

Lawrence Berkeley National Laboratory

LBL Publications

Title

An Effect of the Existence of an Elementary Quark on High Energy Scattering

Permalink

<https://escholarship.org/uc/item/8c45r3gf>

Author

Anderson, Stuart D

Publication Date

1968-05-01

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

eg. 2

University of California
Ernest O. Lawrence
Radiation Laboratory

AN EFFECT OF THE EXISTENCE OF AN ELEMENTARY
QUARK ON HIGH ENERGY SCATTERING

Stuart D. Anderson

May 17, 1968

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

RECEIVE
LAWRENCE
RADIATION LABORA

JUL 2 1968

LIBRARY AND
DOCUMENTS SECTION

Berkeley, California

UCRL-18251 (Rev. 1)
eg. 2

T

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Submitted to the Physical Review

UCRL-18251 Rev. 1
Preprint

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

AEC Contract No. W-7405-eng-48

AN EFFECT OF THE EXISTENCE OF AN ELEMENTARY
QUARK ON HIGH ENERGY SCATTERING

Stuart D. Anderson

May 17, 1968

-iii-

AN EFFECT OF THE EXISTENCE OF AN ELEMENTARY
QUARK ON HIGH ENERGY SCATTERING^{*†}

Stuart D. Anderson

Lawrence Radiation Laboratory
University of California
Berkeley, California

May 17, 1968

ABSTRACT

An elementary quark, if supposed to exist and to couple in quark-antiquark pairs to ordinary two-particle channels, gives rise to a Regge trajectory which would dominate high energy scattering at large momentum transfer. A simple model theory embodying these features is presented. Specifically, the model assumes a single scalar-scalar pair to represent the ordinary two-particle channel which is observed, coupled to a spinor-spinor channel of large mass, and simple pole-type potentials approximated in the effective range method. The Regge trajectory and subtraction generated by this coupling are computed, and arguments given concerning their contribution to the cross section. Comparison with proton-proton scattering suggests that if such an elementary particle exists, its mass must be greater than about 10 GeV.

-1-

I. INTRODUCTION AND OUTLINE

Attempts to verify the existence of the quark by its external production in high energy collisions have to date not met with success.¹ This paper suggests an idea, exemplified here in a simple model, whereby the existence of a massive quark, at least as an internal particle in a collision process, might be made observable. This idea takes its roots in the study of the asymptotic behavior, as generated by the leading Regge singularities of the amplitude, of the scattering of "ordinary" particles (having integral baryonic and electric charge). It is possible that the concept of the elementary quark may be confirmed and better defined should this idea prove useful in explaining ordinary scattering experiments in this high energy region.

Let us consider a scattering amplitude whose leading Regge trajectories fall below angular momentum $J = 0$ for the negative of the squared momentum transfer, t , less than some given (negative) $t_{(0)}$: the amplitude will thus fall asymptotically to zero for large values of the squared energy s . This is apparently an adequate description of a system of "ordinary" composite particles. The same is true of such a system into which are coupled elementary particles of infinite mass, the infinity of the mass effectively negating the coupling. If, however, the mass is finite, coupling to the elementary particle channel is effected, and there appears a new Regge pole which alters this description.

-2-

Let us assume these elementary particles to be our ideal of elementarity: the quarks, conforming to the fundamental spin representation, $J = \frac{1}{2}$, and to the fundamental unitary spin representation $\{3\}$, thus coupling in quark-antiquark pairs to ordinary particle channels. We continue to assume, as the production experiments and the nonrelativistic models suggest, that the quark mass is quite large.

There is a critical difference of character between this new type of trajectory, generated by the introduction of elementary particles, and that type which is already familiar in high energy scattering, and it is this difference which forms the basis of the idea of this work. The familiar Regge poles fall below $J = 0$ for $-t$ large enough-perhaps they fall indefinitely²-whereas this new type of trajectory is bounded from below by $J = s_1 + s_2 - 1$, where s_1 and s_2 are the spins of the elementary particles; i.e., $J = 0$ for the quark pair. Why this is so will be argued immediately below, but the consequence is that the new trajectory is the dominant one and perhaps can be seen as actually dominating-its strength or residue being here the point of attention-for energy and momentum transfer large enough. If the behavior of experimental cross sections can be seen to conform with predictions generated by this new trajectory's presence, properties of the elementary particles, the quarks, in particular the mass, can be obtained. If such modification of present Regge theory is not necessary to obtain

-3-

conformity at the energies and momentum transfers now attainable, we can at least place a lower bound on that mass.

The reason for this difference between the two types of trajectory is, of course, bound to the divergent concepts of the elementary and the composite particle. The presence of a pair of elementary particles gives rise to a term in the partial wave amplitude proportional to the Kronecker delta $\delta_{s_1+s_2-n, J}$, where n is a positive integer.³ To see this, we first note that, for J a given (half-) integral angular momentum, states having helicity $|M| \leq J$, called sense states, are physical states at this J , whereas states having $|M| > J$, nonsense states, are unphysical there. In continuing from high J where all states are physical, sense states, we first encounter such unphysical nonsense states at the (half-)integer $J = s_1 + s_2 - 1$ and subsequently at all (half-) integers $J = s_1 + s_2 - n$, the nonsense (half-)integers. At these values of J the problem of the physical amplitude merely omits such states from consideration, via the Kronecker delta, whereas that of the Regge amplitude, continued to this J value, retains them. A difference between the two solutions is ensured by the Regge continuation's having fictitious elementary particles (the C.D.D. poles) replacing the nonsense states, which are not at all present in the physical partial wave. An immediate consequence of the existence of these Kronecker delta singularities is a

-4-

"hardening" of the asymptotic behavior, the requirement of a subtraction when $s_1 + s_2 - 1 \cong 0$. Out of such a singularity moves a Regge trajectory of this new type as the coupling to the channel is turned on, approaching as $t \rightarrow \infty$ the nonsense (half-)integer from which it emerges, and in the simplest case bounded below by it.

Particles composite in nature will not display such singularities. When a particle can be treated as composite, the Kronecker singularity vanishes in the same way that Mandelstam showed the Amati-Fubini-Stanghellini anomalous cuts to vanish. Thus no subtractions are required if no particles interacting in a system are elementary, and the asymptotic behavior is softened. Any Regge poles which exist are not bound below and we have the familiar nearly linear trajectories well known in high energy physics. But just how to treat an external particle as a composite one is not known. Thus in present theoretical schemes, a scalar-scalar theory has a single subtraction and a pole rising from $J = 0$. So in the model to be presented here, we have chosen to work with a scalar-scalar theory, one not well realized in practice, but one having its leading trajectory falling through $J = 0$, if only to $J = -1$. Had we attempted to work with a theory of a pair of ordinary spinors comparable with the experimentally well known nucleon-nucleon scattering, already there would be a pole covering the one of particular interest, that new one derived from the coupling of this channel to the quark -

-5-

antiquark spinor pair, which rises from and remains near to $J = 0$.

In Section II we will carry out the procedures of calculation entailed in a model which exhibits the above features. This model and those procedures are briefly outlined in the following. We take as our only "ordinary" particle channel in the scattering a pair of scalars—because of the difficulty mentioned above—of equal mass, retaining only a minimum number of parameters in this naive model. These are to be coupled to a quark-antiquark pair, represented by much heavier spin one-half particles in two helicity channels. Simple potentials are chosen and approximated by an effective-range type method. The N over D method, which now reduces from integral equations to numerical equations, is used to obtain the partial wave amplitude, whose poles in J , $\alpha_i(t)$, and reduced residues in the ordinary particle channels, $\gamma_i(t)$, are found. Returning to the scattering amplitude by a Sommerfeld-Watson transformation, we obtain the large-energy amplitude and cross section in the crossed (s) channel. In the concluding part of Section II, the subtraction term is discussed. Finally, in Section III, comparison of the behavior of this imagined scalar pair scattering with experimental nucleon-nucleon scattering, for example, places an order of magnitude lower limit on the mass of the quark.

II. THE MODEL

In the first part of this section, we will set up the problem, proceed through to the solution of the N over D equations in the t channel, and comment briefly on the Sommerfeld-Watson transformation. In the second, we shall find the equations describing the t channel poles, and their residues and contributions to the asymptotic s -channel amplitude. In the third, the subtraction term will be discussed.

A. N Over D Solution

1. Unitarity, the Right-Hand Cut

To begin, we write down the t channel unitarity equation for the amplitude M , normalized appropriately:

$$\frac{M_{ba} - M_{ba}^+}{2i} = \sum_c \int \frac{d\Omega_c}{4\pi} M_{bc}^+ \left\{ \frac{2p_c}{(t)^{\frac{1}{2}}} \theta[t - (M_{c1} + M_{c2})^2] \right\} M_{ca} \quad , \quad (2A.1)$$

in terms of which the cross section is

$$\frac{d\sigma_{ba}}{d\Omega_b} = \frac{p_b}{p_a} \left| \frac{M_{ba}}{E} \right|^2 \quad . \quad (2A.2)$$

Most notations are the customary ones. The particle channels in the problem to be considered are three:

1) a pair of "ordinary" particles which in the model are likened to two pions, having no spin, mass μ , and three-momentum q , but which are to be compared in the final analysis with some particles whose scattering is better known, for instance, the nucleons; and

2) a helicity zero state, and

3) a helicity one state, of a pair of spin one-half antiparticle conjugates of mass M and three-momentum p which are treated as nucleon-antinucleon in the model and which are finally comparable with the very heavy quarks.

We are concerned here only with the dynamical aspects of the model; no isotopic spin (or unitary spin) considerations are put forward.

The angular momentum projection of M in helicity states is

$$m'_{ba}{}^J = \frac{1}{2} \int_{-1}^1 dz d_{\lambda_a \lambda_b}^J(\theta) M_{ba}(t, z) \quad (2A.3)$$

For spinless particles, the factors needed to remove threshold kinematic singularities are well known:

$$m_{11}{}^J = q^{-2J} m'_{11}{}^J = \frac{1}{2\pi} \int_{s_0}^{\infty} ds (q^2)^{-J-1} Q_J\left(1 + \frac{s}{2q^2}\right) [M_{11}]_s \quad (2A.4)$$

For spinless particles, further kinematic singularities must be removed. From Frazer and Fulco we find⁴

$$m'_{21}{}^J = \frac{2}{p} \left\{ -p^2 A_J + \frac{M}{2J+1} (pq) [(J+1) B_{J+1} - J B_{J-1}] \right\} \quad (2A.5a)$$

and

$$m'_{31}{}^J = \frac{2E}{p} (pq) \frac{[J(J+1)]^{\frac{1}{2}}}{2J+1} (B_{J-1} - B_{J+1}) \quad (2A.5b)$$

Note that certain linear combinations of these amplitudes are those of defined parity and orbital angular momentum, not these quantities themselves. Writing $a_J = (pq)^{-J} A_J$, etc., for the projected invariant amplitudes with threshold kinematics removed, we see that

$$m_{21}^J = \frac{p}{M}(pq)^{-J} m'_{21} = 2 \left\{ \frac{p^2}{M} a_J + \frac{1}{2J+1} [(J+1)(pq)^2 b_{J+1} - J b_{J-1}] \right\} \quad (2A.6a)$$

and

$$m_{31}^J = \frac{p}{E}(pq)^{-J} m'_{31} = 2 \frac{[J(J+1)]^{\frac{1}{2}}}{2J+1} [b_{J-1} - (pq)^2 b_{J+1}] \quad (2A.6b)$$

are free of kinematic singularities, and that in terms of these amplitudes the unitarity equation becomes an equation of the discontinuity across the right-hand cut:

$$\frac{m_{ba}^{J+} - m_{ba}^{J-}}{2i} = \sum_c m_{bc}^{J\pm} \rho_c^J m_{ca}^{J\mp}, \text{ or } [m^{-1}]_R = -\rho^J, \quad (2A.7a)$$

where ρ_c^J is a diagonal matrix with elements

$$(\rho^1, \rho^2, \rho^3) = \left[\frac{q^{2J+1}}{E} \theta(t - 4\mu^2), \frac{M^2 p^{2J-1}}{E} \theta(t - 4M^2), p^{2J-1} E \theta(t - 4M^2) \right] \quad (2A.7b)$$

2. The Potential, the Left-Hand Cut

The potential is chosen in the usual simplistic fashion: double-spectral contributions are ignored, and only the poles are saved (see Fig. 1):

$$[M_{11}]_s = (g\mu)^2 \pi \delta(s - \mu^2) \quad , \quad (2.8a)$$

$$[B]_s = (g')^2 \pi \delta(s - M^2) \quad . \quad (2A.8b)$$

We have assumed a scalar type of coupling for the imagined pions, with coupling constant $g\mu$ (μ inserted for convenience). In the coupling of the spin one-half particles to the spinless one, we assume that the coupling constant g' defined in analogy with that of nucleons to the pion remains of order $M^0 = 1$ as the mass M becomes large; how to modify this assumption for other behavior in this mass parameter is obvious. It is also quite easy to modify the assumption made in the interest of economy of parameters that the masses of the exchanged particles are the same as those of the external "pion" and "nucleon."

The corresponding integrals over the left-hand cuts, the potential in the N over D method, are therefore

$$\{m_{11}^J\}_L = g^2 \frac{1}{2} \mu^2 (q^2)^{-J-1} Q_J(z_\pi) \quad , \quad (2A.9a)$$

$$\{m_{21}^J\}_L = g'^2 (pq)^{-J} z Q_J(z) \quad , \quad (2A.9b)$$

and

$$\{m_{31}^J\}_L = -g'^2 \frac{1}{[J(J+1)]^{\frac{1}{2}}} (pq)^{-J} (z^2 - 1) Q_J'(z) \quad , \quad (2A.9c)$$

where

$$z_{\pi} = 1 + \frac{\mu^2}{2q^2} \quad \text{and} \quad z = \frac{p^2 + q^2 + M^2}{2pq} \quad . \quad (2A.9d)$$

Now each part of the potential is approximated by a pole, a familiar scheme called the effective range approximation. The method we have chosen in to find the position and residue of the pole for all J by fitting at the channel 1 threshold ($q^2 \rightarrow 0$) to the value and derivative of the original form. For example:

| | Original Form | Approximating Form |
|-----------------------------|---|-------------------------------|
| | $\frac{1}{2} \mu^2 (q^2)^{-J-1} Q_J(z_{\pi})$ | $\frac{a}{t - t_a}$ |
| $() _{q^2=0}$ | $\frac{1}{2} (\mu^2)^{-J} \pi^{\frac{1}{2}} \frac{\Gamma(J+1)}{\Gamma(J+\frac{3}{2})}$ | $\frac{a}{4\mu^2 - t_a}$ |
| $\frac{d}{dt} () _{q^2=0}$ | $\frac{1}{2} (\mu^2)^{-J} \pi^{\frac{1}{2}} \frac{\Gamma(J+1)}{\Gamma(J+\frac{3}{2})} \left(-\frac{J+1}{2\mu^2} \right)$ | $-\frac{a}{(4\mu^2 - t_a)^2}$ |

So we have that

$$\frac{1}{2} \mu^2 (q^2)^{-J-1} Q_J(z_{\pi})$$

is approximated by

$$\frac{2\pi^{\frac{1}{2}} \Gamma(J+1)}{(\mu^2)^{J-1} \Gamma(J+\frac{3}{2})} \cdot \frac{1}{J+1} \cdot \frac{1}{4\mu^2 \left(x-1+\frac{1}{J+1}\right)},$$

where $x = t/4\mu^2$. The other parts of the potential are approximated in a similar fashion.

In summary, we have approximated the potential by

$$\begin{aligned} \{m^J\}_L = V^J(t) &= G \left[\frac{a_1}{t-t_1} \begin{pmatrix} 100 \\ 000 \\ 000 \end{pmatrix} + \frac{a_2}{t-t_2} \begin{pmatrix} 010 \\ 100 \\ 000 \end{pmatrix} + \frac{a_3}{t-t_3} \begin{pmatrix} 001 \\ 000 \\ 100 \end{pmatrix} \right] \\ &= G \sum_{i=1}^3 \frac{a_i v_i}{t-t_i}, \end{aligned} \quad (2A.10a)$$

where

$$G = 4\mu^2 g^2 \left(\frac{1}{2} \cdot \frac{\pi^{\frac{1}{2}} \Gamma(J+1)}{\mu^{2J} \Gamma(J+\frac{3}{2})} \right); \quad (2A.10b)$$

$$a_1 = \frac{1}{J+1}, \quad x_1 = \frac{t_1}{4\mu^2} = 1 - \frac{1}{J+1} = \frac{2J+1}{2(J+1)}; \quad (2A.10c)$$

$$\begin{aligned} a_2 &= \left(\frac{g'}{g}\right)^2 \frac{\left(\frac{3}{2}+J\right)\lambda^2}{(J+1)(J+2) + (J^2-2)\lambda^2}, \\ x_2 &= \frac{t_2}{4\mu^2} = 1 - \frac{\left(\frac{3}{2}+J\right)\lambda^2}{(J+1)(J+2) + (J^2-2)\lambda^2}; \end{aligned} \quad (2A.10d)$$

and

$$a_3 = \left(\frac{g'}{g}\right)^2 \frac{\left(\frac{3}{2} + J\right)\lambda^2}{J[J+1 + (3J+5)\lambda^2]} \left(\frac{J+1}{J}\right)^{\frac{1}{2}},$$

$$x_3 = \frac{t_3}{4\mu^2} = 1 - \frac{\left(\frac{3}{2} + J\right)\lambda^2}{J[J+1 + (3J+5)\lambda^2]}; \quad (2A.10e)$$

and where $\lambda = \mu/M$ is the anticipatedly small ratio of the masses.

3. N Over D Solution

We have now discussed the quantities which are to be put into the N over D method for m^J : Eqs. (2A.7), (2A.10), where $m^J = ND^{-1}$. The primary purpose in approximating the potential terms by poles is that the integral equation for N now reduces to a numerical equation. Inserting Eq. (2A.10a) into this equation for N,

$$N(t) = V(t) + \frac{1}{\pi} \int_R dt' \frac{V(t') - V(t)}{t' - t} \rho(t') N(t'), \quad (2A.11)$$

we may write the solution

$$N(t) = G \sum_{i=1}^3 \frac{a_i v_i}{t - t_i}, \quad (2A.11a)$$

with the numerical system of equations for the matrices v_i ,

$$v_i = v_i \left(1 - \sum_{j=1}^3 \rho_{ij} a_j v_j \right) \quad \text{or} \quad \sum_{j=1}^3 (\delta_{ij} + v_i \rho_{ij} a_j) v_j = v_i. \quad (2A.12)$$

This is a system of 3×3 matrices in the pole index, the elements of which are 3×3 in the channel number. We define here a matrix whose symmetrical elements in the pole subscript space are

$$\begin{aligned} \rho_{ij} &= \frac{G}{\pi} \int_R dt' \frac{\rho(t')}{(t' - t_i)(t' - t_j)} \\ &= \frac{\frac{G}{\pi} \int_R dt' \frac{\rho(t')}{(t' - t_i)}}{t_i} - \frac{\frac{G}{\pi} \int_R dt' \frac{\rho(t')}{(t' - t_j)}}{t_j} = \frac{\rho_i - \rho_j}{t_i - t_j}, \end{aligned} \quad (2A.13a)$$

whose diagonal elements in the channel number space are

$$(\rho_{ij})_{\alpha\beta} = \delta_{\alpha\beta} \rho_{ij}^{\alpha} = \delta_{\alpha\beta} \frac{\rho_i^{\alpha} - \rho_j^{\alpha}}{t_i - t_j}, \quad (2A.13b)$$

where

$$(\rho_i)_{\alpha\beta} = \delta_{\alpha\beta} \rho_i^{\alpha} = \delta_{\alpha\beta} \frac{G}{\pi} \int_R dt' \frac{\rho^{\alpha}(t')}{(t' - t_i)}. \quad (2A.13c)$$

Note that in the numbers ρ_{ij}^{α} or ρ_i^{α} , the subscripts denote the index of the approximating pole and the superscripts, the channel number. Explicitly these quantities are

$$\rho_i^1 = -4\mu^2 \frac{g^2}{\pi} \frac{\pi}{\sin \pi J} F(1, -J; \frac{3}{2}; x_i), \quad (2A.14a)$$

$$\rho_i^2 = + 4\mu^2 \frac{g^2}{\pi} \frac{\pi}{\sin \pi J} \lambda^{-2J} \left(\frac{J}{J + \frac{1}{2}} \right) F(1, 1 - J; \frac{3}{2}; x_i \lambda^2) , \quad (2A.14b)$$

and

$$\rho_i^3 = - 4\mu^2 \frac{g^2}{\pi} \frac{\pi}{\sin \pi J} \lambda^{-2J} \left(\frac{J}{J + \frac{1}{2}} \right) F(1, -J; \frac{1}{2}; x_i \lambda^2) . \quad (2A.14c)$$

$F(a, b; c; x)$ is the hypergeometric function.

The system (2A.12), which is 9×9 , is immediately reduced to a 5×5 system by virtue of the fact that four rows of v_i are zero, and is solved in terms of the 3×3 determinant d ,

$$d = \begin{vmatrix} 1 + a_1 \rho_{11}^1 & - a_2 \rho_{12}^1 \rho_{22}^2 & - a_3 \rho_{13}^1 \rho_{33}^3 \\ a_1 \rho_{12}^1 & 1 - a_2 \rho_{22}^1 \rho_{22}^2 & - a_3 \rho_{23}^1 \rho_{33}^3 \\ a_1 \rho_{13}^1 & - a_2 \rho_{23}^1 \rho_{22}^2 & 1 - a_3 \rho_{33}^1 \rho_{33}^3 \end{vmatrix} , \quad (2A.15)$$

and its cofactor matrix elements d_{ij} (the determinant derived from d in this form by setting the i th row and j th column to zero, except for the (i, j) element which is set to one). Thus it is found that

$$v_1 = \frac{1}{d} \begin{bmatrix} d_{11} + d_{21} + d_{31} & \frac{a_2}{a_1} d_{12} & \frac{a_3}{a_1} d_{13} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} , \quad (2A.16a)$$

$$v_2 = \frac{1}{d} \begin{bmatrix} -a_2 \rho_{22}^2 (d_{12} + d_{22} + d_{32}) & d_{22} & \frac{a_3}{a_2} d_{23} \\ d_{12} + d_{22} + d_{32} & \frac{d - d_{22}}{a_2 \rho_{22}^2} & -\frac{d_{32}}{a_3 \rho_{33}^3} \\ 0 & 0 & 0 \end{bmatrix}, \quad (2A.16b)$$

and

$$v_3 = \frac{1}{d} \begin{bmatrix} -a_3 \rho_{33}^3 (d_{13} + d_{23} + d_{33}) & \frac{a_2}{a_3} d_{32} & d_{33} \\ 0 & 0 & 0 \\ d_{13} + d_{23} + d_{33} & -\frac{d_{23}}{a_2 \rho_{22}^2} & \frac{d - d_{33}}{a_3 \rho_{33}^3} \end{bmatrix}. \quad (2A.16c)$$

The denominator function $D(t)$ is found by merely integrating:

$$D(t) = 1 - \frac{1}{\pi} \int_R dt' \frac{\rho(t')}{t' - t} N(t'), \quad (2A.17)$$

which is

$$D(t) = 1 - \sum_{i=1}^3 a_i \rho_{0i} v_i \quad (2A.17a)$$

$$= \frac{1}{d} \begin{bmatrix} d - a_1 \rho_{10}^1 (d_{11} + d_{21} + d_{31}) & - a_2 (\rho_{10}^1 d_{12} & - a_3 (\rho_{10}^1 d_{13} \\ + a_2 \rho_{20}^1 \rho_{22}^2 (d_{12} + d_{22} + d_{32}) & + \rho_{20}^1 d_{22} & + \rho_{20}^1 d_{23} \\ + a_3 \rho_{30}^1 \rho_{33}^3 (d_{13} + d_{23} + d_{33}) & + \rho_{30}^1 d_{32}) & + \rho_{30}^1 d_{33}) \\ - a_2 \rho_{20}^2 (d_{12} + d_{22} + d_{32}) & d - \frac{\rho_{20}^2}{\rho_{22}^2} (d - d_{22}) & \frac{a_3}{a_2} \frac{\rho_{20}^2}{\rho_{22}^2} d_{23} \\ - a_3 \rho_{30}^3 (d_{13} + d_{23} + d_{33}) & \frac{a_2}{a_3} \frac{\rho_{30}^3}{\rho_{33}^3} d_{32} & d - \frac{\rho_{30}^3}{\rho_{33}^3} (d - d_{33}) \end{bmatrix},$$

(2A.17b)

where we use the subscript 0 to denote $t_i = t = t_0$. Its determinant is found to be

$$|D| = \frac{\mathcal{D}}{d}, \tag{2A.18}$$

$$\begin{aligned}
 \phi = & \left| \begin{array}{ccc}
 1 + a_1(\rho_{11}^1 - \rho_{10}^1) & - a_2(\rho_{12}^1 - \rho_{20}^1)(\rho_{22}^2 - \rho_{20}^2) & - a_3(\rho_{13}^1 - \rho_{30}^1) \\
 & & \times (\rho_{33}^3 - \rho_{30}^3) \\
 a_1(\rho_{12}^1 - \rho_{10}^1) & 1 - a_2(\rho_{22}^1 - \rho_{20}^1)(\rho_{22}^2 - \rho_{20}^2) & - a_3(\rho_{23}^1 - \rho_{30}^1) \\
 & & \times (\rho_{33}^3 - \rho_{30}^3) \\
 a_1(\rho_{13}^1 - \rho_{10}^1) & - a_2(\rho_{23}^1 - \rho_{20}^1)(\rho_{22}^2 - \rho_{20}^2) & 1 - a_3(\rho_{33}^1 - \rho_{30}^1) \\
 & & \times (\rho_{33}^3 - \rho_{30}^3)
 \end{array} \right| .
 \end{aligned}$$

In order to eliminate factors of d , we define

$$N = \frac{1}{d} \eta , \tag{2A.19}$$

and \mathcal{D}^{-1} to be the cofactor matrix of D ; i.e., $D^{-1} = \mathcal{D}^{-1}/|D|$;

hence

$$D = \frac{\phi}{d} \mathcal{D} \tag{2A.17c}$$

and

$$m^J = \frac{\eta \mathcal{D}^{-1}}{\phi} . \tag{2A.20}$$

If λ is small, or zero, due to large, or infinite, quark mass, and J is not near zero, a good approximation is $a_2 = a_3 = 0$.

Then

$$\not{D} = 1 + a_1(\rho_{11}^1 - \rho_{10}^1) \quad (2A.18a)$$

and

$$(\not{\eta} \not{D}^{-1})_{11} = \frac{G}{4\mu^2} \frac{a_1}{x - x_1} \quad (2A.21a)$$

If λ is small, and if J is near zero, the approximation to be made is $a_2 = 0$, and in this case

$$\not{D} = \begin{vmatrix} 1 + a_1(\rho_{11}^1 - \rho_{10}^1) & -a_3^2(\rho_{13}^1 - \rho_{30}^1)(\rho_{33}^3 - \rho_{30}^3) \\ a_1(\rho_{13}^1 - \rho_{10}^1) & 1 - a_3^2(\rho_{33}^1 - \rho_{30}^1)(\rho_{33}^3 - \rho_{30}^3) \end{vmatrix}$$

and

$$(\not{\eta} \not{D}^{-1})_{11} = \left\{ \frac{G}{4\mu^2} \frac{a_1}{x - x_1} - a_3^2(\rho_{33}^3 - \rho_{30}^3) \left[\frac{1}{x - x_3} + \frac{a_1(\rho_{33}^1 - \rho_{13}^1)}{x - x_1} + \frac{a_1(\rho_{11}^1 - \rho_{13}^1)}{x - x_3} \right] \right\} \quad (2A.21b)$$

This approximation of large quark mass is taken in the following as an assumption to be verified by the final results, in comparison with experiment, and will be seen then to be justified.

4. The Sommerfeld-Watson Transform

The procedure to be followed from here on is well known. One solves the equation $\delta = 0$ for the Regge poles, $J = \alpha_i(t)$, and their reduced residues,

$$\gamma_i(t) = \left. \frac{(\eta \delta^{-1})_{11}}{\frac{\partial \delta}{\partial J}} \right|_{J=\alpha_i(t)}, \quad (2A.22)$$

so that by the Sommerfeld-Watson transform the amplitude is, for large s ,

$$M = \sum_i -2\pi^{\frac{1}{2}} \frac{\Gamma(\alpha_i + \frac{3}{2})}{\Gamma(\alpha_i + \frac{1}{2})} \frac{\gamma_i(-s)^{\alpha_i}}{\sin \pi\alpha_i}$$

A simplifying factorization is made by defining

$$\gamma_i = \left[\frac{\frac{1}{2} \pi^{\frac{1}{2}} \Gamma(\alpha_i + 1)}{\mu^{2\alpha_i} \Gamma(\alpha_i + \frac{3}{2})} \right] \bar{\gamma}_i, \quad (2A.23)$$

this factor being included in η via the factor in G , from Eq.(2A.10b).

With this inserted,

$$\begin{aligned} M &= \sum_i - \frac{\pi}{\sin \pi\alpha_i} \bar{\gamma}_i \left(-\frac{s}{\mu} \right)^{\alpha_i} = \sum_i - \pi \bar{\gamma}_i \left(\frac{s}{\mu} \right)^{\alpha_i} (\cot \pi\alpha_i + 1) \\ &= \sum_i M_i. \end{aligned} \quad (2A.24)$$

We note in passing that for a pole whose position and residue are small and of the same order of magnitude, the contribution to M is

$$M_i = -\frac{\bar{\gamma}_i}{\alpha_i} - \bar{\gamma}_i \left[\ln\left(\frac{s}{\mu^2}\right) + i\pi \right] \quad (2A.24a)$$

By crossing, M is the Reggeized asymptotic amplitude for scattering in the crossed (s) channel, for which

$$\frac{d\sigma}{dt} = \frac{4\pi}{s q_s^2} |M|^2 \quad (2A.25)$$

B. Regge Poles, Residues, and Asymptotic Contributions

1. Uncoupled Case

We first study the case of infinite quark mass for which $M = \infty$, or $\lambda = a_2 = a_3 = 0$. The pole to be found in this case will, of course, be present in the finite mass case, but perturbed by the small but finite λ . Here, setting Eq. (2A.18a) to zero, we have the equation of the trajectory $J = \alpha_I(x)$. Explicitly this is

$$0 = 1 + \frac{g^2}{\pi} \frac{\pi}{\sin \pi \alpha_I} \frac{1}{\alpha_I + 1} \left[\frac{F(1, -\alpha_I; \frac{3}{2}; x) - F(1, -\alpha_I; \frac{3}{2}; x_1)}{x - x_1} - F'(1, -\alpha_I; \frac{3}{2}; x_1) \right], \quad (2B.1)$$

with $x_1 = x_1(\alpha_I) = 1 - [(1/2)/(\alpha_I + 1)]$. Since $F(1, -\alpha; 3/2; x)$ does not take a closed form except for α equal to any half-integer,

for simplicity we shall find the pole positions in x for $\alpha_I = -1$, $-1/2$, and 0 and interpolate.

$$\text{At } \alpha_I = -1, \quad x \text{ approaches } -\frac{1}{2}/(\alpha_I + 1). \quad (2B.2)$$

At $\alpha_I = -\frac{1}{2}$, Eq. (2B.1) becomes

$$1 + \left(\frac{1}{3} + \frac{1}{g^2}\right)x = F\left(1, \frac{1}{2}; \frac{3}{2}; x\right) = \frac{1}{2x^{\frac{1}{2}}} \ln \left(\frac{1 + x^{\frac{1}{2}}}{1 - x^{\frac{1}{2}}}\right). \quad (2B.3)$$

Thus, numerically or graphically, one can find $x_{(-\frac{1}{2})}$, the pole position in energy for angular momentum $\alpha_I = -\frac{1}{2}$ for any coupling g .

At the limits, $x_{(-\frac{1}{2})} = 1$ for $g = 0$ and $x_{(-\frac{1}{2})} = 0$ for $g = \infty$.

At $\alpha_I = 0$, the $0/0$ form is easily reduced, since

$$F\left(1, -\alpha; \frac{3}{2}; x\right) = 1 - \frac{2}{3}\alpha x F\left(1, 1; \frac{5}{2}; x\right) = 1 - 2\alpha(1 - \mu(x)) \quad (2B.4)$$

near $\alpha = 0$, and

$$\mu(x) = \left(\frac{1-x}{-x}\right)^{\frac{1}{2}} \ln \left[(-x)^{\frac{1}{2}} + (1-x)^{\frac{1}{2}}\right] = \left(\frac{1-x}{x}\right)^{\frac{1}{2}} \arcsin(x^{\frac{1}{2}}). \quad (2B.5)$$

The equation (2B.1) at $\alpha_I = 0$ is thus

$$\frac{\pi}{4} + \left[1 - \frac{\pi}{2} - \frac{1}{(g^2/\pi)}\right] \left(x - \frac{1}{2}\right) = \mu(x). \quad (2B.6)$$

Again a numerical or graphical solution is easily obtained for $x_{(0)}$.

The range in x of the pole along $J = 0$ with g varying is from $\frac{1}{2}$ at $g = 0$ to 1 at $g^2/\pi = 1$, and there is no solution for g smaller than this.

Assembling the information from these three values of α_I [for $(g^2/\pi) \geq 1$] and interpolating, we take for the Regge pole, the form

$$x = x_I(J) = \frac{a_I}{J+1} + b_I + c_I(J+1), \quad (2B.7)$$

where

$$a_I = -\frac{1}{2}, \quad b_I = \frac{3}{2} - x_{(0)} + 2x_{(-\frac{1}{2})}, \quad \text{and} \quad c_I = 2[x_{(0)} - x_{(-\frac{1}{2})}] - 1. \quad (2B.7a)$$

Inversely, we have

$$J = \alpha_I(x) = -1 + \frac{x - b_I + [(x - b_I)^2 - 4a_I c_I]^{\frac{1}{2}}}{2c_I}. \quad (2B.8)$$

Thus ϑ may be written

$$\vartheta = \frac{x - x_I(J)}{x_1(J) - x_I(J)}, \quad (2B.9)$$

since $\vartheta = 0$ on the trajectory $x = x_I(J)$, and $\vartheta = 1$ at $x = x_1(J)$.

Hence one may computed

$$\left. \frac{\partial \mathcal{D}}{\partial J} \right|_{J=\alpha_I}$$

and from Eq. (2A.21a) $(\eta \theta^{-1})_{11}$, in order to find $\bar{\gamma}_I$ by

Eq. (2A.22):

$$\bar{\gamma}_I = \frac{\frac{1}{2} \pi \left(\frac{g^2}{\pi} \right)}{[(x - b_I)^2 - 4a_I c_I]^{\frac{1}{2}}} \quad (2B.10)$$

Therefore, for the uncoupled problem, the asymptotic behavior of the amplitude in the s channel is

$$M = -\pi \bar{\gamma}_I \left(\frac{s}{\mu^2} \right)^{\alpha_I} (\cot \pi \alpha_I + i) = M_I \quad (2B.11)$$

For $(g^2/\pi) < 1$, we have

$$J = \alpha_I(x) = -1 + \frac{a_I}{x - b_I} \quad (2B.8a)$$

with

$$a_I = -\frac{1}{2} \quad \text{and} \quad b_I = 1 + x \left(-\frac{1}{2} \right) \quad (2B.8b)$$

and

$$\bar{\gamma}_I = \frac{\frac{1}{2} \pi \left(\frac{g^2}{\pi} \right)}{x - b_I} \quad (2B.10a)$$

In Fig. 2 we have drawn $\alpha_I(x)$ and $\bar{\gamma}_I(x)$, Eqs. (2B.8) and (2B.10), picking arbitrarily the value $(g^2/\pi) = 2$.

2. Coupled Case

Now we shall consider the case when M is not infinite, but is large and finite, $\lambda \ll 1$. Let us note the following general point before continuing on to the details. The elements of ϕ in its second column as written in Eq. (2A.18) containing $a_2^2(\rho_{22}^2 - \rho_{20}^2)$ are proportional to λ^8 (λ^4 near $J = -1$) and those in the third column containing $a_3^2(\rho_{33}^3 - \rho_{30}^3)$ to λ^8 (λ^8/J^3 near $J = 0$). Thus the pole of the uncoupled case, $\alpha_I(x)$, is only negligibly perturbed, and this is also a pole of the coupled case, except near $J = 0$, $x = x_{(0)}$ where the solution is, as yet, undetermined. A second zero of ϕ occurs for J near enough to zero that terms containing $a_3^2(\rho_{33}^3 - \rho_{30}^3)$ become considerable. Terms containing $a_2^2(\rho_{22}^2 - \rho_{20}^2)$ remain always negligibly small and we write as in Eq. (2A.18a),

$$\phi = \begin{vmatrix} \phi_{11} & \phi_{13} \\ \phi_{31} & \phi_{33} \end{vmatrix} = \phi_{11} \left(\phi_{33} - \frac{\phi_{13} \phi_{31}}{\phi_{11}} \right)$$

The zero of $\phi/\phi_{11} = \phi_{33} - \phi_{13} \phi_{31}/\phi_{11}$, $\alpha_{III}(x)$, will be treated below, and it is a pole of the coupled case, except near $x = x_{(0)}$, where apparently the two poles will cross each other, for $(g^2/\pi) \geq 1$.

Actually, the poles will not cross each other in this anticipated fashion, but break and rejoin near $x = x_{(0)}$, $J = 0$, so as to avoid crossing while passing very close by.⁵ Since in what follows we are concerned only with the region somewhat below this point, we may take as the approximate poles of the coupled problem $\alpha_I(x)$ and $\alpha_{III}(x)$, defined as the zeros of \mathcal{D}_1 and $\mathcal{D}/\mathcal{D}_1$.

With the approximations $\lambda \ll 1$ and $|J| \ll 1$ inserted to lowest order into \mathcal{D} , we now find α_{III} as the zero of $\mathcal{D}/\mathcal{D}_1$:

$$\mathcal{D}_1 = 1 + \frac{g^2}{\pi} \left(\frac{\mu_0 - \mu_1}{x_0 - x_1} - \mu_1' \right), \quad (2B.12a)$$

$$\mathcal{D}_3 = 1 + \left(\frac{g^2}{\pi} \right)^2 \lambda^{-2J} 6 \frac{\lambda^8}{J^3} (x_0 - x_3) \left(\frac{\mu_0 - \mu_3}{x_0 - x_3} - \mu_3' \right), \quad (2B.12b)$$

$$\mathcal{D}_{13} \mathcal{D}_{31} = \left(\frac{g^2}{\pi} \right) \left(\frac{g^2}{\pi} \right)^2 \lambda^{-2J} 6 \frac{\lambda^8}{J^3} (x_0 - x_3) \left(\frac{\mu_0 - \mu_3}{x_0 - x_3} - \frac{\mu_1 - \mu_3}{x_1 - x_3} \right) \left(\frac{\mu_0 - \mu_1}{x_0 - x_1} - \frac{\mu_3 - \mu_1}{x_3 - x_1} \right), \quad (2B.12c)$$

where

$$\mu_1 = \mu(x_1), \text{ etc.}, \text{ and } \mu_1' = \left. \frac{d}{dx} \mu(x) \right|_{x=x_1}, \text{ etc.},$$

and

$$x_1 = \frac{1}{2} \quad \text{and} \quad x_3 = 1 - \frac{3\lambda^2}{2J} .$$

From this it is apparent that $J = \alpha_{\text{III}}(x)$ will be of order $\lambda^{8/3}$, that is, not only are α_{III} and λ both small, but as well, $|\alpha_{\text{III}}| \ll \lambda^2$, and x_3 is large and negative. Inserting this further approximation we have, to lowest order in J/λ^2 ,

$$x_3 = -\frac{3\lambda^2}{2J} ,$$

$$\mu_3 = -\frac{1}{2} \ln(J/6\lambda^2) ,$$

and

$$\mu_3' = -\frac{J}{3\lambda^2} ,$$

and for $|x_0| \ll |x_3|$,

$$\frac{\mu_0 - \mu_3}{x_0 - x_3} = \frac{J}{3\lambda^2} \left[\ln\left(\frac{J}{6\lambda^2}\right) + 2\mu_0 \right] .$$

Writing $J/\lambda^{8/3} = j$, we have as the numerically solvable equation for $j_{\text{III}}(x) = \lambda^{-8/3} \alpha_{\text{III}}(x)$, good to order $\lambda^{8/3}$,

$$\psi/\psi_1 = 1 + \left(\frac{g'^2}{\pi}\right)^2 \frac{1}{j_{\text{III}}(x)} [\ln(j_{\text{III}}^3(x)) + h(x)] = 0, \quad (2B.13a)$$

with

$$\begin{aligned} h(x) &= 3 \left[\ln\left(\frac{\lambda^{2/3}}{6}\right) + 2\mu_0 + 1 - 2 \left(\frac{g'^2}{\pi}\right) \frac{(\mu_0 - \mu_1)^2 / (x_0 - x_1)}{\psi_1(x)} \right] \\ &= 3 \left\{ \ln\left(\frac{\lambda^{2/3}}{6}\right) + 2\mu_1 + 1 + 2 \frac{(\mu_0 - \mu_1) [1 - \left(\frac{g'^2}{\pi}\right) \mu_1']}{\psi_1} \right\}, \end{aligned} \quad (2B.13b)$$

valid for $|x_0| \ll \lambda^{-2/3}$ and below (or above) the region of the zero of ψ_1 , $x = x_0$. As $\lambda \rightarrow 0$,

$$j_{\text{III}}(x) \rightarrow \left(\frac{g'^2}{\pi}\right)^{2/3} \ln^{1/3}\left(\frac{1}{\lambda}\right) \rightarrow \infty$$

and

$$\alpha_{\text{III}}(x) \rightarrow \left(\frac{g'^2}{\pi}\right)^{2/3} \lambda^{8/3} \ln^{1/3}\left(\frac{1}{\lambda}\right) \rightarrow 0.$$

One quantity of interest in computing the residue of this pole is

$$\left. \frac{\partial \psi}{\partial j} \right|_{j=\alpha_{\text{III}}} = \lambda^{-8/3} \left. \frac{\partial \psi}{\partial j} \right|_{j=j_{\text{III}}} = \frac{3\lambda^{-8/3} \psi_1(x)}{j_{\text{III}}(x)} \left[1 + \frac{\left(\frac{g'^2}{\pi}\right)^2}{j_{\text{III}}^3(x)} \right].$$

The other is, from Eq. (2A.21b),

$$(\eta s^{-1})_{11} |_{J=\alpha_{\text{III}}} = \frac{\frac{1}{2} \pi \left(\frac{g^2}{\pi} \right)}{x - \frac{1}{2}} \left[\frac{1}{2} \frac{\pi^{\frac{1}{2}}}{2\alpha_{\text{III}}} \frac{\Gamma(\alpha_{\text{III}} + 1)}{\Gamma(\alpha_{\text{III}} + \frac{3}{2})} \right] \\
 \left\{ 1 + \left(\frac{g'^2}{\pi} \right)^2 \frac{1}{j_{\text{III}}^3} \left[\ln(j_{\text{III}}^3) + k \right] \right\} ,$$

where analogously to Eq. (2B.13b),

$$k(x) = 3 \left\{ \ln \left(\frac{\lambda^{2/3}}{6} \right) + 2\mu_1 + 1 + 2 \frac{(x_0 - x_1) \left[1 - \left(\frac{g^2}{\pi} \right) \mu_1' \right]}{(g^2/\pi)} \right\} .$$

Thus, in summary of the properties of this pole,

$$\bar{\gamma}_{\text{III}}(x) = \alpha_{\text{III}} \frac{\frac{1}{6} \pi \left(\frac{g^2}{\pi} \right)}{(x - \frac{1}{2}) \psi_1} \left[\frac{1 + \left(\frac{g'^2}{\pi} \right)^2 \frac{k}{j_{\text{III}}^3} + \left(\frac{g'^2}{\pi} \right)^2 \frac{\ln(j_{\text{III}}^3)}{j_{\text{III}}^3}}{1 + \left(\frac{g'^2}{\pi} \right)^2 \frac{1}{j_{\text{III}}^3}} \right] , \tag{2B.14}$$

and $\alpha_{\text{III}}(x) = \lambda^{8/3} j_{\text{III}}(x)$, which is defined by Eq. (2B.13a). The contribution of this pole to the Reggeized asymptotic amplitude in the s channel is, by Eq. (2A.24a),

$$M_{\text{III}} = - \frac{\bar{\gamma}_{\text{III}}}{\alpha_{\text{III}}} - \bar{\gamma}_{\text{III}} \left[\ln \left(\frac{s}{\mu^2} \right) + i\pi \right] . \tag{2B.15}$$

In Fig. 3 we have drawn $j_{\text{III}}(x) = \lambda^{-8/3} \alpha_{\text{III}}(x)$ and

$\bar{\gamma}'_{\text{III}}(x) = \lambda^{-8/3} \bar{\gamma}_{\text{III}}(x)$, from Eqs. (2B.13a) and (2B.14), choosing

arbitrarily the values $(g^2/\pi) = 2$ and $(g'^2/\pi) = 1$.

Finally we note that all approximations but one appear to be correct to order of magnitude in the mass M , or λ , and that one is that the behavior of the coupling g' , being unknown as a function of λ , was assumed to be constant. If we allow (g'^2/π) to go to $(g'^2/\pi) \lambda^\delta$, in most instance $\lambda^{8/3} \rightarrow \lambda^{(8+\delta)/3}$, and all the structure of the model remains intact.

C. The Subtraction Term

Within the assumptions of the model we have now computed the Reggeized asymptotic amplitude in the s channel,

$$M_r = M_I + M_{III} \quad (2C.1)$$

There yet remains to compute the corresponding physical amplitude whose square is proportional to the cross section of the physical process. These two cannot be the same; they differ by a constant in s , a subtraction term, engendered by a Kronecker delta singularity in the J plane at $J = 0$:

$$m_p = m_r + \delta_{J,0}(\Delta m) \quad (2C.2)$$

To obtain the physical amplitude one must omit from the Reggeized amplitude all contributions arising from terms in the potential connecting sense channels to nonsense at all nonsense (half-) integers - here

$V_{13} = V_{31}$ and $V_{23} = V_{32}$ at $J = 0$. If one attempts the calculation

of Δm within the confines of the model, it appears that the subtraction is zero, due to divergences encountered at $J = 0$. This, however, cannot in fact be so, and we now argue that something can be known about the subtraction term, though the model denies us exact knowledge of it. The pole α_{III} emerges from the Kronecker singularity as the coupling g'^2/π or λ , is turned on, and it must exactly cancel it at zero coupling. A pole can cancel a delta function if its residue and position are limitingly small of the same order of magnitude and their ratio is equal (and opposite) to the coefficient of the delta function:

$$\lim_{\alpha_{III} \sim \bar{\gamma}_{III} \rightarrow 0} \frac{\bar{\gamma}_{III}}{J - \alpha_{III}} = -\frac{\bar{\gamma}_{III}}{\alpha_{III}} \delta_{J,0} \quad , \quad (2C.3)$$

and $\bar{\gamma}_{III}/\alpha_{III}$ is finite at zero coupling, of order λ^0 . Thus the real term Δm must equal $(\bar{\gamma}_{III}/\alpha_{III}) + M_s$, where M_s , a function of t only, is of finite positive order in the coupling λ . This leaves us with the physical amplitude:

$$M_p = M_I + M_s - \bar{\gamma}_{III} \left[\ln\left(\frac{s}{\mu^2}\right) + i\pi \right] \quad . \quad (2C.4)$$

We can say no more of the subtraction term M_s , so let us proceed to see what can be said even in this ignorance.

III. COMPARISON OF THE MODEL WITH EXPERIMENT

To make a comparison of the results of this model with experiment, we allow that, of the terms contained in M_p , M_I represent the contribution of ordinary falling Regge trajectories as determined by present Regge fits; we do not attempt to allow that our α_I pole as calculated above has any physical reality. The remaining terms we retain intact as representing in some fashion the contributions of the quark-induced pole and subtraction. When and if energies and momentum transfers are reached at which contributions as these become visible, data would be fitted to a theory similar to but more nearly correct than the model presented here, rendering a determination of the quark's mass and coupling.

The experimental situation with which we have chosen to compare this model in qualitative fashion is proton-proton scattering at large angles. Immediately after these considerations, a crudely quantitative comparison will be made with the data at 90° in the center of mass. Huang and Pinsky⁶ have observed that the effective trajectory, α_e , may be analyzed into two parts (see the schematic diagram Fig. 4): the pomeron, α_p , and the dipomeron cut, α_{pp} . The trajectory α_e falls from one to below zero as α_p , then rises slightly to the α_{pp} trajectory, and then falls off again as the α_{pp} to the experimental

limits on $-t$ in the present data. Thus, below $t \approx -3.7$, where all ordinary Regge poles (or cuts) have dropped below zero, α_{III} is the conventional "dominant" pole-but not the dominating pole, since α_e shows no signs of rising again to zero, as we might expect if a pole at or near $J = 0$ were actually to dominate. So with present data there is no reason to suppose that a quark, as we have described it, should exist; our task is now to estimate (to order of magnitude only) what is the least mass that would be compatible with the present data, supposing the quark exists.

Since the imaginary part of the scattering amplitude, $M_{Imp} = M_{ImI} - i\pi\bar{\gamma}_{III}$, is effectively M_{ImI} , $\pi^2\bar{\gamma}_{III}^2 \ll M_{ImI}^2 \approx M_{Imp}^2$. But $|M_p|^2 \geq M_{Imp}^2$. Thus we derive the inequality which must be satisfied:

$$\frac{d\sigma}{dt} \gg \frac{4\pi}{sq_s^2} \pi^2 \bar{\gamma}_{III}^2 \equiv \frac{d\sigma}{dt} \Big|_{ImIII} \quad (3.1a)$$

Similarly,

$$\frac{d\sigma}{dt} \gg \frac{4\pi}{sq_s^2} \left[\bar{\gamma}_{III} \ln\left(\frac{s}{\mu^2}\right) - M_s \right]^2 \equiv \frac{d\sigma}{dt} \Big|_{ReIII} \quad (3.1b)$$

The latter is of little use until we can compute M_s , but in the following its right-hand side with $M_s = 0$ will be displayed.

In Fig. 5 the data of Akerlof et al.,⁷ for proton-proton scattering at 90° in the center of mass are compared with $\frac{d\sigma}{dt} \Big|_{ImIII, 90^\circ}$

and $\left. \frac{d\sigma}{dt} \right|_{\text{ReIII}, 90^\circ}$ (for $M_s = 0$). We have arbitrarily chosen $\left(\frac{g^2}{\pi}\right) = 2$
 $\left(\frac{g'^2}{\pi}\right) = 1$ and calculated these quantities at $\lambda = 0.2, 0.1,$ and 0.05 ,
i.e., $M = 5, 10,$ and 20 times the mass of the proton. In order to
satisfy the inequality, Eq. (3.1), M would certainly have to be
greater than about 10 GeV, and one may guess this to be a decent order
of magnitude lower bound on this mass as may be gleaned from these
data, under the many suppositions of the model.

IV. CONCLUSION

It appears that whenever approximations are made on our model, they are expected to be correct to order of magnitude in the mass M of the quark, except that the unknown behavior of the coupling constant g' as a function of λ , was assumed to be constant. Thus we can expect our result to be correct only to order of magnitude, and none of the consequences of subtle details, like the misrepresentation of a spinor by a scalar, to be portrayed with any fidelity. It is in this spirit that the result $M \gtrsim 10 \text{ GeV}$ is stated.

It is not intended that the model presented here be more than a naive first effort towards an analytic approach to the confrontation of idea and fact. The experimental facts, the data, are there; the idea may be well stated in its theoretical context: it is the mapping between the two worlds which is lacking, the model which must be improved. The minor improvements which could be made on this model are many, other similar models could be suggested, or more radical means of relating the idea to the data may be tried. The idea remains an attractive one: that there is a crucial difference of character between composite particles and elementary ones, that if the latter exist, observable effects must arise, and that these may be manifestations of the operation of higher symmetries.

ACKNOWLEDGMENTS

The author expresses his appreciation of the guidance of his research advisor, Professor S. Mandelstam. The main body of this work was done under the auspices of the United States Atomic Energy Commission, and was begun with the assistance of the National Science Foundation.

FOOTNOTES

- * This work was supported in part by the United State Atomic Energy Commission.
- + This paper is based on author's Ph.D. thesis UCRL-18251.
- 1. H. Kasha, L. B. Leipuner, T. P. Wangler, J. Alspectro, and R. K. Adair, Phys. Rev. 154, 1263 (1967): and references.
- 2. S. Mandelstam, in Elementary Particle Physics, Part II, G. Takeda and A. Fujii, editors (Benjamin and Syokabō, New York and Tokyo (1967); and Phys. Rev. 166, 1539 (1968).
- 3. S. Mandelstam, Phys. Rev. 137, 949 (1965).
- 4. W. R. Fraser and J. R. Fulco, Phys. Rev. 117, 1603 (1960).
- 5. J. Hartle and C. E. Jones, Phys. Rev. 140B, 90 (1965).
- 6. K. Huang and S. Pinsky, Massachusetts Institute of Technology preprint (1968).
- 7. C. W. Akerlof, R. H. Hieber, A. D. Krish, K. W. Edwards, L. D. Ratner, and K. Ruddick, Phys. Rev. 159, 1138 (1967).

FIGURE CAPTIONS

- Fig. 1. The potential chosen in the model.
- Fig. 2. $\alpha_I(x)$ and $\bar{\gamma}_I(x)$ for $\frac{g^2}{\pi} = 2$.
- Fig. 3a. $j_{III}(x)$ for $\frac{g^2}{\pi} = 2$, $\frac{g'^2}{\pi} = 1$, and $\lambda = 0.2, 0.1, 0.05$.
- Fig. 3b. $\bar{\gamma}'_{III}(x)$ under the same conditions.
- Fig. 4. Schematic diagram of the trajectories of proton-proton scattering.
- Fig. 5. Comparison of the model with experiment for proton-proton scattering at 90° in the center of mass. The curve marked "proton-proton" is a close fit to the data of Akerlof et al., the broken-line portion denoting preliminary results.
- The three pairs of curves are $\left. \frac{d\sigma}{dt} \right|_{\text{ImIII}, 90^\circ}$ and $\left. \frac{d\sigma}{dt} \right|_{\text{ReIII}, 90^\circ, M_S=0}$
- for $\lambda = 0.2, 0.1, 0.05$, with $\frac{g^2}{\pi} = 2$, $\frac{g'^2}{\pi} = 1$ and $\mu = m_{\text{proton}}$.




Fig. 1

XBL685-2735




Fig. 2

XBL685-2736




Fig. 3a

XBL685-2737




Fig. 3b

XBL685-2738




Fig. 4

XBL685-2739




Fig. 5

XBL685-2740

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

