

Lawrence Berkeley National Laboratory

Recent Work

Title

A NEW TERBIUM ISOTOPE, Tb152

Permalink

<https://escholarship.org/uc/item/8c82v7xm>

Authors

Toth, Kenneth S.
Faler, Kenneth T.
Rasmussen, John O.

Publication Date

1958-10-01

UNIVERSITY OF
CALIFORNIA

*Radiation
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

For publication in Physical Review

UCRL-8446

UNIVERSITY OF CALIFORNIA

Radiation Laboratory
Berkeley, California

Contract No. W-7405-eng-48

A NEW TERBIUM ISOTOPE, Tb^{152}

Kenneth S. Toth, Kenneth T. Faler, and John O. Rasmussen

October 1958

Printed for the U. S. Atomic Energy Commission

A NEW TERBIUM ISOTOPE, Tb^{152*}

Kenneth S. Toth,[†] Kenneth T. Faler, and John O. Rasmussen

Radiation Laboratory and Department of Chemistry
University of California, Berkeley, California

October, 1958

Terbium-152 has not been previously reported despite the fact that Tb¹⁵¹, Tb¹⁵³, and Tb¹⁵⁴ have been known for some time. In this study Tb¹⁵² has been produced and mass-assigned in a series of bombardments using alpha particles accelerated in the Berkeley 60-inch cyclotron. The half-life has been found to be 18.5 ± 0.5 hr. The similarity of the half-lives of two neighboring isotopes, Tb¹⁵¹ (19 hr) and Tb¹⁵⁴ (17 hr) makes it clear why this isotope has been previously overlooked.

A series of bombardments were undertaken using the enriched isotopes, Eu¹⁵¹ (91.9% Eu¹⁵¹, 8.1% Eu¹⁵³) and Eu¹⁵³ (5.0% Eu¹⁵¹, 95.0% Eu¹⁵³) with bombarding energies of 48 Mev and 37 Mev. At the higher energy the most probable reaction is ($\alpha,4n$), while the lower energy is below the threshold for this reaction and produced mostly ($\alpha,3n$).

After bombardment the elements were chemically separated by means of a cation-exchange method described elsewhere.¹ Gamma rays emitted by the terbium isotopes were studied using 3 x 3 inch and 1 x 1-1/2 inch NaI(Tl) crystal scintillation counters in conjunction with a 100-channel pulse-height analyzer. Accurate gamma ray energies were obtained from conversion electron spectra taken in 180-degree-focusing permanent-magnet electron spectrographs described by Smith and Hollander.²

In figure 1, curve A shows the gamma spectrum of the chemically separated terbium isotopes following 37 Mev alpha-particle bombardment of Eu¹⁵³. This sample should consist mostly of Tb¹⁵⁴ with smaller amounts of

* This work was performed under the auspices of the U. S. Atomic Energy Commission.

[†] Present address: University Institute for Theoretical Physics, Copenhagen, Denmark.

Tb¹⁵⁵ and very little Tb¹⁵³ and Tb¹⁵². The three peaks have energies (determined precisely by conversion electron measurements) of 123.2, 248.1, and 347.1 keV and have previously been ascribed to Tb¹⁵⁴ decay.^{3,4}

Curve B of figure 1 shows the gamma spectrum following 37 MeV alpha-particle bombardment of Eu¹⁵¹. Here the majority of product should be Tb¹⁵² with smaller amounts of Tb¹⁵³ and Tb¹⁵⁴. Again a peak is seen at 340 keV and its rate of decay as obtained from a series of spectra taken at various times after bombardment indicates a half-life nearly the same as that of Tb¹⁵⁴. That it is not, in fact, the same gamma as the 347.1 keV one of Tb¹⁵⁴ is readily seen when its intensity is compared with the intensities of the other peaks seen in the Tb¹⁵⁴ spectrum in curve A. In curve B the relative intensities of the 120 and 240 keV gammas are greatly reduced. This evidence indicates the presence of a new isotope whose decay includes a prominent 340 keV gamma. Since Tb¹⁵³ is known to have a half-life of 60 hr,^{3,5} the isotope must be Tb¹⁵² or perhaps Tb¹⁵¹ which also has nearly the same half-life. Since this bombardment was at an energy below the threshold for production of Tb¹⁵¹ it is most probable that the gamma belongs to Tb¹⁵², and additional evidence is obtained by comparison with curve C of figure 1. This is the spectrum of gamma rays following a 48 MeV bombardment of Eu¹⁵¹ which produced an abundance of Tb¹⁵¹. This spectrum was obtained with a 1 x 1-1/2 inch scintillation crystal and thus in a visual intensity comparison with the upper curves, the different counting efficiencies must be borne in mind. It is possible, however, to obtain relative intensities of gamma rays and correct these for differences in counting efficiencies in order to make comparisons between curves B and C. This having been done, the corrected intensity ratio of the 340 keV gamma to the 110 keV gamma (assuming all of the 110 keV peaks in curves B and C to be due to Tb¹⁵¹) is 9.5 to 1 in curve B and 5.8 to 1 in curve C. This variation strongly supports assignment of the 340 keV gamma to Tb¹⁵².

The spectrum shown in curve B was obtained after an elapsed time of 32 hours following bombardment and thus some of the peaks shown are due to gamma rays of longer-lived nuclides. The peak at 212.2 keV is the prominent gamma seen in the decay of 62 hr Tb¹⁵³. The peaks near 100 keV are complex, consisting of the 97.3 and 103.1 keV gammas from the decay of Gd¹⁵³ and

several weak gammas from Tb^{153} decay. It is also possible that there is a gamma of about 90 keV in Tb^{152} . No gammas are seen which can be assigned to Tb^{151} or Tb^{154} . The rate of decay of the peaks at 180 and 265 keV indicate that these also belong to the decay of Tb^{152} . A spectrum taken at higher energies shows a large number of low intensity gammas which could not be resolved and assigned with certainty. From the high-energy side of the 340 keV peak may be resolved a gamma of 415 keV energy with an intensity about 20% that of the 340. This is doubtless the 413 keV gamma seen in the decay of Eu^{152} to the same daughter nuclide.


Several permanent magnet spectrograph sources were made from the terbium fraction of the sample obtained in the 48 MeV alpha particle bombardment of Eu^{151} . Conversion electron energies were compared with energies observed in the decay of Eu^{152} on the same instrument and corresponded exactly to those of the 344.1 keV gamma seen here and elsewhere.

In addition to the two gamma rays discussed above, there are peaks at 180 and 265 keV in spectrum B which decayed with about the correct half-life to indicate that they belonged to Tb^{152} decay, and there was some indication that a gamma around 90 keV also had a 19 hour half-life but this was less certain. An additional gamma was seen at 125 keV which decayed very rapidly and whose identity or half-life was not determined.

Finally, alpha particles from Tb^{152} decay were sought in the sample obtained from the 37 MeV bombardment of Eu^{151} . No alpha counts were detected, and estimating the disintegration rate of Tb^{152} from the K x-ray peak together with the statistical uncertainty in the alpha counter background allowed an alpha-to-electron capture branching ratio to be set at a limit of $< 10^{-7}$. This number is to be compared with the branching ratio of 3×10^{-6} for Tb^{151} which has been recently determined experimentally by Toth and Rasmussen.⁵

REFERENCES

1. Thompson, Harvey, Choppin, and Seaborg, J. Am. Chem. Soc. 76, 6229 (1954).
2. W. G. Smith and J. M. Hollander, Phys. Rev. 101, 746 (1956).
3. Mihelich, Harmatz, and Handley, Phys. Rev. 108, 989 (1957).
4. K. S. Toth and J. O. Rasmussen, University of California Radiation Laboratory Report UCRL-8373 (to be published, Phys. Rev.).
5. K. S. Toth and J. O. Rasmussen, unpublished data.



MJ-16212

Fig. 1. Gamma spectra of terbium isotopes by alpha-particle bombardment of enriched europium isotopes. (A) 37-Mev alphas on Eu^{153} , 3 x 3-inch NaI(Tl) detector; (B) 37-Mev alphas on Eu^{151} , 3 x 3-inch detector; and (C) 48-Mev alphas on Eu^{151} , 1.5 x 1-inch detector.