

UC Davis

UC Davis Previously Published Works

Title

Finished Genome Sequence of the Indole-3-Acetic Acid-Catabolizing Bacterium
Pseudomonas putida 1290

Permalink

<https://escholarship.org/uc/item/8cm9j84x>

Journal

Microbiology Resource Announcements, 8(28)

ISSN

2576-098X

Authors

Laird, Tyler S
Leveau, Johan HJ

Publication Date

2019-07-11


DOI

10.1128/mra.00519-19

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed



Finished Genome Sequence of the Indole-3-Acetic Acid-Catabolizing Bacterium *Pseudomonas putida* 1290

Tyler S. Laird,^a Johan H. J. Leveau^a

^aDepartment of Plant Pathology, University of California, Davis, Davis, California, USA

ABSTRACT Use of indole-3-acetic acid (IAA) as a carbon, nitrogen, and energy source by *Pseudomonas putida* 1290 is linked to the possession of a gene cluster that codes for conversion to catechol. Here, we present the genomic context of this *iac* gene cluster, which includes genes for IAA chemotaxis/transport and catechol catabolism.

Originally isolated from the pear phyllosphere (1), *Pseudomonas putida* 1290 was the first bacterium for which the ability to grow at the expense of the plant hormone indole-3-acetic acid (IAA) was linked to a gene cluster designated *iacABCDEFGH-I*, which allows the enzymatic conversion of IAA to catechol for further channeling into central metabolism via the beta-ketoadipate pathway (2). *P. putida* 1290 also exhibits positive chemotaxis toward IAA (3). To better understand the genomic context within which the *iac* gene cluster operates in this strain, we sequenced the genome of *P. putida* 1290.

Genomic DNA was extracted from a single-colony culture of *P. putida* 1290 grown overnight in LB, using a DNeasy blood and tissue kit (Qiagen, Valencia, CA) to construct a 10-kb library for PacBio RS II sequencing on 3 single-molecule real-time (SMRT) cells using P4-C2 chemistry at the UC Davis Genome Center. The total number of reads was 160,877, covering a total of 597,416,768 bp (average subread length, 3,713 bp; N_{50} subread length, 5,224 bp). *De novo* assembly was accomplished with Hierarchical Genome Assembly Process (HGAP) version 3 within SMRT Analysis software v2.2.0 (Pacific Biosciences) (4). Default parameters were used for this and all other software. Assembly correction was performed with Quiver and Gepard v1.30 (5), resulting in two circularized contigs, a 6,495,886-bp chromosome (G+C content, 63.14%; 62.26× coverage) and a 114,265-bp, major facilitator superfamily (MFS)-type I plasmid (6) designated pPp1290 (G+C content, 54.59%; 25.17× coverage). Pairwise average nucleotide identity values based on fastANI (7) showed the closest genomic match (87.4%) with soil isolate *P. putida* DRA525 (8). The plasmid most similar to pPp1290 in the PLSDb database (9) was pGRT1 from *P. putida* DOT-T1E (10). Gene prediction by RAST (11) revealed 6,048 coding sequences, 77 tRNA genes, and 20 rRNA genes on the chromosome and 123 coding sequences on pPp1290.

We identified an open reading frame (ORF), directly upstream of the *iacABCDEFGH-I* gene cluster in *P. putida* 1290, coding for a LysR-type transcriptional regulator, a partial duplication of the *iac* gene cluster (*iacCDF*), genes predicted to code for an OprD-like outer membrane protein, an ATP-binding protein belonging to the ABC family of transporters, and an MFS-type transporter similar to lact1 from *Paraburkholderia phytofirmans* PsJN, implicated in the transport of an IAA degradation pathway intermediate (12). Also upstream of the *iac* gene cluster, an ORF predicted to code for a methyl-accepting chemotaxis protein possibly involved in the chemotaxis of *P. putida* toward IAA (3) and a *catR-catBCA* gene cluster for catechol catabolism (3) were found. Interestingly, the *iac* gene cluster appears to be part of an approximately 30-kb

Citation Laird TS, Leveau JHJ. 2019. Finished genome sequence of the indole-3-acetic acid-catabolizing bacterium *Pseudomonas putida* 1290. Microbiol Resour Announc 8:e00519-19. <https://doi.org/10.1128/MRA.00519-19>.

Editor David A. Baltus, University of Arizona

Copyright © 2019 Laird and Leveau. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Johan H. J. Leveau, jleveau@ucdavis.edu.

Received 6 May 2019

Accepted 19 June 2019

Published 11 July 2019

genomic island predicted by the IslandPick algorithm in IslandViewer v4 (13) and encompassing multiple genes for the utilization of plant-derived compounds other than IAA, including methylamine (14), phenylacetaldehyde (15), and opines (16). The presence of *aldA* and *aldB* homologs (17) on the chromosome is consistent with IAA production reported for *P. putida* 1290 (2). Plasmid pPp1290 harbors multiple insertion sequence (IS)-like elements, as well as genes related to aromatic compound degradation, stress tolerance, fluoride efflux, and conjugal transfer.

Data availability. The *P. putida* 1290 chromosome and plasmid sequences are available under GenBank accession numbers [CP039371](#) and [CP039372](#), respectively. Raw reads from the three SMRT cells are available under the SRA accession number [SRP195595](#).

ACKNOWLEDGMENTS

We thank Joseph Fass (UC Davis Genome Center) for his help with genome assembly.

This study was funded by USDA-NIFA grant number 2011-67017-30024. We also acknowledge financial support from UC Davis to T.S.L. in the form of a Dean's Distinguished Graduate Fellowship. The funders had no role in the study design, data collection and interpretation, or the decision to submit the work for publication.

REFERENCES

- Leveau JHJ, Lindow SE. 2005. Utilization of the plant hormone indole-3-acetic acid for growth by *Pseudomonas putida* strain 1290. *Appl Environ Microbiol* 71:2365–2371. <https://doi.org/10.1128/AEM.71.5.2365-2371.2005>.
- Leveau JHJ, Gerards S. 2008. Discovery of a bacterial gene cluster for catabolism of the plant hormone indole 3-acetic acid. *FEMS Microbiol Ecol* 65:238–250. <https://doi.org/10.1111/j.1574-6941.2008.00436.x>.
- Scott JC, Greenhut IV, Leveau JHJ. 2013. Functional characterization of the bacterial *iac* genes for degradation of the plant hormone indole-3-acetic acid. *J Chem Ecol* 39:942–951. <https://doi.org/10.1007/s10886-013-0324-x>.
- Chin C-S, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, Clum A, Copeland A, Huddleston J, Eichler EE, Turner SW, Korlach J. 2013. Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. *Nat Methods* 10:563–569. <https://doi.org/10.1038/nmeth.2474>.
- Krumsiek J, Arnold R, Rattei T. 2007. Gepard: a rapid and sensitive tool for creating dotplots on genome scale. *Bioinformatics* 23:1026–1028. <https://doi.org/10.1093/bioinformatics/btm039>.
- Robertson J, Nash JHE. 2018. MOB-suite: software tools for clustering, reconstruction and typing of plasmids from draft assemblies. *Microb Genom* 4:e000206. <https://doi.org/10.1099/mgen.0.000206>.
- Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S. 2018. High throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat Commun* 9:5114. <https://doi.org/10.1038/s41467-018-07641-9>.
- Drace K, Giangli S, LeSar C, Kiefer AM. 2018. Draft genome sequence of mercury-resistant *Pseudomonas putida* strain DRA525. *Genome Announc* 6:e00370-18. <https://doi.org/10.1128/genomeA.00370-18>.
- Galata V, Fehlmann T, Backes C, Keller A. 2019. PLSDB: a resource of complete bacterial plasmids. *Nucleic Acids Res* 47:D195–D202. <https://doi.org/10.1093/nar/gky1050>.
- Molina L, Duque E, Gómez MJ, Krell T, Lacal J, García-Puente A, García V, Matilla MA, Ramos J-L, Segura A. 2011. The pGRT1 plasmid of *Pseudomonas putida* DOT-T1E encodes functions relevant for survival under harsh conditions in the environment. *Environ Microbiol* 13:2315–2327. <https://doi.org/10.1111/j.1462-2920.2011.02492.x>.
- Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <https://doi.org/10.1186/1471-2164-9-75>.
- Donoso R, Leiva-Novoa P, Zúñiga A, Timmermann T, Recabarren-Gajardo G, González B. 2017. Biochemical and genetic bases of indole-3-acetic acid (auxin phytohormone) degradation by the plant-growth-promoting rhizobacterium *Paraburkholderia phytofirmans* PsJN. *Appl Environ Microbiol* 83:e01991-16. <https://doi.org/10.1128/AEM.01991-16>.
- Bertelli C, Laird MR, Williams KP, Simon Fraser University Research Computing Group, Lau BY, Hoard G, Winsor GL, Brinkman FSL. 2017. IslandViewer 4: expanded prediction of genomic islands for larger-scale datasets. *Nucleic Acids Res* 45:W30–W35. <https://doi.org/10.1093/nar/gkx343>.
- Shiraishi K, Oku M, Kawaguchi K, Uchida D, Yurimoto H, Sakai Y. 2015. Yeast nitrogen utilization in the phyllosphere during plant lifespan under regulation of autophagy. *Sci Rep* 5:9719. <https://doi.org/10.1038/srep09719>.
- Dudareva N, Klempien A, Muhlemann JK, Kaplan I. 2013. Biosynthesis, function and metabolic engineering of plant volatile organic compounds. *New Phytol* 198:16–32. <https://doi.org/10.1111/nph.12145>.
- Moore LW, Chilton WS, Canfield ML. 1997. Diversity of opines and opine-catabolizing bacteria isolated from naturally occurring crown gall tumors. *Appl Environ Microbiol* 63:201–207.
- McClerkin SA, Lee SG, Harper CP, Nwumeh R, Jez JM, Kunkel BN. 2018. Indole-3-acetaldehyde dehydrogenase-dependent auxin synthesis contributes to virulence of *Pseudomonas syringae* strain DC3000. *PLoS Pathog* 14:e1006811. <https://doi.org/10.1371/journal.ppat.1006811>.