# Lawrence Berkeley National Laboratory

**Recent Work** 

## Title

THE DISCOVERY OF A SECOND NARROW RESONANCE IN e+e- ANNIHILATION

## Permalink

https://escholarship.org/uc/item/8cx6g66t

## Authors

Abrams, G.S. Briggs, D. Chinowsky, W. <u>et al.</u>

## **Publication Date**

1974-11-01

THE DISVOERY OF A SECOND NARROW RESONANCE IN e<sup>+</sup>e<sup>-</sup> ANNIHILATION

- "%



**\***?

5

2

ా

0

ung**e** ⊰

3

Э

Э



LBL-3605

**C**.

. 3

#### DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California. × SIAC-PUB-1510 /LBL-3605 November 1974 (T/E)

#### THE DISCOVERY OF A SECOND NARROW RESONANCE IN e<sup>+</sup>e<sup>-</sup> ANNIHILATION\*

G. S. Abrams, D. Briggs, W. Chinowsky, C. E. Friedberg,
G. Goldhaber, R. J. Hollebeek, J. A. Kadyk, A. Litke, B. Lulu,
F. Pierre<sup>1</sup>, B. Sadoulet, G. H. Trilling, J. S. Whitaker,
J. Wiss, and J. E. Zipse

Lawrence Berkeley Laboratory and Department of Physics University of California, Berkeley, California 94720

J.-E. Augustin, A. M. Boyarski, M. Breidenbach, F. Bulos, G. J. Feldman, G. E. Fischer, D. Fryberger, G. Hanson, B. Jean-Marie, R. R. Larsen, V. Luth, H. L. Lynch, D. Lyon, C. C. Morehouse, J. M. Paterson, M. L. Perl, B. Richter, P. Rapidis, R. F. Schwitters, W. Tanenbaum, and F. Vannucci‡

> Stanford Linear Accelerator Center Stanford University, Stanford, California 94305

#### ABSTRACT

We have observed a second sharp peak in the cross section for  $e^+e^- \rightarrow$  hadrons at a center-of-mass energy of 3.695  $\stackrel{+}{}$  0.004 GeV. The upper limit of the full width at half maximum is 2.7 MeV.

(Submitted to Phys. Rev. Letters)

\* Work supported by the U. S. Atomic Energy Commission † CEN, Saclay, France

M)

٠.

0

 $\bigcirc$ 

M

٦.

naniy Lucius

 $\Box$ 

3

‡ Laboratoire de l'Accélérateur Linéaire, Centre d'Orsay de l'Université de Paris, 91 Orsay, France

FInstitut de Physique Nucléaire, Orsay, France

The recent discovery of a very narrow resonant state coupled to leptons and hadrons<sup>1,2,3</sup> has raised the obvious question of the existence of other narrow resonances also coupled to leptons and hadrons. We therefore began a systematic search of the mass region accessible with the SLAC e<sup>+</sup>e<sup>-</sup> storage ring SPEAR and quickly found a second narrow resonance decaying to hadrons. The parameters of the new state (which we suggest calling  $\psi(3695)$ )are

 $\odot$ 

where the mass uncertainty reflects the uncertainty in the absolute energy calibration of the storage ring.

The  $\psi(3695)$ , like the  $\psi(3105)$ , was found using the SLAC-LBL magnetic detector at SPEAR.<sup>4</sup> The luminosity monitoring, event acceptance criteria, and storage ring energy determination have been described previously.<sup>1</sup>

The new feature of this run is the search procedure used to hunt for narrow  $e^+e^-$  resonances. In the search mode the storage ring energy is increased in about 1 MeV steps ( $E_{CM} = 2 \times E_{BEAM}$ ) every three minutes. The data taken during each step are analyzed in real time and the relative cross sections computed at the end of each step. Figure 1a shows the search mode data taken during a calibration scan over the previously discovered  $\psi(3105)$ . Figure 1b shows the data taken during the first scan which began at a ring energy of 1.8 GeV. A clear indication of a narrow resonance with a mass of about 3.70 GeV is seen. It should be emphasized that we have not yet scanned any mass region other than that between 3.6 and 3.71 GeV.

-2-

#### LEGAL NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights. TECHNICAL INFORMATION DIVISION LAWRENCE BERKELEY LABORATORY UNIVERSITY OF CALIFORNIA BERKELEY, CALIFORNIA 94720