UCLA

UCLA Previously Published Works

Title

Examining the Impact of Separate Components of a Multicomponent Intervention Designed to Reduce At-Risk Drinking Among Older Adults: The Project SHARE Study

Permalink

<https://escholarship.org/uc/item/8d53c00v>

Journal

Alcoholism: Clinical and Experimental Research, 39(7)

ISSN

0145-6008

Authors

Duru, OK
Xu, H
Moore, AA
et al.

Publication Date

2015-07-01

DOI

10.1111/acer.12754

Peer reviewed

Examining the Impact of Separate Components of a Multicomponent Intervention Designed to Reduce At-Risk Drinking Among Older Adults: The Project SHARE Study

Obidiugwu K. Duru, Haiyong Xu, Alison A. Moore, Michelle Mirkin, Alfonso Ang, Louise Tallen, Chi-Hong Tseng, and Susan L. Ettner

Background: Health promotion interventions often include multiple components and several patient contacts. The objective of this study was to examine how participation within a multicomponent intervention (Project SHARE) is associated with changes in at-risk drinking among older adults.

Methods: We analyzed observational data from a cluster-randomized trial of 31 primary care physicians and their patients aged ≥ 60 years, at a community-based practice with 7 clinics. Recruitment occurred between 2005 and 2007. At-risk drinkers in a particular physician's practice were randomly assigned as a group to usual care ($n = 640$ patients) versus intervention ($n = 546$ patients). The intervention included personalized reports, educational materials, drinking diaries, in-person physician advice, and telephone counseling by health educators (HEs). The primary outcome was at-risk drinking at follow-up, defined by scores on the Comorbidity Alcohol Risk Evaluation Tool (CARET). Predictors included whether a physician-patient alcohol risk discussion occurred, HE call occurred, drinking agreement with the HE was made, and patients self-reported keeping a drinking diary as suggested by the HE.

Results: At 6 months, there was no association of at-risk drinking with having had a physician-patient discussion. Compared to having had no HE call, the odds of at-risk drinking at 6 months were lower if an agreement was made or patients reported keeping a diary (odds ratio [OR] 0.58, 95% confidence interval [CI] 0.37 to 0.90), or if an agreement was made and patients reported keeping a diary (OR 0.52, CI 0.28 to 0.97). At 12 months, a physician-patient discussion (OR 0.61, CI 0.38 to 0.98) or an agreement and reported use of a diary (OR 0.45, CI 0.25) were associated with lower odds of at-risk drinking.

Conclusions: Within the Project SHARE intervention, discussing alcohol risk with a physician, making a drinking agreement, and/or self-reporting the use of a drinking diary were associated with lower odds of at-risk drinking at follow-up. Future studies targeting at-risk drinking among older adults should consider incorporating both intervention components.

Key Words: Alcohol Use, Older Adults, Physician-Patient Discussion, Drinking Diary, Drinking Agreement.

APPROXIMATELY 30 to 40% of American adults aged 65 years and over drink alcohol regularly (Blazer and Wu, 2011; Kirchner et al., 2007). While heavy drinking is less common with advancing age, light or moderate drink-

ing has different risks and benefits for older adults compared with younger populations (Chan et al., 2009; Cooper et al., 2009; Reid et al., 2002). Several observational studies have found that older adults who drink moderately have better cognitive function, a reduction in cardiovascular disease, and decreased mortality compared with nondrinkers (Byles et al., 2006; Karlamangla et al., 2009; Lang et al., 2007a,b; Reid et al., 2006; Townsend et al., 2009). Conversely, age-related physiologic changes in older adults elevate blood alcohol levels per amount of alcohol consumed and increase brain sensitivity to the effects of alcohol, thereby increasing the potential risk of injury (Gilbertson et al., 2009; Vestal et al., 1977). Compared with younger persons, older adults also have more comorbid conditions and use more medications, including those that may increase risks with alcohol (e.g., nonsteroidal anti-inflammatories and warfarin) (Moore et al., 2007). Current practice guidelines recommend that adults be asked about their alcohol consumption as part of routine examinations as well as prior to prescribing medications that interact with alcohol (U.S. Department of Health & Human Services National Institute of Health National

From the Division of General Internal Medicine & Health Services Research (OKD, HX, MM, AA, LT, C-HT, SLE), Department of Medicine, David Geffen School of Medicine at University of California, Los Angeles, Los Angeles, California; Division of Geriatrics (AAM), Department of Medicine, University of California, Los Angeles, Los Angeles, California; Ahava Center for Spiritual Living (LT), Lexington, Kentucky; and Fielding School of Public Health (SLE), University of California, Los Angeles, California.

Received for publication July 8, 2014; accepted April 14, 2015.

Reprint requests: Obidiugwu K. Duru, MD, MSHS, Associate Professor of Medicine, David Geffen School of Medicine at University of California, Los Angeles, 10940 Wilshire Blvd., Suite 700, Los Angeles, CA 90024; Tel.: 310-794-8138; Fax: 310-794-0723; E-mail: kduru@mednet.ucla.edu

Copyright © 2015 by the Research Society on Alcoholism.

DOI: 10.1111/acer.12754

Institute on Alcohol Abuse and Alcoholism, 2007). However, physicians infrequently screen their existing patients for alcohol use and misuse, because they have limited time or because they simply may not remember to do so (Aira et al., 2003; Friedmann et al., 2000; Johansson et al., 2005; McCormick et al., 2006). As a result, there has been significant interest in developing effective interventions to reduce at-risk drinking in this population, and several intervention protocols have been implemented with varying degrees of success (Fink et al., 2005; Jonas et al., 2012; Lin et al., 2010a,b; Moore et al., 2011; Saitz et al., 2003; Whitlock et al., 2004).

While the evidence suggests that multicomponent interventions for at-risk adult drinkers are more successful than single brief interventions in reducing risky alcohol use (Jonas et al., 2012; Whitlock et al., 2004), there is significant heterogeneity in how these interventions have been designed. These interventions typically involve between 2 and 4 contacts with the patient over a period of several months and often combine several different components such as personalized counseling, drinking diaries to quantify alcohol consumption, and goal setting to reduce drinking to less harmful levels. However, some of these interventions are delivered entirely by primary care physicians, while others include nonphysicians, typically nurses or health educators (HEs). Some are limited to face-to-face patient contacts, while others include telephone calls. These reports do not frequently describe whether specific, individual components of these interventions are each linked to reductions in at-risk drinking. Distinguishing whether different intervention components (e.g., patient-physician discussions, keeping a drinking diary, etc.) are linked to reductions in at-risk drinking may help to identify the essential intervention components for future efforts to translate clinical trials into practice.

We recently conducted a large cluster-randomized clinical trial, designed to evaluate the effectiveness of a multicomponent intervention designed to reduce at-risk drinking among older adults, and defined by scores on the Comorbidity Alcohol Risk Evaluation Tool (CARET) (Moore et al., 2011). Project SHARE (Senior Health and Alcohol Risk Education) found a statistically significant decrease in at-risk drinking among older adults at 12-month follow-up (Ettner et al., 2014). While the intervention included several different components, including a physician component that involved physician-patient discussions of alcohol-related risk, as well as an HE component that involved HE telephone calls and encouragement to keep a drinking diary and/or make an agreement to cut down on alcohol intake, the specific mechanism/s that resulted in reductions in at-risk drinking are not clear. Therefore, we conducted an analysis to investigate to what extent participation in the different intervention components may be associated with the observed behavior change.

We hypothesized that Project SHARE patients with greater participation in the physician intervention component would have lower odds of being classified as at-risk drinkers at 6 months and at 12 months of follow-up, as compared to patients who did not participate in this intervention

component. We further hypothesized that patients who participated in the HE component, specifically those who (i) received a call from the HE but did not keep a drinking diary or make a drinking agreement, (ii) either made an agreement or self-reported that they kept a diary, and (iii) both made an agreement and self-reported that they kept a diary, would have lower odds of being classified as at-risk drinkers at 6 and at 12 months of follow-up, as compared to patients who did not participate at all in the HE intervention component.

MATERIALS AND METHODS

Study Design/Setting

The Project SHARE study population was drawn from primary care physicians, and their patients aged 60 years and over at Sansum Clinic, a community-based group practice with 7 clinics in and near Santa Barbara, California. The Project SHARE study was approved by the Institutional Review Board of the University of California, Los Angeles.

Sample—Primary Care Providers and Their Patients

All 43 Sansum Clinic physicians were approached about the study, and 31 (72%) agreed to participate. As shown in Fig. 1, Sansum Clinic staff then identified patients ($n = 12,573$) of these 31 physicians who were aged 60 years and older. From these lists, providers indicated which patients should be excluded due to severe cognitive impairment, terminal illness, whether they were known to be moving out of the area within the next year, or inability to speak and understand English. Of the remaining patients not excluded by the physicians, 9,476 were mailed recruitment letters and 6,919 (73%) agreed to participate. They were then screened by telephone to verify the self-reported absence of these same exclusion criteria and to confirm that they were current drinkers (consumed at least 1 drink in the prior 3 months).

Potentially eligible patients then completed a written survey including demographic and clinical information as well as the CARET. The CARET, an updated and revised version of the Alcohol-Related Problems Survey (Fink et al., 2002; Moore et al., 2011), identifies older adults at risk for harm from their alcohol consumption based on their responses to a set of questions about their clinical status, medication use, symptoms consistent with increased risk, and drinking habits (see the Appendix). Patients who completed the baseline survey and were classified as at-risk drinkers ($n = 1,186$) by the CARET were included in the Project SHARE intervention study and were randomized at the level of the primary care physician into experimental and control groups (Fig. 1). All patients, both control and experimental, were asked to complete follow-up surveys at 6 and 12 months that included the CARET items. All surveys in the Project SHARE study were mailed to patients.

Intervention

Experimental patients received a multicomponent intervention that included individualized patient risk reports provided to patients and to their physicians, as well as telephone-based counseling from an HE (Table 1). Both at baseline and at 6-month follow-up, CARET software was used to generate 2 risk reports with reasons for the classification as an at-risk drinker (e.g., taking blood thinning medications that could interact adversely with alcohol, symptoms of depression that may be exacerbated by alcohol, driving after drinking, etc.) and the potential harms that could result from the patient's at-risk drinking practices. One report was designed for and mailed to patients. The other was

Fig. 1. Participant flow in Project SHARE.

designed for physicians and was attached to the front of each experimental patient’s chart prior to each scheduled visit to an experimental physician.

Physicians received a single 2-hour training session from the study investigators prior to the patient recruitment, which provided guidance on how to interpret these reports and use them to initiate an alcohol risk discussion with the patient. The physicians

were asked to discuss the baseline and/or 6-month reports with the patient during regularly scheduled visits during the 12 months postbaseline, focusing on the relationship between risky drinking behavior and medication use and/or health conditions. After the appointment, the physician filled out a visit log confirming that a risk-related discussion did or did not take place and either describing the content of the discussion if it did take place or

Table 1. Components of the Project SHARE Intervention

Reports and educational materials for patients	<p>Materials mailed to patients:</p> <ol style="list-style-type: none"> 1. Comorbidity Alcohol Risk Evaluation Tool-generated individualized risk report with drinking classification (at-risk vs. not at-risk), reasons for the classification, and potential harms from the patient's at-risk drinking practices 2. Educational booklet ("Healthy Drinking as You Age") 3. Drinking diary ("Healthy Living as You Age") 4. Tip sheets (could include any combination of the following based on patient risk factors: automobile safety, drinking sensibly, diabetes, depression, preventing falls and fractures, gastritis, gout, heartburn, medications, forgetfulness, ulcers, liver disease, and sleep)
Telephone health educator (HE) intervention	<p>Five activities took place during HE telephone calls:</p> <ol style="list-style-type: none"> 1. Assessment and direct feedback <ul style="list-style-type: none"> • HE expressed concern regarding the patient's drinking pattern, linking it to the mailed risk report 2. Negotiation and goal setting 3. Behavioral modification techniques <ul style="list-style-type: none"> • HE helped patient identify high-risk situations in which drinking would likely occur, and familiarized the patient with coping techniques and with ways for establishing a support network to help in this process 4. Self-help-directed bibliotherapy (educational booklet discussed for reinforcement) 5. Follow-up and reinforcement
Brief provider intervention	<ol style="list-style-type: none"> 1. Providers discussed the individualized risk report with the patient, focusing on any risky drinking behavior and its relation to medication use and/or health conditions 2. Providers then set goals with patients for reducing their drinking 3. After each appointment, the physician completed a visit log detailing what s/he discussed with the patient

explaining why it did not take place, for example, due to lack of time.

The telephone-based HE intervention was based on the same risk report. HEs called each experimental patient 3 times: within 2 weeks after both the baseline and 6-month reports were mailed, as well as at the 3-month time period. The baseline and 6-month calls included explanation of the contents of the report, as well as reinforcement of the potential health consequences of the patient's drinking. The HE intervention component was essentially independent of the physician intervention component, because HEs were not aware of whether patients had already discussed alcohol use with their physicians, and vice-versa. The HE encouraged patients to keep a drinking diary detailing their daily alcohol intake and also encouraged them to make an agreement that they would cut down on their drinking to a specific negotiated goal. The HE also focused on behavior modification, such as identifying high-risk situations in which drinking would likely occur, and familiarized the patient with coping techniques for managing these situations. The 3-month calls were designed to check on patients' progress with their initial goals. After each call, the HE completed a form documenting the length of each call and content discussed. This form also included information on whether or not a drinking agreement was actually made and whether or not the patient reported that they kept a drinking diary.

Analysis of Participation in Different Intervention Components

To assess changes in at-risk drinking among patients who participated in different intervention components, we created separate variables to measure the separate components. For the physician component, using the records from the physician visit logs, we classified patients as either having had, or not had, an alcohol-related discussion with their physician at any time during the 12-month follow-up period. We created 4 categories to measure participation in the HE component: (i) "no HE intervention," defined as not receiving any HE calls, (ii) receiving at least 1 HE call but not making a drinking agreement or reporting that they kept a drinking diary, (iii) either making a drinking agreement or reporting that they kept a drinking diary, and (iv) making a drinking agreement and reporting that they kept a drinking diary. Notably, patients who were originally randomized to the experimental group in the Project SHARE study were classified in this analysis as having received no physician intervention if they did not have an alcohol-related discussion with their physician, and as having received no HE intervention if they did not receive any HE calls. Because all Project SHARE patients received a personalized risk report of their at-risk drinking as well as educational materials discussing at-risk drinking, we did not include these intervention components as predictor variables.

Other Measures

The dependent variable for this analysis, whether or not a patient was an at-risk drinker, was determined at 6 and 12 months based on patient responses to the CARET survey items. Covariates were drawn from the baseline survey and included gender, race, ethnicity, marital status, education, income, home ownership, and whether or not alcohol counseling from any provider took place in the 12 months prior to Project SHARE.

Statistical Methods

We used chi-squared tests for unadjusted comparisons between groups. We then constructed separate logistic regressions that included random effects for physicians, at 6-month follow-up and at 12-month follow-up. All analyses were completed using STATA Version 10.1 (Stata Corp, College Station, TX). Adjusted odds ratios (ORs) for at-risk drinking and 95% confidence intervals (CIs) are presented in the tables; in addition, predicted probabilities associated with statistically significant findings are presented in the text. The predicted probabilities were calculated using the "predict" command in STATA, which sets the physician random effect equal to zero.

We also examined several sensitivity analyses, first assuming that all of the patients who dropped out of the study and therefore had missing outcome values were at-risk drinkers at 6-month follow-up and 12-month follow-up, and then assuming that none of these patients were at-risk drinkers at follow-up. These sensitivity analyses did not produce different results from the main, complete-case analyses, and therefore, we only report results from the main analyses.

RESULTS

A total of 1,186 older adults met criteria for at-risk drinking, were enrolled in Project SHARE, and completed the baseline data collection survey. A subsample of 1,073 older adults remained in the study and completed the questionnaire at 6-month follow-up, and 1,049 remained in the study and completed the questionnaire at 12-month follow-up. Among the 419 patients in the intervention arm, 392 (94%), 315 (75%), and 364 (87%) received the baseline, 3-month,

Table 2. Unadjusted Participant Characteristics, by Whether or Not Physician-Level Intervention (Alcohol-Related Risk Discussion) Occurred During Project SHARE (*n* = 997)

	Risk discussion occurred during 12 months postbaseline (<i>n</i> = 300) (%)	Risk discussion did not occur (<i>n</i> = 697) (%)	<i>p</i> -Value
Female gender	36.0	33.9	0.51
Latino ethnicity	5.7	5.3	0.82
Age			
60 to 64 years	18.0	24.1	0.12
65 to 69 years	30.0	28.1	
70 to 74 years	17.7	19.7	
75 to 79 years	17.3	14.9	
80 years or greater	17.3	13.2	
Race			
White	95.3	98.1	0.06
Black	0.67	0.29	
Asian	2.33	0.57	
American Indian	1.7	1.7	
Marital status			
Married	78.5	71.3	0.11
Divorced or separated	8.9	12.3	
Widowed	13.3	10.3	
Never married	3.0	2.3	
Education level			
Less than high school	2.7	3.2	0.62
High school or GED	11.0	9.0	
Some college	28.3	25.5	
Graduated college	25.7	25.8	
Professional or graduate degree	32.3	36.4	
Annual household income			
<\$30,000	12.7	8.9	0.25
>\$30,000 to \$40,000	10.0	7.9	
>\$40,000 to \$60,000	17.7	17.4	
>\$60,000 to \$80,000	16.7	16.4	
>\$80,000 to \$100,000	13.3	18.4	
>\$100,000 to \$200,000	21.7	21.5	
>\$200,000	8.0	9.6	
Own home	87.0	89.2	0.31
Patient received alcohol counseling in the past from any provider			
Never	50.7	60.4	
During the last year	24.0	15.1	
More than 1 year ago	17.3	16.6	0.005
Mean # of alcoholic drinks/wk (SD)	13.5 (8.0)	13.7 (7.9)	
Categories of alcohol-related risk			
Taking medications that increase risk	62.3	59.7	0.43
Symptoms consistent with risk	63.0	60.8	0.52
Behaviors that increase risk	65.0	62.4	0.44

Results with *p*-value < 0.05 in bold font.

and 6-month HE calls. Furthermore, 37 received 1 call from the HE, 67 received 2 calls, and 300 received 3 calls, while 164 had 1 alcohol-related risk discussion with their physician, 65 had 2, 51 had 3, and 20 had 4 or more. We had <5% missing data for covariates and limited the multivariate analyses to the 997 patients for whom we had complete data at both follow-up time points.

Greater than 95% of the sample was white, and most patients were men (Tables 2 and 3). More than half of the

sample were college graduates and had annual household incomes exceeding \$60,000. On average, the patients in the sample consumed 13.7 alcoholic drinks per week. Each of 3 relatively common risk factors for at-risk drinking (taking medications that increase risk, reporting symptoms consistent with risk, or engaging in behaviors that increase risk) was present in approximately 60% of the sample. Compared to patients who did not have a discussion of alcohol risk during the 12-month study window, those who had a discussion were more likely (*p* = 0.005) to have already received some alcohol counseling in the past 12 months (24% vs. 15%) or over 12 months ago (17.3% vs. 16.6%); we controlled for these differences in the multivariate analyses. There were no other statistically significant differences between patients who had a risk discussion with their physician during the study window and those who did not (Table 2), or between patients by level of HE intervention (Table 3).

In adjusted multivariate analyses, the odds of being an at-risk drinker at 6 months did not vary significantly by whether or not a patient-physician alcohol risk discussion took place (Table 4). Compared to patients who did not receive an HE intervention, those who either made a drinking agreement or reported that they kept a drinking diary had a lower probability of being an at-risk drinker at 6 months (OR = 0.58, 95% CI 0.37 to 0.90; number needed to treat [NNT] of 10). Patients who made a drinking agreement and reported that they kept a drinking diary also had a lower odds of being an at-risk drinker at 6 months (OR = 0.52, 95% CI 0.28 to 0.97; NNT of 9). The predicted probability of at-risk drinking at 6 months would be reduced from 71% if patients did not receive an HE intervention to 60% if patients kept a drinking diary or made a drinking agreement, and would be further reduced to 58% if patients did both.

As shown in Table 4, compared with patients who did not have a discussion, patients who had an alcohol-related risk discussion with their physician had lower odds of being an at-risk drinker at 12 months (OR 0.61, 95% CI 0.38 to 0.98, NNT of 10). The predicted probability of at-risk drinking at 12 months would be reduced from 67% if patients did not have an alcohol-related discussion with their physician to 54% if they did have such a discussion. No difference was seen in the probability of being an at-risk drinker at 12 months between patients with no HE calls and those who either only received an HE call without making a drinking agreement or reporting that they kept a drinking diary, or those who either made an agreement or reported that they kept a diary. However, patients who both made a drinking agreement and reported that they kept a drinking diary had lower odds of being an at-risk drinker at 12 months (OR 0.45, 95% CI 0.25 to 0.82, NNT of 7) compared with patients who did not receive an HE intervention. The predicted probability of at-risk drinking at 12 months would be reduced from 68% if patients did not receive an HE intervention to 49% if they kept a drinking diary and made a drinking agreement.

Table 3. Adjusted Project SHARE Participant Characteristics, by Use of Components of the Health Educator (HE)-Level Intervention (*n* = 997)

	No phone call from HE (classified as no HE intervention) (<i>n</i> = 593) (%)	Patient received a phone call from HE			<i>p</i> -Value
		No drinking diary, no drinking agreement (<i>n</i> = 124) (%)	Drinking diary or drinking agreement (<i>n</i> = 184) (%)	Drinking diary and drinking agreement (<i>n</i> = 96) (%)	
Female gender	32.0	37.1	37.0	41.7	0.20
Latino ethnicity	5.7	4.0	4.9	6.3	0.85
Age					
60 to 64 years	23.8	25.0	16.3	20.8	0.05
65 to 69 years	27.5	35.5	28.8	26.0	
70 to 74 years	20.9	11.3	16.3	22.9	
75 to 79 years	14.2	16.1	18.5	18.8	
80 years or greater	13.7	12.1	20.1	11.5	
Race					
White	97.8	97.6	95.1	97.9	0.36
Black	0.3	0.0	1.1	0.0	
Asian	0.5	1.6	2.2	2.1	
American Indian	1.4	0.8	1.6	0.0	
Marital status					
Married	79.9	73.4	67.4	75.0	0.005
Divorced or separated	8.1	16.1	12.0	9.4	
Widowed	9.4	7.3	18.5	13.5	
Never married	2.5	3.2	2.2	2.1	
Education level					
Less than high school	3.0	1.6	3.3	4.2	0.79
High school or GED	9.8	7.3	10.3	10.4	
Some college	24.8	30.7	29.4	25.0	
Graduated college	25.1	29.8	23.4	29.2	
Professional or graduate degree	37.3	30.7	33.7	31.3	
Annual household income					
<\$30,000	8.6	11.3	13.6	10.4	0.14
>\$30,000 to \$40,000	7.6	8.1	12.5	7.3	
>\$40,000 to \$60,000	18.0	16.9	17.4	14.6	
>\$60,000 to \$80,000	15.2	20.2	12.5	27.1	
>\$80,000 to \$100,000	18.0	18.6	15.2	10.4	
>\$100,000 to \$200,000	22.4	18.6	20.7	21.9	
>\$200,000	10.1	6.5	8.2	8.3	
Own home	87.9	91.9	88.0	89.6	0.61
Patient received alcohol counseling in the past					
Never	58.9	54.8	56.0	55.2	0.13
During the last year	15.2	17.7	24.5	20.8	
More than 1 year ago	17.2	21.0	12.0	18.8	
Unknown	8.8	6.5	7.6	5.2	
Categories of alcohol-related risk					
Taking medications that increase risk	60.2	54.8	64.1	62.5	0.41
Symptoms consistent with risk	61.7	63.0	61.5	64.6	0.88
Behaviors that increase risk	65.1	61.3	57.6	64.6	0.30

Results with *p*-value < 0.05 in bold font.

DISCUSSION

This study examining data from Project SHARE supported our hypothesis that both the occurrence of a patient–physician risk discussion, and making a drinking agreement and/or self-report of keeping a drinking diary, were associated with lower rates of at-risk drinking at study follow-up. This suggests that both aspects of the intervention are important in reducing at-risk drinking and that efforts to reduce drinking in primary care settings should include both physicians and HEs.

We found that use of intervention components associated with the HE (drinking agreement, self-report of keeping a drinking diary) was associated with lower rates of at-risk drinking as compared to patients who did not

receive any HE calls, at both study end points. A prior analysis reported a benefit of receiving HE calls within a larger intervention study, specifically a reduction in the rates of at-risk drinking at 3 months if 3 calls were received within that time period (Maisto et al., 2001). These investigators did not find an effect of receiving HE calls at 12-month follow-up. They were unable to specifically examine the association between at-risk drinking and having completed a drinking agreement during these phone calls, because few patients had completed a drinking agreement. Our results indicate that completion of a drinking agreement, as well as self-reported use of a drinking diary, may be important in achieving larger decreases in at-risk drinking that are sustained over a longer period.

Table 4. Adjusted Odds of At-Risk Drinking Among Project SHARE Participants, at 6 and 12 Months (*n* = 997)

	Odds ratio (95% CI)	<i>p</i> -Value for difference
<i>6-month follow-up</i>		
Physician-level intervention		
Physician-patient alcohol risk discussion did not occur (REFERENCE)	1.0	
Physician-patient alcohol risk discussion occurred	0.81 (0.52 to 1.25)	0.34
Health educator (HE)-level intervention		
No HE intervention (REFERENCE)	1.0	
Patient received HE phone calls ONLY	0.72 (0.44 to 1.18)	0.19
Patient kept a drinking diary or made a drinking agreement	0.58 (0.37 to 0.90)	0.02
Patient kept a drinking diary and made a drinking agreement	0.52 (0.28 to 0.97)	0.04
<i>12-month follow-up</i>		
Physician-level intervention		
Physician-patient alcohol risk discussion did not occur (REFERENCE)	1.0	
Physician-patient alcohol risk discussion occurred	0.61 (0.38 to 0.98)	0.04
HE-level intervention		
No HE intervention (REFERENCE)	1.0	
Patient received HE phone calls ONLY	1.15 (0.67 to 1.99)	0.60
Patient kept a drinking diary or made a drinking agreement	0.75 (0.45 to 1.27)	0.29
Patient kept a drinking diary and made a drinking agreement	0.45 (0.25 to 0.82)	0.01

Results are derived from logistic regression analyses. Adjusted for gender, race, ethnicity, marital status, education, income, home ownership, whether or not alcohol counseling took place prior to Project SHARE, use of medications that increase alcohol risk, presence of symptoms that increase alcohol risk, behaviors that increase alcohol risk.

Results with *p*-value < 0.05 in bold font.

However, we did not assess outcomes beyond 12 months, and the literature indicates that extended interventions or repeated “booster” sessions may be required to maintain these effects and may in fact be cost-effective over the long term (Lash et al., 2011; McKay, 2005; Stout et al., 1999).

We found that having had a physician discussion about alcohol risk was associated with lower rates of at-risk drinking at 12 months but the association was not significant at the intermediate 6-month point. Many earlier intervention studies have focused on the 12-month end point, although the significant intervention effect is generally observed by 6 months (Curry et al., 2003; Fink et al., 2005; Fleming et al., 1997; Lin et al., 2010a,b; Maisto et al., 2001; Richmond et al., 1995). The specific explanation for the lack of a significant effect at 6 months in our study is not entirely clear. However, because physicians were asked to engage in alcohol risk discussions with intervention patients at each visit during the 12-month follow-up period, only 62% of the alcohol risk discussions occurred in the first 6 months. The other 38% occurred in the second 6 months, and it is possible that the specific patients who had their initial discussion (or a follow-up discussion building on a prior discussion)

between months 6 and 12 were more likely to change their behavior.

Studies of alcohol risk reduction have largely focused on describing the success of multicomponent interventions, with less attention paid to the specific components within the intervention that may be most important. A recent study asked 17 primary care physicians who frequently deliver alcohol reduction interventions for young adults to rate the most effective approaches they include (Grossberg et al., 2010). Of 24 possible intervention components, 5 were identified as the most useful, specifically summarizing the patient’s drinking level, making a drinking agreement, keeping a drinking diary, discussing drinking likes and dislikes, and discussing life goals. While the target population is different in terms of age, the results of our study support these clinical observations. The ultimate goal of this line of research was to support the translation of effective interventions into widespread use in nonresearch situations. However, translated interventions are often modified or abbreviated to fit the capacity of different “real-world” delivery settings (Carvalho et al., 2013; Napoles et al., 2013). Additional studies investigating which specific components of alcohol risk reduction interventions are critical to include in translation, and by whom the intervention can be delivered, will help to advance the field.

This study has several limitations, including the possibility that unmeasured differences in patient motivation or self-efficacy may lead to selection bias and different levels of intervention uptake. For example, patients who perceived their alcohol use as a more significant problem may have been more likely to engage with the HEs. However, in results not shown here, we examined the relationship between use of different intervention components and several baseline survey items that address patient self-efficacy, including confidence in being able to cut down on drinking if it was worsening one’s health and confidence in being able to ask physicians or other healthcare personnel for information about alcohol. The distribution of these survey items did not vary across patients who kept a drinking diary, made a drinking agreement, did both, or did neither, suggesting that unmeasured differences between patients were probably not the primary reason for the results of our main analyses. Also, due to limited statistical power, we were unable to examine interactions between patient participation in the physician intervention and each of the different aspects of the HE intervention. Other limitations include that fact that risk factors used in the CARET, such as drinking frequency and quantity, were self-reported. However, evidence from the literature suggests that patient self-reported alcohol consumption tends to be reliable and valid (Del Boca and Darkes, 2003). We also relied on patient self-report that they kept a drinking diary and did not require patients to show the diary to the HEs. Finally, this study included primarily white, well-educated patients in a relatively affluent area and may not be generalizable to other settings.

In summary, we found that use of several components of the Project SHARE intervention, including the physician-delivered as well as the HE-delivered approaches, was associated with lower rates of at-risk drinking at follow-up. This suggests that both components are important in achieving the desired effect, and future studies targeting at-risk drinking among older adults should consider incorporating both elements of the intervention.

APPENDIX. Comorbidity Alcohol Risk Evaluation Tool (CARET©UC Regents)

Item	Amount of drinking considered at-risk
<i>Alcohol use and behaviors in the last 12 months</i>	
Number of drinks and frequency of drinking	≥5/d at any frequency, 4/d at least 2 times/mon, 3/d at least 4 times/wk
Four or more drinks on 1 occasion (heavy episodic drinking)	At least 1 time/wk
Driving within 2 hours of drinking 3 or more drinks	Any frequency
Someone concerned about participant's alcohol use	Any amount
Someone concerned about participant's alcohol use more than 12 months ago	≥4/d at any frequency, 2 to 3/d at least 4 times/wk
<i>Alcohol use and medications taken at least 3 to 4 times per week currently</i>	
Medications that may cause bleeding, dizziness, sedation	≥4/d at any frequency, 2 to 3/d at least 4 times/wk
Medications used for gastroesophageal reflux, ulcer disease, depression	≥4/d at any frequency, 2 to 3/d at least 4 times/wk
Medications for hypertension	≥5/d at any frequency, 4/d at least 2 times/wk, 3/d at least 4 times/wk
<i>Alcohol use and comorbidities in the past 12 months</i>	
Liver disease, pancreatitis	Any amount
Gout, depression	≥4/d at any frequency, 3/d at least 2 times/wk, 2/d at least 4 times/wk
High blood pressure, diabetes	5/d at any frequency, 4/d at least 2 times/mon
Sometimes have problems with sleeping, falling	3/d at least 4 times/wk
Memory problems, heartburn, stomach pain, nausea, vomiting, or feel sad/blue	≥5/d at any frequency, 4/d at least 2 times/mon, 3/d at least 2 times/wk
Often have problems with sleeping, falling, memory, heartburn, stomach pain, nausea, vomiting, or feel sad/blue	≥4/d at any frequency, 2 to 3/d at least 2 times/wk

ACKNOWLEDGMENTS

This project was funded by National Institute on Alcohol Abuse and Alcoholism (NIAAA) grant 1R01AA013990 (Principal investigator: Susan L. Ettner). Alison A. Moore's time was additionally supported by NIAAA grants R01 AA15957, K24 AA15957, P30 AG028748 and P30 AG021684 (Principal investigator: Alison A. Moore). O. Kenrik Duru's time was supported by National Institute on Aging grants 5P30AG021684-12 and 5K08AG033630-05. The authors thank the following:

the leadership, research staff, and information technology staff of Sansum Clinic, in particular Mr. Paul Jaconette, Ms. Chris McNamara, Ms. Linda Chapman, and Mr. Tom Colbert; Ms. Deborah Marshall and the University of California at Los Angeles research staff; the Project SHARE (Senior Health and Alcohol Risk Education) health educators; and the members of our research advisory board. Finally, we are indebted to the Sansum Clinic patients and physicians who participated in Project SHARE, without whom the study could not have been conducted.

REFERENCES

- Aira M, Kauhanen J, Larivaara P, Rautio P (2003) Factors influencing inquiry about patients' alcohol consumption by primary health care physicians: qualitative semi-structured interview study. *Fam Pract* 20:270–275.
- Blazer DG, Wu LT (2011) The epidemiology of alcohol use disorders and subthreshold dependence in a middle-aged and elderly community sample. *Am J Geriatr Psychiatry* 19:685–694.
- Byles J, Young A, Furuya H, Parkinson L (2006) A drink to healthy aging: the association between older women's use of alcohol and their health-related quality of life. *J Am Geriatr Soc* 54:1341–1347.
- Carvalho ML, Honeycutt S, Escoffery C, Glanz K, Sabbs D, Kegler MC (2013) Balancing fidelity and adaptation: implementing evidence-based chronic disease prevention programs. *J Public Health Manag Pract* 4:348–356.
- Chan AM, von Muhlen D, Kritiz-Silverstein D, Barrett-Connor E (2009) Regular alcohol consumption is associated with increasing quality of life and mood in older men and women: The Rancho Bernardo Study. *Maturitas* 62:294–300.
- Cooper C, Bebbington P, Meltzer H, Jenkins R, Brugha T, Lindesay JE, Livingston G (2009) Alcohol in moderation, premorbid intelligence and cognition in older adults: results from the psychiatric morbidity survey. *J Neurol Neurosurg Psychiatry* 80:1236–1239.
- Curry SJ, Ludman EJ, Grothaus LC, Donovan D, Kim E (2003) A randomized trial of a brief primary-care-based intervention for reducing at-risk drinking practices. *Health Psychol* 22:156–165.
- Del Boca FK, Darkes J (2003) The validity of self-reports of alcohol consumption: state of the science and challenges for research. *Addiction* 98 (Suppl 2):1–12.
- Ettner SL, Xu H, Duru OK, Ang A, Tseng CH, Tallen L, Barnes A, Mirkin M, Ransohoff K, Moore AA (2014) The effect of an educational intervention on alcohol consumption, at-risk drinking, and health care utilization in older adults: The Project SHARE study. *J Stud Alcohol Drugs* 75:447–457.
- Fink A, Elliott MN, Tsai M, Beck JC (2005) An evaluation of an intervention to assist primary care physicians in screening and educating older patients who use alcohol. *J Am Geriatr Soc* 53:1937–1943.
- Fink A, Morton SC, Beck JC, Hays RD, Spritzer K, Oishi S, Moore AA (2002) The alcohol-related problems survey: identifying hazardous and harmful drinking in older primary care patients. *J Am Geriatr Soc* 50:1717–1722.
- Fleming MF, Barry KL, Manwell LB, Johnson K, London R (1997) Brief physician advice for problem alcohol drinkers. A randomized controlled trial in community-based primary care practices. *JAMA* 277:1039–1045.
- Friedmann PD, McCullough D, Chin MH, Saitz R (2000) Screening and intervention for alcohol problems. A national survey of primary care physicians and psychiatrists. *J Gen Intern Med* 15:84–91.
- Gilbertson R, Ceballos NA, Prather R, Nixon SJ (2009) Effects of acute alcohol consumption in older and younger adults: perceived impairment versus psychomotor performance. *J Stud Alcohol Drugs* 70:242–252.

- Grossberg P, Halperin A, Mackenzie S, Gisslow M, Brown D, Fleming M (2010) Inside the physician's black bag: critical ingredients of brief alcohol interventions. *Subst Abus* 31:240–250.
- Johansson K, Bendtsen P, Akerlind I (2005) Factors influencing GPs' decisions regarding screening for high alcohol consumption: a focus group study in Swedish primary care. *Public Health* 119:781–788.
- Jonas DE, Garbutt JC, Amick HR, Brown JM, Brownley KA, Council CL, Viera AJ, Wilkins TM, Schwartz CJ, Richmond EM, Yeatts J, Evans TS, Wood SD, Harris RP (2012) Behavioral counseling after screening for alcohol misuse in primary care: a systematic review and meta-analysis for the US Preventive Services Task Force. *Ann Intern Med* 157:645–654.
- Karlamangla AS, Sarkisian CA, Kado DM, Dedes H, Liao DH, Kim S, Reuben DB, Greendale GA, Moore AA (2009) Light to moderate alcohol consumption and disability: variable benefits by health status. *Am J Epidemiol* 169:96–104.
- Kirchner JE, Zubritsky C, Cody M, Coakley E, Chen H, Ware JH, Oslin DW, Sanchez HA, Durai UN, Miles KM, Llorente MD, Costantino G, Levkoff S (2007) Alcohol consumption among older adults in primary care. *J Gen Intern Med* 22:92–97.
- Lang I, Guralnik J, Wallace RB, Melzer D (2007a) What level of alcohol consumption is hazardous for older people? Functioning and mortality in US and English national cohorts. *J Am Geriatr Soc* 55:49–57.
- Lang I, Wallace RB, Huppert FA, Melzer D (2007b) Moderate alcohol consumption in older adults is associated with better cognition and well-being than abstinence. *Age Ageing* 36:256–261.
- Lash SJ, Timko C, Curran GM, McKay JR, Burden JL (2011) Implementation of evidence-based substance use disorder continuing care interventions. *Psychol Addict Behav* 25:238–251.
- Lin JC, Karno MP, Barry KL, Blow FC, Davis JW, Tang L, Moore AA (2010a) Determinants of early reductions in drinking in older at-risk drinkers participating in the intervention arm of a trial to reduce at-risk drinking in primary care. *J Am Geriatr Soc* 58:227–233.
- Lin JC, Karno MP, Tang L, Barry KL, Blow FC, Davis JW, Ramirez KD, Welgreen S, Hoffing M, Moore AA (2010b) Do health educator telephone calls reduce at-risk drinking among older adults in primary care? *J Gen Intern Med* 25:334–339.
- Maisto SA, Conigliaro J, McNeil M, Kraemer K, Conigliaro RL, Kelley ME (2001) Effects of two types of brief intervention and readiness to change on alcohol use in hazardous drinkers. *J Stud Alcohol* 62:605–614.
- McCormick KA, Cochran NE, Back AL, Merrill JO, Williams EC, Bradley KA (2006) How primary care providers talk to patients about alcohol: a qualitative study. *J Gen Intern Med* 21:966–972.
- McKay JR (2005) Is there a case for extended interventions for alcohol and drug use disorders? *Addiction* 100:1594–1610.
- Moore AA, Blow FC, Hoffing M, Welgreen S, Davis JW, Lin JC, Ramirez KD, Liao DH, Tang L, Gould R, Gill M, Chen O, Barry KL (2011) Primary care-based intervention to reduce at-risk drinking in older adults: a randomized controlled trial. *Addiction* 106:111–120.
- Moore AA, Whiteman EJ, Ward KT (2007) Risks of combined alcohol/medication use in older adults. *Am J Geriatr Pharmacother* 5:64–74.
- Napoles AM, Santoyo-Olsson J, Stewart AL (2013) Methods for translating evidence-based behavioral interventions into health disparity communities. *Prev Chronic Dis* 10:E193.
- Reid MC, Boutros NN, O'Connor PG, Cadariu A, Concato J (2002) The health-related effects of alcohol use in older persons: a systematic review. *Subst Abus* 23:149–164.
- Reid MC, Van Ness PH, Hawkins KA, Towle V, Concato J, Guo Z (2006) Light to moderate alcohol consumption is associated with better cognitive function among older male veterans receiving primary care. *J Geriatr Psychiatry Neurol* 19:98–105.
- Richmond R, Heather N, Wodak A, Kehoe L, Webster I (1995) Controlled evaluation of a general practice-based brief intervention for excessive drinking. *Addiction* 90:119–132.
- Saitz R, Horton NJ, Sullivan LM, Moskowitz MA, Samet JH (2003) Addressing alcohol problems in primary care: a cluster randomized, controlled trial of a systems intervention. The screening and intervention in primary care (sip) study. *Ann Intern Med* 138:372–382.
- Stout RL, Rubin A, Zwick W, Zywiak W, Bellino L (1999) Optimizing the cost-effectiveness of alcohol treatment: a rationale for extended case monitoring. *Addict Behav* 24:17–35.
- Townsend MK, Devore E, Kang JH, Grodstein F (2009) The relation between moderate alcohol consumption and cognitive function in older women with type 2 diabetes. *Diabetes Res Clin Pract* 85:322–327.
- U.S. Department of Health & Human Services National Institute of Health National Institute on Alcohol Abuse and Alcoholism (2007). Helping patients with alcohol problems: a health practitioner's guide. Available at: <http://pubs.niaaa.nih.gov/publications/Practitioner/CliniciansGuide2005/guide.pdf>. Accessed January 12, 2015.
- Vestal RE, McGuire EA, Tobin JD (1977) Aging and ethanol metabolism in man. *Clin Pharmacol Ther* 21:343–345.
- Whitlock EP, Polen MR, Green CA, Orleans T, Klein J (2004) Behavioral counseling interventions in primary care to reduce risky/harmful alcohol use by adults: a summary of the evidence for the U.S. Preventive Services Task Force. *Ann Intern Med* 140:557–568.