

UC San Diego

UC San Diego Previously Published Works

Title

Cathelicidin Host Defence Peptide Augments Clearance of Pulmonary Pseudomonas aeruginosa Infection by Its Influence on Neutrophil Function In Vivo

Permalink

<https://escholarship.org/uc/item/8dn455w8>

Journal

PLOS ONE, 9(6)

ISSN

1932-6203

Authors

Beaumont, Paula E
McHugh, Brian
Findlay, Emily Gwyer
et al.

Publication Date

2014

DOI

10.1371/journal.pone.0099029

Peer reviewed



Cathelicidin Host Defence Peptide Augments Clearance of Pulmonary *Pseudomonas aeruginosa* Infection by Its Influence on Neutrophil Function *In Vivo*

Paula E. Beaumont¹, Brian McHugh¹, Emily Gwyer Findlay¹, Annie Mackellar¹, Karen J. Mackenzie¹, Richard L. Gallo², John R. W. Govan³, A. John Simpson^{1,4}, Donald J. Davidson^{1*}

1 MRC Centre for Inflammation Research, Queen's Medical Research Institute, The University of Edinburgh, Edinburgh, United Kingdom, **2** Division of Dermatology, Department of Medicine, University of California at San Diego and VA San Diego Health Care System, San Diego, California, United States of America, **3** Edinburgh Infectious Diseases, The Chancellor's Building, New Royal Infirmary, University of Edinburgh, Edinburgh, Scotland, United Kingdom, **4** Institute of Cellular Medicine, Medical School, Newcastle University, Newcastle upon Tyne, United Kingdom

Abstract

Cathelicidins are multifunctional cationic host-defence peptides (CHDP; also known as antimicrobial peptides) and an important component of innate host defence against infection. In addition to microbicidal potential, these peptides have properties with the capacity to modulate inflammation and immunity. However, the extent to which such properties play a significant role during infection *in vivo* has remained unclear. A murine model of acute *P. aeruginosa* lung infection was utilised, demonstrating cathelicidin-mediated enhancement of bacterial clearance *in vivo*. The delivery of exogenous synthetic human cathelicidin LL-37 was found to enhance a protective pro-inflammatory response to infection, effectively promoting bacterial clearance from the lung in the absence of direct microbicidal activity, with an enhanced early neutrophil response that required both infection and peptide exposure and was independent of native cathelicidin production. Furthermore, although cathelicidin-deficient mice had an intact early cellular inflammatory response, later phase neutrophil response to infection was absent in these animals, with significantly impaired clearance of *P. aeruginosa*. These findings demonstrate the importance of the modulatory properties of cathelicidins in pulmonary infection *in vivo* and highlight a key role for cathelicidins in the induction of protective pulmonary neutrophil responses, specific to the infectious milieu. In addition to their physiological roles, CHDP has been proposed as future antimicrobial therapeutics. Elucidating and utilising the modulatory properties of cathelicidins has the potential to inform the development of synthetic peptide analogues and novel therapeutic approaches based on enhancing innate host defence against infection with or without direct microbicidal targeting of pathogens.

Citation: Beaumont PE, McHugh B, Gwyer Findlay E, Mackellar A, Mackenzie KJ, et al. (2014) Cathelicidin Host Defence Peptide Augments Clearance of Pulmonary *Pseudomonas aeruginosa* Infection by Its Influence on Neutrophil Function *In Vivo*. PLoS ONE 9(6): e99029. doi:10.1371/journal.pone.0099029

Editor: Samithamby Jayaseelan, Louisiana State University, United States of America

Received: February 5, 2014; **Accepted:** May 9, 2014; **Published:** June 2, 2014

Copyright: © 2014 Beaumont et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: PEB was supported by an Asthma UK/MRC Joint PhD Studentship (S08/001). KJM was supported by a MRC Clinical Research Training Fellowship (G0701350). RLG was supported by NIH grants R01-AR052728, R01-AI052453 and R01 AI0833358. DJD was supported by a MRC Senior Non-clinical Fellowship (G1002046). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: donald.davidson@ed.ac.uk

Introduction

Cationic host-defence peptides (CHDP; also known as antimicrobial peptides or AMPs) are important components of early innate host defences. In addition to their physiological roles, these peptides and their derivatives have been proposed as future antimicrobial therapeutics, relatively unaffected by the development of sustained microbial resistance [1]. Although initially characterised as directly microbicidal agents, it is now clear that many CHDP also have multiple functions as modulators of inflammation and immunity [2,3,4], with emerging roles in diseases affecting multiple organs including the lung, skin and gastrointestinal tract. Human clinical trials using analogues of CHDP modified to maximise direct microbicidal function have achieved only moderate efficacy [5], perhaps due to failure to recognise the importance of the immunomodulatory functions of the native peptides. Interestingly, studies using non-microbicidal analogues of naturally-occurring CHDP that retained other

bioactive functions, have demonstrated effective host defence augmentation in mice [6,7]. These studies raise questions about the relative roles of microbicidal and immunomodulatory properties of naturally-occurring CHDP in infections.

Cathelicidins are multipotent immunomodulatory CHDP [8]. The sole human cathelicidin Human Cationic Antimicrobial Peptide of 18 kDa (hCAP-18; encoded by *CAMP*) is expressed by multiple cell types including neutrophils, where it is stored in specific granules and proteolytically cleaved following release, to produce a 37 amino acid mature peptide fragment named LL-37 [9,10]. hCAP-18/LL-37 is upregulated in pulmonary infections [11] and, in children with RSV bronchiolitis, low serum cathelicidin is correlated with more severe disease [12]. Mice, like humans, express only a single cathelicidin gene; *Camp* (encoding the mCRAMP peptide) [13], with similar patterns of expression, which is cleaved to produce an active 34 amino acid peptide [14]. Mice deficient in mCRAMP (*Camp*^{-/-}) have increased suscep-

tibility to bacterial infections of the skin [15], intestinal tract [16], cornea [17] and urinary tract [18]. These *Camp*^{-/-} mice also have impaired host defence against lung infection [19,20], while therapeutic use of LL-37 and/or mCRAMP in wild type mice is protective in models of pulmonary infection with *P. aeruginosa* [21] or influenza virus [22]. These studies demonstrate a critical, non-redundant role for endogenous cathelicidin in host defence against lung infection and the therapeutic potential of the unmodified peptides, but the mechanisms by which pulmonary host defence is enhanced *in vivo* remains unclear. Although generally presented as being primarily a consequence of direct microbicidal activity, this is not fully consistent with *in vivo* concentrations and microbicidal properties in a physiological environment. However, the extent to which any of the plethora of immunomodulatory properties ascribed to cathelicidins play a significant role during infection has never been demonstrated *in vivo*. Understanding the critical modulatory roles of native CHDP and how these contribute to innate host defence against infection, may prove to be vital in development of specific pathogen-targeted analogues of these peptides for therapeutic use.

Respiratory diseases are among the most common causes of morbidity and account for 1 in 5 deaths in the UK [23]. A third of mortalities are due to acute respiratory infections, influenza or pneumonia and pathogens resistant to conventional therapeutics represent an increasing clinical challenge. *Pseudomonas aeruginosa* is the primary cause of nosocomial pulmonary infections and pulmonary colonisation with this pathogen is considered to be responsible for the fatal deterioration of lung function in patients with cystic fibrosis (CF) (reviewed in [24]). This opportunistic pathogen is difficult to treat because of its widespread resistance to multiple antibiotics [25], with the limited number of effective antimicrobial treatments reduced further by the emergence of carbapenem- and polymyxin-B resistant isolates [26]. A greater understanding of the natural host defence mechanisms involved in pulmonary defence against this organism is required in order to develop novel therapeutic approaches. Cathelicidins can alter susceptibility to pulmonary infection with *P. aeruginosa* in murine models [19,20,21], despite this pathogen being resistant to the directly microbicidal effects of these peptides in the presence of physiologically relevant levels of cations *in vitro* [27,28,29]. Thus, the *in vivo* roles of endogenous cathelicidin in host defence against *P. aeruginosa*, the relative effects of microbicidal and modulatory properties, and the consequences of therapeutic targeting of cathelicidin expression or exogenous delivery of peptide remain unknown.

We demonstrate that therapeutically administered synthetic LL-37 peptide can enhance the clearance of *P. aeruginosa* from the murine lung, in the absence of demonstrable direct microbicidal effects, and can induce an upregulation of the early neutrophil response to pathogen in the lungs that is dependent both upon the presence of the peptide and the pathogen. We show that despite a normal early neutrophil response, second phase pulmonary neutrophil influx was deficient in *Camp*^{-/-} mice, with impaired clearance of pulmonary *P. aeruginosa*. Delivery of LL-37 to these cathelicidin-deficient mice enhanced the neutrophil response and restore bacterial clearance, demonstrating proof of principle for therapeutic use of LL-37 in cathelicidin deficiency. These studies indicate that the protective effects of cathelicidins in *P. aeruginosa* infection *in vivo* can result from modulatory effects in innate immune responses, synergising with infectious stimuli to enhance a protective neutrophil response.

Materials and Methods

Peptide

LL-37 (LLGDFFRKSKEKIGKEFKRIVQRIKDFLRNLPV-RTES; MW 4493.33) was either synthesised by N-(9-fluorenyl) methoxycarbonyl chemistry at the Nucleic Acid/Protein Service unit at the University of British Columbia (UBC; Vancouver, Canada), or custom synthesised by Almac (East Lothian, Scotland) using Fmoc solid phase synthesis and reversed phase HPLC purification. Peptide identity was confirmed by electrospray mass spectrometry, purity (>95% area) by RP-HPLC and net peptide content determined by amino acid analysis. Lyophilised peptides were reconstituted in endotoxin free water at 5 mg/ml stock concentration and determined to be endotoxin-free using a Limulus Amebocyte Lysate Chromogenic Endotoxin Quantitation Kit (Thermo Scientific, UK). Peptide functionality was confirmed by assessing anti-endotoxic activity [30].

Bacteria

Pseudomonas aeruginosa strain PAO1 was grown in Luria Bertani (LB) broth at 37°C in an orbital shaker (250 rpm) overnight to achieve a stationary-phase suspension. Bacterial cultures were then diluted 1:10 in fresh LB broth and incubated at 37°C for 90 min to reach logarithmic phase. Standardisation was performed by dilution with LB broth to an optical density of 0.1 at 595 nm using spectrophotometry (WPA UV 1101, Biotech Photometer), then bacteria were centrifuged at 1,500 ×g for 15 min and resuspended in PBS for use.

Murine infection model

Mouse experiments were performed in accordance with Home Office UK project licence 60/4216, under the Animal (Scientific Procedures) Act 1986. Wild type control mice were all C57Bl/6 strain, supplied by Charles River Laboratories, UK, and housed at the University of Edinburgh for at least two weeks before use, or were animals bred from those mice. *Camp*^{-/-} mice [15] were bred to congenicity on a C57Bl/6 strain background and were the offspring of homozygous mutant matings raised in the same facility at the University of Edinburgh. Both male and female mice were used, between 8–12 weeks old, housed in individually ventilated cages and randomly assigned to treatment groups (no significant difference were found in end points between male and female mice). Mice were weighed, given a general anaesthetic (isoflurane) in a category 2 biosafety hood, then held vertically by scruffing over the front of the thorax and inoculated by an intranasal delivery up to a total of 50 µl volume. Mice were inoculated with 3 × 10⁷ colony forming units (cfu) of PAO1 or the same volume of PBS, and 10 µg LL-37 peptide in PBS or PBS only control. PBS alone (carrier for both bacteria and peptide) was used as a control (rather than scrambled peptide, which previous pulmonary infection studies indicated had no effects ([22] and unpublished data), in order that the wild type control infected animals were appropriate controls both for the LL-37-treated infected wild types and for the infected *Camp*^{-/-} animals (in which no peptide was delivered). Mice were then returned to cages, placed on a heat mat to maintain body temperature, and monitored and scored for signs of infection every 2 hours, with peak of illness occurring at 6–8 hours post infection, followed by recovery with diminishing severity score. Mice were re-weighed and culled at selected timepoints (0, 6 or 24 hours), culled by pentobarbital injection and lungs and trachea exposed by dissection. Lungs were lavaged in 1 ml sterile PBS via intramedic polyethylene tubing (Sigma Aldrich, UK) inserted into the trachea, and bronchoalveolar lavage fluid (BALF) stored on ice. Following

lavage, lungs were either homogenised in 2 ml sterile PBS for cfu counts or were perfused by PBS injection into the heart, then removed and frozen for RNA preparation.

CFU counts

BALF or homogenised lungs were serially diluted in PBS, plated on *Pseudomonas* Isolation agar (Becton Dickinson Difco, Oxford, UK), incubated overnight at 37°C and bacterial colonies counted using a Stuart SC6 colony counter. Total colonies on the lowest dilution plate countable were multiplied by the appropriate dilution factors to determine the total CFU count of the lung tissue or BALF sample.

Cytospins and counts

BALF was centrifuged at 200×g for 5 minutes, and supernatant was removed for cytokine measurement. Pelleted cells were resuspended and counted by NucleoCounter YC-100 (Chemo-Metec, Allerød, Denmark) automated cell number counting. 100 µl of cell suspension was then loaded onto a glass slide using a disposable sample funnel and cytocentrifuged at 10×g for 3 minutes in a Shandon Cytospin 2 centrifuge. Slides were air dried for 20 minutes, fixed in methanol for 20 minutes, stained with Diff Quik (Fisher Scientific, Loughborough, UK), and mounted in DPX Mountant (Fluka BioChemika/Sigma Aldrich, UK). Differential counts for neutrophils and monocytes were then performed by light microscopy at 20× magnification using an EVOS FL microscope (Peqlab, Sarisbury Green, UK).

ELISAs

BALF was used to measure cytokine levels by ELISA according to manufacturer's instructions, for KC, MIP-2 alpha (R&D Systems, UK) or using a cytometric bead assay mouse inflammation kit (BD Biosciences, UK) for TNF, IL-6, MCP-1, IL-10, IFN-gamma, IL-12p70.

qRT-PCR

Mouse lung tissue was homogenized in Qiagen RLT buffer (Qiagen, Manchester, UK) using a Precellys 24 homogeniser with Precellys-Keramik-kit ceramic beads (PeqLab). RNA was then prepared from homogenised mouse lung tissue using RNeasy mini kits (Qiagen), according to the manufacturer's instructions. After DNase treatment with RQ1 DNase (Promega, Southampton, UK), cDNA was prepared from RNA using TaqMan reverse transcriptase reagents and random hexamer primers (Life Technologies Ltd, Paisley, UK), according to the manufacturer's instructions. Quantitative Real Time PCR was performed on a StepOne Real Time PCR machine (Life Technologies), using Gene Expression Mastermix and TaqMan gene expression assays for *Camp* (assay I.D. Mm00438285_m1) and 18S (assay I.D. Mm03928990_g1). Relative quantitation of *Camp* was calculated using the $\Delta\Delta C_T$ method.

Analysis of mCRAMP protein expression

Harvested lung tissue was placed in 600 µl M-PER Mammalian Protein Extraction Reagent with Complete Protease Inhibitor Cocktail (Roche Applied Science, Burgess Hill, UK) added, and homogenised using a Precellys 24 homogeniser with Precellys-Keramik-kit ceramic beads (PeqLab). Homogenised tissue was shaken on an IKA-Vibramax-VXR (Sigma Aldrich, UK) for 20 minutes at 4°C and lysates were subsequently centrifuged at 15,000×g for 10 minutes at 4°C to pellet insoluble material. Protein concentration in lysates was measured by Pierce BCA assay (Thermo Scientific), according to manufacturer's instruc-

tions. Lysate concentrations were equalised with lysis buffer, and subsequently boiled at 96°C for 5 minutes in the presence of loading buffer and reducing agent (Life Technologies), then run on Novex NuPAGE 4–12% Bis-Tris pre-cast gels, in MOPS buffer (Life Technologies), and subsequently transferred to Novex 0.2 µm pore Nitrocellulose Membrane. mCRAMP was detected with rabbit anti-mouse mCRAMP antibody (R-170, Santa Cruz Biotechnology, Heidelberg, Germany), followed by staining with IRDye 800CW anti-rabbit secondary antibody, with subsequent detection using a LI-COR Odyssey Infrared Imaging System. Rabbit pan-actin antibody (Cell Signalling Technology, Danvers, MA, USA) staining was used as a loading control, detected with anti-rabbit secondary antibody and infrared imaging as above. Mouse CRAMP staining was then quantitated using LI-COR Odyssey software.

Statistics

Statistical analyses were performed using Graphpad Prism version 5.04 for Windows. Bacterial counts were normalised by logarithmic transformation before analysis by 2 way ANOVA with Bonferroni's post tests where appropriate. Cell counts and cytokine concentrations were analysed by Mann Whitney test. Differences were considered statistically significant at $P < 0.05$.

Results

LL-37-mediated microbicidal activity against *P. aeruginosa* PAO1 is not observed *in vivo*

LL-37 and other CHDP were initially described as having rapid direct microbicidal properties based on *in vitro* studies conducted under favourable ionic environments (reviewed in [27]). However, the activity of LL-37 can be inhibited by physiological levels of divalent cations [27,28], serum apolipoprotein, f-actin and DNA [31,32,33]. Thus, the relative contributions of microbicidal versus modulatory properties in conferring the protective effects of this peptide *in vivo* remain uncertain. We and others have shown that LL-37 has negligible microbicidal activity against *P. aeruginosa in vitro* [27,28,29]. In order to evaluate the contribution of any early direct microbicidal properties of exogenously delivered LL-37 peptide in a murine pulmonary *P. aeruginosa* infection model, mice were culled immediately after the intranasal delivery of bacteria with peptide or carrier-only control to the cohort. Lungs were removed, homogenised and plated to evaluate the number of viable bacteria in the lungs. These samples are referred to as $t = 0$, however homogenisation did not occur until 60 minutes after inoculation, during which time interaction between peptide and bacteria could occur. No significant difference was observed between infected mice receiving LL-37 and control infected animals (Fig 1a) demonstrating that LL-37, under these conditions, had no discernable early microbicidal affects.

Therapeutic delivery of LL-37 is protective against *P. aeruginosa* infection *in vivo*

In order to evaluate the protective antimicrobial properties of LL-37 against *P. aeruginosa* PAO1 *in vivo*, mice were infected with or without concomitant delivery of LL-37. All mice lost ~4% body weight in the first 6 hours post procedure (data not shown), with infected animals continuing to lose weight over 24 hours, but no significant effect of LL-37 treatment was observed (Fig 1b). Mice were culled 6 and 24 hours post-infection and the total number of viable bacteria in the BALF (bronchoalveolar lavage fluid) and lung homogenate was assessed (Fig 1 c–f). At 6 hours post-infection, treatment with LL-37 showed no statistically significant effect on the total number of bacteria in the lung homogenates, but




Figure 1. Exogenous LL-37 enhances pulmonary clearance of *P. aeruginosa*. Wild type C57Bl/6 mice were weighed, then inoculated with 3×10^7 cfu of *P. aeruginosa* PAO1 or PBS and 10 μ g LL-37 peptide or PBS by intranasal delivery. a) Immediately after inoculation of all mice, a subset (called 0 hr; $n=3$ per group) were culled and their lungs homogenised (60 minutes after initial inoculation), or b–f) 6 or 24 hours after inoculation mice were re-weighed and culled, and their lungs were lavaged before homogenisation. BALF and lung homogenates were serially diluted, plated and incubated overnight at 37°C before bacterial colonies were counted and corrected for volume. Mean PAO1 cfu \pm SEM in the lung homogenate (a, c & e) or BALF (d & f) for infected animals ($n \geq 9$ per condition) are displayed. No bacteria were detected in samples from uninfected mice. b) Data show mean percentage weight loss \pm SEM. For statistical analyses bacterial counts were normalised by logarithmic transformation. Analyses were conducted using 2 way ANOVA with Bonferroni's post tests; * $p < 0.05$, ** $p < 0.01$. doi:10.1371/journal.pone.0099029.g001

did result in significantly lower levels of bacteria in the BALF (Fig 1c/d). By 24 hours post-infection, LL-37 treatment had significantly enhanced pathogen clearance from the lungs, compared to controls, leaving only a residual infection,

inaccessible to BAL (Fig 1e/f). These data demonstrate the capacity of LL-37 to enhance pulmonary bacterial clearance in the absence of early microbicidal properties.

Therapeutic delivery of LL-37 enhances neutrophil responses in infected animals

Cathelicidins have been proposed to have multiple inflammatory properties that could modulate the clearance of infection *in vivo* [2], including direct chemotactic activity of LL-37 for neutrophils and monocytes [34,35,36,37,38]. Differential cytospin cell counts were performed on the BALF from LL-37-treated and control-treated infected and uninfected mice at 6 and 24 hours after infection. LL-37 treatment resulted in a significantly upregulated neutrophil response to infection (over 2 fold increase in median cell number) compared to control infected animals at 6 hours post-infection (Fig 2a). No neutrophils were detected in the BALF from uninfected mice, regardless of LL-37 treatment, at this timepoint (Fig 2b). Whereas a larger second phase neutrophil response was observed by 24 hours in control-treated infected mice, the mean neutrophil number in the LL-37-treated infected animals was actually significantly lower than the controls at this timepoint, in line with the enhanced earlier clearance of the infection (Fig 2c). A degree of neutrophil influx was observed in response to LL-37 alone at 24 hours in uninfected mice (Fig 2d). In contrast, no significant LL-37-mediated effects on pulmonary monocyte numbers were observed at 6 or 24 hours in infected (Fig 2 e/f) or uninfected (data not shown) mice.

Therapeutic delivery of LL-37 does not affect pulmonary cytokine responses in infected animals

In order to determine whether LL-37-mediated enhanced neutrophil responses were secondary to modulation of pulmonary cytokine and chemokine responses, CBA (Cytometric Bead Array) and ELISA assays were performed on the BALF from LL-37-treated and control-treated infected mice at 6 and 24 hours after infection, to determine the concentrations of TNF, IL-6, MIP-2, KC, MCP-1, IL-10, IFN γ and IL-12. Although TNF, IL-6, MIP-2, KC and MCP-1 were all highly expressed in response to infection at 6 hours (Fig 3a–e) compared to baseline levels in uninfected mice (data not shown), and resolving by 24 hours (Fig 3f–j), treatment with LL-37 had no significant effect on any of the cytokines measured. In contrast, IL-10, IFN γ and IL-12 were not detected in significant quantities.

Pulmonary infection with *P. aeruginosa* induces *Camp* expression in the murine lung

In this model, therapeutic administration of LL-37 was in addition to any effects of endogenous murine cathelicidin mCRAMP produced in the murine lungs in response to infection. mCRAMP has also been shown to have neutrophil chemotactic properties *in vivo*, in an air pouch model [36]. Thus, in order to establish the temporal expression pattern of *Camp* in *P. aeruginosa* infected mice, qRT-PCR and western immunoblot analyses were performed on lung homogenates at 0, 2, 6 and 24 hours post-infection. *Camp* transcription was not detected at 0 hour, but was detectable at very low levels by 2 hours after infection. Transcription was dramatically increased at 6 and 24 hours after infection (upregulated 1886-fold, +/-137, and 1124-fold, +/-66 respectively, relative to the 2 hour timepoint), with mCRAMP protein clearly detectable at these timepoints (data not shown) in keeping with previously published data [20]. Thus, the inflammatory responses were potentially modified by cathelicidin from around 6 hours post-infection in all mice, but additionally modified by cathelicidin within the first few hours in mice receiving an intranasal bolus of LL-37.

Endogenous mCRAMP is protective against *P. aeruginosa* infection *in vivo*

In order to evaluate the protective antimicrobial properties of induced endogenous mCRAMP against *P. aeruginosa* PAO1 in this model, *Camp* $-/-$ mice were infected and compared to wild type control animals. The profile of weight loss in infected *Camp* $-/-$ mice was not significantly different from the wild type control (data not shown). Mice were culled 6 and 24 hours post-infection and the total number of viable bacteria in the BALF and lung homogenate was assessed (Fig 4 a–d). At 6 hours post-infection *Camp* $-/-$ mice showed no statistically significant difference in the total number of bacteria in the lung homogenates or BALF when compared to infected wild type controls (Fig 4 a/b). However, by 24 hours post-infection, wild type mice had more effectively cleared the bacteria, with significantly higher pathogen loads found in both the lung homogenates and BALF from the *Camp* $-/-$ mice (Fig 4c/d). These data demonstrate the capacity of endogenous mCRAMP to enhance pulmonary bacterial clearance, occurring after inducible *Camp* expression is detectable in the lungs of wild-type mice.

Endogenous mCRAMP enhances neutrophil responses in infected animals

The extent to which endogenous mCRAMP might play a role in the pulmonary neutrophil response to *P. aeruginosa* infection was examined by comparing BALF cytospin differential cell counts from infected *Camp* $-/-$ mice and wild type mice at 6 and 24 hours after infection. No significant differences were observed in neutrophil or monocyte counts at 6 hours (Fig 5a/b), with an initial neutrophil influx occurring similarly in both genotypes (Fig 5a). However, *Camp* $-/-$ mice failed to further upregulate this response, demonstrating a significantly less elevated neutrophil count than wild type controls at 24 hours (Fig 5c). A trend towards fewer monocytes was also observed, but did not reach significance (Fig 5d). These data indicate that endogenous mCRAMP is not involved in the first phase of neutrophil influx, but is required, following induction, for the second phase neutrophil response to *P. aeruginosa* infection. TNF, IL-6, MIP-2, KC and MCP-1 were all highly expressed in response to infection in *Camp* $-/-$ mice at 6 hours (Fig 6a–e), and resolving by 24 hours (Fig 6f–j), but were not significantly different from the responses quantified in infected wild type mice.

Therapeutic delivery of LL-37 can restore protection against *P. aeruginosa* infection in cathelicidin deficient mice

Cathelicidin insufficiency has been associated with increased susceptibility to infection in humans [12,39,40]. Thus, in order to determine whether therapeutic use of synthetic human LL-37 could restore cathelicidin-mediated protective antimicrobial function in cathelicidin deficiency, *Camp* $-/-$ mice were infected with *P. aeruginosa* PAO1 *in vivo*, with or without concomitant delivery of LL-37. The profile of weight loss in infected *Camp* $-/-$ mice was not significantly altered by delivery of LL-37 (data not shown). Mice were culled 6 and 24 hours post-infection and the total number of viable bacteria in the BALF (bronchoalveolar lavage fluid) and lung homogenate was assessed (Fig 7 a–d). At 6 hours post-infection, treatment with LL-37 showed no significant effect on the total number of bacteria in the lung homogenates (despite a trend towards enhanced clearance), but led to significantly lower levels of bacteria in the BALF (Fig 7a/b). By 24 hours post-infection, LL-37 treatment had very significantly enhanced pathogen clearance from the lungs, compared to control-treated




Figure 2. Exogenous LL-37 promotes an early neutrophil response to *P. aeruginosa*. Wild type C57Bl/6 mice were inoculated with 3×10^7 cfu of *P. aeruginosa* PAO1 or PBS and 10 μ g LL-37 peptide or PBS by intranasal delivery. At 6 hours (a, b & e) or 24 hours (c, d & f) after inoculation mice were culled and their lungs were lavaged. BALF was cytocentrifuged and differential counts were conducted for neutrophils (a–d) and monocytes (e & f). Data show Tukey box and whiskers plots for infected (a, c, e & f) ($n \geq 9$ per condition) and uninfected (b & d) animals ($n \geq 5$ per condition). Analyses were conducted using the Mann Whitney test; * $p < 0.05$. ND denotes “not detected”. doi:10.1371/journal.pone.0099029.g002

Camp^{−/−} (Fig 7c/d). These data demonstrate that delivery of exogenous synthetic LL-37 can enhance host defence against infection by mechanisms that do not require endogenous host cathelicidin production, and show cross-species functionality of these peptides.

Therapeutic delivery of LL-37 promotes an early neutrophil response to *P. aeruginosa* infection, associated with enhanced clearance

Cathelicidin-mediated enhancement of bacterial clearance was associated with upregulated neutrophil influx in LL-37-treated infected wild type mice (compared to untreated infected controls). However, in addition, endogenous cathelicidin clearly also had a critical role in the induction of a maximal neutrophil responses to

infection. Thus, in order to determine whether LL-37-mediated enhanced neutrophil responses were independent of endogenous cathelicidin production, BALF cytopspin differential cell counts were also evaluated from LL-37-treated and control infected *Camp*^{−/−} mice at 6 and 24 hours after infection. As also observed in LL-37-treated wild type mice (Fig 2a), early, infection-induced neutrophil influx (at 6 hours) was significantly greater in LL-37-treated mice (Fig 8a), but this early effect of the therapeutic bolus was lost by 24 hours (Fig 8b). However, whereas wild type mice showed a robust later neutrophil response to infection regardless of peptide treatment (Fig 2c), this second phase neutrophil response failed to occur in infected *Camp*^{−/−} mice, irrespective of peptide treatment (Fig 8b), demonstrating the dependence of this later response upon pathogen-induced *Camp* expression. No significant




Figure 3. *P. aeruginosa*, but not exogenous LL-37, induces pulmonary cytokine responses. Wild type C57Bl/6 mice were inoculated with 3×10^7 cfu of *P. aeruginosa* PAO1 and 10 μ g LL-37 peptide or PBS by intranasal delivery. At 6 hours (a–e) or 24 hours (f–j) after inoculation, mice were culled and their lungs were lavaged. BALF was centrifuged to remove cells and levels of TNF (a, f), IL-6 (b, g), MIP-2 (c, h), KC (d, i) and MCP-1 (e, j) were determined. Data show Tukey box and whiskers plots for $n \geq 9$ animals per condition. Analyses were conducted using the Mann Whitney test. doi:10.1371/journal.pone.0099029.g003




Figure 4. Cathelicidin-deficient mice display impaired pulmonary clearance of *P. aeruginosa*. *Camp*^{-/-} mice and wild type controls were inoculated with 3×10^7 cfu of *P. aeruginosa* PAO1 or PBS by intranasal delivery. At 6 or 24 hours after inoculation mice were culled and their lungs were lavaged before homogenisation. BALF and lung homogenates were serially diluted, plated and incubated overnight at 37°C before bacterial colonies were counted and corrected for volume. Mean PAO1 cfu \pm SEM in the lung homogenate (a & c) or BALF (b & d) for infected animals ($n \geq 10$ per condition) are displayed. No bacteria were detected in samples from uninfected mice. For statistical analyses bacterial counts were normalised by logarithmic transformation. Analyses were conducted using 2 way ANOVA with Bonferroni's post tests; * $p < 0.05$. doi:10.1371/journal.pone.0099029.g004




Figure 5. Cathelicidin-deficient mice display impaired late neutrophil responses to *P. aeruginosa*. Camp^{-/-} mice and wild type controls were inoculated with 3 × 10⁷ cfu of *P. aeruginosa* PAO1 by intranasal delivery. At 6 hours (a & b) or 24 hours (c & d) after inoculation mice were culled and their lungs were lavaged. BALF was cytocentrifuged and differential counts were conducted for neutrophils (a & c) and monocytes (b & d). Data show Tukey box and whiskers plots for n ≥ 8 animals per condition. Analyses were conducted using the Mann Whitney test; * p < 0.05. doi:10.1371/journal.pone.0099029.g005

effects upon monocyte counts were observed (data not shown). These data demonstrate that the early infection-mediated neutrophil response, enhanced by the bolus of LL-37, was independent of

endogenous cathelicidin expression and associated with enhanced clearance of pulmonary *P. aeruginosa*.




Figure 6. *P. aeruginosa*, but not cathelicidin sufficiency, induces pulmonary cytokine responses. Camp^{-/-} mice and wild type controls were inoculated with 3 × 10⁷ cfu of *P. aeruginosa* PAO1 by intranasal delivery. At 6 hours (a–e) or 24 hours (f–j) after inoculation, mice were culled and their lungs were lavaged. BALF was centrifuged to remove cells and levels of TNF (a, f), IL-6 (b, g), MIP-2 (c, h), KC (d, i) and MCP-1 (e, j) were determined. Data show Tukey box and whiskers plots for n ≥ 8 animals per condition. Analyses were conducted using the Mann Whitney test. doi:10.1371/journal.pone.0099029.g006




Figure 7. Exogenous LL-37 enhances pulmonary clearance of *P. aeruginosa* in cathelicidin-deficient mice. *Camp*^{-/-} mice were inoculated with 3×10^7 cfu of *P. aeruginosa* PAO1 or PBS and 10 μ g LL-37 peptide or PBS by intranasal delivery. At 6 or 24 hours after inoculation mice were re-weighed and culled, and their lungs were lavaged before homogenisation. BALF and lung homogenates were serially diluted, plated and incubated overnight at 37°C before bacterial colonies were counted and corrected for volume. Mean PAO1 cfu \pm SEM in the lung homogenate (a & c) or BALF (b & d) for infected animals ($n \geq 6$ per condition at 6 hours and $n \geq 10$ per condition at 24 hours) are displayed. No bacteria were detected in samples from uninfected mice. For statistical analyses bacterial counts were normalised by logarithmic transformation. Analyses were conducted using 2 way ANOVA with Bonferroni's post tests; ** $p < 0.01$, *** $p < 0.001$. doi:10.1371/journal.pone.0099029.g007

Discussion

Cathelicidins are recognised as key multifunctional modulators of innate immunity and host defence against infection, and offer possible novel therapeutic templates. In addition to directly microbicidal potential, these peptides have been described as having a broad range of inflammomodulatory and immunomod-

ulatory properties [2]. However, no clear evidence exists for these functions being involved in cathelicidin-mediated enhanced host defence against pulmonary infection *in vivo*, with the relative significance of microbicidal potential and modulatory functions remaining unclear. Using a murine model of acute *P. aeruginosa* lung infection, we demonstrate cathelicidin-mediated enhancement of bacterial clearance *in vivo* in the absence of direct early




Figure 8. Exogenous LL-37 promotes an early neutrophil response to *P. aeruginosa* in cathelicidin-deficient mice. *Camp*^{-/-} mice were inoculated with 3×10^7 cfu of *P. aeruginosa* PAO1 and 10 μ g LL-37 peptide or PBS by intranasal delivery. At 6 hours (a) or 24 hours (b) after inoculation mice were culled and their lungs were lavaged. BALF was cytocentrifuged and differential counts were conducted for neutrophils. Data show Tukey box and whiskers plots for $n \geq 8$ animals per condition. Analyses were conducted using the Mann Whitney test; * $p < 0.05$. doi:10.1371/journal.pone.0099029.g008

microbicidal activity. Administration of synthetic LL-37 promoted an upregulation of the early neutrophil response that was dependent upon both infection and peptide, but was independent of native cathelicidin production, and enhanced bacterial clearance from the lung. Mice deficient in endogenous mCRAMP had a normal initial neutrophil response to infection, during a period in which *Camp* was not initially expressed in the wild type lung. However, these *Camp*^{-/-} mice were deficient in the larger second phase neutrophil response observed in *Camp*-expressing infected wild type mice and had an impaired capacity to clear the infection. To our knowledge, this is the first demonstration of cathelicidins enhancing host defence against infection through primarily modulatory mechanisms *in vivo* and suggests induction of a cathelicidin-mediated protective proinflammatory response.

The associations between hCAP-18/LL-37 expression and susceptibility to infection in humans [39][40][11][12] suggest an important, but as yet undefined role for hCAP-18/LL-37 in innate host defence against infection in humans. Although this peptide has microbicidal potential, its activity is poor against many microorganisms in physiologically relevant environments at the low concentrations found *in vivo* in most systems [27,28,29]. However, cathelicidins have additionally been shown to have multiple modulatory activities, including chemotactic function [34,35,36,37,38], the ability to modulate chemokine, cytokine and cellular responses [30,41,42,43], the capacity to alter leukocyte differentiation and function [44,45,46] and cell death modulating properties [28,47,48]. Critical to understanding these peptides and utilising their properties therapeutically is the need to clarify their modes of action *in vivo* in infectious contexts.

In this study, an acute murine pulmonary infection model with *P. aeruginosa* was utilised in order to evaluate the capacity of cathelicidins to enhance host defence against infection with a microbe which is largely resistant to these peptides under physiological conditions *in vitro* [27,28,29]. Under favourable *in vitro* conditions in which microbicidal properties are evident for LL-37, this peptide has been shown to permeabilise bacterial membranes within minutes [49]. However, we found no evidence for direct microbicidal activity against *P. aeruginosa* after co-incubation with LL-37 *in vivo*, yet exogenously delivered LL-37 was found to significantly enhance pathogen clearance over 24 hours. Although we cannot exclude some alternative form of late direct microbicidal activity of LL-37, even by 6 hours after infection no significant impact on bacterial load of the whole lung could be demonstrated in response to LL-37 treatment, although interestingly a therapeutic bolus of peptide was found to diminish the number of live bacteria accessible to bronchoalveolar lavage at this time point. The reason for this is unclear, but may relate to early removal of the most accessible bacteria by the enhanced neutrophil influx observed. A previous study using adenoviral vectors carrying the cDNA for hCAP18/LL-37, to overexpress the human cathelicidin in the murine lung over the 5 days prior to infection, resulted in the promotion of a small, but significant enhancement of *P. aeruginosa* clearance from the murine lung over a 24 hour period [21]. This was observed to be accompanied by decreased pulmonary TNF levels, but the mechanism underpinning this therapeutic effect was not evaluated and was assumed to be microbicidal. In contrast, we found no evidence to support a microbicidal effect, but demonstrate a peptide-mediated enhanced early neutrophil influx *in vivo*.

Prior research has demonstrated the capacity for cathelicidins to have direct chemotactic activity for human neutrophils and monocytes and murine leukocytes *in vitro* [34,36] and for murine leukocytes in an experimentally-formed murine air pouch model [36]. In that *in vivo* model, injection of 2 μ M LL-37 or mCRAMP

into the air pouch significantly enhanced the influx of neutrophils and monocytes within a 4 hour period. This is in contrast to the complete absence of neutrophils observed in our studies in the murine lung 6 hours after instillation of LL-37 alone (Figure 2b). In addition, LL-37 was not found to mediate any significant effects on the number of monocytes in the BALF, in contrast to the previously published findings in other systems [35,36]. A small, but significant neutrophil response was observed in the lungs of LL-37-treated uninfected mice at 24 hours after instillation (Figure 2d), demonstrating some LL-37-mediated neutrophil influx. However, indirect effects cannot be excluded and LL-37 has also been shown to enhance the production of neutrophil chemokines by other cells [30,50]. Nevertheless, despite this absence of any substantial response to LL-37 alone, a significantly enhanced pulmonary neutrophilia was observed in response to LL-37 upon concomitant infection (Figure 2a). With regard to this apparent contradiction to previous findings, it is worth noting that even the control air pouches in the previous report yielded a substantial number of neutrophils [36], indicating that this was already an inflamed environment and may in fact be analogous to infected lungs in our study. The requirement for concomitant infection in order to establish the early enhanced LL-37-mediated neutrophil response suggests a synergy with inflammatory mediators that remain to be identified, with no peptide-mediated modulation having been observed in the levels of the chemokines studied (including KC, MIP2, MCP-1, TNF and IL-6). Nevertheless, these data provide clear evidence for the *in vivo* capacity of exogenous LL-37 to modulate the innate cellular immune response in the context of pulmonary infection, enhancing pathogen clearance in the absence of microbicidal activity and having potential therapeutic implications.

In addition to the potential therapeutic roles of exogenous cathelicidins, the primary roles of endogenous peptides in pulmonary infection remain unclear. Studies using *Camp*^{-/-} mice have demonstrated a deficiency in the clearance of both pulmonary *Klebsiella pneumoniae* and *P. aeruginosa* infections [19,20]. *K. pneumoniae* promoted a later induction of pulmonary *Camp* expression than *P. aeruginosa* and mCRAMP appeared to have a more potent effect on *K. pneumoniae*, with *Camp*^{-/-} mice having a significant and severe clearance defect at 24 hours, resulting in more florid inflammation by 48 hours in the absence of endogenous cathelicidin and increased mortality [20]. Given that, in the case of *K. pneumoniae*, mCRAMP was reported to have microbicidal effects at a relatively modest 1 μ M, it is possible that the phenotype in this particular infection was influenced by loss of a relevant microbicidal agent in the face of a lethal infectious dose. Indeed, the late induction of *Camp* in wild type mice infected with *K. pneumoniae* may suggest less relevance for the peptide in the inflammatory response to this particular infection. In contrast, we demonstrate that *P. aeruginosa* infection can ultimately be controlled and cleared even in the absence of mCRAMP, and the earlier expression in wild type mice may indicate a more important role in the inflammatory response. The previous study reported that *Camp*^{-/-} mice infected with *P. aeruginosa* (at a lower infectious dose than in our study) has a significantly impaired bacterial clearance at 48 hours, with a decreased neutrophil response at 24 hrs [20]. These observations are compatible with our study, but were attributed to a loss of a direct chemotactic response to endogenous cathelicidin. Our new data, examining earlier timepoints, indicate that the initial murine pulmonary neutrophil response to *P. aeruginosa* infection precedes induction of and is independent of mCRAMP, and thus proceeded normally in *Camp*^{-/-} mice. However, second phase neutrophil influx was dependent upon mCRAMP expression, which may synergise with

infection-induced factors as yet unidentified, and was therefore defective in *Camp*^{-/-} mice, in whom impaired pathogen clearance then occurred. These data suggest that in individuals with impaired endogenous cathelicidin production, an effective, protective pulmonary inflammatory response will be suboptimal.

hCAP-18/LL-37 in humans is pre-formed in neutrophil granules, but can also be induced in a vitamin-D dependent manner in epithelial cells and macrophages [51,52,53,54]. In addition, strategies to induce LL-37 expression are under development, including the use of compounds such as 4-phenylbutyrate (reviewed in [55]), which can effectively upregulate hCAP-18/LL-37 expression *in vitro*, including in airway epithelial cells [56], and *in vivo* in a model of Shigella infection [57]. Such approaches may be of value in enhancing protective cathelicidin expression in humans, particularly in vitamin-D insufficient seasonal conditions. However, it was also important to consider whether therapeutic application of cathelicidin could provide rapid short term improvement of host defence in the absence of effective endogenous cathelicidin expression. In this regard, our studies demonstrate that the delivery of LL-37 to *P. aeruginosa* infected mice could promote an early neutrophil response and enhanced pathogen clearance in *Camp*^{-/-} mice as effectively as in wild type mice. These data indicate that this protective effect was in response to the exogenous LL-37 delivered, independent of endogenous mCRAMP expression and of native cathelicidin release from incoming PMN and supports the potential for the use of exogenous peptides in infection.

Thus, using a murine model of acute *P. aeruginosa* lung infection, we demonstrate cathelicidin-mediated enhancement of bacterial

clearance *in vivo* in the absence of direct microbicidal activity. The delivery of exogenous cathelicidin functioned to enhance a protective pro-inflammatory response to infection, promoting bacterial clearance from the lung, with an infection- and peptide-dependent early neutrophil response that was independent of native cathelicidin production. Furthermore, although *Camp*^{-/-} mice had an intact early cellular inflammatory response (which was comparable to cathelicidin-sufficient animals in the period preceding the induction of mCRAMP expression), they had significantly impaired bacterial clearance and absence of a second phase neutrophil response to infection. These findings demonstrate the importance of the inflammomodulatory properties of cathelicidins in pulmonary infection *in vivo* and highlight the significance of understanding and utilising these properties in the development of novel therapeutic approaches.

Acknowledgments

The authors would like to thank Peter Barlow, Robert Gray, Mark Marsden, Pat Swan and Sharon Hannah for advice and assistance.

Author Contributions

Conceived and designed the experiments: PEB BM JRWG AJS DJD. Performed the experiments: PEB BM EGF AM KJM. Analyzed the data: PEB BM DJD. Contributed reagents/materials/analysis tools: RLG JRWG. Wrote the paper: BM DJD. Expert revision of the manuscript: RLG AJS.

References

- Zasloff M (2002) Antimicrobial peptides of multicellular organisms. *Nature* 415: 389–395.
- Beaumont PE, Li H, Davidson DJ (2013) LL-37: An Innate Immunomodulatory Antimicrobial Host Defence Peptide. In: Hiemstra PS, Zaat SAJ, Antimicrobial peptides and Innate Immunity. Basel: Springer pp. 97–122.
- Semple F, Dorin JR (2012) beta-Defensins: Multifunctional Modulators of Infection, Inflammation and More? *J Innate Immun* 4: 337–348.
- Bowdish DME, Davidson DJ, Hancock REW (2005) A re-evaluation of the role of host defence peptides in mammalian immunity. *Curr Protein Pept Sci* 6: 35–51.
- Lipsky BA, Holroyd KJ, Zasloff M (2008) Topical versus systemic antimicrobial therapy for treating mildly infected diabetic foot ulcers: a randomized, controlled, double-blinded, multicenter trial of pexiganan cream. *Clin Infect Dis* 47: 1537–1545.
- Scott MG, Dullaghan E, Mookherjee N, Glavas N, Waldbrook M, et al. (2007) An anti-infective peptide that selectively modulates the innate immune response. *Nat Biotechnol* 25: 465–472.
- Rivas-Santiago B, Castaneda-Delgado JE, Rivas Santiago CE, Waldbrook M, Gonzalez-Curiel I, et al. (2013) Ability of Innate Defence Regulator Peptides IDR-1002, IDR-HH2 and IDR-1018 to Protect against Mycobacterium tuberculosis Infections in Animal Models. *PLoS One* 8: e59119.
- Zanetti M (2005) The role of cathelicidins in the innate host defenses of mammals. *Curr Issues Mol Biol* 7: 179–196.
- Gudmundsson GH, Agerberth B, Odeberg J, Bergman T, Olsson B, et al. (1996) The human gene FALL39 and processing of the cathelin precursor to the antibacterial peptide LL-37 in granulocytes. *Eur J Biochem* 238: 325–332.
- Sorensen OE, Follin P, Johnsen AH, Calafat J, Tjallingii GS, et al. (2001) Human cathelicidin, hCAP-18, is processed to the antimicrobial peptide LL-37 by extracellular cleavage with proteinase 3. *Blood* 97: 3951–3959.
- Schaller-Bals S, Schulze A, Bals R (2002) Increased Levels of Antimicrobial Peptides in Tracheal Aspirates of Newborn Infants during Infection. *Am J Respir Crit Care Med* 165: 992–995.
- Mansbach JM, Piedra PA, Borregaard N, Martineau AR, Neuman MI, et al. (2012) Serum cathelicidin level is associated with viral etiology and severity of bronchiolitis. *J Allergy Clin Immunol* 130: 1007–1008.e1001.
- Gallo RL, Kim KJ, Bernfield M, Kozak CA, Zanetti M, et al. (1997) Identification of CRAMP, a cathelin-related antimicrobial peptide expressed in the embryonic and adult mouse. *J Biol Chem* 272: 13088–13093.
- Pestonjamas VK, Huttner KH, Gallo RL (2001) Processing site and gene structure for the murine antimicrobial peptide CRAMP. *Peptides* 22: 1643–1650.
- Nizet V, Ohtake T, Lauth X, Trowbridge J, Rudisill J, et al. (2001) Innate antimicrobial peptide protects the skin from invasive bacterial infection. *Nature* 414: 454–457.
- Iimura M, Gallo RL, Hase K, Miyamoto Y, Eckmann L, et al. (2005) Cathelicidin Mediates Innate Intestinal Defense against Colonization with Epithelial Adherent Bacterial Pathogens. *J Immunol* 174: 4901–4907.
- Huang LC, Reins RY, Gallo RL, McDermott AM (2007) Cathelicidin-deficient (*Cnlp*^{-/-}) mice show increased susceptibility to *Pseudomonas aeruginosa* keratitis. *Invest Ophthalmol Vis Sci* 48: 4498–4508.
- Chromek M, Slamova Z, Bergman P, Kovacs L, Podracka L, et al. (2006) The antimicrobial peptide cathelicidin protects the urinary tract against invasive bacterial infection. *Nat Med* 12: 636–641.
- Yu FS, Cornicelli MD, Kovach MA, Newstead MW, Zeng X, et al. (2010) Flagellin Stimulates Protective Lung Mucosal Immunity: Role of Cathelicidin-Related Antimicrobial Peptide. *J Immunol* 185: 1142–1149.
- Kovach MA, Ballinger MN, Newstead MW, Zeng X, Bhan U, et al. (2012) Cathelicidin-Related Antimicrobial Peptide Is Required for Effective Lung Mucosal Immunity in Gram-Negative Bacterial Pneumonia. *J Immunol* 189: 304–311.
- Bals R, Weiner DJ, Moscioni AD, Meegalla RL, Wilson JM (1999) Augmentation of innate host defense by expression of a cathelicidin antimicrobial peptide. *Infect Immun* 67: 6084–6089.
- Barlow PG, Svoboda P, Mackellar A, Nash AA, York IA, et al. (2011) Antiviral Activity and Increased Host Defense against Influenza Infection Elicited by the Human Cathelicidin LL-37. *PLoS One* 6: e25333.
- British Thoracic Society (2006) The burden of lung disease: a statistical report from the British Thoracic Society. 2nd ed London: British Thoracic Society.
- Davidson DJ, Currie AJ, Speert DP (2003) *Pseudomonas aeruginosa* infections in individuals with cystic fibrosis: North American perspective. In: Hause A and Rello J, Severe infections caused by *Pseudomonas aeruginosa*. Norwell: Kluwer Academic Publishers. pp. 71–89.
- Carmeli Y, Troillet N, Eliopoulos GM, Samore MH (1999) Emergence of antibiotic-resistant *Pseudomonas aeruginosa*: comparison of risks associated with different antipseudomonal agents. *Antimicrob Agents Chemother* 43: 1379–1382.
- Hermes DM, Pommann Pitt C, Lutz L, Teixeira AB, Ribeiro VB, et al. (2013) Evaluation of heteroresistance to polymyxin B among carbapenem-susceptible and -resistant *Pseudomonas aeruginosa*. *J Med Microbiol* 62: 1184–1189.
- Bowdish DME, Davidson DJ, Lau YE, Lee K, Scott MG, et al. (2005) Impact of LL-37 on anti-infective immunity. *J Leukoc Biol* 77: 451–459.

28. Barlow PG, Beaumont PE, Cossecau C, Mackellar A, Wilkinson TS, et al. (2010) The Human Cathelicidin LL-37 Preferentially Promotes Apoptosis of Infected Airway Epithelium. *Am J Respir Cell Mol Biol* 43: 692–702.
29. Pompilio A, Scocchi M, Pomponio S, Guida F, Di Primio A, et al. (2011) Antibacterial and anti-biofilm effects of cathelicidin peptides against pathogens isolated from cystic fibrosis patients. *Peptides* 32: 1807–1814.
30. Scott MG, Davidson DJ, Gold MR, Bowdish DME, Hancock REW (2002) The Human Antimicrobial Peptide LL-37 Is a Multifunctional Modulator of Innate Immune Responses. *J Immunol* 169: 3883–3891.
31. Wang Y, Agerberth B, Johansson J (1998) Structure and activity of cathelicidin antibacterial proteins. *J Protein Chem* 17: 522–523.
32. Weiner DJ, Bucki R, Janmey PA (2003) The antimicrobial activity of the cathelicidin LL37 is inhibited by F-actin bundles and restored by gelsolin. *Am J Respir Cell Mol Biol* 28: 738–745.
33. Bucki R, Byfield FJ, Janmey PA (2007) Release of the antimicrobial LL37 peptide from DNA/F-actin bundles in CF sputum. *Eur Respir J* 29: 624–632.
34. Yang D, Chen Q, Schmidt AP, Anderson GM, Wang JM, et al. (2000) LL-37, the neutrophil granule- and epithelial cell-derived cathelicidin, utilizes formyl peptide receptor-like 1 (FPRL1) as a receptor to chemoattract human peripheral blood neutrophils, monocytes, and T cells. *J Exp Med* 192: 1069–1074.
35. Wantha S, Alard JE, Megens RT, van der Does AM, Doring Y, et al. (2013) Neutrophil-derived cathelicidin promotes adhesion of classical monocytes. *Circ Res* 112: 792–801.
36. Kurosaka K, Chen Q, Yarovsky F, Oppenheim JJ, Yang D (2005) Mouse cathelin-related antimicrobial Peptide chemoattracts leukocytes using formyl Peptide receptor-like 1/mouse formyl Peptide receptor-like 2 as the receptor and acts as an immune adjuvant. *J Immunol* 174: 6257–6265.
37. Zhang Z, Cherryholmes G, Chang F, Rose DM, Schraufstatter I, et al. (2009) Evidence that cathelicidin peptide LL-37 may act as a functional ligand for CXCR2 on human neutrophils. *Eur J Immunol* 39: 3181–3194.
38. Tjabringa GS, Ninaber DK, Drijfhout JW, Rabe KF, Hiemstra PS (2006) Human Cathelicidin LL-37 Is a Chemoattractant for Eosinophils and Neutrophils That Acts via Formyl-Peptide Receptors. *Int Arch Allergy Immunol* 140: 103–112.
39. Putsep K, Carlsson G, Boman HG, Andersson M (2002) Deficiency of antibacterial peptides in patients with morbus Kostmann: an observation study. *Lancet* 360: 1144–1149.
40. Schaubert J, Gallo RL (2008) Antimicrobial peptides and the skin immune defense system. *J Allergy Clin Immunol* 122: 261–266.
41. Mookherjee N, Brown KL, Bowdish DME, Doria S, Falsafi R, et al. (2006) Modulation of the TLR-Mediated Inflammatory Response by the Endogenous Human Host Defense Peptide LL-37. *J Immunol* 176: 2455–2464.
42. Nagaoka I, Hirota S, Niyosaba F, Hirata M, Adachi Y, et al. (2001) Cathelicidin family of antibacterial peptides CAP18 and CAP11 inhibit the expression of TNF-alpha by blocking the binding of LPS to CD14(+) cells. *J Immunol* 167: 3329–3338.
43. Lande R, Gregorio J, Facchinetti V, Chatterjee B, Wang YH, et al. (2007) Plasmacytoid dendritic cells sense self-DNA coupled with antimicrobial peptide. *Nature* 449: 564–569.
44. Davidson DJ, Currie AJ, Reid GS, Bowdish DME, MacDonald KL, et al. (2004) The cationic antimicrobial peptide LL-37 modulates dendritic cell differentiation and dendritic cell-induced T cell polarization. *J Immunol* 172: 1146–1156.
45. van der Does AM, Beekhuizen H, Ravensbergen B, Vos T, Ottenhoff TH, et al. (2010) LL-37 directs macrophage differentiation toward macrophages with a proinflammatory signature. *J Immunol* 185: 1442–1449.
46. Alalwani MS, Sierigk J, Herr C, Pinkenburg O, Gallo R, et al. (2010) The antimicrobial peptide LL-37 modulates the inflammatory and host defense response of human neutrophils. *Eur J Immunol* 40: 1118–1126.
47. Li HN, Barlow PG, Bylund J, Mackellar A, Bjorstad A, et al. (2009) Secondary necrosis of apoptotic neutrophils induced by the human cathelicidin LL-37 is not proinflammatory to phagocytosing macrophages. *J Leukoc Biol* 86: 891–902.
48. Mader JS, Mookherjee N, Hancock RE, Bleackley RC (2009) The Human Host Defense Peptide LL-37 Induces Apoptosis in a Calpain- and Apoptosis-Inducing Factor-Dependent Manner Involving Bax Activity. *Mol Cancer Res* 7: 689–702.
49. Turner J, Cho Y, Dinh NN, Waring AJ, Lehrer RI (1998) Activities of LL-37, a cathelin-associated antimicrobial peptide of human neutrophils. *Antimicrob Agents Chemother* 42: 2206–2214.
50. Tjabringa GS, Aarbiou J, Ninaber DK, Drijfhout JW, Sorensen OE, et al. (2003) The Antimicrobial Peptide LL-37 Activates Innate Immunity at the Airway Epithelial Surface by Transactivation of the Epidermal Growth Factor Receptor. *J Immunol* 171: 6690–6696.
51. Wang TT, Nestel FP, Bourdeau V, Nagai Y, Wang Q, et al. (2004) Cutting edge: 1,25-dihydroxyvitamin D3 is a direct inducer of antimicrobial peptide gene expression. *J Immunol* 173: 2909–2912.
52. Gombart AF, Borregaard N, Koeffler HP (2005) Human cathelicidin antimicrobial peptide (CAMP) gene is a direct target of the vitamin D receptor and is strongly up-regulated in myeloid cells by 1,25-dihydroxyvitamin D3. *FASEB J* 19: 1067–1077.
53. Liu PT, Stenger S, Li H, Wenzel L, Tan BH, et al. (2006) Toll-like receptor triggering of a vitamin D-mediated human antimicrobial response. *Science* 311: 1770–1773.
54. Yim S, Dhawan P, Ragunath C, Christakos S, Diamond G (2007) Induction of cathelicidin in normal and CF bronchial epithelial cells by 1,25-dihydroxyvitamin D(3). *J Cyst Fibros* 6: 403–410.
55. van der Does AM, Bergman P, Agerberth B, Lindbom L (2012) Induction of the human cathelicidin LL-37 as a novel treatment against bacterial infections. *J Leukoc Biol* 92: 735–742.
56. Steinmann J, Halldorsson S, Agerberth B, Gudmundsson GH (2009) Phenylbutyrate induces antimicrobial peptide expression. *Antimicrob Agents Chemother* 53: 5127–5133.
57. Sarker P, Ahmed S, Tiash S, Rekha RS, Stromberg R, et al. (2011) Phenylbutyrate counteracts Shigella mediated downregulation of cathelicidin in rabbit lung and intestinal epithelia: a potential therapeutic strategy. *PLoS One* 6: e20637.