UC Irvine

UC Irvine Previously Published Works

Title

Coherent-state Hall effect in the heavy fermions CeCu₆ and U₂PtC₂ (abstract)

Permalink

<https://escholarship.org/uc/item/8qb3b5v9>

Journal

Journal of Applied Physics, 61(8)

ISSN

0021-8979

Authors

Penney, T
Milliken, FP
Holtzberg, F
et al.

Publication Date

1987-04-15

DOI

10.1063/1.338435

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Coherent-state Hall effect in the heavy fermions CeCu₆ and U₂PtC₂ (abstract)

T. Penney, F. P. Milliken, and F. Holtzberg

IBM T. J. Watson Research Center, P. O. Box 218, Yorktown Heights, New York 10598

Z. Fisk

Los Alamos National Laboratory, Los Alamos, New Mexico 87545

G. P. Meisner

General Motors Research Laboratories, Warren, Michigan 48090

The Hall effect is found to set the scale for coherence in the normal state of CeCu₆ and U₂PtC₂. CeCu₆ is a nonmagnetic, nonsuperconducting heavy fermion system.¹⁻⁴ Its resistivity at high temperature is like that of a collection of incoherent Kondo scatterers. At low temperature the resistivity smoothly decreases to a very small value, indicating that scattering has become coherent. The Hall effect has two strong extrema in its temperature dependence which define a high-temperature incoherent scattering region, a transition region, and a low temperature coherent region. Although U₂PtC₂ is superconducting below 1.5 K,⁵ the Hall results in the normal state show two extrema similar to those of CeCu₆. However, these features are scaled to higher temperatures, consistent with the smaller low-temperature electronic specific heat and higher Fermi temperature.

Work at Los Alamos supported by the U. S. Dept. of Energy.

¹H. R. Ott *et al.*, Solid State Commun. **53**, 235 (1985).

²Y. Onuki *et al.*, J. Phys. Soc. Jpn. **54**, 2804 (1985).

³J. Flouquet *et al.*, J. Magn. Magn. Mater. **52**, 85 (1985).

⁴T. Penney *et al.*, J. Magn. Magn. Mater. **54-57**, 370 (1986); Phys. Rev. B **34**, 5959 (1986).

⁵G. P. Meisner *et al.*, Phys. Rev. Lett. **53**, 1829 (1984).

Hall effect in heavy fermion compounds (abstract)

P. M. Levy

Department of Physics, New York University, New York, New York 10003

A. Fert

Laboratoire de Physique des Solides, Université de Paris-Sud 91405 Orsay, France

As a result of a recent analysis of data on the Hall effect in heavy fermion compounds¹ we evaluated the higher-order resonant scattering contributions to the skew scattering of conduction electrons. In the single-site approximation we have calculated the *t* matrix elements for skew scattering correct to fourth order in the Anderson mixing interaction, i.e., we have neglected pair correlation effects. By including this fourth-order correction in our calculation of the Hall resistivity we find it is positive at high temperatures which is in agreement with data on heavy fermion compounds. Prior calculations of this scattering in these compounds were limited to second order in the mixing interaction above *T_K*, and predicted a negative Hall constant at high temperatures.² By using a phase shift analysis of the skew scattering we have extended our calculation to the incoherent (single-site) strong coupling regime. In the temperature ranges we considered the Hall resistivity is proportional to the product of the resistivity and magnetic susceptibility. This is in agreement with data on heavy fermion compounds.

¹M. Hadžić-Leroux, A. Hamzić, A. Fert, P. Haen, F. Lapierre, and O. Laborde, Europhys. Lett. **1**, 579 (1986).

²P. Coleman, P. W. Anderson, and T. V. Ramakrishnan, Phys. Rev. Lett. **55**, 414 (1985).

Journal of Applied Physics is copyrighted by the American Institute of Physics (AIP). Redistribution of journal material is subject to the AIP online journal license and/or AIP copyright. For more information, see <http://ojps.aip.org/japo/japcr/jsp>
Copyright of Journal of Applied Physics is the property of American Institute of Physics and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.

Journal of Applied Physics is copyrighted by the American Institute of Physics (AIP). Redistribution of journal material is subject to the AIP online journal license and/or AIP copyright. For more information, see <http://ojps.aip.org/japo/japcr/jsp>