

UCSF

UC San Francisco Previously Published Works

Title

Rationale and Design of the Registry for Stones of the Kidney and Ureter (ReSKU): A Prospective Observational Registry to Study the Natural History of Urolithiasis Patients

Permalink

<https://escholarship.org/uc/item/8h2421b6>

Journal

Journal of Endourology, 30(12)

ISSN

0892-7790

Authors

Chang, Helena C
Tzou, David T
Usawachintachit, Manint
[et al.](#)

Publication Date

2016-12-01

DOI

10.1089/end.2016.0648

Peer reviewed

Rationale and Design of the Registry for Stones of the Kidney and Ureter (ReSKU): A Prospective Observational Registry to Study the Natural History of Urolithiasis Patients

Helena C. Chang, MD,¹ David T. Tzou, MD,¹ Manint Usawachintachit, MD,^{1,2} Brian D. Duty, MD,³
Ryan S. Hsi, MD,¹ Jonathan D. Harper, MD,⁴ Mathew D. Sorensen, MD,⁴ Marshall L. Stoller, MD,¹
Roger L. Sur, MD,⁵ and Thomas Chi, MD¹

Abstract

Objectives: Registry-based clinical research in nephrolithiasis is critical to advancing quality in urinary stone disease management and ultimately reducing stone recurrence. A need exists to develop Health Insurance Portability and Accountability Act (HIPAA)-compliant registries that comprise integrated electronic health record (EHR) data using prospectively defined variables. An EHR-based standardized patient database—the Registry for Stones of the Kidney and Ureter (ReSKUTM)—was developed, and herein we describe our implementation outcomes.

Materials and Methods: Interviews with academic and community endourologists in the United States, Canada, China, and Japan identified demographic, intraoperative, and perioperative variables to populate our registry. Variables were incorporated into a HIPAA-compliant Research Electronic Data Capture database linked to text prompts and registration data within the Epic EHR platform. Specific data collection instruments supporting New patient, Surgery, Postoperative, and Follow-up clinical encounters were created within Epic to facilitate automated data extraction into ReSKU.

Results: The number of variables within each instrument includes the following: New patient—60, Surgery—80, Postoperative—64, and Follow-up—64. With manual data entry, the mean times to complete each of the clinic-based instruments were (minutes) as follows: New patient— 12.06 ± 2.30 , Postoperative— 7.18 ± 1.02 , and Follow-up— 8.10 ± 0.58 . These times were significantly reduced with the use of ReSKU structured clinic note templates to the following: New patient— 4.09 ± 1.73 , Postoperative— 1.41 ± 0.41 , and Follow-up— 0.79 ± 0.38 . With automated data extraction from Epic, manual entry is obviated.

Conclusions: ReSKU is a longitudinal prospective nephrolithiasis registry that integrates EHR data, lowering the barriers to performing high quality clinical research and quality outcome assessments in urinary stone disease.

Keywords: prospective studies, registries, urolithiasis

Introduction

HIGH QUALITY PROSPECTIVE data on long-term outcomes have been lacking in urolithiasis research. Without it, many basic aspects of urinary stone disease remain poorly understood, including the natural history of stone recurrence and ne-

phrolithiasis treatment practice trends and quality outcomes. Longitudinal observational registries provide an unparalleled opportunity for high quality research to support evidence-based care.¹ Within the field of urology, multiple examples exist of effective registries, including the Surveillance, Epidemiology, End-Results Program, Cancer of the Prostate Strategic Urologic

¹Department of Urology, University of California San Francisco, San Francisco, California.

²Division of Urology, Faculty of Medicine, Chulalongkorn University and King Chulalongkorn Memorial Hospital, Bangkok, Thailand.

³Department of Urology, Oregon Health and Science University, Portland, Oregon.

⁴Department of Urology, University of Washington, Seattle, Washington.

⁵Department of Urology, University of California San Diego, La Jolla, California.

Research Endeavor (CaPSURE™), Prostate Cancer Outcomes Study, and the American Urological Association (AUA) Quality (AQUA) Registry. These registries have been used to evaluate disease prevalence, treatment outcomes, national practice trends, and health service utilization across multiple diseases leading to advances in patient care.² Unfortunately, most registry efforts still rely on manual data entry, which results in expensive, labor-intensive long-term maintenance.³

Data entry for traditional registries is time-consuming and labor intensive. We hypothesized that by designing a registry tied to an electronic health record (EHR), we could significantly decrease the time of data extraction. Our goal was to design the Registry for Stones of the Kidney and Ureter (ReSKU™) as a longitudinal, observational prospective registry of patients with urinary stone disease to address the limitations of traditional registries. To do so, inpatient and outpatient data in all phases of care would be extracted from the EHR prospectively in an automated manner and then organized in a secure, web-based application portal designed to support data capture for research studies. This approach will improve the quality of care for patients with urinary stone disease by prospectively tracking data regarding patient characteristics, surgical outcomes, follow-up management, and stone recurrence. One primary aim for ReSKU is to lower the barriers to participating in and maintaining a registry for urinary stone patients for any practice setting. In this study, we describe the design approach and initial experience with building and implementing ReSKU.

Materials and Methods

Evaluation of registry options

In developing ReSKU, it was important to first recognize how current registries operate and how an ideal

modern registry would compare since traditional registry design and data entry are accompanied by challenges that hinder their implementation. These include labor-intensive data entry, cost of registry maintenance, patient confidentiality with meeting Health Insurance Portability and Accountability Act (HIPAA) requirements and human subjects research regulations, manpower for maintaining and organizing a database, concerns for how data will be stored and used, and interference with day to day clinical care³⁻⁶ (Table 1).

A modern registry should ideally be easy to populate through integration with existing EHR systems and have a low maintenance cost, all while recording accurate, detailed patient clinical information, documenting surgeon outcomes, and tracking quality of care.⁷ By leveraging the EHR, many of the challenges facing patient registries can be overcome.

Secondarily, the EHR would ideally be coupled with a data warehouse to allow information to be extracted from the EHR and housed securely in an easily accessible manner. For ReSKU, we elected to utilize a Research Electronic Data Capture (REDCap) database. REDCap is a mature, HIPAA-compliant, secure web application for building and managing online surveys and databases. It provides the following: (1) an intuitive interface for validated data entry; (2) audit trails for tracking data manipulation and export procedures; (3) automated export procedures for seamless data downloads to common statistical packages; and (4) procedures for importing data from external sources.⁸ The Epic platform (Verona, WI) was the initial system selected to build the EHR tools for ReSKU, given that it has consistently had the largest market share of any EHR vendor.⁹ These two software systems were selected as they are widely utilized in the medical and research communities and contain native features that make integration readily achievable.¹⁰

TABLE 1. COMPARISON OF BENEFITS AND CHALLENGES OF DIFFERENT REGISTRY APPROACHES

	<i>Traditional registry</i>	<i>Registry using REDCap and without EHR integration</i>	<i>Registry using REDCap and with EHR integration</i>
Benefits	<ul style="list-style-type: none"> • Database design and management • Works with both EHR and paper charts 	<ul style="list-style-type: none"> • Works with both EHR and paper charts • HIPAA compliant data management • REDCap makes database design and updates easy • Electronic web-based registry 	<ul style="list-style-type: none"> • Works with both EHR and paper charts • HIPAA compliant data management • REDCap makes database design and updates easy • Electronic web-based registry • Automated data entry • Automated EHR data extraction • Dropdown lists in clinic notes • Data quality is as good as clinical data
Challenges	<ul style="list-style-type: none"> • Manual data entry • Interference with clinical flow • Data quality dependent on data entry • Need for additional personnel • Patient confidentiality • Increased paperwork • Updating spreadsheets or data management software 	<ul style="list-style-type: none"> • Manual data entry • Interference with clinical flow • Data quality dependent on data entry • May need additional personnel 	<ul style="list-style-type: none"> • Upfront investment in programming structured data and data extraction with analysts

EHR = electronic health record; HIPAA = Health Insurance Portability and Accountability Act; REDCap = Research Electronic Data Capture.

TABLE 2. VARIABLES CONSIDERED FOR INCLUSION IN THE REGISTRY

	Encounter in which specific variables are captured			
	New Patient variables	Surgery variables	Postoperative Follow up variables	Follow up variables
Included in ReSKU™	<ul style="list-style-type: none"> • Race • Gender • Age • Education • Past medical history • Past surgical history • Family history • Alcohol, tobacco, substance use • Biometrics • Anatomic anomalies (congenital and acquired) • Stone radiologic characteristics 	<ul style="list-style-type: none"> • Serum labs • Urine culture • ASA score • Stone surgical characteristics • Type of surgery • Equipment used • Surgery duration • Urinary drainage tube type • Estimated blood loss • Surgeon characteristics 	<ul style="list-style-type: none"> • Stone analysis • Hospital stay length • Residual stone after surgery • Need for staged procedures • Complications • Urinary drainage tube duration 	<ul style="list-style-type: none"> • 24-hour urine parameters • Follow up imaging • Recurrence management strategy • Intervention compliance • Stone recurrence • Serum labs
Excluded	<ul style="list-style-type: none"> • Income factors • Work exposures • Environmental exposures • Country of origin • Diet and hydration status • Activity level • Climate geography 	<ul style="list-style-type: none"> • Anesthetics used • Equipment usage duration 	<ul style="list-style-type: none"> • Access to care • Insurance coverage • Cost 	<ul style="list-style-type: none"> • Urinalysis • Genetic analysis • Patient quality of life • Patient satisfaction with care • Patient pain metrics • Readmission • Outside hospital ED visits

Data points listed demonstrate conceptualizing the determinants of stone patient outcomes. Included variables in bold are readily collected as part of each patient clinical encounter.

ASA = American Society of Anesthesiologists; BMP = Basic metabolic panel; ReSKU™ = Registry for Stones of the Kidney and Ureter.

Selecting measures to collect for the ReSKU study

The determinants of patient-centered outcomes were broadly conceptualized, relying on both urologist practice patterns and the existing literature to build a framework of variables for collection. To maximize the validity, reproducibility, and quantitative nature of the data collected, current AUA and European Association of Urology guidelines for urolithiasis management were referenced.¹¹⁻¹³ Where possible, validated clinical instruments, including American Society of Anesthesiologists score,¹⁴ Clavien–Dindo score,¹⁵ and modified Guy’s stone score,¹⁶ were integrated. Variables were divided into broad categories: sociodemographic characteristics, clinical and biometric characteristics, stone characteristics, operative factors, process characteristics, and outcomes. The next design phase focused on feasible implementation of ReSKU into daily clinical practice.

While the ideal prospective observational registry would record all potential stone-related determinants of health, this approach is impractical given the time constraints intrinsic to clinical visits. Therefore, we narrowed the measures to be collected down to the most critical variables that could be

feasibly gathered in the usual course of clinical care. After a comprehensive list of measures was constructed, we performed collaborative interviews of 11 academic and community endourologists based in the United States, Canada, China, and Japan to determine which of these clinical variables to collect. This resulted in a condensed set of variables for inclusion in the ReSKU database, separated into instruments specific to each patient encounter (Table 2). The registry design process evolved over the course of 1 year, involving multiple rounds of data instrument pretesting and revision. Particular attention was paid to structuring variables into queries that could be collected in the course of routine patient care with minimal disruption to the clinical encounter. Pre-testing included piloting data collection at two medical centers with assessment of compliance for accuracy of data collection.

Patient selection

After obtaining appropriate institutional review board (IRB) approval, all patients of any age presenting to the urology clinic for evaluation of stone disease who had clinical evidence




FIG. 1. ReSKU™ database structure is based on clinical flow for patient care. *Light blue arrows* indicate clinical flow. *White boxes* represent the type of patient encounter. The *light gray box* represents the EHR. Under “EHR data extraction,” the *colored boxes* indicate structured clinical note templates with dropdown lists of variables for data capture. The *arrows* portray how automated data integration occurs within ReSKU™. The *white arrows* delineate discrete data extraction from the EHR into ReSKU™ (e.g., laboratories and vital signs). The *gray arrows* represent data (e.g., stone symptoms and stone size) extracted from dropdown list selections within the EHR clinic note templates. Under “ReSKU,” the *blue box* represents the collection and storage of data into different variable categories using REDCap. EHR, electronic health record; REDCap=Research Electronic Data Capture; ReSKU™, Registry for Stones of the Kidney and Ureter.

of nephrolithiasis were included. Clinical evidence of nephrolithiasis was defined as any upper tract urinary stone demonstrated by imaging performed within the 12 months before clinic presentation or adjudicated passage (i.e., the patient brought their stone to their clinic visit). Patients were included into the registry after providing informed consent for participation. No patients who met these inclusion criteria were excluded from participation in ReSKU.

Results

ReSKU registry variables were incorporated into the REDCap database by dividing all of the variables into four specific data collection instruments to parallel patient encounters: New Outpatient, Surgery, Postoperative, and Follow-Up visits (Table 2). Instruments were structured in a patient-centered manner by following the patient’s phases of care. An encounter is created each time the patient is seen. Within the EHR, clinical note templates related to each encounter were created: “New patient history and physical,” “Operative report,” “Postoperative progress note,” and “Follow-up progress note.” Within each template, dropdown lists were created for each of the registry variables. Using these dropdown lists facilitates storing note data in a structured format, which allows information extraction directly into the registry database, thereby bypassing manual data entry (Fig. 1). During the pilot phase,

compliance with accurate coding of clinical encounters into each of the note templates was assessed by three urologists independently, and where variables were perceived as possibly introducing uncertainty into coding accuracy, templates were adjusted to meet agreement between the three adjudicating urologists.

A total of 268 variables across all encounter types were implemented in ReSKU, with 60 New Patient, 80 Surgery, 64 Postoperative follow-up, and 64 Follow-up variables. Using conventional registry data population (i.e., a research assistant abstracting data from the provider’s clinical note into ReSKU), manual data entry into REDCap required 12.06 ± 2.3 minutes for New patient, 7.18 ± 1.02 minutes for Postoperative follow-up, and 8.10 ± 0.58 minutes for subsequent follow-up clinical encounters. By using ReSKU structured clinical note templates in Epic to document each encounter, the manual data entry times decreased to 4.09 ± 1.73 minutes for New patients, 1.41 ± 0.41 minute for Postoperative follow-up, and 0.79 ± 0.38 minute for subsequent Follow-up patients, respectively. With full EHR integration, use of ReSKU templates to document clinic encounters facilitated automated data extraction, and no added manual data entry time was required (Table 3).

Discussion

Clinical research in nephrolithiasis most often relies on retrospective clinical data collection.¹⁷ The resultant quality

TABLE 3. DATA ENTRY TIMES NEEDED TO POPULATE DATA COLLECTION INSTRUMENTS IN REDCAP

<i>Data abstraction source</i>	<i>New patient</i>	<i>Postoperative</i>	<i>Follow-up</i>
Number of variables	60	64	64
Conventional clinical notes, mean \pm SD (minutes)	12.06 \pm 2.30	7.18 \pm 1.02	8.10 \pm 0.58
Using ReSKU template, mean \pm SD (minutes)	4.09 \pm 1.73	1.41 \pm 0.41	0.79 \pm 0.38
Using ReSKU template with automated data extraction, mean \pm SD (minutes)	0	0	0
<i>p</i>	<0.01	<0.01	<0.01

Manual extraction from conventional unstructured clinic notes (third row) compared with manual extraction from ReSKU™ Epic clinic note templates (fourth row), and lastly implementation of ReSKU™ Epic clinic note templates along with automated data abstraction (fifth row). *p*-Values were calculated with unpaired Student's *t*-test using Stata 14.1 (StataCorp, College Station, TX).

of studies is dependent on the thoroughness of retrospective chart review, accuracy of visit coding, and accuracy of patient records. Prospective high quality registries linked to clinical outcomes are critical for forward progress in understanding the natural history of kidney stones.

Some prospective registries exist in urinary stone disease, but differ from ReSKU in that they are limited to specific procedures, have limited follow-up, or include only patients with rare kidney stones. Examples include the Clinical Research Office of the Endourological Society percutaneous nephrolithotomy and ureteroscopy global registries, the Brushite Kidney Stone Registry, the Rare Kidney Stone Consortium registries, the Percutaneous Nephrolithotomy Registry, the Health-Related Quality of Life in Rare Kidney Stone Registry, and the Vietnam Era Twin Registry.^{18–21} By design, each of these registries has a narrow focus with the intention of bringing attention to particular stone types or interventions. The aim of ReSKU is to capture the larger scope of stone disease burden across all stone types in order for study results to be more generalizable to a broader population of nephrolithiasis patients.

ReSKU's approach is unique in that it captures longitudinal clinical data already being collected by providers during the course of a patient encounter. The ReSKU data collection instruments were designed based on current stone management guidelines, quality care metrics, and endourologist interviews to identify the data points that any provider ought to be asking patients at each clinic visit. By tapping into the power of EHRs, ReSKU eliminates the burden of data entry, making prospective comprehensive inpatient and outpatient data collection on all stone patients a sustainable endeavor.

ReSKU's automated data extraction process drives several key advantages compared with traditional registries that rely on manual data entry. Since manual entry often relies on a research assistant or third party interpreting the provider's clinical notes to extract data, human data entry problems such as typographic errors and variable definition interpretation errors are common. These databases require significant time to rid the data set of erroneous entries. In contrast, ReSKU minimizes these errors since templates are populated by consistent dropdown menus. Moreover, the urologist is the one performing data collection directly into the clinic note, which then populates the data registry. Since the provider is best positioned to understand the patient's clinical status, registry data quality can be expected to be at least as good as the data collected for clinical care. In addition, data collection instruments in both clinic and operative note templates were designed with clinical care as well

as billing charge capture in mind. This allows ReSKU to reflect real world practice with minimal impact on provider documentation time.

Understanding that different practices will have their own set of time and available resources, ReSKU was designed so that providers can participate at different levels of data integration. While the optimum implementation of ReSKU includes full integration with the EHR for automated data capture, this is not the only available option. Based on experience with piloting ReSKU in medical centers without an EHR in place, implementation can still yield excellent data quality results with designated, trained research team members to collect and upload clinical data directly into REDCap (Table 3, row 1). REDCap data capture has been proven to be a robust way to manage instrument design and secure data collection across institutions.⁸ For practices with an EHR present but no infrastructure for automated data capture, ReSKU clinical templates can be built into the EHR and research team members can extract data from the template note into REDCap (Table 3, row 2). As seen by our results, the use of templates significantly reduced data entry time into REDCap. Therefore, any clinical provider who is interested in implementing ReSKU at their institution can do so. ReSKU's implementation strategy is flexible in that it can be modified for different practice settings to account for unique EHR environments and personnel.

Additional benefits of ReSKU will be realized with future clinical and translational research. As a longitudinal, prospective observational registry, ReSKU was designed to reveal patterns in clinical outcomes and practice as well as quality outcomes. However, once data collection is implemented into a provider's daily routine, clinical information will be continuously collected and managed in a HIPAA- and IRB-compliant manner. Such demographic and generalized nephrolithiasis-related clinical data provide a baseline framework on which to layer additional IRB-approved clinical studies. As new clinical and translational research questions arise surrounding validation of biomarkers, clinical endpoints, and technologies over time, ReSKU data can support project aims without the need for additional retrospective data extraction. For example, patients could be randomized into different groups for an interventional study. New data collection instruments can be easily integrated so that only specific data points related to the randomized trial need to be collected in addition to ReSKU data, thereby decreasing the amount of work needed to complete the clinical study. This could be performed on a single- or multi-institutional scale. By lowering the resources needed to effectively complete clinical trials,

ReSKU can expand data collection networks and lower the cost of multicenter trials.

While ReSKU has several advantages over traditional registries, it is not without its own associated challenges. For automated data extraction to be implemented, an upfront investment is required for programming both structured data capture into clinical notes and data extraction from the EHR. Collaboration between the urologist and data analysts at each site is essential to optimize the quality of the data being extracted and verify that the captured data points correspond to the desired data. Meanwhile, ReSKU is also limited in that long-term data collection may be limited by patients lost to follow-up. This is an inherent limitation given ReSKU's infrastructure and reflects the nature of registry studies in general. In addition, ReSKU was designed to integrate with Epic, given its marketshare and relatively widespread use in the United States. This integration relies on tools built specifically for the Epic EMR system. To broaden its applicability to other EMR systems, both within the United States as well as globally, these tools could be adapted for automated data extraction in non-Epic environments. However, in its current state, ReSKU is designed for use with Epic as a data source.

Finally, in the current ethos of the Affordable Care Act and accountability for quality and healthcare value, ReSKU may also be of practical value for providers. EHRs, as mandated by the American Reinvestment and Recovery Act in 2009, provide an opportunity to automate aspects of data entry that were previously accomplished only through labor-intensive chart review and manual data entry.⁶ Participation in the Physician Quality Reporting System (PQRS) through qualified clinical data registries (QCDR) can help providers maintain compliance with Medicare and Medicaid Services requirements.²² Responding to increasing pressure from the federal government and private insurers to report on quality and to meet PQRS reporting, the AUA has led efforts to launch nationwide clinical data registries with the AQUA Registry. Initial efforts for AQUA have focused on urologic oncology, highlighting the importance of physician participation in registries for tracking of quality metrics. Continued AQUA efforts and registries such as ReSKU can help to reduce the burden of participation in QCDR and keep urologists aligned with Centers for Medicare & Medicaid Services mandates.

Conclusions

As a longitudinal, observational prospective registry of patients with urinary stone disease that automates data extraction from EHRs, the ReSKU and Ureter represent an evolution from traditional registries. ReSKU provides a research infrastructure that decreases the burdens of quality metric reporting and lowers the barriers to performing high quality clinical research for urolithiasis.

Acknowledgments

This study was supported by the NIH K12-DK-07-006 (TC), P20-DK-100863 (MLS and TC), and R21-DK-109433 (TC).

Author Disclosure Statement

No competing financial interests exist.

References

1. Dreyer NA, Tunis SR, Berger M, et al. Why observational studies should be among the tools used in comparative effectiveness research. *Health Aff (Millwood)* 2010;29:1818–1825.
2. Hussein AA, Welty CJ, Broerring J, et al. National prostate cancer registries: Contemporary trends of prostate cancer in the United States. *Urol Pract* 2014;1:198–204.
3. Grant A, Ure J, Nicolson DJ, et al. Acceptability and perceived barriers and facilitators to creating a national research register to enable 'direct to patient' enrolment into research: The Scottish Health Research Register (SHARE). *BMC Health Serv Res* 2013;13:422.
4. Harris PA, Lane L, Biaggioni I. Clinical research subject recruitment: The Volunteer for Vanderbilt Research Program www.volunteer.mc.vanderbilt.edu/. *J Am Med Inform Assoc* 2005;12:608–613.
5. Couchoud C, Lassalle M, Cornet R, et al. Renal replacement therapy registries—Time for a structured data quality evaluation programme. *Nephrol Dial Transplant* 2013;28:2215–2220.
6. Gliklich RE, Dreyer NA, Leavy MB, et al. Standards in the conduct of registry studies for patient-centered outcomes research. Report to PCORI [Internet]. 2012 March. 58 p. Available at: www.pcori.org/sites/default/files/Standards-in-the-Conduct-of-Registry-Studies-for-Patient-Centered-Outcomes-Research1.pdf (Last accessed July 11, 2016).
7. Gliklich RE, Dreyer NA, Leavy MB. Interfacing registries with electronic health records. In: Gliklich RE, Dreyer NA, Leavy MB, eds. *Registries for Evaluating Patient Outcomes: A User's Guide* [Internet]. 2014 April. Available at: <http://effectivehealthcareahrq.gov/index.cfm/search-for-guides-reviews-and-reports/?productid=1897&pageaction=displayproduct> (Last accessed July 12, 2016).
8. Harris PA, Taylor R, Thielke R, et al. Research electronic data capture (REDCap)—A metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform* 2009;42:377–381.
9. SK&A. Physician Office Usage of Electronic Health Records Software: Market Insights Report [Internet]. SK&A (US). February 2016. Available at: www.skainfo.com/reports/physician-ehr-software-usage (Last accessed July 17, 2016).
10. Alavi CB, Massman JD III. Selecting an electronic data capture system. *Urol Pract* 2016;3:236–241.
11. Pearle MS, Goldfarb DS, Assimos DG, et al. Medical management of kidney stones: AUA guideline *J Urol* 2014;192:316–324.
12. Turk C, Petrik A, Sarica K, et al. EAU guidelines on diagnosis and conservative management of urolithiasis. *Eur Urol* 2016;69:468–474.
13. Turk C, Petrik A, Sarica K, et al. EAU guidelines on interventional treatment for urolithiasis. *Eur Urol* 2016;69:475–482.
14. Dripps RD. New classification of physical status. *Anesthesiology* 1963;24:111.
15. Dindo D, Demartines N, Clavien PA. Classification of surgical complications: A new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg* 2004;240:205–213.
16. Thomas K, Smith NC, Hegarty N, et al. The Guy's stone score—Grading the complexity of percutaneous nephrolithotomy procedures. *Urology* 2011;78:277–281.

17. Parks JH, Coe FL. Evidence for durable kidney stone prevention over several decades. *BJU Int* 2009;103:1238–1246.
18. Krambeck AE, Handa SE, Evan AP, et al. Profile of the brushite stone former. *J Urol* 2010;184:1367–1371.
19. de la Rosette J, Assimos D, Desai M, et al. The clinical research office of the endourological society percutaneous nephrolithotomy global study: Indications, complications, and outcomes in 5803 patients. *J Endourol* 2011;25:11–17.
20. Edvardsson VO, Goldfarb DS, Lieske JC, et al. Hereditary causes of kidney stones and chronic kidney disease. *Pediatr Nephrol* 2013;28:1923–1942.
21. Goldfarb DS, Fischer ME, Keich Y, et al. A twin study of genetic and dietary influences on nephrolithiasis: A report from the Vietnam Era Twin (VET) Registry. *Kidney Int* 2005;67:1053–1061.
22. Centers for Medicare & Medicaid Services [Internet]. [Place unknown]: the Centers for Medicare & Medicaid Services: 2016. Registry Reporting; 2016. Available at: www.cms.gov/Medicare/Quality-Initiatives-Patient-Assessment-Instruments/PQRS/Registry-Reporting.html (Last accessed July 11, 2016).

Address correspondence to:

Thomas Chi, MD

Department of Urology

University of California, San Francisco

400 Parnassus Avenue

Suite A610 Box 0330

San Francisco, CA 94143

E-mail: tom.chi@ucsf.edu

Abbreviations Used

AQUA = American Urological Association
Quality Registry
AUA = American Urological Association
EHR = electronic health record
IRB = institutional review board
PQRS = Physician Quality Reporting System
QCDR = qualified clinical data registries
REDCap = Research Electronic Data Capture
ReSKU = Registry for Stones of the Kidney and Ureter