

UC Irvine

UC Irvine Previously Published Works

Title

Whole genome sequencing data of multiple individuals of Pakistani descent

Permalink

<https://escholarship.org/uc/item/8jz1b64b>

Journal

Scientific Data, 7(1)

ISSN

2052-4463

Authors

Khan, Shahid Y
Ali, Muhammad
Lee, Mei-Chong W
et al.

Publication Date

2020

DOI

10.1038/s41597-020-00664-2

Peer reviewed



OPEN

DATA DESCRIPTOR

Whole genome sequencing data of multiple individuals of Pakistani descent

Shahid Y. Khan¹, Muhammad Ali¹, Mei-Chong W. Lee², Zhiwei Ma³, Pooja Biswas⁴, Asma A. Khan⁵, Muhammad Asif Naeem⁵, Saima Riazuddin⁶, Sheikh Riazuddin^{5,7,8}, Radha Ayyagari⁴, J. Fielding Hejtmancik³ & S. Amer Riazuddin¹✉

Here we report whole genome sequencing of four individuals (H3, H4, H5, and H6) from a family of Pakistani descent. Whole genome sequencing yielded 1084.92, 894.73, 1068.62, and 1005.77 million mapped reads corresponding to 162.73, 134.21, 160.29, and 150.86 Gb sequence data and 52.49x, 43.29x, 51.70x, and 48.66x average coverage for H3, H4, H5, and H6, respectively. We identified 3,529,659, 3,478,495, 3,407,895, and 3,426,862 variants in the genomes of H3, H4, H5, and H6, respectively, including 1,668,024 variants common in the four genomes. Further, we identified 42,422, 39,824, 28,599, and 35,206 novel variants in the genomes of H3, H4, H5, and H6, respectively. A major fraction of the variants identified in the four genomes reside within the intergenic regions of the genome. Single nucleotide polymorphism (SNP) genotype based comparative analysis with ethnic populations of 1000 Genomes database linked the ancestry of all four genomes with the South Asian populations, which was further supported by mitochondria based haplogroup analysis. In conclusion, we report whole genome sequencing of four individuals of Pakistani descent.

Background & Summary

The completion of Human Genome Project ignited several large scale efforts to characterize variations in the human genome, which led to a comprehensive catalog of the common variants including single-nucleotide polymorphisms (SNPs) and insertions/deletions (indels), across the entire human genome^{1,2}. Population-based genome reference datasets played an important role in elucidation of rare variants in specific populations^{3,4}. So far, comprehensive genome reference datasets have been reported for African, Japanese, Korean, and Chinese populations^{5–8}.

Advancements in next-generation sequencing technologies have impelled the development of a comprehensive catalog of genetic variants from different ethnic populations^{9–15}. The 1000 Genomes Project reports human genetic variation profiles from 26 ethnic populations, including one Pakistani (Punjabi), two Indian (Gujarati and Telugu), one Bangladeshi (Bengali), and one Sri Lankan (Tamil) population—all descendants of the Indian subcontinent¹⁵.

Additionally, independent groups have recently published two Indian and two Pakistani genomes with an overall 25–30× sequencing coverage^{16–19}. Recently, the GenomeAsia 100 K project reported genomes of 1,739 individuals, including 113 individuals of Pakistani origin (<https://browser.genomeasia100k.org>). We previously reported the whole genome sequencing of two Pakistani individuals²⁰. Here, we report whole genome sequencing of four individuals of Pakistani descent.

¹The Wilmer Eye Institute, Johns Hopkins University School of Medicine, Baltimore, MD, 21287, USA. ²Department of Computer Science, San José State University, San José, CA, 95192, USA. ³Ophthalmic Genetics and Visual Function Branch, National Eye Institute, National Institutes of Health, Bethesda, MD, 20892, USA. ⁴Shiley Eye Institute, University of California San Diego, La Jolla, CA, 92093, USA. ⁵National Centre of Excellence in Molecular Biology, University of the Punjab, Lahore, 53700, Pakistan. ⁶Department of Otorhinolaryngology-Head & Neck Surgery, University of Maryland School of Medicine, Baltimore, MD, 21201, USA. ⁷Allama Iqbal Medical College, University of Health Sciences, Lahore, 54550, Pakistan. ⁸Department of Molecular Biology, Shaheed Zulfiqar Ali Bhutto Medical University, Islamabad, 44080, Pakistan. ✉e-mail: riazuddin@jhmi.edu

Methods

Sample collection. The protocol for this study was approved by the Institutional Review Board of the Johns Hopkins University School of Medicine (Baltimore, MD), the National Centre of Excellence in Molecular Biology (Lahore, Pakistan), and the National Eye Institute (Bethesda, MD). The participating members provided informed written consent consistent with the tenets of the Declaration of Helsinki. A small aliquot (~10 ml) of a blood sample was collected from each individual and genomic DNA was extracted as previously described²⁰.

Library preparation and next-generation sequencing. Whole genome sequencing was performed using the Illumina HiSeq X10 (Illumina, San Diego, CA, USA). Briefly, 1.0–2.0 µg of fragmented gDNA was used to prepare paired-end libraries with the TruSeq DNA PCR-Free Library Preparation Kit for four samples (H3, H4, H5, and H6) according to the manufacturer's instructions (Illumina Inc., San Diego, CA). All four libraries were sequenced using Illumina HiSeq X10 in paired-end fashion (2 × 150 bp; Illumina Inc.). The base calls were assigned through Illumina Real-Time Analysis software (Ver. 1.17.20) and binary base call (BCL) files were converted to flat-file format (qseq.txt) using Illumina BCL Converter software (Ver. 1.9.4).

Bioinformatics analysis. Paired-end raw reads were aligned to the human reference genome (GRCh38.p13) using Burrows-Wheeler Aligner-MEM (BWA-MEM; Ver. 0.7.17-r1188) without ALT-aware mode²¹. The quality of the read alignments was examined using CollectAlignmentSummaryMetrics from Picard Tools (Ver. 2.19.0; <http://broadinstitute.github.io/picard>). The duplicate reads were removed from the mapped reads using MarkDuplicates from Picard Tools. The variants including SNPs and indels were called using the Genome Analysis Tool Kit (GATK; Ver. 4.0) best-practices^{12,22}. Briefly, the recalibration of base qualities of input reads was performed using GATK tools (BaseRecalibrator and ApplyBQSR). Subsequently, the SNPs, indels, and genotype of variants were identified using multiple tools i.e. HaplotypeCaller (in GVCF mode), GenotypeGVCFs, and VCFtools (Ver. 0.1.15)²³. Alignment metrics were generated using CollectAlignmentSummaryMetrics and CollectInsertSizeMetrics from Picard Tools. Genome-wide read coverage was generated using Bedtools (Ver. 2.26.0)²⁴.

Variant filtering and annotation. The variants identified through the GATK tool kit were further screened using the high-confidence regions characterized by Genome in a Bottle (GIAB) database²⁵. The variants aligned within the large segmental duplication regions of the human genome were discarded while variants mapped to the high-confidence regions of GIAB were used in downstream analyses including Venn diagram generation using VennPainter²⁶. **Note:** An allele (variant) with a minimum of 40% of the total reads mapped to reference allele is considered authentic. The filtered variants were annotated using clinEff (Ver. 1.0 h; <http://www.dnaminer.com/clineff.html>), a professional version of SNPEff²⁷, designed for the prediction of functional effects of variants.

Variant calling. The CNVnator (Ver. 0.4.1) algorithm was used for the identification of copy number variations (CNVs) with a bin size of 1,000 and 10,000²⁸. The GIAB filtered variants (SNPs) were imported into the CNV analysis pipeline for plotting the B-allele frequency (BAF) along the read depths for all deletion and duplication events.

Ancestry prediction. The ancestral roots of H3, H4, H5, and H6 were examined using the algorithms of Peddy (Ver. 0.3.5)²⁹. The study utilized the high-performance computational capabilities of the Biowulf Linux cluster at the National Institutes of Health, Bethesda, MD. PCA plots were created using SNPs genotype information obtained from VCF (variant call format) files (from whole genome sequencing data of H3, H4, H5, and H6) and comparing it with combined ethnic populations from the 1000 Genomes dataset.

In parallel, ancestral roots of H3, H4, H5, and H6 were examined through a comparative analysis with genomes of five different ethnic populations within the 1000 Genomes database. We randomly selected 96 samples from African, Ad Mixed American, East Asian, European, and South Asian populations for comparative analysis by the bcftools-isec algorithm. These variants from 1000 Genomes database and four genomes in VCF format were converted to BCF using bcftools (Ver. 1.8). The BCF files were converted to PLINK format using PLINK (Ver. 1.90b6.18) and PLINK algorithms were used to filter the variants to generate a list of markers in approximate linkage equilibrium for PCA analysis.

Haplogroup analysis. The mitochondrial sequencing reads were mapped to the revised Cambridge reference sequence (rCRS) of the human mitochondrial genome³⁰. Mitochondrial variants were identified using GATK (Ver. 4.0) best practices and are classified into phylogenetic clusters in the haplogroup analysis using HaploGrep 2 (Ver. 2.1.25)³¹, with Kulczynski measure and Phylotree (build 17). The Y chromosomal haplogroup analysis for all four genomes was performed using Yleaf³².

Data Records

The next-generation whole genome sequencing raw reads of H3, H4, H5, and H6 have been deposited in the NCBI Sequence Read Archive (SRA) with the accession number PRJNA596295³³. The chromosomal distribution of the variants identified in H3, H4, H5, and H6 genomes is available at figshare³⁴.

Technical Validation

The next-generation whole genome sequencing generated 1344.74, 1110.55, 1200.77, and 1142.35 million total reads for H3, H4, H5, and H6, respectively (Table 1)³³. Quality control (QC) examination of the sequencing reads revealed that >99% of the sequencing data yielded a PHRED score of 30 or above (PHRED score of 30 represents the probability of 0.001 that the base call is wrong). Subsequent to QC examination and the removal of PCR duplicates (~10–18% of reads were marked duplicates and subsequently removed in downstream analysis),




Fig. 1 Histogram depicting the sequence coverage of the genomes of H3, H4, H5, and H6. The x- and y-axes represent the genome coverage (sequencing depth) and genome fraction (%), respectively. The blue, yellow, green, and red colors represent the genomes of H3, H4, H5, and H6, respectively.

Sample ID	Total reads (10^6)	Total reads w/o PCR duplication (10^6)	Mapped reads (10^6)	Mapped reads (%)	Sequenced bases (Gb)	Mean depth (x)
H3	1344.74	1091.08	1084.92	99.44	162.73	52.49
H4	1110.55	899.761	894.73	99.44	134.21	43.29
H5	1200.77	1075.53	1068.62	99.36	160.29	51.70
H6	1142.35	1012.29	1005.77	99.36	150.86	48.66

Table 1. Summary of the next-generation whole genome sequencing data.

Genome Coverage	Fraction of H3 genome (%)	Fraction of H4 genome (%)	Fraction of H5 genome (%)	Fraction of H6 genome (%)
0x	5.44	6.17	5.48	6.15
1–10x	1.36	1.61	1.50	1.54
10–20x	2.31	2.30	2.80	1.98
20–30x	6.29	7.58	7.44	5.10
30–40x	9.18	30.6	10.95	15.97
40–50x	26.09	37.66	28.65	36.02
50–60x	32.50	11.56	31.04	25.76
60–70x	13.27	1.38	9.88	5.88
70–80x	2.31	0.27	1.21	0.67
80–90x	0.37	0.17	0.22	0.19
90–100x	0.16	0.12	0.14	0.13
>100x	0.72	0.58	0.69	0.61

Table 2. The genome coverage of whole genome sequencing data.

the majority of the reads (>99% of reads with a PHRED score ≥ 30) mapped to reference human genome (GRCh38.p13; Table 1). Mapping of the paired-end reads identified an estimated mean insert size of 390 bp in all four genomes. Majority of the total mapped reads showed paired-end alignment, resulting in 1084.92, 894.73, 1068.62, and 1005.77 million mapped reads corresponding to 162.73, 134.21, 160.29, and 150.86 Gb sequence data and 52.49x, 43.29x, 51.70x, and 48.66x average coverage for the genomes of H3, H4, H5, and H6, respectively (Table 1).

The evaluation of sequencing reads revealed that a significant fraction of the genomes of H3, H4, H5, and H6 exhibited 30–60x read coverage (Fig. 1 and Table 2). Importantly, 5–6% of the genomes of H3, H4, H5, and H6 were not captured, representing 0x read coverage while approximately, 1%, 2%, and 5% of four genomes exhibited 1–10x, 10–20x, and 20–30x read coverage, respectively (Fig. 1 and Table 2). A minor fraction i.e. <1% of the genomes of H3, H4, H5, and H6 exhibited 80–100x read coverage (Fig. 1 and Table 2).




Fig. 2 Venn diagram illustration of the overlapping variants characterized by the Genome in a Bottle (GIAB) database in the genomes H3, H4, H5, and H6. **(a)** Venn diagram illustrating all variants, **(b)** Venn diagram illustrating all known variants, and **(c)** Venn diagram illustrating all novel variants identified in the high-confidence regions characterized by the GIAB database in H3, H4, H5, and H6 genomes. **Note:** Yellow, green, blue, and pink represent variants in the genomes of H3, H4, H5, and H6, respectively, whereas darker shades represent common variants in these genomes.

Sequence analysis of the genome of H3 revealed a total of 3,529,659 variants including 3,035,369 SNPs and 494,290 indels. The SNPs were annotated against dbSNP (Ver. 150) that identified 7,553 novel variants (0.21% of the total variants) in the genome of H3³⁴. A total of 494,290 indels including 34,869 novel indels (7.05% of the total indels) were identified in the H3 genome³⁴.

Sequence analysis of the genome of H4 identified 3,478,495 total variants including 2,996,403 SNPs and 482,092 indels while annotation of the SNPs identified 6,631 novel SNPs (0.19% of the total variants) in the genome of H4³⁴. A total of 482,092 indels including 33,193 novel indels (6.88% of the total indels) were identified in the genome of H4³⁴. Sequence analysis of the genome of H5 identified 3,407,895 total variants including 2,983,279 SNPs and 424,616 indels while annotation of the SNPs identified 5,560 novel SNPs (0.16% of the total variants) in the genome of H5³⁴. A total of 424,616 indels including 23,039 novel indels (5.42% of the total indels) were identified in the genome of H5³⁴. Finally, sequence analysis of the genome of H6 identified 3,426,862 total




Fig. 3 Examining the ancestral roots of H3, H4, H5, and H6 through SNP-genotype comparative analysis with the combined ethnic populations of the 1000 Genomes dataset. The arrows point to (a) H3, (b) H4, (c) H5 and (d) H6 shown as red circles in the principal component analysis (PCA) plots with South Asian populations in principal components 1 and 3 (PC1 and PC3) and between South Asian, and European populations in principal component 2 (PC2). The x-axis represents PC1 while the y- and the z-axis represent PC2 and PC3, respectively. **Note:** South Asian: SAS; African: AFR; Ad Mixed American: AMR; and East Asian: EAS.

variants including 2,972,863 SNPs and 453,999 indels while annotation of the SNPs identified 6,703 novel SNPs (0.19% of the total variants) in the genome H6³⁴. A total of 453,999 indels including 28,503 novel indels (6.28% of the total indels) were identified in the genome of H6³⁴.

1000 Genomes populations	African			
Sample ID	common	unique	% common	% unique
H3	2758075	277294	90.865	9.135
H4	2716661	245307	90.664	9.336
H5	2744407	238872	91.993	8.007
H6	2699787	273076	90.814	9.186
1000 Genomes populations	South Asian			
Sample ID	common	unique	% common	% unique
H3	2878259	157110	94.824	5.176
H4	2829770	166633	94.439	5.561
H5	2823764	159515	94.653	5.347
H6	2809228	163635	94.496	5.504
1000 Genomes populations	East Asian			
Sample ID	common	unique	% common	% unique
H3	2700367	335002	88.963	11.366
H4	2654822	341581	88.600	11.400
H5	2687681	295598	90.092	9.908
H6	2644954	327909	88.970	11.030
1000 Genomes populations	European			
Sample ID	common	unique	% common	% unique
H3	2794183	213551	92.054	7.946
H4	2753320	208648	91.888	8.112
H5	2778513	206924	93.136	6.864
H6	2733120	208367	91.936	8.064
1000 Genomes populations	Admixed American			
Sample ID	common	unique	% common	% unique
H3	2789486	245883	91.899	8.101
H4	2748836	247567	91.738	8.262
H5	2774780	208499	93.011	6.989
H6	2729535	243328	91.815	8.185

Table 3. The variants present in H3, H4, H5, and H6 genomes overlapping with 1000 Genomes ethnic populations.

Importantly, we identified a total of 1,668,024 variants including 1,666,232 variants reported previously and 1,792 novel SNPs common in the four genomes (Fig. 2a–c). Altogether, the variants common in the four genomes constitute nearly half of the total variants identified in each genome.

We examined the putative effect of the variants based on their location in the genome (exon, intron, etc.), functional impact (high, moderate, and low), and classification (synonymous vs. non-synonymous), etc. The analysis revealed that intergenic regions harbor the majority of SNPs consistent with the GIAB high-confidence variants. Furthermore, in contrast to intergenic variants, fewer variants were identified in the exons, splice site, and untranslated regions (UTRs) of the genome. Furthermore, >3 K, >82 K, and >28 K variants present in all four genomes were predicted to exhibit a putative high, moderate, and low impact, respectively.

We used CNVnator, an algorithm to characterize copy number variations (CNVs), to examine structural variants in the genomes of H3, H4, H5, and H6. The analysis identified a total of 4,269 copy number variation regions (CNVRs) common in four genomes, covering 305.95 Mb (9.53%) of the reference human genome (GRCh38.p13).

Although H3, H4, H5, and H6 belong to the Punjabi ethnic group of Pakistani population suggesting a close ancestral relationship with South Asian populations, we sought of genomic evidence to confirm the ancestral roots of the four genomes. We compared the SNP genotypes of H3, H4, H5, and H6 to the combined population of the 1000 Genomes project by the Peddy algorithm. The analysis localized the all four genomes within South Asian populations in principal component 1 and 3 (PC1 and PC3) (Fig. 3a–d; arrows pointing to samples shown as red circles in PCA plots) and on the edge of the South Asian populations in principal component 2 (PC2) towards the European populations (Fig. 3a–d). The localization of H3, H4, H5, and H6 in PC2 suggests some ancestral link with European populations.

In parallel, we performed an additional comparative analysis of the four genomes with the genomes of five different ethnic populations in the 1000 Genomes database. The analysis identified >94% overlap of variants in the genomes of H3, H4, H5, and H6 with South Asian populations (Table 3) with a small number of variants in the genomes of H3 (157,110), H4 (166,633), H5 (159,515), and H6 (163,635) genomes not present in South Asian populations (Table 3). We identified >92% overlap of variants in the genomes of H3, H4, H5, and H6 with both European and Ad Mixed American populations (Table 3). Likewise, we identified >88% and >90% overlap of variants in the genomes of H3, H4, H5, and H6 with East Asian and African populations, respectively (Table 3). These data support the notion that H3, H4, H5, and H6 have a close ancestral relationship with South Asian populations (Fig. 4a–c).




Fig. 4 Investigating the ancestral origins of H3, H4, H5, and H6 by SNP-genotype comparative analysis with 96 random genomes from five different ethnic groups within the 1000 Genomes dataset. (a–c) Principal component analysis (PCA) plots illustrate three different angles of principal component # 1 (PC1), principal component # 2 (PC2), and principal component # 3 (PC3), respectively. The x-, y-, and z-axes depict the three largest components of the total variation in the percent of PC1, PC2, and PC3, respectively. The analysis illustrates an overlapping of four genomes with the South Asian population. **Note:** South Asian: SAS; African: AFR; Ad Mixed American: AMR; East Asian: EAS; and European: EUR.

To further confirm the results of SNP genotype based analysis, we performed mitochondria and Y chromosome based haplogroup analysis. The mitochondria genome analysis revealed M35b haplogroup in the H3 genome and M6 haplogroup in H4, H5, and H6 genomes. Both mitochondrial haplogroups (M35b and M6) have been mainly identified in South Asian populations^{35,36}. The Y chromosome analysis identified G1a1b2a haplogroup in H3 and H5 genomes, suggesting a Middle Eastern origin. Taken together, the mitochondria haplogroup based analyses support the results of the SNP genotype based analysis and strengthen the notion that H3, H4, H5, and H6 have a close ancestral relationship with South Asian populations.

In conclusion, we have completed next-generation based whole genome sequencing of four individuals from a family of Pakistani descent. Importantly, nearly 1% of the total variants identified in each of the four genomes are novel and have not been reported previously. To the best of our knowledge, this is the first report of whole genome sequencing of four individuals from a family.

Received: 15 January 2020; Accepted: 2 September 2020;

Published online: 13 October 2020

References

- Lander, E. S. *et al.* Initial sequencing and analysis of the human genome. *Nature* **409**, 860–921 (2001).
- Sachidanandam, R. *et al.* A map of human genome sequence variation containing 1.42 million single nucleotide polymorphisms. *Nature* **409**, 928–933 (2001).
- Southam, L. *et al.* Whole genome sequencing and imputation in isolated populations identify genetic associations with medically-relevant complex traits. *Nat. Commun.* **8**, 15606 (2017).
- Ebenesersdottir, S. S. *et al.* Ancient genomes from Iceland reveal the making of a human population. *Science* **360**, 1028–1032 (2018).
- Gurdasani, D. *et al.* The African Genome Variation Project shapes medical genetics in Africa. *Nature* **517**, 327–332 (2015).
- Nagai, A. *et al.* Overview of the BioBank Japan Project: Study design and profile. *J. Epidemiol.* **27**, S2–S8 (2017).
- Kim, J. *et al.* KoVariome: Korean National Standard Reference Variome database of whole genomes with comprehensive SNV, indel, CNV, and SV analyses. *Sci. Rep.* **8**, 5677 (2018).
- Gao, Y. *et al.* PGG.Han: the Han Chinese genome database and analysis platform. *Nucleic Acids Res.* **48**, D971–D976 (2020).
- Li, B. & Leal, S. M. Methods for detecting associations with rare variants for common diseases: application to analysis of sequence data. *Am. J. Hum. Genet.* **83**, 311–321 (2008).
- Tucker, T., Marra, M. & Friedman, J. M. Massively parallel sequencing: the next big thing in genetic medicine. *Am. J. Hum. Genet.* **85**, 142–154 (2009).
- Frazer, K. A., Murray, S. S., Schork, N. J. & Topol, E. J. Human genetic variation and its contribution to complex traits. *Nat. Rev. Genet.* **10**, 241–251 (2009).
- DePristo, M. A. *et al.* A framework for variation discovery and genotyping using next-generation DNA sequencing data. *Nat. Genet.* **43**, 491–498 (2011).
- Li, H. & Durbin, R. Inference of human population history from individual whole-genome sequences. *Nature* **475**, 493–496 (2011).
- Auton, A. *et al.* A global reference for human genetic variation. *Nature* **526**, 68–74 (2015).
- Sudmant, P. H. *et al.* An integrated map of structural variation in 2,504 human genomes. *Nature* **526**, 75–81 (2015).
- Patowary, A. *et al.* Systematic analysis and functional annotation of variations in the genome of an Indian individual. *Hum. Mutat.* **33**, 1133–1140 (2012).
- Gupta, R. *et al.* Sequencing and analysis of a South Asian-Indian personal genome. *BMC. Genomics* **13**, 440 (2012).
- Azim, M. K. *et al.* Complete genome sequencing and variant analysis of a Pakistani individual. *J. Hum. Genet.* **58**, 622–626 (2013).
- Ilyas, M. *et al.* Whole genome sequencing of an ethnic Pathan (Pakhtun) from the north-west of Pakistan. *BMC. Genomics* **16**, 172 (2015).
- Khan, S. Y. *et al.* Whole genome sequencing data for two individuals of Pakistani descent. *Sci. Data* **5**, 180174 (2018).
- Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics*. **25**, 1754–1760 (2009).
- Van der Auwera, G. A. *et al.* From FastQ data to high confidence variant calls: the Genome Analysis Toolkit best practices pipeline. *Curr. Protoc. Bioinformatics*. **43**, 11 (2013).
- Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics*. **27**, 2156–2158 (2011).
- Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics*. **26**, 841–842 (2010).
- Zook, J. M. *et al.* Integrating human sequence data sets provides a resource of benchmark SNP and indel genotype calls. *Nat. Biotechnol.* **32**, 246–251 (2014).
- Lin, G. *et al.* VennPainter: A Tool for the Comparison and Identification of Candidate Genes Based on Venn Diagrams. *PLoS. One*. **11**, e0154315 (2016).
- Cingolani, P. *et al.* A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaster* strain w1118; iso-2; iso-3. *Fly. (Austin.)* **6**, 80–92 (2012).
- Abyzov, A., Urban, A. E., Snyder, M. & Gerstein, M. CNVnator: an approach to discover, genotype, and characterize typical and atypical CNVs from family and population genome sequencing. *Genome Res.* **21**, 974–984 (2011).
- Pedersen, B. S. & Quinlan, A. R. Who's Who? Detecting and Resolving Sample Anomalies in Human DNA Sequencing Studies with Peddy. *Am. J. Hum. Genet.* **100**, 406–413 (2017).
- Andrews, R. M. *et al.* Reanalysis and revision of the Cambridge reference sequence for human mitochondrial DNA. *Nat. Genet.* **23**, 147 (1999).
- Weissensteiner, H. *et al.* HaploGrep 2: mitochondrial haplogroup classification in the era of high-throughput sequencing. *Nucleic Acids Res.* **44**, W58–W63 (2016).
- Ralf, A., Montiel, G. D., Zhong, K. & Kayser, M. Yleaf: Software for Human Y-Chromosomal Haplogroup Inference from Next-Generation Sequencing Data. *Mol. Biol. Evol.* **35**, 1291–1294 (2018).
- NCBI Sequence Read Archive <https://identifiers.org/ncbi/insdc.sra:SRP237990> (2020).
- Khan, S. Y. *et al.* Whole-genome sequencing of multiple individuals of Pakistani descent. *figshare* <https://doi.org/10.6084/m9.figshare.c.4787124> (2020).
- Metspalu, M. *et al.* Most of the extant mtDNA boundaries in south and southwest Asia were likely shaped during the initial settlement of Eurasia by anatomically modern humans. *BMC. Genet.* **5**, 26 (2004).
- Malyarchuk, B. A. *et al.* Mitochondrial DNA variability in Slovaks, with application to the Roma origin. *Ann. Hum. Genet.* **72**, 228–240 (2008).

Acknowledgements

We are thankful to individuals H3, H4, H5, and H6 for their participation in this study. This study was supported by the National Eye Institute Grant R01EY022714 (SAR).

Author contributions

S.Y.K. and S.A.R.: conceived and designed the experiments; M.W.L., S. R., S.R., R.A., J.F.H. and S.A.R.: contributed reagents, materials, and analytical tools; S.Y.K., M.A., M.W.L., Z.M., P.B. and M.A.N.: performed experiments; S.Y.K., M.A., M.W.L., Z.M., P.B., A.A.K., M.A.N., S. R., S.R., R.A., J.F.H. and S.A.R.: analyzed the data; S.Y.K., M.A., M.W.L., S. R., S.R., R.A., J.F.H. and S.A.R.: contributed to writing the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to S.A.R.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

The Creative Commons Public Domain Dedication waiver <http://creativecommons.org/publicdomain/zero/1.0/> applies to the metadata files associated with this article.

© The Author(s) 2020