

UC Irvine

UC Irvine Previously Published Works

Title

Crystallographic characterization of rare-earth cyanotriphenylborate complexes and the cyanoborates [NCBPh₃]¹⁻, [NCBPh₂Me]¹⁻, and [NCBPh₂(μ-O)BPh₂]¹⁻

Permalink

<https://escholarship.org/uc/item/8kv1m9gc>

Journal

Acta Crystallographica Section E: Crystallographic Communications, 77(Pt 8)

ISSN

2056-9890

Authors

Dumas, Megan T
White, Jessica RK
Ziller, Joseph W
[et al.](#)

Publication Date

2021-08-01

DOI

10.1107/s2056989021006861

Peer reviewed

Crystallographic characterization of rare-earth cyanotriphenylborate complexes and the cyanoborates $[\text{NCBPh}_3]^{1-}$, $[\text{NCBPh}_2\text{Me}]^{1-}$, and $[\text{NCBPh}_2(\mu\text{-O})\text{BPh}_2]^{1-}$

Megan T. Dumas, Jessica R. K. White, Joseph W. Ziller and William J. Evans*

Received 19 May 2021

Accepted 3 July 2021

Department of Chemistry, University of California, Irvine, CA 92697-2025, USA. *Correspondence e-mail: wevans@uci.edu

Edited by S. Parkin, University of Kentucky, USA

Keywords: lanthanide; rare earth; cyanide; cyanotriphenylborate; crystal structure.

CCDC references: 2094137; 2094136; 2094135; 2094134; 2094133


Supporting information: this article has supporting information at journals.iucr.org/e

The investigation of the coordination chemistry of rare-earth metal complexes with cyanide ligands led to the isolation and crystallographic characterization of the Ln^{III} cyanotriphenylborate complexes dichlorido(cyanotriphenylborato- κN)tetrakis(tetrahydrofuran- κO)lanthanide(III), $[\text{LnCl}_2(\text{C}_{19}\text{H}_{15}\text{BN})(\text{C}_4\text{H}_8\text{O})_4]$ [lanthanide (Ln) = dysprosium (Dy) and yttrium Y)] from reactions of LnCl_3 , KCN, and NaBPh_4 . Attempts to independently synthesize the tetraethylammonium salt of $(\text{NCBPh}_3)^-$ from BPh_3 and $[\text{NEt}_4][\text{CN}]$ in THF yielded crystals of the phenyl-substituted cyclic borate, tetraethylazanium 2,2,4,6-tetraphenyl-1,3,5,2 λ^4 ,4,6-trioxatriborinan-2-ide, $\text{C}_8\text{H}_{20}\text{N}^+\cdot\text{C}_{24}\text{H}_{20}\text{B}_3\text{O}_3^-$ or $[\text{NEt}_4][\text{B}_3(\mu\text{-O})_3(\text{C}_6\text{H}_5)_4]$. The mechanochemical reaction of BPh_3 and $[\text{NEt}_4][\text{CN}]$ without solvent produced crystals of tetraethylazanium cyanodiphenyl- λ^4 -boranyl diphenylborinate, $\text{C}_8\text{H}_{20}\text{N}^+\cdot\text{C}_{25}\text{H}_{20}\text{B}_2\text{NO}^-$ or $[\text{NEt}_4][\text{NCBPh}_2(\mu\text{-O})\text{BPh}_2]$. Reaction of BPh_3 and KCN in THF in the presence of 2.2.2-cryptand (crypt) led to a crystal of bis[(2.2.2-cryptand)potassium] 2,2,4,6-tetraphenyl-1,3,5,2 λ^4 ,4,6-trioxatriborinan-2-ide cyanomethyl-diphenylborate tetrahydrofuran disolvate, $2\text{C}_{18}\text{H}_{36}\text{KN}_2\text{O}_6^+\cdot\text{C}_{24}\text{H}_{20}\text{B}_3\text{O}_3^-\cdot\text{C}_{14}\text{H}_{13}\text{BN}^-\cdot 2\text{C}_4\text{H}_8\text{O}$ or $[\text{K}(\text{crypt})]_2[\text{B}_3(\mu\text{-O})_3(\text{C}_6\text{H}_5)_4][\text{NCBPh}_2\text{Me}]\cdot 2\text{THF}$. The $[\text{NCBPh}_2(\mu\text{-O})\text{BPh}_2]^{1-}$ and $(\text{NCBPh}_2\text{Me})^{1-}$ anions have not been structurally characterized previously. The structure of **1-Y** was refined as a two-component twin with occupancy factors 0.513 (1) and 0.487 (1). In **4**, one solvent molecule was disordered and included using multiple components with partial site-occupancy factors.


1. Chemical context

Attempts to make cationic rare-earth metal cyanide complexes of the type $[\text{Ln}(\text{CN})_2(\text{THF})_x][\text{BPh}_4]$ by combining LnCl_3 with sodium tetraphenylborate and potassium cyanide led to the isolation of the cyanotriphenylborate complexes $\text{LnCl}_2(\text{THF})_4(\text{NCBPh}_3)$, **1-Ln** ($\text{Ln} = \text{Dy}, \text{Y}$). Previously, transition-metal complexes of $(\text{NCBPh}_3)^{-}$ have been known to form from $\text{RhCl}(\text{PPh}_3)_3$, KCN, and BPh_3 (Pankowski *et al.*, 1996; Carlton *et al.*, 1998; Fernandes *et al.*, 2002) and from $[\text{Et}_4\text{N}][\text{Cr}(\text{CN})_6]$ and BPh_3 (Schelter *et al.*, 2005).

Efforts to independently synthesize the tetraethylammonium salt of the $(\text{NCBPh}_3)^{-}$ ligand generated a borate anion and two new cyanophenylborate anions that, to our knowledge, have not been structurally characterized. Specifically, the reaction of BPh_3 and $[\text{NEt}_4][\text{CN}]$ in THF led to crystals of the cyclic borate, $[\text{NEt}_4][\text{B}_3(\mu\text{-O})_3(\text{C}_6\text{H}_5)_4]$, **2**. When the analogous reaction was tried mechanochemically without solvent, the cyanoborate, $[\text{NEt}_4][\text{NCBPh}_2(\mu\text{-O})\text{BPh}_2]^{1-}$



O)BPh₂], **3**, was obtained. Reaction of BPh₃ with KCN in the presence of 2.2.2-cryptand (crypt) gave crystals of the double salt [K(crypt)]₂[B₃(μ-O)₃Ph₄][NCBPh₂Me], **4**. The cyanoborate anions in **3** and **4** have not been previously characterized by X-ray crystallography. The ChemDraw representations of **1-Ln** (Ln = Dy, Y), **2**, **3**, and **4** are depicted in the scheme below.



2. Structural commentary

The displacement ellipsoid plot of $\text{LnCl}_2(\text{THF})_4(\text{NCBPh}_3)$ **1-Ln** (Ln = Dy, Y) is depicted in Fig. 1 and the structural parameters are tabulated in Table 1. **1-Dy** and **1-Y** crystallize in the $\bar{P}1$ space group and are isomorphous. The geometry around the Ln^{III} ions is distorted pentagonal bipyramidal. The C1–N1 bond distances are 1.141 (3) and 1.144 (4) Å in **1-Dy** and **1-Y**, respectively. These distances are consistent with a C≡N triple bond (Allen *et al.*, 1987). The 178.7 (3) and 178.4 (3)° N1–C1–B1 bond angles in **1-Dy** and **1-Y**, respectively, are also consistent with a C≡N triple bond. The C1–N1–Ln1 angles are 163.92 (19) and 164.6 (3)° in **1-Dy** and **1-Y**, respectively. In comparison, the titanium complex, $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Ti}(\eta^2\text{-COR})(\text{NCBPh}_3)]$, has a C–N distance of 1.14 (2) Å, an N–C–B angle of 176.8 (16)°, and a C–N–Ti




Figure 1
Displacement ellipsoid plot of $\text{DyCl}_2(\text{THF})_4(\text{NCBPh}_3)$, **1-Dy**, drawn at the 30% probability level.

Table 1
Selected bond lengths and angles (Å, °) for **1-Dy** and **1-Y**.

	1-Dy	1-Y
Ln1–O1	2.3790 (18)	2.369 (2)
Ln1–O2	2.3838 (17)	2.370 (2)
Ln1–O3	2.4022 (16)	2.390 (2)
Ln1–O4	2.3932 (17)	2.382 (2)
Ln1–N1	2.431 (2)	2.420 (3)
Ln1–Cl1	2.5888 (6)	2.5803 (9)
Ln1–Cl2	2.5835 (6)	2.5730 (8)
N1–C1	1.141 (3)	1.144 (4)
B1–C1	1.621 (3)	1.629 (4)
B1–C18	1.625 (3)	1.630 (5)
B1–C24	1.632 (4)	1.626 (5)
B1–C30	1.641 (3)	1.642 (5)
C1–N1–Ln1	163.92 (19)	164.6 (3)
N1–C1–B1	178.7 (3)	178.4 (3)

angle of 169.1 (13)° (Pankowski *et al.*, 1996). The chromium complex $[\text{Et}_4\text{N}]_3[\text{Cr}(\text{NCBPh}_3)_6]$ has C–N distances of 1.127 (5), 1.100 (5), and 1.150 (5) Å, N–C–B angles of 178.9 (4), 176.8 (4), and 179.8 (5)°, and Cr–N–C angles of 174.8 (3), 175.4 (3), and 173.5 (3)° (Schelter *et al.*, 2005). The B–C(CN) and B–C(phenyl) distances in **1-Dy** and **1-Y**, respectively, are similar to those reported in transition-metal complexes with $[\text{NCBPh}_3]^{1-}$ ligands (Pankowski *et al.*, 1996; Fernandes *et al.*, 2002).

The displacement ellipsoid plots of $[\text{NEt}_4][\text{B}_3(\mu\text{-O})_3(\text{C}_6\text{H}_5)_4]$, **2**, $[\text{NEt}_4][\text{NCBPh}_2(\mu\text{-O})\text{BPh}_2]$, **3**, and $[\text{K}(\text{crypt})]_2[\text{B}_3(\mu\text{-O})_3(\text{C}_6\text{H}_5)_4][\text{NCBPh}_2\text{Me}]$, **4**, are shown below in Figs. 2, 3 and 4, respectively, and the structural parameters are tabulated in Table 1. Both **2** and **4** contain a non-planar anionic tetraphenylboroxinate ring as reported previously in the tetramethylammonium salt, $[\text{NMe}_4][\text{B}_3(\mu\text{-O})_3(\text{C}_6\text{H}_5)_4]$ (Kliegel *et al.*, 1986). While complex **2** is a tetraethylammonium salt of the tetraphenylboroxinate anion, complex **4** is a double salt with the second anion being




Figure 2
Displacement ellipsoid plot of **2**, $[\text{NEt}_4][\text{B}_3(\mu\text{-O})_3(\text{C}_6\text{H}_5)_4]$, drawn at the 30% probability level. The disorder in the $[\text{NEt}_4]^+$ cation is omitted for clarity.




Figure 3
Displacement ellipsoid plot of **3**, [NET₄][NCBPh₂(μ -O)BPh₂], drawn at the 50% probability level.

[NCBPh₂Me]¹⁻. There are no crystallographically characterized examples of this anion in the literature to our knowledge. Complex **3** contains a [Ph₂B(μ -O)BPh₂CN]¹⁻ anion, differing both from the cyclic B₃O₃ phenylboroxinate anions found in **2** and **4** and from the [NCBPh₂Me]¹⁻ anion in **4**. We found no crystallographically characterized examples of this anion in the literature.

The O–B–O and B–O–B angles range from 110.4 (1) to 122.5 (2)° for **2** and range from 109.9 (5) to 123.3 (5)° for **4**, both ranges are similar to those in [NMe₄][B₃(μ -O)₃(C₆H₅)₄]. The 1.497 (2) and 1.504 (2) Å B–O bonds involving four-coordinate B1 in **2** are longer than the other four B–O bond distances involving three-coordinate boron, which range from 1.333 (2) Å to 1.399 (2) Å for **2**. Similar distances are found in **4**.

Interestingly, though the B–C(CN) and B–C(phenyl) distances for **4** are similar to those of the [NCBPh₃]¹⁻ anion in **1-Ln**, the N–C–B angle of 171.6 (9)° in **4** is less linear than the N–C–B angles in **1-Ln** and in **3**, 178 and 177.10 (13)°, respectively.




Figure 4
Displacement ellipsoid plot of **4**, [K(crypt)]₂[B₃(μ -O)₃(C₆H₅)₄][NCBPh₂Me], drawn at the 50% probability level.

Both the C–N and B–C(CN) lengths in **3** and **4** are similar to those in **1-Ln**. The 1.488 (10) Å B₄–C(Me) distance in **4** is much shorter than the reported B–C(Me) distances in [BMe₄]¹⁻ [1.639 (2) to 1.648 (2) Å] (Zhu & Kochi, 1999), [BPh₃Me]¹⁻ [1.653 (2) Å] (Zhu & Kochi, 1999), and [FcB(Mes^F)(Me)(CN)]¹⁻ [1.628 (5) Å] (Broomsgrrove *et al.*, 2010). The B₄–C(Me) distance is also shorter than the 1.636 (2) and 1.614 (10) Å B–C(CN) distances in **3** and **4**, respectively. The 1.633 (11) and 1.648 (11) Å B₄–C(phenyl) distances are similar to the other B–C(phenyl) distances in the structures reported here.

3. Supramolecular features

There are no notable supramolecular features in **1-Ln**, **2**, **3**, or **4**.

4. Database survey

The Cambridge Structural Database contains 14 known structures of transition metal complexes with the [NCBPh₃]¹⁻ ligand found in **1-Ln** or related [NCB(C₆F₅)₃]¹⁻ ligands, including titanium (Pankowski *et al.*, 1996, CSD refcode: TEXLEN), chromium (Schelter *et al.*, 2005, CSD Refcode: XAKCUI), iron (Vei *et al.*, 2003, CSD Refcode: TAGKUI), nickel (Brunkan *et al.*, 2004, CSD Refcode: AVOKAX), copper (Nazarenko *et al.*, 1996, CSD Refcode: REYHEI) zirconium (Zhou *et al.*, 2001, CSD Refcode: YEQZYZ), and rhodium (Fernandes *et al.*, 2002, CSD Refcode: XUTDIZ; Cornock *et al.*, 1977, CSD Refcode: CBORRH). There is also a crystallographically-characterized cerium cyanotriphenylborate complex, [Ce(L_{OEt})₂(NCBPh₃)₂] (L_{OEt}¹⁻ = [Co(η^5 -C₅H₅){P(O)(OEt)₂]₃]¹⁻) (Au-Yeung *et al.*, 2016, CSD Refcode: EYAZOV).

Other crystallographically characterized tetraphenylboroxinates of the type found in **2** that are reported in the Cambridge Structural Database include [Me₃NCH₂CH₂OH][B₃(μ -O)₃(C₆H₅)₄] (Beckett *et al.*, 2006, CSD Refcode: ICUWAF), [NET₃H][B₃(μ -O)₃(C₆H₅)₄] (Kratzert & Krossing, 2018, CSD Refcode: HERJUM01), [(^tBu₃PAu)₄P][B₃(μ -O)₃(C₆H₅)₄] (Zeller *et al.*, 1993, CSD Refcode: PEVZOF), [C₆H₁₁NMe₃][B₃(μ -O)₃(C₆H₅)₄] (Beckett *et al.*, 2018, CSD Refcode: VEKVIT), [Ph₂B{OCH₂CH₂N(Me)(CH₂)_n}]₂[B₃(μ -O)₃(C₆H₅)₄] (*n* = 4, 5) (Beckett *et al.*, 2010, CSD Refcode: VUTGUN), and [(η^5 -C₅H₅)Ni(η^6 -C₆H₆)Ni(η^5 -C₅H₅)]₂[B₃(μ -O)₃(C₆F₅)₅] (Priego *et al.*, 2000, CSD Refcode: MEKLAP).

There are no crystallographically characterized examples of the [Ph₂B(μ -O)BPh₂CN]¹⁻ and [NCBPh₂Me]¹⁻ anions found in **3** and **4**, respectively, in the literature.

5. Synthesis and crystallization

DyCl₂(THF)₄(NCBPh₃), **1-Dy**. In an argon-filled glovebox, KCN (42 mg, 0.642 mmol) was added to a stirred slurry of DyCl₃ (75 mg, 0.279 mmol) in THF (10 mL). NaBPh₄ (96 mg,

Table 2
Experimental details.

	1-Dy	1-Y	2	3	4
Crystal data					
Chemical formula	[DyCl ₂ (C ₁₉ H ₁₅ BN)- (C ₄ H ₈ O) ₄]	[YCl ₂ (C ₁₉ H ₁₅ BN)- (C ₄ H ₈ O) ₄]	C ₈ H ₂₀ N ⁺ ·C ₂₄ H ₂₀ B ₃ O ₃ ⁻	C ₈ H ₂₀ N ⁺ · C ₂₅ H ₂₀ B ₂ NO ⁻	2C ₁₈ H ₃₆ KN ₂ O ₆ ⁺ · C ₂₄ H ₂₀ B ₃ O ₃ ⁻ · C ₁₄ H ₁₅ BN ⁻ · 2C ₄ H ₈ O
<i>M_r</i>	789.94	716.35	519.08	502.29	1570.27
Crystal system, space group	Triclinic, <i>P</i> $\bar{1}$	Triclinic, <i>P</i> $\bar{1}$	Monoclinic, <i>P</i> 2 ₁ / <i>n</i>	Monoclinic, <i>P</i> 2 ₁ / <i>n</i>	Orthorhombic, <i>Pca</i> 2 ₁
Temperature (K)	158	173	133	88	88
<i>a</i> , <i>b</i> , <i>c</i> (Å)	12.0043 (17), 12.5261 (17), 13.4913 (19)	12.0404 (9), 12.5428 (9), 13.4654 (10)	9.7245 (7), 18.3021 (13), 16.7716 (12)	11.0269 (8), 13.9387 (11), 18.8488 (14)	27.193 (2), 14.5520 (11), 21.2218 (16)
α , β , γ (°)	114.7161 (13), 101.2230 (16), 94.2256 (16)	114.6967 (8), 101.3142 (9), 94.3882 (9)	90, 101.5456 (9), 90	90, 100.6357 (10), 90	90, 90, 90
<i>V</i> (Å ³)	1780.0 (4)	1782.7 (2)	2924.6 (4)	2847.3 (4)	8397.7 (11)
<i>Z</i>	2	2	4	4	4
Radiation type	Mo <i>K</i> α	Mo <i>K</i> α	Mo <i>K</i> α	Mo <i>K</i> α	Mo <i>K</i> α
μ (mm ⁻¹)	2.29	1.82	0.07	0.07	0.18
Crystal size (mm)	0.39 × 0.33 × 0.26	0.32 × 0.20 × 0.13	0.37 × 0.33 × 0.23	0.28 × 0.27 × 0.22	0.45 × 0.33 × 0.28
Data collection					
Diffraction	Bruker SMART APEXII CCD	Bruker SMART APEXII CCD	Bruker SMART APEXII CCD	Bruker SMART APEXII CCD	Bruker SMART APEXII CCD
Absorption correction	Numerical (<i>SADABS</i> ; Krause <i>et al.</i> , 2015)	Multi-scan (<i>TWINABS</i> ; Sheldrick, 2012)	Multi-scan (<i>SADABS</i> ; Krause <i>et al.</i> , 2015)	Multi-scan (<i>SADABS</i> ; Krause <i>et al.</i> , 2015)	Multi-scan (<i>SADABS</i> ; Krause <i>et al.</i> , 2015)
<i>T</i> _{min} – <i>T</i> _{max}	0.558, 0.696	0.576, 0.746	0.834, 0.862	0.715, 0.746	0.825, 0.862
No. of measured, independent and observed [<i>I</i> > 2 σ (<i>I</i>)] reflections	21440, 8460, 7562	7988, 7988, 6188	31741, 5959, 5030	34768, 7242, 5500	85621, 15958, 14236
<i>R</i> _{int}	0.019	–	0.029	0.042	0.038
(<i>sin</i> θ / λ) _{max} (Å ⁻¹)	0.680	0.650	0.625	0.685	0.610
Refinement					
<i>R</i> [<i>F</i> ² > 2 σ (<i>F</i> ²)], <i>wR</i> (<i>F</i> ²), <i>S</i>	0.024, 0.061, 1.04	0.047, 0.102, 1.04	0.064, 0.179, 1.02	0.048, 0.119, 1.04	0.074, 0.209, 1.06
No. of reflections	8460	7988	5959	7242	15958
No. of parameters	406	425	442	347	988
No. of restraints	0	0	0	0	1
H-atom treatment	H-atom parameters constrained	H-atom parameters constrained	H-atom parameters constrained	H-atom parameters constrained	H-atom parameters constrained
$\Delta\rho_{\max}$, $\Delta\rho_{\min}$ (e Å ⁻³)	1.34, -0.89	0.64, -0.81	1.06, -0.47	0.41, -0.24	0.78, -0.34

Computer programs: *APEX2* (Bruker, 2014), *SAINTE* (Bruker, 2013), *SHELXT2014/4* (Sheldrick, 2015a), *SHELXL2014/7* (Sheldrick, 2015b), and *SHELXTL* (Sheldrick, 2008).

0.279 mmol) was added to the stirred slurry. The cloudy white solution was stirred overnight. The volatiles were removed under vacuum. The product was extracted into THF (10 mL) and centrifuged to remove white solids. The clear colorless solution had its volatiles removed under vacuum. The product was isolated as a colorless powder. Colorless crystals of DyCl₂(THF)₄(NCBPh₃), **1-Dy**, suitable for X-ray diffraction, were isolated from a vapor diffusion of hexane into a concentrated THF solution at room temperature after 6 d.

YCl₂(THF)₄(NCBPh₃), 1-Y. In an argon-filled glovebox, KCN (34 mg, 0.522 mmol) was stirred in THF (10 mL) to form a cloudy white slurry. After 4 h, YCl₃ (51 mg, 0.260 mmol) was tapped into the stirred white slurry. After 5 min, a solution of NaBPh₄ (89 mg, 0.260 mmol) in THF (8 mL) was added to the stirred slurry. The cloudy white slurry was stirred overnight. The white slurry was centrifuged. The clear, colorless supernatant was collected, and the centrifuge pellet was washed with THF (5 mL), and the wash was combined with the

supernatant. The colorless solution had its volatiles removed under vacuum. The product was isolated as a colorless solid (107 mg). X-ray quality crystals were isolated from a vapor diffusion of hexane into a concentrated THF solution at room temperature after 10 d.

[NEt₄][B₃(μ -O)₃(C₆H₅)₄], 2. In an argon-filled glovebox, BPh₃ (78 mg, 0.320 mmol) was tapped into a stirred slurry of Et₄NCN (50 mg, 0.320 mmol) in THF (5 mL). THF (10 mL) was added to the slurry, and the solution was heated lightly on a hot plate to encourage the Et₄NCN to dissolve. After the sample was heated lightly for about an hour, most of the white solids had dissolved. The solution was allowed to stir at room temperature overnight. The volatiles were removed under vacuum and the white solids were washed with toluene (2 mL) twice. The thick tacky colorless solids were extracted into THF. Colorless X-ray quality crystals of [NEt₄][B₃(μ -O)₃(C₆H₅)₄], **2**, were grown from a slow evaporation of a THF solution at room temperature.

[NEt₄][NCBPh₂(μ -O)BPh₂], 3. In an argon-filled glovebox, BPh₃ and NEt₄CN were added to a BMT-20-S tube drive along with 40 steel balls (6 mm). The reaction mixture was ball milled together for 40 minutes using an Ultra-Turrax Tube Drive at the maximum speed setting. After this time, the colorless solids were extracted into toluene and THF. The volatiles were removed under vacuum. X-ray quality colorless crystals of [NEt₄][NCBPh₂(μ -O)BPh₂], **3**, were grown from a slow evaporation of a concentrated THF solution at 258 K after a few days.

[K(crypt)]₂[B₃(μ -O)₃(C₆H₅)₄][NCBPh₂Me], 4. In an argon-filled glovebox, 2,2,2-cryptand (156 mg, 0.413 mmol) and KCN (30 mg, 0.460 mmol) were stirred in THF (10 mL) for 2 h, which allowed most of the white solids to dissolve. A solution of BPh₃ (100 mg, 0.413 mmol) in THF (5 mL) was added to this mixture and the combination was allowed to stir for 3 d. The volatiles were removed under vacuum from the clear and colorless solution. The sample was extracted into THF (20 mL) and volatiles were removed under vacuum. The product was isolated as a colorless solid (423 mg). Colorless X-ray quality crystals of [K(crypt)]₂[B₃(μ -O)₃(C₆H₅)₄][NCBPh₂Me], **4**, were obtained from a vapor diffusion of pentane into a concentrated THF solution at room temperature.

6. Refinement

General Structure Solution and Refinement. The analytical scattering factors (Wilson, 1992) for neutral atoms were used throughout the analysis. Hydrogen atoms were included using a riding model. **DyCl₂(THF)₄(NCBPh₃), 1-Dy:** Data were collected using a 15 sec/frame scan time. There were no systematic absences nor any diffraction symmetry other than the Friedel condition. Atom C5 was disordered and included using multiple components with partial site-occupancy factors. **YCl₂(THF)₄(NCBPh₃), 1-Y:** Data were collected using a 30 sec/frame scan time. There were no systematic absences nor any diffraction symmetry other than the Friedel condition. Disordered atoms were included using multiple components with partial site-occupancy-factors. The structure was refined as a two-component twin with occupancy factors 0.513 (1) and 0.487 (1). **[NEt₄][B₃(μ -O)₃(C₆H₅)₄], 2:** Data were collected using a 20 sec/frame scan time. The tetraethylammonium ion was fully disordered. The disordered atoms were included using multiple components with partial site-occupancy factors. **[NEt₄][NCBPh₂(μ -O)BPh₂], 3:** Data were collected using a 30 sec/frame scan time. **[K(crypt)]₂[B₃(μ -O)₃(C₆H₅)₄][NCBPh₂Me], 4:** Data were collected using a 60 sec/frame scan time. There were two molecules of tetrahydrofuran solvent present. One solvent molecule was disordered and included using multiple components with partial site-occupancy factors. Crystal data, data collection and structure refinement details are summarized in Table 2.

Acknowledgements

We thank Dr Michael Wojnar for assistance with X-ray crystallography.

Funding information

Funding for this research was provided by: National Science Foundation (grant No. CHE-1855328).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, **2**, S1–S19.
- Au-Yeung, K. C., So, Y. M., Sung, H. H. Y., Williams, I. D. & Leung, W. H. (2016). *Dalton Trans.* **45**, 18163–18170.
- Beckett, M. A., Bennett, E. L., Horton, P. N. & Hursthouse, M. B. (2010). *J. Organomet. Chem.* **695**, 1080–1083.
- Beckett, M. A., Coles, S. J., Horton, P. N., Jones, C. L., Marshall, E. V. & Perry, T. (2018). *J. Organomet. Chem.* **865**, 72–79.
- Beckett, M. A., Coles, S. J., Light, M. E., Fischer, L., Stiefvater-Thomas, B. M. & Varma, K. S. (2006). *Polyhedron*, **25**, 1011–1016.
- Broomsgrave, A. E. J., Addy, D. A., Di Paolo, A., Morgan, I. R., Bresner, C., Chislett, V., Fallis, I. A., Thompson, A. L., Vidovic, D. & Aldridge, S. (2010). *Inorg. Chem.* **49**, 157–173.
- Bruker (2013). *SAINT*. Bruker AXS, Inc., Madison, Wisconsin, USA.
- Bruker (2014). *APEX2*. Bruker AXS, Inc., Madison, Wisconsin, USA.
- Brunkan, N. M., Brestensky, D. M. & Jones, W. D. (2004). *J. Am. Chem. Soc.* **126**, 3627–3641.
- Carlton, L., Weber, R. & Leventis, D. C. (1998). *Inorg. Chem.* **37**, 1264–1271.
- Cornock, M. C., Robertson, D. R., Stephenson, T. A., Jones, C. L., Milburn, G. H. W. & Sawyer, L. (1977). *J. Organomet. Chem.* **135**, C50–C52.
- Fernandes, M. A., Cîrcu, V., Weber, R., Varnali, T. & Carlton, L. (2002). *J. Chem. Crystallogr.* **32**, 273–278.
- Kliegel, W., Motzkus, H.-W., Rettig, S. J. & Trotter, J. (1986). *Can. J. Chem.* **64**, 507–513.
- Kratzert, D. & Krossing, I. (2018). *J. Appl. Cryst.* **51**, 928–934.
- Krause, L., Herbst-Irmer, R., Sheldrick, G. M. & Stalke, D. (2015). *J. Appl. Cryst.* **48**, 3–10.
- Nazarenko, A. Y., Belsky, V. K. & Streltsova, N. R. (1996). *Zh. Neorg. Khim.* **41**, 1119–1123.
- Pankowski, M., Cabestaing, C. & Jaouen, G. (1996). *J. Organomet. Chem.* **516**, 11–16.
- Priego, J. L., Doerrer, L. H., Rees, L. H. & Green, M. L. H. (2000). *Chem. Commun.* pp. 779–780.
- Schelter, E. J., Shatruk, M., Heintz, R. A., Galán-Mascarós, J. R. & Dunbar, K. R. (2005). *Chem. Commun.* pp. 1417–1419.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Sheldrick, G. M. (2012). *TWINABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2015a). *Acta Cryst. A* **71**, 3–8.
- Sheldrick, G. M. (2015b). *Acta Cryst. C* **71**, 3–8.
- Vei, I. C., Pascu, S. I., Green, M. L. H., Green, J. C., Schilling, R. E., Anderson, G. D. W. & Rees, L. H. (2003). *Dalton Trans.* pp. 2550–2557.
- Wilson, A. J. C. (1992). *International Tables for Crystallography*, Vol. C. Dordrecht: Kluwer Academic Publishers.
- Zeller, E., Beruda, H. & Schmidbaur, H. (1993). *Chem. Ber.* **126**, 2033–2036.
- Zhou, J., Lancaster, S. J., Walker, D. A., Beck, S., Thornton-Pett, M. & Bochmann, M. (2001). *J. Am. Chem. Soc.* **123**, 223–237.
- Zhu, D. & Kochi, J. K. (1999). *Organometallics*, **18**, 161–172.

supporting information

Acta Cryst. (2021). E77, 799-803 [https://doi.org/10.1107/S2056989021006861]

Crystallographic characterization of rare-earth cyanotriphenylborate complexes and the cyanoborates [NCBPh₃]¹⁻, [NCBPh₂Me]¹⁻, and [NCBPh₂(μ-O)BPh₂]¹⁻

Megan T. Dumas, Jessica R. K. White, Joseph W. Ziller and William J. Evans

Computing details

For all structures, data collection: *APEX2* (Bruker, 2014); cell refinement: *SAINTE* (Bruker, 2013); data reduction: *SAINTE* (Bruker, 2013); program(s) used to solve structure: *SHELXT2014/4* (Sheldrick, 2015a); program(s) used to refine structure: *SHELXL2014/7* (Sheldrick, 2015); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

Dichlorido(cyanotriphenylborato-κN)tetrakis(tetrahydrofuran-κO)dysprosium(III) (1-Dy)

Crystal data

[DyCl₂(C₁₉H₁₅BN)(C₄H₈O)₄]

M_r = 789.94

Triclinic, *P*1

a = 12.0043 (17) Å

b = 12.5261 (17) Å

c = 13.4913 (19) Å

α = 114.7161 (13)°

β = 101.2230 (16)°

γ = 94.2256 (16)°

V = 1780.0 (4) Å³

Z = 2

F(000) = 802

D_x = 1.474 Mg m⁻³

Mo *K*α radiation, λ = 0.71073 Å

Cell parameters from 9908 reflections

θ = 2.3–28.8°

μ = 2.29 mm⁻¹

T = 158 K

Irregular, colorless

0.39 × 0.33 × 0.26 mm

Data collection

Bruker SMART APEXII CCD
diffractometer

Radiation source: fine-focus sealed tube

φ and ω scans

Absorption correction: numerical
(SADABS; Krause et al., 2015)

T_{min} = 0.558, *T_{max}* = 0.696

21440 measured reflections

8460 independent reflections

7562 reflections with *I* > 2σ(*I*)

R_{int} = 0.019

θ_{max} = 28.9°, θ_{min} = 1.7°

h = -16→16

k = -16→16

l = -18→18

Refinement

Refinement on *F*²

Least-squares matrix: full

R[*F*² > 2σ(*F*²)] = 0.024

wR(*F*²) = 0.061

S = 1.04

8460 reflections

406 parameters

0 restraints

Primary atom site location: dual space

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

w = 1/[σ²(*F_o*²) + (0.0251*P*)² + 2.2673*P*]

where *P* = (*F_o*² + 2*F_c*²)/3

$$(\Delta/\sigma)_{\max} = 0.002$$

$$\Delta\rho_{\max} = 1.34 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.89 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. A colorless crystal of approximate dimensions 0.255 x 0.332 x 0.391 mm was mounted in a cryoloop and transferred to a Bruker SMART APEX II diffractometer. The APEX2 program package was used to determine the unit-cell parameters and for data collection (15 sec/frame scan time for a sphere of diffraction data). The raw frame data was processed using SAINT and SADABS to yield the reflection data file. Subsequent calculations were carried out using the SHELXTL program. There were no systematic absences nor any diffraction symmetry other than the Friedel condition. The centrosymmetric triclinic space group P-1 was assigned and later determined to be correct.

The structure was solved by dual space methods and refined on F2 by full-matrix least-squares techniques. The analytical scattering factors for neutral atoms were used throughout the analysis. Hydrogen atoms were included using a riding model. Atom C(5) was disordered and included using multiple components with partial site-occupancy-factors.

Least-squares analysis yielded $wR2 = 0.0609$ and $Goof = 1.042$ for 406 variables refined against 8460 data (0.74Å), $R1 = 0.0239$ for those 7562 data with $I > 2.0\sigma(I)$.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Dy1	0.24691 (2)	0.24994 (2)	0.49865 (2)	0.01889 (4)	
C11	0.23261 (5)	0.07841 (6)	0.55751 (5)	0.03227 (13)	
C12	0.25400 (5)	0.40466 (5)	0.41894 (5)	0.03005 (12)	
O1	0.45097 (15)	0.26894 (16)	0.55203 (16)	0.0342 (4)	
O2	0.30435 (16)	0.12131 (15)	0.33601 (14)	0.0316 (4)	
O3	0.07375 (14)	0.14589 (15)	0.35225 (14)	0.0287 (4)	
O4	0.08323 (15)	0.30171 (15)	0.57396 (14)	0.0288 (4)	
N1	0.31345 (17)	0.42077 (18)	0.68261 (17)	0.0284 (4)	
B1	0.3578 (2)	0.6482 (2)	0.8573 (2)	0.0236 (5)	
C1	0.33280 (19)	0.5144 (2)	0.75535 (19)	0.0247 (5)	
C2	0.5339 (2)	0.3715 (3)	0.5732 (3)	0.0376 (6)	
H2A	0.5001	0.4452	0.5982	0.045*	
H2B	0.5601	0.3596	0.5047	0.045*	
C3	0.6319 (3)	0.3779 (3)	0.6651 (4)	0.0676 (12)	
H3A	0.6208	0.4295	0.7402	0.081*	
H3B	0.7064	0.4100	0.6582	0.081*	
C4	0.6293 (3)	0.2511 (3)	0.6487 (3)	0.0465 (7)	
H4A	0.6249	0.2447	0.7186	0.056*	0.65
H4B	0.6985	0.2207	0.6256	0.056*	0.65
H4C	0.6521	0.2006	0.5793	0.056*	0.35
H4D	0.6784	0.2464	0.7143	0.056*	0.35
C5	0.5180 (4)	0.1813 (4)	0.5532 (5)	0.0455 (11)	0.65
H5A	0.5371	0.1397	0.4797	0.055*	0.65
H5B	0.4761	0.1217	0.5693	0.055*	0.65
C5B	0.5075 (7)	0.2203 (9)	0.6388 (9)	0.046 (2)	0.35
H5B1	0.4880	0.2609	0.7122	0.055*	0.35
H5B2	0.4831	0.1330	0.6084	0.055*	0.35

C6	0.2852 (4)	-0.0100 (3)	0.2908 (3)	0.0578 (9)
H6A	0.2018	-0.0433	0.2635	0.069*
H6B	0.3200	-0.0342	0.3493	0.069*
C7	0.3429 (3)	-0.0522 (3)	0.1953 (3)	0.0465 (7)
H7A	0.3031	-0.1318	0.1356	0.056*
H7B	0.4249	-0.0558	0.2218	0.056*
C8	0.3317 (3)	0.0430 (3)	0.1538 (3)	0.0513 (8)
H8A	0.3889	0.0440	0.1102	0.062*
H8B	0.2532	0.0309	0.1066	0.062*
C9	0.3554 (3)	0.1549 (3)	0.2616 (2)	0.0432 (7)
H9A	0.4394	0.1836	0.2935	0.052*
H9B	0.3195	0.2185	0.2495	0.052*
C10	-0.0179 (2)	0.0684 (2)	0.3591 (2)	0.0378 (6)
H10A	-0.0791	0.1127	0.3866	0.045*
H10B	0.0125	0.0328	0.4097	0.045*
C11	-0.0631 (2)	-0.0268 (2)	0.2381 (2)	0.0406 (6)
H11A	-0.0149	-0.0899	0.2198	0.049*
H11B	-0.1440	-0.0637	0.2240	0.049*
C12	-0.0541 (3)	0.0436 (3)	0.1712 (3)	0.0540 (9)
H12A	-0.0318	-0.0047	0.1009	0.065*
H12B	-0.1288	0.0678	0.1515	0.065*
C13	0.0369 (3)	0.1510 (3)	0.2465 (2)	0.0454 (7)
H13A	0.1028	0.1504	0.2122	0.054*
H13B	0.0053	0.2249	0.2580	0.054*
C14	0.0192 (3)	0.3867 (3)	0.5532 (3)	0.0421 (7)
H14A	-0.0553	0.3449	0.4974	0.051*
H14B	0.0636	0.4307	0.5232	0.051*
C15	-0.0006 (4)	0.4690 (5)	0.6599 (3)	0.0839 (15)
H15A	-0.0720	0.5015	0.6480	0.101*
H15B	0.0652	0.5359	0.7031	0.101*
C16	-0.0113 (4)	0.3935 (5)	0.7179 (4)	0.0923 (17)
H16A	-0.0926	0.3553	0.6992	0.111*
H16B	0.0152	0.4419	0.8005	0.111*
C17	0.0639 (3)	0.2994 (3)	0.6771 (3)	0.0493 (8)
H17A	0.1378	0.3196	0.7340	0.059*
H17B	0.0240	0.2198	0.6610	0.059*
C18	0.3477 (2)	0.7417 (2)	0.8026 (2)	0.0248 (5)
C19	0.3067 (2)	0.7120 (2)	0.6885 (2)	0.0298 (5)
H19A	0.2836	0.6304	0.6356	0.036*
C20	0.2988 (2)	0.7998 (3)	0.6504 (2)	0.0370 (6)
H20A	0.2713	0.7770	0.5722	0.044*
C21	0.3305 (2)	0.9190 (3)	0.7249 (3)	0.0395 (6)
H21A	0.3247	0.9783	0.6986	0.047*
C22	0.3708 (2)	0.9512 (2)	0.8386 (3)	0.0383 (6)
H22A	0.3921	1.0331	0.8912	0.046*
C23	0.3800 (2)	0.8635 (2)	0.8755 (2)	0.0313 (5)
H23A	0.4095	0.8871	0.9536	0.038*
C24	0.2584 (2)	0.6512 (2)	0.92518 (19)	0.0254 (5)

C25	0.1699 (2)	0.7173 (2)	0.9233 (2)	0.0337 (5)
H25A	0.1687	0.7633	0.8823	0.040*
C26	0.0836 (2)	0.7187 (3)	0.9790 (3)	0.0412 (6)
H26A	0.0251	0.7650	0.9756	0.049*
C27	0.0830 (3)	0.6528 (3)	1.0393 (3)	0.0425 (7)
H27A	0.0247	0.6537	1.0780	0.051*
C28	0.1687 (3)	0.5852 (3)	1.0425 (2)	0.0398 (6)
H28A	0.1690	0.5392	1.0835	0.048*
C29	0.2542 (2)	0.5840 (2)	0.9863 (2)	0.0318 (5)
H29A	0.3117	0.5364	0.9891	0.038*
C30	0.4912 (2)	0.67005 (19)	0.9294 (2)	0.0267 (5)
C31	0.5262 (2)	0.7043 (2)	1.0449 (2)	0.0330 (6)
H31A	0.4696	0.7137	1.0869	0.040*
C32	0.6433 (3)	0.7251 (2)	1.1005 (2)	0.0414 (7)
H32A	0.6646	0.7483	1.1792	0.050*
C33	0.7263 (2)	0.7123 (2)	1.0424 (3)	0.0429 (7)
H33A	0.8052	0.7258	1.0803	0.051*
C34	0.6955 (2)	0.6797 (3)	0.9283 (3)	0.0424 (7)
H34A	0.7530	0.6712	0.8874	0.051*
C35	0.5798 (2)	0.6594 (2)	0.8735 (2)	0.0351 (6)
H35A	0.5599	0.6373	0.7950	0.042*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Dy1	0.02183 (6)	0.01583 (6)	0.01804 (6)	0.00100 (4)	0.00370 (4)	0.00748 (4)
Cl1	0.0365 (3)	0.0313 (3)	0.0369 (3)	0.0028 (2)	0.0059 (2)	0.0243 (3)
Cl2	0.0376 (3)	0.0231 (3)	0.0322 (3)	0.0012 (2)	0.0053 (2)	0.0169 (2)
O1	0.0244 (8)	0.0269 (9)	0.0423 (10)	0.0024 (7)	0.0038 (7)	0.0090 (8)
O2	0.0381 (10)	0.0284 (9)	0.0259 (9)	0.0075 (7)	0.0121 (7)	0.0076 (7)
O3	0.0256 (8)	0.0307 (9)	0.0256 (8)	-0.0050 (7)	0.0001 (7)	0.0128 (7)
O4	0.0305 (9)	0.0291 (9)	0.0296 (9)	0.0063 (7)	0.0119 (7)	0.0136 (7)
N1	0.0272 (10)	0.0246 (10)	0.0259 (10)	0.0018 (8)	0.0032 (8)	0.0061 (8)
B1	0.0278 (12)	0.0190 (11)	0.0197 (12)	0.0026 (9)	0.0018 (10)	0.0064 (10)
C1	0.0231 (11)	0.0264 (12)	0.0236 (11)	0.0030 (9)	0.0035 (9)	0.0114 (10)
C2	0.0244 (12)	0.0460 (16)	0.0486 (16)	-0.0008 (11)	0.0057 (11)	0.0292 (14)
C3	0.0453 (19)	0.0471 (19)	0.095 (3)	-0.0098 (15)	-0.0298 (19)	0.040 (2)
C4	0.0424 (16)	0.0489 (18)	0.0525 (18)	0.0083 (13)	0.0032 (14)	0.0301 (15)
C5	0.030 (2)	0.034 (2)	0.074 (4)	0.0091 (18)	0.011 (2)	0.026 (2)
C5B	0.028 (4)	0.062 (6)	0.066 (6)	0.004 (4)	0.000 (4)	0.050 (5)
C6	0.084 (3)	0.0343 (16)	0.056 (2)	0.0188 (16)	0.0311 (19)	0.0130 (15)
C7	0.0495 (17)	0.0352 (15)	0.0406 (16)	0.0061 (13)	0.0187 (13)	0.0005 (13)
C8	0.059 (2)	0.0543 (19)	0.0323 (15)	0.0081 (16)	0.0211 (14)	0.0070 (14)
C9	0.0504 (17)	0.0402 (16)	0.0393 (15)	0.0044 (13)	0.0254 (13)	0.0123 (13)
C10	0.0328 (13)	0.0355 (14)	0.0384 (15)	-0.0075 (11)	0.0026 (11)	0.0149 (12)
C11	0.0354 (14)	0.0273 (13)	0.0456 (16)	-0.0037 (11)	-0.0010 (12)	0.0096 (12)
C12	0.061 (2)	0.0490 (18)	0.0325 (15)	-0.0120 (15)	-0.0106 (14)	0.0131 (14)
C13	0.0461 (17)	0.0491 (17)	0.0352 (15)	-0.0084 (13)	-0.0088 (12)	0.0241 (14)

C14	0.0383 (15)	0.0452 (16)	0.0558 (18)	0.0190 (13)	0.0187 (13)	0.0296 (15)
C15	0.090 (3)	0.108 (4)	0.050 (2)	0.071 (3)	0.017 (2)	0.024 (2)
C16	0.076 (3)	0.173 (5)	0.082 (3)	0.075 (3)	0.059 (3)	0.083 (4)
C17	0.0502 (18)	0.070 (2)	0.0479 (18)	0.0171 (16)	0.0252 (15)	0.0390 (17)
C18	0.0245 (11)	0.0263 (11)	0.0255 (11)	0.0049 (9)	0.0063 (9)	0.0131 (9)
C19	0.0277 (12)	0.0361 (13)	0.0260 (12)	0.0054 (10)	0.0061 (9)	0.0145 (10)
C20	0.0308 (13)	0.0594 (18)	0.0356 (14)	0.0129 (12)	0.0094 (11)	0.0335 (14)
C21	0.0312 (13)	0.0515 (17)	0.0590 (18)	0.0146 (12)	0.0155 (13)	0.0432 (16)
C22	0.0389 (14)	0.0310 (13)	0.0511 (17)	0.0080 (11)	0.0114 (13)	0.0236 (13)
C23	0.0377 (13)	0.0269 (12)	0.0300 (13)	0.0056 (10)	0.0068 (10)	0.0140 (10)
C24	0.0320 (12)	0.0189 (10)	0.0195 (10)	0.0008 (9)	0.0037 (9)	0.0047 (9)
C25	0.0355 (13)	0.0343 (13)	0.0364 (14)	0.0083 (11)	0.0094 (11)	0.0198 (12)
C26	0.0359 (14)	0.0423 (16)	0.0524 (17)	0.0136 (12)	0.0165 (13)	0.0241 (14)
C27	0.0428 (16)	0.0451 (16)	0.0454 (16)	0.0057 (13)	0.0207 (13)	0.0217 (14)
C28	0.0503 (17)	0.0371 (15)	0.0388 (15)	0.0045 (12)	0.0145 (13)	0.0224 (13)
C29	0.0403 (14)	0.0238 (12)	0.0313 (13)	0.0048 (10)	0.0096 (11)	0.0120 (10)
C30	0.0327 (12)	0.0144 (10)	0.0262 (11)	-0.0002 (9)	-0.0018 (9)	0.0071 (9)
C31	0.0423 (14)	0.0218 (11)	0.0271 (12)	0.0062 (10)	-0.0012 (10)	0.0074 (10)
C32	0.0518 (17)	0.0247 (12)	0.0311 (14)	0.0065 (12)	-0.0122 (12)	0.0065 (11)
C33	0.0317 (14)	0.0248 (13)	0.0570 (18)	0.0016 (10)	-0.0095 (13)	0.0131 (13)
C34	0.0300 (13)	0.0360 (15)	0.0563 (18)	0.0002 (11)	0.0028 (12)	0.0201 (14)
C35	0.0316 (13)	0.0351 (14)	0.0367 (14)	-0.0007 (11)	0.0016 (11)	0.0180 (12)

Geometric parameters (Å, °)

Dy1—O1	2.3790 (18)	C7—C8	1.519 (5)
Dy1—O2	2.3838 (17)	C8—C9	1.495 (4)
Dy1—O4	2.3932 (17)	C10—C11	1.521 (4)
Dy1—O3	2.4022 (16)	C11—C12	1.514 (4)
Dy1—N1	2.431 (2)	C12—C13	1.498 (4)
Dy1—C12	2.5835 (6)	C14—C15	1.456 (5)
Dy1—C11	2.5888 (6)	C15—C16	1.470 (7)
O1—C5	1.411 (5)	C16—C17	1.522 (5)
O1—C2	1.455 (3)	C18—C19	1.397 (3)
O1—C5B	1.597 (8)	C18—C23	1.398 (3)
O2—C9	1.454 (3)	C19—C20	1.396 (4)
O2—C6	1.476 (4)	C20—C21	1.377 (4)
O3—C13	1.439 (3)	C21—C22	1.384 (4)
O3—C10	1.455 (3)	C22—C23	1.386 (4)
O4—C14	1.445 (3)	C24—C25	1.397 (3)
O4—C17	1.465 (3)	C24—C29	1.407 (3)
N1—C1	1.141 (3)	C25—C26	1.390 (4)
B1—C1	1.621 (3)	C26—C27	1.382 (4)
B1—C18	1.625 (3)	C27—C28	1.385 (4)
B1—C24	1.632 (4)	C28—C29	1.387 (4)
B1—C30	1.641 (3)	C30—C31	1.396 (3)
C2—C3	1.505 (4)	C30—C35	1.402 (4)
C3—C4	1.507 (4)	C31—C32	1.407 (4)

C4—C5B	1.450 (9)	C32—C33	1.363 (5)
C4—C5	1.557 (6)	C33—C34	1.380 (4)
C6—C7	1.499 (4)	C34—C35	1.390 (4)
O1—Dy1—O2	73.01 (6)	O1—C2—C3	104.0 (2)
O1—Dy1—O4	142.46 (6)	C2—C3—C4	105.0 (3)
O2—Dy1—O4	143.59 (6)	C5B—C4—C3	98.6 (4)
O1—Dy1—O3	145.88 (6)	C3—C4—C5	103.7 (3)
O2—Dy1—O3	73.03 (6)	O1—C5—C4	105.1 (3)
O4—Dy1—O3	70.83 (6)	C4—C5B—O1	101.3 (5)
O1—Dy1—N1	73.26 (7)	O2—C6—C7	105.1 (3)
O2—Dy1—N1	144.41 (7)	C6—C7—C8	102.4 (2)
O4—Dy1—N1	71.76 (6)	C9—C8—C7	102.1 (3)
O3—Dy1—N1	140.64 (6)	O2—C9—C8	104.8 (2)
O1—Dy1—Cl2	94.03 (5)	O3—C10—C11	103.4 (2)
O2—Dy1—Cl2	85.89 (5)	C12—C11—C10	102.5 (2)
O4—Dy1—Cl2	96.37 (4)	C13—C12—C11	105.6 (2)
O3—Dy1—Cl2	86.87 (4)	O3—C13—C12	107.3 (2)
N1—Dy1—Cl2	85.47 (5)	O4—C14—C15	107.6 (3)
O1—Dy1—Cl1	87.73 (5)	C14—C15—C16	102.9 (4)
O2—Dy1—Cl1	89.32 (5)	C15—C16—C17	106.5 (3)
O4—Dy1—Cl1	85.53 (4)	O4—C17—C16	104.3 (3)
O3—Dy1—Cl1	88.55 (4)	C19—C18—C23	116.1 (2)
N1—Dy1—Cl1	100.36 (5)	C19—C18—B1	126.0 (2)
Cl2—Dy1—Cl1	174.169 (19)	C23—C18—B1	117.9 (2)
C5—O1—C2	104.8 (2)	C20—C19—C18	121.4 (2)
C2—O1—C5B	105.3 (3)	C21—C20—C19	120.8 (2)
C5—O1—Dy1	129.0 (2)	C20—C21—C22	119.2 (2)
C2—O1—Dy1	125.50 (15)	C21—C22—C23	119.7 (3)
C5B—O1—Dy1	119.5 (3)	C22—C23—C18	122.8 (2)
C9—O2—C6	108.8 (2)	C25—C24—C29	115.6 (2)
C9—O2—Dy1	127.40 (16)	C25—C24—B1	122.3 (2)
C6—O2—Dy1	123.67 (17)	C29—C24—B1	122.1 (2)
C13—O3—C10	107.00 (19)	C26—C25—C24	122.8 (2)
C13—O3—Dy1	128.45 (15)	C27—C26—C25	120.0 (3)
C10—O3—Dy1	124.48 (15)	C26—C27—C28	119.1 (3)
C14—O4—C17	107.9 (2)	C27—C28—C29	120.5 (3)
C14—O4—Dy1	120.89 (15)	C28—C29—C24	122.1 (2)
C17—O4—Dy1	126.17 (16)	C31—C30—C35	115.8 (2)
C1—N1—Dy1	163.92 (19)	C31—C30—B1	125.4 (2)
C1—B1—C18	108.07 (19)	C35—C30—B1	118.6 (2)
C1—B1—C24	105.67 (19)	C30—C31—C32	121.6 (3)
C18—B1—C24	111.50 (19)	C33—C32—C31	120.5 (3)
C1—B1—C30	106.34 (19)	C32—C33—C34	119.8 (3)
C18—B1—C30	108.93 (19)	C33—C34—C35	119.6 (3)
C24—B1—C30	115.9 (2)	C34—C35—C30	122.7 (3)
N1—C1—B1	178.7 (3)		

Dichlorido(cyanotriphenylborato- κ N)tetrakis(tetrahydrofuran- κ O)ytterbium(III) (1-Y)*Crystal data*[YCl₂(C₁₉H₁₅BN)(C₄H₈O)₄] $M_r = 716.35$ Triclinic, $P\bar{1}$ $a = 12.0404$ (9) Å $b = 12.5428$ (9) Å $c = 13.4654$ (10) Å $\alpha = 114.6967$ (8)° $\beta = 101.3142$ (9)° $\gamma = 94.3882$ (9)° $V = 1782.7$ (2) Å³ $Z = 2$ $F(000) = 748$ $D_x = 1.334$ Mg m⁻³Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 9606 reflections

 $\theta = 2.3$ – 27.1 ° $\mu = 1.82$ mm⁻¹ $T = 173$ K

Prism, colorless

 $0.32 \times 0.20 \times 0.13$ mm*Data collection*Bruker SMART APEXII CCD
diffractometer

Radiation source: fine-focus sealed tube

 φ and ω scansAbsorption correction: multi-scan
(*TWINABS*; Sheldrick, 2012) $T_{\min} = 0.576$, $T_{\max} = 0.746$

7988 measured reflections

7988 independent reflections

6188 reflections with $I > 2\sigma(I)$ $\theta_{\max} = 27.5$ °, $\theta_{\min} = 1.7$ ° $h = -15$ → 15 $k = -16$ → 14 $l = 0$ → 17 *Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.047$ $wR(F^2) = 0.102$ $S = 1.04$

7988 reflections

425 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0438P)^2 + 0.9643P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.64$ e Å⁻³ $\Delta\rho_{\min} = -0.80$ e Å⁻³*Special details*

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. A colorless crystal of approximate dimensions 0.126 x 0.198 x 0.324 mm was mounted on a glass fiber and transferred to a Bruker SMART APEX II diffractometer system. The APEX2 program package and the CELL_NOW were used to determine the unit-cell parameters. Data was collected using a 30 sec/frame scan time. The raw frame data was processed using SAINT3 and TWINABS to yield the reflection data file (HKL5 format). Subsequent calculations were carried out using the SHELXTL program package. There were no systematic absences nor any diffraction symmetry other than the Friedel condition. The centrosymmetric triclinic space group P-1 was assigned and later determined to be correct.

The structure was solved by direct methods and refined on F2 by full-matrix least-squares techniques. The analytical scattering factors for neutral atoms were used throughout the analysis. Hydrogen atoms were included using a riding model. Disordered atoms were included using multiple components with partial site-occupancy-factors.

Least-squares analysis yielded $wR2 = 0.1023$ and $Goof = 1.037$ for 425 variables refined against 7988 data (0.77), $R1 = 0.0470$ for those 6188 with $I > 2.0\sigma(I)$. The structure was refined as a two-component twin, $BASF = 0.4868$.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Y1	0.24658 (3)	0.24946 (3)	0.49747 (3)	0.01941 (8)	
Cl1	0.23233 (7)	0.07981 (7)	0.55775 (7)	0.0330 (2)	
Cl2	0.25375 (7)	0.40279 (7)	0.41725 (7)	0.03054 (19)	
O1	0.44949 (18)	0.2696 (2)	0.5508 (2)	0.0338 (6)	
O2	0.30323 (19)	0.1206 (2)	0.33612 (18)	0.0304 (5)	
O3	0.07439 (18)	0.1455 (2)	0.35125 (18)	0.0291 (5)	
O4	0.08446 (18)	0.3011 (2)	0.57256 (18)	0.0291 (5)	
N1	0.3129 (2)	0.4202 (2)	0.6805 (2)	0.0273 (6)	
B1	0.3570 (3)	0.6480 (3)	0.8568 (3)	0.0242 (7)	
C1	0.3324 (3)	0.5138 (3)	0.7540 (3)	0.0249 (7)	
C2	0.5324 (3)	0.3724 (3)	0.5732 (3)	0.0368 (8)	
H2A	0.4986	0.4459	0.5980	0.044*	
H2B	0.5594	0.3611	0.5052	0.044*	
C3	0.6292 (4)	0.3788 (4)	0.6661 (4)	0.0612 (14)	
H3A	0.6173	0.4296	0.7411	0.073*	
H3B	0.7039	0.4117	0.6605	0.073*	
C4	0.6265 (3)	0.2520 (4)	0.6489 (4)	0.0481 (10)	
H4A	0.6216	0.2449	0.7185	0.058*	0.65
H4B	0.6958	0.2223	0.6263	0.058*	0.65
H4C	0.6516	0.2028	0.5806	0.058*	0.35
H4D	0.6740	0.2474	0.7154	0.058*	0.35
C5	0.5161 (5)	0.1826 (5)	0.5525 (6)	0.0425 (15)	0.65
H5A	0.5358	0.1414	0.4793	0.051*	0.65
H5B	0.4739	0.1227	0.5679	0.051*	0.65
C5B	0.5064 (10)	0.2189 (15)	0.6354 (15)	0.065 (4)	0.35
H5B1	0.4853	0.2562	0.7083	0.078*	0.35
H5B2	0.4832	0.1312	0.6023	0.078*	0.35
C6	0.2840 (4)	-0.0103 (3)	0.2910 (4)	0.0563 (12)	
H6A	0.2006	-0.0436	0.2636	0.068*	
H6B	0.3185	-0.0342	0.3500	0.068*	
C7	0.3416 (3)	-0.0537 (3)	0.1950 (3)	0.0466 (10)	
H7A	0.3008	-0.1328	0.1350	0.056*	
H7B	0.4230	-0.0582	0.2216	0.056*	
C8	0.3321 (4)	0.0423 (4)	0.1539 (3)	0.0509 (11)	
H8A	0.3900	0.0435	0.1113	0.061*	
H8B	0.2543	0.0305	0.1058	0.061*	
C9	0.3553 (3)	0.1534 (3)	0.2621 (3)	0.0435 (10)	
H9A	0.4392	0.1819	0.2949	0.052*	
H9B	0.3204	0.2171	0.2497	0.052*	
C10	-0.0181 (3)	0.0683 (3)	0.3584 (3)	0.0392 (9)	
H10A	-0.0787	0.1131	0.3859	0.047*	
H10B	0.0120	0.0329	0.4093	0.047*	
C11	-0.0638 (3)	-0.0268 (3)	0.2377 (3)	0.0417 (9)	
H11A	-0.1448	-0.0630	0.2237	0.050*	
H11B	-0.0165	-0.0904	0.2196	0.050*	

C12	-0.0539 (4)	0.0431 (4)	0.1702 (3)	0.0558 (12)	
H12A	-0.0326	-0.0059	0.0996	0.067*	
H12B	-0.1278	0.0682	0.1507	0.067*	
C13	0.0381 (3)	0.1494 (4)	0.2447 (3)	0.0469 (10)	
H13A	0.1039	0.1472	0.2102	0.056*	
H13B	0.0080	0.2236	0.2555	0.056*	
C14	0.0212 (3)	0.3876 (3)	0.5531 (3)	0.0429 (9)	
H14A	0.0648	0.4292	0.5205	0.052*	0.65
H14B	-0.0542	0.3462	0.4986	0.052*	0.65
H14C	-0.0332	0.3522	0.4774	0.052*	0.35
H14D	0.0731	0.4585	0.5633	0.052*	0.35
C15	0.0041 (7)	0.4723 (7)	0.6573 (6)	0.0430 (17)	0.65
H15A	-0.0664	0.5055	0.6446	0.052*	0.65
H15B	0.0708	0.5384	0.6986	0.052*	0.65
C16	-0.0078 (6)	0.3991 (8)	0.7203 (6)	0.0461 (17)	0.65
H16A	-0.0895	0.3649	0.7055	0.055*	0.65
H16B	0.0231	0.4485	0.8026	0.055*	0.65
C15B	-0.0431 (15)	0.4171 (17)	0.6509 (18)	0.071 (5)	0.35
H15C	-0.0002	0.4898	0.7188	0.085*	0.35
H15D	-0.1215	0.4309	0.6262	0.085*	0.35
C16B	-0.0492 (13)	0.3163 (16)	0.6758 (13)	0.061 (4)	0.35
H16C	-0.0646	0.3364	0.7505	0.073*	0.35
H16D	-0.1071	0.2475	0.6167	0.073*	0.35
C17	0.0642 (4)	0.2970 (4)	0.6745 (3)	0.0507 (11)	
H17A	0.1378	0.3126	0.7304	0.061*	0.65
H17B	0.0203	0.2185	0.6565	0.061*	0.65
H17C	0.0732	0.2184	0.6726	0.061*	0.35
H17D	0.1195	0.3598	0.7425	0.061*	0.35
C18	0.3475 (3)	0.7411 (3)	0.8012 (2)	0.0234 (6)	
C19	0.3062 (3)	0.7111 (3)	0.6868 (3)	0.0300 (7)	
H19A	0.2829	0.6295	0.6338	0.036*	
C20	0.2984 (3)	0.7984 (4)	0.6489 (3)	0.0373 (8)	
H20A	0.2706	0.7754	0.5705	0.045*	
C21	0.3304 (3)	0.9175 (4)	0.7232 (3)	0.0402 (9)	
H21A	0.3250	0.9764	0.6964	0.048*	
C22	0.3703 (3)	0.9503 (3)	0.8368 (3)	0.0377 (8)	
H22A	0.3914	1.0323	0.8893	0.045*	
C23	0.3795 (3)	0.8627 (3)	0.8743 (3)	0.0303 (7)	
H23A	0.4087	0.8865	0.9527	0.036*	
C24	0.2577 (3)	0.6505 (3)	0.9236 (3)	0.0258 (7)	
C25	0.1695 (3)	0.7157 (3)	0.9213 (3)	0.0336 (8)	
H25A	0.1685	0.7615	0.8801	0.040*	
C26	0.0831 (3)	0.7171 (4)	0.9762 (3)	0.0437 (9)	
H26A	0.0243	0.7626	0.9720	0.052*	
C27	0.0831 (3)	0.6515 (3)	1.0372 (3)	0.0435 (9)	
H27A	0.0250	0.6525	1.0761	0.052*	
C28	0.1685 (3)	0.5845 (3)	1.0411 (3)	0.0395 (9)	
H28A	0.1689	0.5390	1.0825	0.047*	

C29	0.2534 (3)	0.5837 (3)	0.9850 (3)	0.0314 (8)
H29A	0.3108	0.5365	0.9881	0.038*
C30	0.4906 (3)	0.6702 (3)	0.9288 (3)	0.0264 (7)
C31	0.5250 (3)	0.7033 (3)	1.0442 (3)	0.0345 (8)
H31A	0.4683	0.7113	1.0858	0.041*
C32	0.6417 (3)	0.7251 (3)	1.1004 (3)	0.0446 (10)
H32A	0.6629	0.7491	1.1795	0.053*
C33	0.7251 (3)	0.7120 (3)	1.0424 (4)	0.0441 (10)
H33A	0.8037	0.7245	1.0804	0.053*
C34	0.6941 (3)	0.6806 (3)	0.9286 (4)	0.0426 (10)
H34A	0.7511	0.6724	0.8874	0.051*
C35	0.5787 (3)	0.6610 (3)	0.8744 (3)	0.0355 (8)
H35A	0.5588	0.6401	0.7959	0.043*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Y1	0.02100 (14)	0.01684 (13)	0.01973 (13)	0.00180 (10)	0.00476 (10)	0.00791 (10)
Cl1	0.0363 (5)	0.0321 (4)	0.0387 (5)	0.0042 (4)	0.0077 (4)	0.0244 (4)
Cl2	0.0377 (5)	0.0236 (4)	0.0332 (4)	0.0024 (3)	0.0066 (4)	0.0168 (3)
O1	0.0216 (12)	0.0270 (12)	0.0441 (14)	0.0039 (10)	0.0048 (10)	0.0092 (11)
O2	0.0347 (13)	0.0268 (13)	0.0268 (12)	0.0075 (10)	0.0124 (10)	0.0068 (10)
O3	0.0243 (12)	0.0325 (13)	0.0268 (12)	-0.0043 (10)	0.0019 (9)	0.0131 (10)
O4	0.0300 (12)	0.0300 (13)	0.0299 (12)	0.0062 (10)	0.0125 (10)	0.0133 (10)
N1	0.0262 (14)	0.0236 (14)	0.0257 (14)	0.0023 (11)	0.0047 (11)	0.0059 (12)
B1	0.0294 (19)	0.0165 (16)	0.0219 (17)	0.0023 (15)	0.0024 (15)	0.0062 (13)
C1	0.0224 (16)	0.0287 (17)	0.0249 (16)	0.0055 (13)	0.0052 (13)	0.0134 (14)
C2	0.0264 (18)	0.042 (2)	0.047 (2)	0.0009 (16)	0.0076 (16)	0.0252 (18)
C3	0.043 (2)	0.043 (3)	0.084 (3)	-0.0033 (19)	-0.019 (2)	0.032 (2)
C4	0.041 (2)	0.054 (3)	0.054 (3)	0.0083 (19)	0.0030 (19)	0.033 (2)
C5	0.026 (3)	0.036 (3)	0.065 (4)	0.009 (2)	0.009 (3)	0.024 (3)
C5B	0.023 (6)	0.099 (12)	0.108 (12)	0.001 (7)	0.002 (7)	0.086 (11)
C6	0.082 (3)	0.029 (2)	0.055 (3)	0.017 (2)	0.028 (2)	0.010 (2)
C7	0.047 (2)	0.036 (2)	0.044 (2)	0.0079 (18)	0.0209 (19)	0.0007 (18)
C8	0.058 (3)	0.053 (3)	0.032 (2)	0.008 (2)	0.0207 (19)	0.0056 (19)
C9	0.049 (2)	0.040 (2)	0.043 (2)	0.0062 (18)	0.0268 (19)	0.0123 (18)
C10	0.0304 (19)	0.039 (2)	0.043 (2)	-0.0083 (16)	0.0030 (16)	0.0193 (17)
C11	0.0314 (19)	0.0271 (18)	0.053 (2)	-0.0030 (16)	-0.0018 (17)	0.0111 (17)
C12	0.060 (3)	0.056 (3)	0.036 (2)	-0.009 (2)	-0.009 (2)	0.018 (2)
C13	0.048 (2)	0.051 (3)	0.037 (2)	-0.0078 (19)	-0.0073 (18)	0.025 (2)
C14	0.037 (2)	0.045 (2)	0.061 (3)	0.0201 (18)	0.0213 (19)	0.031 (2)
C15	0.039 (4)	0.055 (5)	0.039 (3)	0.022 (3)	0.013 (3)	0.020 (3)
C16	0.044 (4)	0.065 (5)	0.040 (4)	0.026 (4)	0.022 (3)	0.026 (4)
C15B	0.049 (10)	0.082 (14)	0.110 (15)	0.039 (9)	0.059 (11)	0.048 (13)
C16B	0.056 (9)	0.093 (12)	0.053 (9)	0.027 (9)	0.037 (7)	0.038 (9)
C17	0.051 (3)	0.076 (3)	0.049 (2)	0.020 (2)	0.029 (2)	0.041 (2)
C18	0.0194 (15)	0.0272 (16)	0.0249 (15)	0.0058 (13)	0.0066 (13)	0.0121 (13)
C19	0.0272 (18)	0.0378 (19)	0.0295 (17)	0.0084 (15)	0.0098 (14)	0.0175 (15)

C20	0.0279 (19)	0.063 (3)	0.0351 (18)	0.0153 (18)	0.0099 (16)	0.0333 (19)
C21	0.031 (2)	0.052 (2)	0.064 (3)	0.0170 (17)	0.0179 (18)	0.047 (2)
C22	0.035 (2)	0.0295 (19)	0.053 (2)	0.0081 (15)	0.0103 (17)	0.0228 (17)
C23	0.0336 (18)	0.0278 (17)	0.0301 (17)	0.0058 (15)	0.0062 (15)	0.0143 (14)
C24	0.0309 (17)	0.0204 (16)	0.0219 (16)	0.0030 (13)	0.0045 (13)	0.0065 (13)
C25	0.035 (2)	0.034 (2)	0.0369 (19)	0.0062 (16)	0.0082 (16)	0.0209 (16)
C26	0.038 (2)	0.046 (2)	0.056 (2)	0.0147 (18)	0.0178 (19)	0.0264 (19)
C27	0.043 (2)	0.048 (2)	0.046 (2)	0.0050 (19)	0.0234 (19)	0.0219 (19)
C28	0.050 (2)	0.036 (2)	0.039 (2)	0.0043 (18)	0.0143 (18)	0.0217 (17)
C29	0.038 (2)	0.0248 (18)	0.0322 (18)	0.0053 (15)	0.0121 (15)	0.0123 (15)
C30	0.0324 (18)	0.0124 (15)	0.0272 (17)	0.0005 (13)	-0.0014 (14)	0.0065 (13)
C31	0.044 (2)	0.0237 (18)	0.0276 (18)	0.0065 (15)	-0.0010 (15)	0.0077 (15)
C32	0.057 (3)	0.0235 (18)	0.0333 (19)	0.0053 (17)	-0.0145 (18)	0.0056 (15)
C33	0.033 (2)	0.0239 (19)	0.058 (3)	0.0026 (15)	-0.0088 (19)	0.0116 (18)
C34	0.033 (2)	0.033 (2)	0.058 (3)	0.0006 (16)	0.0059 (18)	0.0201 (19)
C35	0.0307 (19)	0.0341 (19)	0.040 (2)	0.0002 (15)	0.0047 (16)	0.0176 (16)

Geometric parameters (Å, °)

Y1—O1	2.369 (2)	C10—C11	1.515 (5)
Y1—O2	2.370 (2)	C11—C12	1.518 (5)
Y1—O4	2.382 (2)	C12—C13	1.493 (5)
Y1—O3	2.390 (2)	C14—C15	1.429 (8)
Y1—N1	2.420 (3)	C14—C15B	1.583 (16)
Y1—C12	2.5730 (8)	C15—C16	1.503 (11)
Y1—C11	2.5803 (9)	C16—C17	1.578 (8)
O1—C5	1.408 (6)	C15B—C16B	1.44 (2)
O1—C2	1.452 (4)	C16B—C17	1.408 (14)
O1—C5B	1.594 (12)	C18—C19	1.396 (4)
O2—C9	1.451 (4)	C18—C23	1.396 (4)
O2—C6	1.473 (4)	C19—C20	1.390 (5)
O3—C13	1.438 (4)	C20—C21	1.376 (5)
O3—C10	1.463 (4)	C21—C22	1.377 (5)
O4—C14	1.453 (4)	C22—C23	1.391 (5)
O4—C17	1.460 (4)	C24—C25	1.392 (5)
N1—C1	1.144 (4)	C24—C29	1.407 (5)
B1—C24	1.626 (5)	C25—C26	1.388 (5)
B1—C1	1.629 (4)	C26—C27	1.384 (5)
B1—C18	1.630 (5)	C27—C28	1.383 (5)
B1—C30	1.642 (5)	C28—C29	1.384 (5)
C2—C3	1.504 (5)	C30—C35	1.388 (5)
C3—C4	1.505 (6)	C30—C31	1.393 (5)
C4—C5B	1.428 (12)	C31—C32	1.404 (5)
C4—C5	1.552 (7)	C32—C33	1.368 (6)
C6—C7	1.506 (5)	C33—C34	1.376 (6)
C7—C8	1.523 (6)	C34—C35	1.387 (5)
C8—C9	1.492 (5)		

O1—Y1—O2	72.94 (8)	C5B—C4—C3	99.2 (7)
O1—Y1—O4	142.45 (8)	C3—C4—C5	103.6 (4)
O2—Y1—O4	143.68 (8)	O1—C5—C4	105.3 (4)
O1—Y1—O3	145.89 (8)	C4—C5B—O1	102.3 (8)
O2—Y1—O3	73.07 (8)	O2—C6—C7	105.5 (3)
O4—Y1—O3	70.93 (8)	C6—C7—C8	102.2 (3)
O1—Y1—N1	73.27 (8)	C9—C8—C7	102.2 (3)
O2—Y1—N1	144.47 (9)	O2—C9—C8	105.4 (3)
O4—Y1—N1	71.65 (8)	O3—C10—C11	103.5 (3)
O3—Y1—N1	140.55 (9)	C10—C11—C12	102.4 (3)
O1—Y1—C12	93.71 (6)	C13—C12—C11	105.8 (3)
O2—Y1—C12	86.20 (6)	O3—C13—C12	107.4 (3)
O4—Y1—C12	96.39 (6)	C15—C14—O4	109.7 (4)
O3—Y1—C12	86.81 (6)	O4—C14—C15B	98.3 (7)
N1—Y1—C12	85.32 (7)	C14—C15—C16	102.8 (6)
O1—Y1—C11	88.01 (6)	C15—C16—C17	105.4 (5)
O2—Y1—C11	89.21 (6)	C16B—C15B—C14	107.1 (11)
O4—Y1—C11	85.42 (6)	C17—C16B—C15B	97.5 (11)
O3—Y1—C11	88.77 (6)	C16B—C17—O4	106.2 (7)
N1—Y1—C11	100.28 (7)	O4—C17—C16	103.9 (4)
C12—Y1—C11	174.39 (3)	C19—C18—C23	116.3 (3)
C5—O1—C2	104.8 (3)	C19—C18—B1	126.0 (3)
C2—O1—C5B	105.3 (5)	C23—C18—B1	117.6 (3)
C5—O1—Y1	128.8 (3)	C20—C19—C18	121.2 (3)
C2—O1—Y1	125.9 (2)	C21—C20—C19	121.0 (3)
C5B—O1—Y1	119.6 (4)	C20—C21—C22	119.3 (3)
C9—O2—C6	108.5 (3)	C21—C22—C23	119.5 (3)
C9—O2—Y1	127.3 (2)	C22—C23—C18	122.6 (3)
C6—O2—Y1	124.1 (2)	C25—C24—C29	115.5 (3)
C13—O3—C10	107.0 (2)	C25—C24—B1	122.6 (3)
C13—O3—Y1	128.5 (2)	C29—C24—B1	121.9 (3)
C10—O3—Y1	124.49 (19)	C26—C25—C24	123.2 (3)
C14—O4—C17	107.8 (3)	C27—C26—C25	119.4 (4)
C14—O4—Y1	121.12 (19)	C28—C27—C26	119.4 (3)
C17—O4—Y1	126.3 (2)	C27—C28—C29	120.2 (3)
C1—N1—Y1	164.6 (3)	C28—C29—C24	122.2 (3)
C24—B1—C1	105.7 (3)	C35—C30—C31	115.7 (3)
C24—B1—C18	111.6 (3)	C35—C30—B1	119.3 (3)
C1—B1—C18	107.8 (2)	C31—C30—B1	125.0 (3)
C24—B1—C30	116.6 (3)	C30—C31—C32	121.4 (4)
C1—B1—C30	106.0 (3)	C33—C32—C31	120.6 (3)
C18—B1—C30	108.6 (3)	C32—C33—C34	119.4 (3)
N1—C1—B1	178.4 (3)	C33—C34—C35	119.4 (4)
O1—C2—C3	103.9 (3)	C34—C35—C30	123.5 (3)
C2—C3—C4	105.0 (3)		

Tetraethylazanium 2,2,4,6-tetraphenyl-1,3,5,2 λ ⁴,4,6-trioxatriborinan-2-ide (2)*Crystal data*C₈H₂₀N⁺·C₂₄H₂₀B₃O₃⁻ $M_r = 519.08$ Monoclinic, $P2_1/n$ $a = 9.7245$ (7) Å $b = 18.3021$ (13) Å $c = 16.7716$ (12) Å $\beta = 101.5456$ (9)° $V = 2924.6$ (4) Å³ $Z = 4$ $F(000) = 1112$ $D_x = 1.179$ Mg m⁻³Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 9884 reflections

 $\theta = 2.4$ – 28.3 ° $\mu = 0.07$ mm⁻¹ $T = 133$ K

Prism, colorless

 $0.37 \times 0.33 \times 0.23$ mm*Data collection*Bruker SMART APEXII CCD
diffractometer

Radiation source: fine-focus sealed tube

 φ and ω scansAbsorption correction: multi-scan
(SADABS; Krause et al., 2015) $T_{\min} = 0.834$, $T_{\max} = 0.862$

31741 measured reflections

5959 independent reflections

5030 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.029$ $\theta_{\max} = 26.4$ °, $\theta_{\min} = 1.7$ ° $h = -12$ → 12 $k = -22$ → 22 $l = -20$ → 20 *Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.064$ $wR(F^2) = 0.179$ $S = 1.02$

5959 reflections

442 parameters

0 restraints

Primary atom site location: dual space

Secondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0909P)^2 + 2.9541P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} = 0.001$ $\Delta\rho_{\max} = 1.06$ e Å⁻³ $\Delta\rho_{\min} = -0.47$ e Å⁻³*Special details*

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. A colorless crystal of approximate dimensions 0.228 x 0.331 x 0.367 mm was mounted on a glass fiber and transferred to a Bruker SMART APEX II diffractometer. The APEX2 program package was used to determine the unit-cell parameters and for data collection (20 sec/frame scan time for a sphere of diffraction data). The raw frame data was processed using SAINT and SADABS to yield the reflection data file. Subsequent calculations were carried out using the SHELXTL program. The diffraction symmetry was 2/m and the systematic absences were consistent with the monoclinic space group $P2_1/n$ that was later determined to be correct.

The structure was solved by dual space methods and refined on F^2 by full-matrix least-squares techniques. The analytical scattering factors for neutral atoms were used throughout the analysis. Hydrogen atoms were included using a riding model. The tetraethylammonium ion was fully disordered. The disordered atoms were included using multiple components with partial site-occupancy-factors.

Least-squares analysis yielded $wR2 = 0.1785$ and $\text{Goof} = 1.017$ for 442 variables refined against 5959 data (0.80 Å), $R1 = 0.0639$ for those 5030 data with $I > 2.0\sigma(I)$.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
B1	0.4762 (2)	0.56789 (11)	0.77924 (12)	0.0193 (4)	
B2	0.6190 (2)	0.62127 (11)	0.68613 (12)	0.0177 (4)	
B3	0.6440 (2)	0.66930 (11)	0.82091 (13)	0.0190 (4)	
O1	0.52579 (14)	0.57190 (7)	0.69997 (7)	0.0215 (3)	
O2	0.67303 (14)	0.67482 (7)	0.74274 (8)	0.0226 (3)	
O3	0.56271 (14)	0.61696 (7)	0.84114 (7)	0.0212 (3)	
C1	0.3139 (2)	0.59569 (10)	0.76601 (11)	0.0208 (4)	
C2	0.2638 (2)	0.63277 (11)	0.82703 (12)	0.0257 (4)	
H2A	0.3257	0.6410	0.8777	0.031*	
C3	0.1269 (2)	0.65811 (11)	0.81642 (13)	0.0295 (5)	
H3A	0.0968	0.6832	0.8595	0.035*	
C4	0.0340 (2)	0.64703 (12)	0.74362 (14)	0.0310 (5)	
H4A	-0.0598	0.6644	0.7361	0.037*	
C5	0.0796 (2)	0.61020 (14)	0.68181 (14)	0.0384 (5)	
H5A	0.0170	0.6021	0.6314	0.046*	
C6	0.2174 (2)	0.58499 (13)	0.69332 (13)	0.0318 (5)	
H6A	0.2467	0.5596	0.6502	0.038*	
C7	0.49230 (18)	0.48395 (10)	0.81153 (11)	0.0191 (4)	
C8	0.5228 (2)	0.46709 (11)	0.89406 (12)	0.0291 (5)	
H8A	0.5386	0.5058	0.9326	0.035*	
C9	0.5309 (3)	0.39533 (13)	0.92183 (14)	0.0368 (5)	
H9A	0.5528	0.3856	0.9786	0.044*	
C10	0.5071 (2)	0.33809 (11)	0.86710 (14)	0.0325 (5)	
H10A	0.5101	0.2891	0.8859	0.039*	
C11	0.4790 (2)	0.35292 (11)	0.78472 (14)	0.0295 (5)	
H11A	0.4645	0.3139	0.7466	0.035*	
C12	0.4719 (2)	0.42482 (11)	0.75758 (12)	0.0245 (4)	
H12A	0.4525	0.4341	0.7007	0.029*	
C13	0.67346 (18)	0.62034 (10)	0.60354 (11)	0.0189 (4)	
C14	0.6536 (2)	0.55854 (11)	0.55359 (12)	0.0253 (4)	
H14A	0.6069	0.5172	0.5698	0.030*	
C15	0.7006 (3)	0.55642 (13)	0.48118 (13)	0.0368 (5)	
H15A	0.6873	0.5136	0.4485	0.044*	
C16	0.7671 (3)	0.61664 (15)	0.45617 (14)	0.0402 (6)	
H16A	0.7992	0.6152	0.4063	0.048*	
C17	0.7867 (2)	0.67887 (14)	0.50387 (13)	0.0368 (5)	
H17A	0.8312	0.7205	0.4865	0.044*	
C18	0.7410 (2)	0.68035 (12)	0.57715 (12)	0.0265 (4)	
H18A	0.7560	0.7230	0.6100	0.032*	
C19	0.71090 (19)	0.72806 (10)	0.88572 (11)	0.0215 (4)	
C20	0.8255 (2)	0.77103 (11)	0.87687 (13)	0.0275 (4)	
H20A	0.8646	0.7654	0.8297	0.033*	
C21	0.8835 (2)	0.82185 (12)	0.93569 (15)	0.0358 (5)	
H21A	0.9633	0.8495	0.9293	0.043*	
C22	0.8252 (2)	0.83221 (12)	1.00344 (14)	0.0336 (5)	

H22A	0.8639	0.8674	1.0433	0.040*	
C23	0.7105 (2)	0.79122 (12)	1.01311 (13)	0.0335 (5)	
H23A	0.6695	0.7985	1.0594	0.040*	
C24	0.6552 (2)	0.73938 (12)	0.95518 (12)	0.0289 (4)	
H24A	0.5773	0.7109	0.9630	0.035*	
N4	0.4859 (11)	0.8966 (7)	0.6646 (6)	0.0247 (17)	0.567 (3)
C25	0.4659 (5)	0.9730 (2)	0.6325 (4)	0.0550 (15)	0.567 (3)
H25A	0.5493	0.9879	0.6111	0.066*	0.567 (3)
H25B	0.4569	1.0065	0.6775	0.066*	0.567 (3)
C26	0.3277 (5)	0.9795 (3)	0.5612 (3)	0.0479 (12)	0.567 (3)
H26A	0.3213	1.0291	0.5387	0.072*	0.567 (3)
H26B	0.2441	0.9691	0.5835	0.072*	0.567 (3)
H26C	0.3339	0.9443	0.5180	0.072*	0.567 (3)
C27	0.4859 (6)	0.8388 (3)	0.5960 (3)	0.0553 (13)	0.567 (3)
H27A	0.5212	0.7923	0.6225	0.066*	0.567 (3)
H27D	0.3871	0.8306	0.5686	0.066*	0.567 (3)
C28	0.5623 (8)	0.8535 (3)	0.5362 (3)	0.0687 (18)	0.567 (3)
H28A	0.5610	0.8106	0.5011	0.103*	0.567 (3)
H28B	0.6595	0.8651	0.5618	0.103*	0.567 (3)
H28C	0.5204	0.8952	0.5034	0.103*	0.567 (3)
C29	0.6298 (5)	0.8933 (3)	0.7230 (5)	0.082 (3)	0.567 (3)
H29A	0.6458	0.8425	0.7431	0.098*	0.567 (3)
H29D	0.7033	0.9047	0.6917	0.098*	0.567 (3)
C30	0.6475 (10)	0.9406 (5)	0.7909 (5)	0.114 (4)	0.567 (3)
H30A	0.7443	0.9374	0.8214	0.170*	0.567 (3)
H30B	0.5829	0.9263	0.8262	0.170*	0.567 (3)
H30C	0.6273	0.9909	0.7722	0.170*	0.567 (3)
C31	0.3724 (4)	0.8759 (3)	0.7034 (3)	0.0452 (12)	0.567 (3)
H31A	0.2815	0.8906	0.6692	0.054*	0.567 (3)
H31B	0.3825	0.9007	0.7568	0.054*	0.567 (3)
C32	0.3769 (6)	0.7839 (3)	0.7165 (4)	0.0343 (11)	0.567 (3)
H32A	0.2809	0.7649	0.7076	0.051*	0.567 (3)
H32B	0.4257	0.7723	0.7720	0.051*	0.567 (3)
H32C	0.4268	0.7615	0.6774	0.051*	0.567 (3)
N4B	0.4989 (18)	0.9025 (10)	0.6867 (8)	0.030 (3)	0.433 (3)
C25B	0.6237 (6)	0.9310 (3)	0.6647 (4)	0.0357 (12)	0.433 (3)
H25C	0.6315	0.9100	0.6114	0.043*	0.433 (3)
H25D	0.7057	0.9136	0.7052	0.043*	0.433 (3)
C26B	0.6326 (4)	1.0148 (2)	0.6590 (3)	0.0206 (9)	0.433 (3)
H26D	0.7147	1.0282	0.6365	0.031*	0.433 (3)
H26E	0.6415	1.0361	0.7134	0.031*	0.433 (3)
H26F	0.5474	1.0334	0.6235	0.031*	0.433 (3)
C27B	0.3630 (5)	0.9241 (3)	0.6138 (3)	0.0358 (12)	0.433 (3)
H27B	0.2776	0.9014	0.6265	0.043*	0.433 (3)
H27C	0.3499	0.9778	0.6139	0.043*	0.433 (3)
C28B	0.3771 (8)	0.9007 (4)	0.5299 (4)	0.0544 (17)	0.433 (3)
H28D	0.2900	0.9117	0.4911	0.082*	0.433 (3)
H28E	0.3954	0.8481	0.5297	0.082*	0.433 (3)

H28F	0.4552	0.9271	0.5141	0.082*	0.433 (3)
C29B	0.4530 (6)	0.9384 (3)	0.7578 (3)	0.0360 (12)	0.433 (3)
H29B	0.3618	0.9178	0.7644	0.043*	0.433 (3)
H29C	0.4403	0.9915	0.7474	0.043*	0.433 (3)
C30B	0.5623 (6)	0.9263 (3)	0.8356 (3)	0.0295 (11)	0.433 (3)
H30D	0.5333	0.9518	0.8809	0.044*	0.433 (3)
H30E	0.6532	0.9453	0.8284	0.044*	0.433 (3)
H30F	0.5707	0.8738	0.8476	0.044*	0.433 (3)
C31B	0.4989 (5)	0.8204 (3)	0.6900 (3)	0.0312 (11)	0.433 (3)
H31C	0.5934	0.8007	0.7127	0.037*	0.433 (3)
H31D	0.4633	0.7986	0.6358	0.037*	0.433 (3)
C32B	0.3869 (8)	0.8069 (5)	0.7537 (5)	0.050 (2)	0.433 (3)
H32D	0.3422	0.7590	0.7430	0.075*	0.433 (3)
H32E	0.3148	0.8451	0.7448	0.075*	0.433 (3)
H32F	0.4387	0.8089	0.8101	0.075*	0.433 (3)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
B1	0.0231 (10)	0.0195 (10)	0.0164 (9)	-0.0042 (8)	0.0062 (8)	-0.0017 (7)
B2	0.0160 (9)	0.0192 (10)	0.0177 (9)	0.0028 (7)	0.0030 (7)	0.0022 (7)
B3	0.0163 (9)	0.0193 (10)	0.0209 (10)	0.0016 (7)	0.0023 (8)	-0.0002 (8)
O1	0.0255 (7)	0.0231 (7)	0.0169 (6)	-0.0053 (5)	0.0065 (5)	-0.0020 (5)
O2	0.0242 (7)	0.0237 (7)	0.0217 (7)	-0.0065 (5)	0.0089 (5)	-0.0046 (5)
O3	0.0249 (7)	0.0210 (7)	0.0177 (6)	-0.0045 (5)	0.0047 (5)	-0.0023 (5)
C1	0.0258 (9)	0.0151 (8)	0.0224 (9)	-0.0025 (7)	0.0069 (7)	0.0022 (7)
C2	0.0303 (10)	0.0269 (10)	0.0198 (9)	0.0009 (8)	0.0049 (8)	0.0004 (7)
C3	0.0332 (11)	0.0280 (10)	0.0301 (11)	0.0064 (8)	0.0132 (9)	0.0000 (8)
C4	0.0254 (10)	0.0293 (11)	0.0389 (12)	0.0042 (8)	0.0080 (9)	0.0007 (9)
C5	0.0301 (11)	0.0470 (14)	0.0342 (12)	0.0054 (10)	-0.0028 (9)	-0.0108 (10)
C6	0.0288 (11)	0.0397 (12)	0.0263 (10)	0.0044 (9)	0.0037 (8)	-0.0099 (9)
C7	0.0152 (8)	0.0220 (9)	0.0205 (9)	-0.0023 (7)	0.0047 (7)	-0.0009 (7)
C8	0.0378 (11)	0.0272 (10)	0.0217 (10)	-0.0039 (9)	0.0041 (8)	0.0009 (8)
C9	0.0447 (13)	0.0349 (12)	0.0286 (11)	-0.0009 (10)	0.0022 (10)	0.0118 (9)
C10	0.0288 (11)	0.0208 (10)	0.0489 (13)	0.0039 (8)	0.0101 (10)	0.0108 (9)
C11	0.0286 (10)	0.0204 (10)	0.0429 (12)	-0.0007 (8)	0.0151 (9)	-0.0064 (8)
C12	0.0276 (10)	0.0233 (9)	0.0238 (9)	-0.0027 (8)	0.0080 (8)	-0.0019 (7)
C13	0.0143 (8)	0.0241 (9)	0.0179 (8)	0.0021 (7)	0.0022 (7)	0.0015 (7)
C14	0.0288 (10)	0.0248 (10)	0.0229 (9)	0.0035 (8)	0.0062 (8)	0.0000 (7)
C15	0.0431 (13)	0.0420 (13)	0.0273 (11)	0.0055 (10)	0.0118 (10)	-0.0083 (9)
C16	0.0364 (12)	0.0659 (17)	0.0220 (10)	-0.0008 (11)	0.0149 (9)	-0.0032 (10)
C17	0.0315 (11)	0.0531 (14)	0.0275 (11)	-0.0125 (10)	0.0097 (9)	0.0066 (10)
C18	0.0252 (10)	0.0309 (10)	0.0233 (10)	-0.0055 (8)	0.0046 (8)	0.0008 (8)
C19	0.0201 (9)	0.0212 (9)	0.0224 (9)	0.0005 (7)	0.0021 (7)	-0.0009 (7)
C20	0.0226 (10)	0.0277 (10)	0.0332 (11)	-0.0030 (8)	0.0082 (8)	-0.0070 (8)
C21	0.0248 (10)	0.0324 (12)	0.0499 (14)	-0.0088 (9)	0.0071 (10)	-0.0133 (10)
C22	0.0323 (11)	0.0293 (11)	0.0355 (12)	-0.0028 (9)	-0.0020 (9)	-0.0144 (9)
C23	0.0409 (12)	0.0339 (12)	0.0257 (10)	-0.0037 (9)	0.0071 (9)	-0.0090 (9)

C24	0.0317 (11)	0.0297 (11)	0.0259 (10)	-0.0084 (8)	0.0071 (8)	-0.0052 (8)
N4	0.016 (2)	0.026 (3)	0.030 (5)	-0.0004 (19)	-0.001 (3)	0.001 (3)
C25	0.044 (3)	0.020 (2)	0.115 (4)	0.0086 (17)	0.048 (3)	0.020 (2)
C26	0.048 (3)	0.048 (3)	0.049 (3)	0.011 (2)	0.013 (2)	0.007 (2)
C27	0.066 (3)	0.042 (3)	0.057 (3)	0.011 (2)	0.009 (3)	-0.010 (2)
C28	0.115 (5)	0.052 (3)	0.038 (3)	0.022 (3)	0.014 (3)	0.004 (2)
C29	0.031 (3)	0.030 (3)	0.162 (7)	0.010 (2)	-0.032 (3)	-0.018 (3)
C30	0.108 (7)	0.106 (6)	0.093 (6)	0.054 (5)	-0.060 (5)	-0.043 (5)
C31	0.0221 (19)	0.057 (3)	0.058 (3)	0.0066 (18)	0.0125 (18)	0.037 (2)
C32	0.031 (2)	0.026 (2)	0.048 (3)	-0.0033 (18)	0.013 (2)	0.006 (2)
N4B	0.044 (5)	0.018 (3)	0.027 (6)	-0.007 (3)	0.005 (4)	-0.002 (4)
C25B	0.028 (3)	0.026 (3)	0.054 (3)	0.004 (2)	0.011 (2)	0.007 (2)
C26B	0.0155 (19)	0.015 (2)	0.033 (2)	-0.0087 (15)	0.0080 (17)	-0.0005 (16)
C27B	0.025 (2)	0.036 (3)	0.042 (3)	0.001 (2)	-0.001 (2)	0.011 (2)
C28B	0.061 (4)	0.057 (4)	0.039 (3)	-0.017 (3)	-0.004 (3)	0.002 (3)
C29B	0.036 (3)	0.027 (3)	0.045 (3)	-0.001 (2)	0.010 (2)	-0.006 (2)
C30B	0.036 (3)	0.029 (2)	0.022 (2)	-0.004 (2)	0.002 (2)	0.0007 (19)
C31B	0.027 (2)	0.021 (2)	0.045 (3)	0.0010 (18)	0.005 (2)	-0.001 (2)
C32B	0.043 (4)	0.048 (5)	0.066 (5)	0.021 (4)	0.028 (4)	0.036 (4)

Geometric parameters (Å, °)

B1—O3	1.497 (2)	C15—C16	1.385 (4)
B1—O1	1.504 (2)	C16—C17	1.383 (4)
B1—C7	1.626 (3)	C17—C18	1.388 (3)
B1—C1	1.631 (3)	C19—C24	1.395 (3)
B2—O1	1.333 (2)	C19—C20	1.396 (3)
B2—O2	1.392 (2)	C20—C21	1.391 (3)
B2—C13	1.579 (3)	C21—C22	1.380 (3)
B3—O3	1.329 (2)	C22—C23	1.380 (3)
B3—O2	1.399 (2)	C23—C24	1.387 (3)
B3—C19	1.574 (3)	N4—C31	1.441 (13)
C1—C2	1.394 (3)	N4—C25	1.497 (14)
C1—C6	1.395 (3)	N4—C29	1.541 (11)
C2—C3	1.388 (3)	N4—C27	1.561 (11)
C3—C4	1.381 (3)	C25—C26	1.615 (8)
C4—C5	1.382 (3)	C27—C28	1.390 (8)
C5—C6	1.394 (3)	C29—C30	1.414 (9)
C7—C8	1.391 (3)	C31—C32	1.697 (7)
C7—C12	1.399 (3)	N4B—C25B	1.435 (19)
C8—C9	1.391 (3)	N4B—C31B	1.504 (18)
C9—C10	1.381 (3)	N4B—C29B	1.506 (15)
C10—C11	1.381 (3)	N4B—C27B	1.659 (16)
C11—C12	1.390 (3)	C25B—C26B	1.541 (6)
C13—C18	1.396 (3)	C27B—C28B	1.504 (9)
C13—C14	1.398 (3)	C29B—C30B	1.525 (8)
C14—C15	1.381 (3)	C31B—C32B	1.688 (9)

O3—B1—O1	110.41 (14)	C15—C14—C13	121.3 (2)
O3—B1—C7	109.43 (15)	C14—C15—C16	120.0 (2)
O1—B1—C7	108.31 (15)	C17—C16—C15	119.9 (2)
O3—B1—C1	107.53 (15)	C16—C17—C18	119.8 (2)
O1—B1—C1	109.70 (15)	C17—C18—C13	121.3 (2)
C7—B1—C1	111.47 (15)	C24—C19—C20	117.19 (18)
O1—B2—O2	122.37 (16)	C24—C19—B3	120.14 (17)
O1—B2—C13	120.31 (17)	C20—C19—B3	122.67 (17)
O2—B2—C13	117.32 (16)	C21—C20—C19	121.36 (19)
O3—B3—O2	122.23 (17)	C22—C21—C20	120.1 (2)
O3—B3—C19	120.06 (17)	C23—C22—C21	119.76 (19)
O2—B3—C19	117.70 (16)	C22—C23—C24	119.9 (2)
B2—O1—B1	122.45 (15)	C23—C24—C19	121.68 (19)
B2—O2—B3	117.97 (15)	C31—N4—C25	110.6 (8)
B3—O3—B1	122.39 (15)	C31—N4—C29	112.5 (8)
C2—C1—C6	115.91 (18)	C25—N4—C29	107.5 (8)
C2—C1—B1	121.51 (17)	C31—N4—C27	105.6 (8)
C6—C1—B1	122.57 (17)	C25—N4—C27	112.8 (7)
C3—C2—C1	122.39 (19)	C29—N4—C27	107.9 (8)
C4—C3—C2	120.37 (19)	N4—C25—C26	111.4 (5)
C3—C4—C5	118.9 (2)	C28—C27—N4	118.7 (7)
C4—C5—C6	120.1 (2)	C30—C29—N4	115.9 (6)
C5—C6—C1	122.3 (2)	N4—C31—C32	108.4 (6)
C8—C7—C12	116.51 (18)	C25B—N4B—C31B	112.3 (12)
C8—C7—B1	121.88 (17)	C25B—N4B—C29B	116.4 (10)
C12—C7—B1	121.58 (16)	C31B—N4B—C29B	113.9 (11)
C9—C8—C7	122.0 (2)	C25B—N4B—C27B	108.2 (10)
C10—C9—C8	120.2 (2)	C31B—N4B—C27B	105.1 (9)
C11—C10—C9	119.29 (19)	C29B—N4B—C27B	99.2 (11)
C10—C11—C12	120.06 (19)	N4B—C25B—C26B	116.0 (8)
C11—C12—C7	121.96 (19)	C28B—C27B—N4B	114.4 (7)
C18—C13—C14	117.63 (17)	N4B—C29B—C30B	110.3 (7)
C18—C13—B2	122.09 (17)	N4B—C31B—C32B	100.0 (8)
C14—C13—B2	120.28 (17)		

Tetraethylazanium cyanodiphenyl- λ^4 -boranyl diphenylborinate (3)

Crystal data

$\text{C}_8\text{H}_{20}\text{N}^+\cdot\text{C}_{25}\text{H}_{20}\text{B}_2\text{NO}^-$

$M_r = 502.29$

Monoclinic, $P2_1/n$

$a = 11.0269$ (8) Å

$b = 13.9387$ (11) Å

$c = 18.8488$ (14) Å

$\beta = 100.6357$ (10)°

$V = 2847.3$ (4) Å³

$Z = 4$

$F(000) = 1080$

$D_x = 1.172$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 9716 reflections

$\theta = 2.2\text{--}29.1^\circ$

$\mu = 0.07$ mm⁻¹

$T = 88$ K

Irregular, colorless

$0.28 \times 0.27 \times 0.22$ mm

Data collection

Bruker SMART APEXII CCD
diffractometer
Radiation source: fine-focus sealed tube
 φ and ω scans
Absorption correction: multi-scan
(SADABS; Krause et al., 2015)
 $T_{\min} = 0.715$, $T_{\max} = 0.746$
34768 measured reflections

7242 independent reflections
5500 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.042$
 $\theta_{\max} = 29.1^\circ$, $\theta_{\min} = 1.8^\circ$
 $h = -14 \rightarrow 14$
 $k = -18 \rightarrow 18$
 $l = -25 \rightarrow 25$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.048$
 $wR(F^2) = 0.119$
 $S = 1.04$
7242 reflections
347 parameters
0 restraints
Primary atom site location: dual space

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0477P)^2 + 1.1994P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.41 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.24 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. A colorless crystal of approximate dimensions 0.216 x 0.265 x 0.280 mm was mounted in a cryoloop and transferred to a Bruker SMART APEX II diffractometer. The APEX2 program package was used to determine the unit-cell parameters and for data collection (30 sec/frame scan time for a sphere of diffraction data). The raw frame data was processed using SAINT and SADABS to yield the reflection data file. Subsequent calculations were carried out using the SHELXTL program. The diffraction symmetry was 2/m and the systematic absences were consistent with the monoclinic space group P21/n that was later determined to be correct.

The structure was solved by dual space methods and refined on F2 by full-matrix least-squares techniques. The analytical scattering factors for neutral atoms were used throughout the analysis. Hydrogen atoms were included using a riding model.

Least-squares analysis yielded $wR2 = 0.1188$ and $\text{Goof} = 1.038$ for 347 variables refined against 7242 data (0.73 Å), $R1 = 0.0476$ for those 5500 data with $I > 2.0\sigma(I)$.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.54584 (8)	0.23892 (7)	0.71613 (5)	0.0169 (2)
N1	0.65451 (11)	0.48002 (9)	0.69950 (7)	0.0223 (3)
B1	0.57013 (13)	0.29746 (11)	0.65455 (8)	0.0150 (3)
B2	0.48921 (13)	0.24852 (11)	0.77251 (8)	0.0164 (3)
C1	0.61696 (12)	0.40504 (10)	0.68153 (7)	0.0164 (3)
C2	0.68248 (12)	0.24787 (9)	0.62196 (7)	0.0155 (3)
C3	0.74007 (12)	0.29439 (10)	0.57114 (7)	0.0167 (3)
H3	0.7132	0.3569	0.5555	0.020*
C4	0.83524 (13)	0.25223 (10)	0.54282 (7)	0.0193 (3)
H4	0.8716	0.2856	0.5081	0.023*
C5	0.87694 (13)	0.16128 (10)	0.56543 (8)	0.0229 (3)

H5	0.9419	0.1321	0.5463	0.028*
C6	0.82301 (14)	0.11343 (11)	0.61610 (9)	0.0271 (3)
H6	0.8516	0.0515	0.6322	0.033*
C7	0.72688 (13)	0.15612 (10)	0.64344 (8)	0.0229 (3)
H7	0.6903	0.1219	0.6777	0.028*
C8	0.44546 (12)	0.30775 (9)	0.59338 (7)	0.0155 (3)
C9	0.42486 (12)	0.38370 (10)	0.54401 (7)	0.0178 (3)
H9	0.4882	0.4299	0.5440	0.021*
C10	0.31446 (13)	0.39366 (10)	0.49485 (7)	0.0210 (3)
H10	0.3036	0.4461	0.4621	0.025*
C11	0.22064 (13)	0.32728 (11)	0.49356 (7)	0.0243 (3)
H11	0.1445	0.3347	0.4610	0.029*
C12	0.23942 (14)	0.24983 (12)	0.54039 (8)	0.0279 (3)
H12	0.1765	0.2030	0.5392	0.033*
C13	0.35024 (13)	0.24034 (11)	0.58922 (8)	0.0234 (3)
H13	0.3615	0.1865	0.6206	0.028*
C14	0.42967 (12)	0.34557 (10)	0.79522 (7)	0.0179 (3)
C15	0.35810 (12)	0.40696 (10)	0.74516 (7)	0.0205 (3)
H15	0.3443	0.3909	0.6953	0.025*
C16	0.30703 (13)	0.49057 (10)	0.76672 (8)	0.0234 (3)
H16	0.2571	0.5299	0.7318	0.028*
C17	0.32839 (13)	0.51700 (10)	0.83869 (8)	0.0240 (3)
H17	0.2951	0.5752	0.8531	0.029*
C18	0.39858 (13)	0.45811 (10)	0.88957 (8)	0.0228 (3)
H18	0.4135	0.4757	0.9391	0.027*
C19	0.44713 (13)	0.37332 (10)	0.86811 (7)	0.0195 (3)
H19	0.4935	0.3329	0.9037	0.023*
C20	0.48423 (12)	0.15502 (9)	0.82032 (7)	0.0160 (3)
C21	0.39985 (12)	0.14284 (10)	0.86704 (7)	0.0171 (3)
H21	0.3452	0.1939	0.8725	0.021*
C22	0.39383 (13)	0.05835 (10)	0.90564 (7)	0.0199 (3)
H22	0.3345	0.0515	0.9360	0.024*
C23	0.47508 (13)	-0.01578 (10)	0.89946 (7)	0.0204 (3)
H23	0.4719	-0.0735	0.9259	0.024*
C24	0.56085 (13)	-0.00558 (10)	0.85476 (7)	0.0210 (3)
H24	0.6171	-0.0561	0.8510	0.025*
C25	0.56500 (13)	0.07811 (10)	0.81549 (7)	0.0187 (3)
H25	0.6237	0.0837	0.7846	0.022*
N2	0.51337 (10)	0.80126 (8)	0.65697 (6)	0.0166 (2)
C26	0.53989 (13)	0.69500 (10)	0.65074 (8)	0.0228 (3)
H26A	0.5995	0.6869	0.6178	0.027*
H26B	0.5797	0.6709	0.6988	0.027*
C27	0.42736 (13)	0.63381 (11)	0.62331 (8)	0.0253 (3)
H27A	0.3910	0.6533	0.5740	0.038*
H27B	0.3666	0.6424	0.6547	0.038*
H27C	0.4517	0.5662	0.6235	0.038*
C28	0.42039 (13)	0.81887 (10)	0.70616 (7)	0.0206 (3)
H28A	0.4178	0.8886	0.7159	0.025*

H28B	0.3376	0.7998	0.6801	0.025*
C29	0.44632 (14)	0.76619 (11)	0.77769 (8)	0.0252 (3)
H29A	0.4397	0.6969	0.7690	0.038*
H29B	0.3862	0.7859	0.8072	0.038*
H29C	0.5298	0.7817	0.8031	0.038*
C30	0.63326 (12)	0.85169 (10)	0.68896 (7)	0.0200 (3)
H30A	0.6145	0.9194	0.6986	0.024*
H30B	0.6676	0.8214	0.7359	0.024*
C31	0.73123 (13)	0.84982 (12)	0.64196 (8)	0.0256 (3)
H31A	0.6986	0.8798	0.5953	0.038*
H31B	0.7541	0.7832	0.6343	0.038*
H31C	0.8041	0.8852	0.6660	0.038*
C32	0.46006 (14)	0.84029 (11)	0.58214 (8)	0.0258 (3)
H32A	0.5154	0.8217	0.5487	0.031*
H32B	0.3792	0.8093	0.5649	0.031*
C33	0.44268 (16)	0.94792 (13)	0.57844 (9)	0.0359 (4)
H33A	0.4017	0.9662	0.5297	0.054*
H33B	0.5233	0.9795	0.5899	0.054*
H33C	0.3918	0.9678	0.6133	0.054*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0185 (5)	0.0189 (5)	0.0140 (4)	0.0017 (4)	0.0045 (4)	0.0036 (4)
N1	0.0202 (6)	0.0224 (6)	0.0246 (6)	-0.0005 (5)	0.0043 (5)	-0.0017 (5)
B1	0.0171 (7)	0.0156 (7)	0.0127 (6)	-0.0001 (5)	0.0034 (5)	0.0017 (5)
B2	0.0143 (7)	0.0180 (7)	0.0163 (7)	-0.0005 (5)	0.0008 (5)	0.0015 (6)
C1	0.0137 (6)	0.0214 (7)	0.0141 (6)	0.0027 (5)	0.0028 (5)	0.0022 (5)
C2	0.0160 (6)	0.0164 (6)	0.0137 (6)	-0.0023 (5)	0.0018 (5)	-0.0023 (5)
C3	0.0200 (6)	0.0169 (6)	0.0129 (6)	-0.0007 (5)	0.0019 (5)	0.0006 (5)
C4	0.0211 (7)	0.0241 (7)	0.0136 (6)	-0.0030 (5)	0.0055 (5)	-0.0008 (5)
C5	0.0235 (7)	0.0225 (7)	0.0249 (7)	-0.0005 (6)	0.0100 (6)	-0.0067 (6)
C6	0.0317 (8)	0.0158 (7)	0.0369 (9)	0.0025 (6)	0.0147 (7)	0.0002 (6)
C7	0.0264 (7)	0.0179 (7)	0.0275 (7)	-0.0012 (6)	0.0127 (6)	0.0033 (6)
C8	0.0168 (6)	0.0175 (6)	0.0131 (6)	0.0010 (5)	0.0051 (5)	-0.0010 (5)
C9	0.0193 (6)	0.0178 (6)	0.0166 (6)	0.0001 (5)	0.0043 (5)	-0.0008 (5)
C10	0.0251 (7)	0.0209 (7)	0.0166 (6)	0.0060 (6)	0.0028 (5)	0.0007 (5)
C11	0.0184 (7)	0.0359 (8)	0.0174 (7)	0.0031 (6)	-0.0001 (5)	-0.0033 (6)
C12	0.0222 (7)	0.0361 (9)	0.0241 (7)	-0.0107 (6)	0.0011 (6)	0.0019 (6)
C13	0.0248 (7)	0.0258 (8)	0.0188 (7)	-0.0057 (6)	0.0015 (6)	0.0054 (6)
C14	0.0168 (6)	0.0178 (6)	0.0199 (7)	-0.0023 (5)	0.0057 (5)	0.0027 (5)
C15	0.0199 (7)	0.0241 (7)	0.0187 (7)	0.0003 (5)	0.0070 (5)	0.0022 (6)
C16	0.0223 (7)	0.0220 (7)	0.0268 (7)	0.0031 (6)	0.0068 (6)	0.0099 (6)
C17	0.0253 (7)	0.0163 (7)	0.0336 (8)	0.0010 (6)	0.0135 (6)	-0.0009 (6)
C18	0.0263 (7)	0.0227 (7)	0.0199 (7)	-0.0033 (6)	0.0058 (6)	-0.0024 (6)
C19	0.0213 (7)	0.0183 (7)	0.0190 (7)	-0.0014 (5)	0.0038 (5)	0.0011 (5)
C20	0.0173 (6)	0.0165 (6)	0.0134 (6)	0.0008 (5)	0.0007 (5)	-0.0013 (5)
C21	0.0162 (6)	0.0171 (6)	0.0171 (6)	0.0019 (5)	0.0004 (5)	-0.0009 (5)

C22	0.0204 (7)	0.0241 (7)	0.0156 (6)	-0.0021 (5)	0.0041 (5)	-0.0007 (5)
C23	0.0274 (7)	0.0159 (7)	0.0171 (6)	-0.0011 (5)	0.0018 (5)	0.0023 (5)
C24	0.0269 (7)	0.0167 (7)	0.0191 (7)	0.0041 (5)	0.0035 (6)	-0.0008 (5)
C25	0.0221 (7)	0.0205 (7)	0.0139 (6)	0.0025 (5)	0.0046 (5)	0.0000 (5)
N2	0.0149 (5)	0.0198 (6)	0.0150 (5)	-0.0009 (4)	0.0027 (4)	-0.0014 (4)
C26	0.0193 (7)	0.0205 (7)	0.0289 (8)	0.0013 (5)	0.0050 (6)	-0.0069 (6)
C27	0.0234 (7)	0.0226 (7)	0.0292 (8)	-0.0040 (6)	0.0031 (6)	-0.0065 (6)
C28	0.0192 (7)	0.0221 (7)	0.0220 (7)	0.0015 (5)	0.0081 (5)	-0.0015 (6)
C29	0.0309 (8)	0.0265 (8)	0.0197 (7)	-0.0026 (6)	0.0089 (6)	-0.0002 (6)
C30	0.0180 (6)	0.0224 (7)	0.0189 (7)	-0.0031 (5)	0.0020 (5)	-0.0033 (5)
C31	0.0185 (7)	0.0330 (8)	0.0259 (7)	-0.0043 (6)	0.0056 (6)	-0.0045 (6)
C32	0.0216 (7)	0.0379 (9)	0.0171 (7)	-0.0031 (6)	0.0014 (5)	0.0040 (6)
C33	0.0328 (9)	0.0409 (10)	0.0355 (9)	0.0058 (7)	0.0102 (7)	0.0193 (8)

Geometric parameters (Å, °)

O1—B2	1.3350 (17)	C14—C15	1.4036 (19)
O1—B1	1.4831 (16)	C14—C19	1.4059 (19)
N1—C1	1.1518 (18)	C15—C16	1.387 (2)
B1—C8	1.6293 (19)	C16—C17	1.383 (2)
B1—C2	1.6343 (19)	C17—C18	1.385 (2)
B1—C1	1.636 (2)	C18—C19	1.388 (2)
B2—C20	1.591 (2)	C20—C21	1.4045 (18)
B2—C14	1.596 (2)	C20—C25	1.4074 (18)
C2—C3	1.4019 (18)	C21—C22	1.3923 (19)
C2—C7	1.4021 (19)	C22—C23	1.386 (2)
C3—C4	1.3923 (19)	C23—C24	1.385 (2)
C4—C5	1.389 (2)	C24—C25	1.3869 (19)
C5—C6	1.386 (2)	N2—C26	1.5184 (18)
C6—C7	1.395 (2)	N2—C30	1.5202 (17)
C8—C13	1.4004 (19)	N2—C28	1.5239 (17)
C8—C9	1.4001 (18)	N2—C32	1.5249 (18)
C9—C10	1.3940 (19)	C26—C27	1.5158 (19)
C10—C11	1.385 (2)	C28—C29	1.515 (2)
C11—C12	1.386 (2)	C30—C31	1.5185 (19)
C12—C13	1.394 (2)	C32—C33	1.512 (2)
B2—O1—B1	138.29 (11)	C15—C14—C19	116.60 (13)
O1—B1—C8	110.58 (11)	C15—C14—B2	122.99 (12)
O1—B1—C2	108.69 (10)	C19—C14—B2	120.41 (12)
C8—B1—C2	111.74 (10)	C16—C15—C14	121.56 (13)
O1—B1—C1	110.69 (10)	C17—C16—C15	120.41 (13)
C8—B1—C1	108.42 (10)	C16—C17—C18	119.56 (13)
C2—B1—C1	106.67 (10)	C17—C18—C19	119.93 (13)
O1—B2—C20	116.03 (12)	C18—C19—C14	121.90 (13)
O1—B2—C14	125.05 (12)	C21—C20—C25	116.67 (12)
C20—B2—C14	118.91 (11)	C21—C20—B2	123.39 (12)
C3—C2—C7	116.17 (12)	C25—C20—B2	119.92 (12)

C3—C2—B1	122.11 (11)	C22—C21—C20	122.06 (12)
C7—C2—B1	121.72 (12)	C23—C22—C21	119.52 (13)
C4—C3—C2	122.36 (12)	C24—C23—C22	119.93 (13)
C5—C4—C3	119.86 (13)	C23—C24—C25	120.30 (13)
C6—C5—C4	119.42 (13)	C24—C25—C20	121.50 (13)
C5—C6—C7	120.09 (14)	C26—N2—C30	108.41 (10)
N1—C1—B1	177.10 (13)	C26—N2—C28	111.41 (10)
C6—C7—C2	122.08 (13)	C30—N2—C28	108.52 (10)
C13—C8—C9	116.23 (12)	C26—N2—C32	108.87 (11)
C13—C8—B1	120.44 (12)	C30—N2—C32	111.12 (10)
C9—C8—B1	123.31 (12)	C28—N2—C32	108.52 (10)
C10—C9—C8	122.12 (13)	C27—C26—N2	114.71 (11)
C11—C10—C9	120.21 (13)	C29—C28—N2	115.19 (11)
C10—C11—C12	119.08 (13)	C31—C30—N2	114.89 (11)
C11—C12—C13	120.29 (14)	C33—C32—N2	114.87 (13)
C12—C13—C8	122.02 (13)		

Bis[(2.2.2-cryptand)potassium] 2,2,4,6-tetraphenyl-1,3,5,2 λ^4 ,4,6-trioxatriborinan-2-ide cyanomethyldiphenylborate tetrahydrofuran disolvate (4)

Crystal data

$2\text{C}_{18}\text{H}_{36}\text{KN}_2\text{O}_6^+ \cdot \text{C}_{24}\text{H}_{20}\text{B}_3\text{O}_3^- \cdot \text{C}_{14}\text{H}_{13}\text{BN}^- \cdot 2\text{C}_4\text{H}_8\text{O}$

$M_r = 1570.27$

Orthorhombic, $Pca2_1$

$a = 27.193$ (2) Å

$b = 14.5520$ (11) Å

$c = 21.2218$ (16) Å

$V = 8397.7$ (11) Å³

$Z = 4$

$F(000) = 3368$

$D_x = 1.242$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 9334 reflections

$\theta = 2.3$ – 25.3°

$\mu = 0.18$ mm⁻¹

$T = 88$ K

Irregular, colorless

$0.45 \times 0.33 \times 0.28$ mm

Data collection

Bruker SMART APEXII CCD
diffractometer

Radiation source: fine-focus sealed tube

φ and ω scans

Absorption correction: multi-scan
(SADABS; Krause et al., 2015)

$T_{\min} = 0.825$, $T_{\max} = 0.862$

85621 measured reflections

15958 independent reflections

14236 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.038$

$\theta_{\max} = 25.7^\circ$, $\theta_{\min} = 1.6^\circ$

$h = -33 \rightarrow 33$

$k = -17 \rightarrow 17$

$l = -25 \rightarrow 25$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.074$

$wR(F^2) = 0.209$

$S = 1.06$

15958 reflections

988 parameters

1 restraint

Primary atom site location: dual space

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.1366P)^2 + 5.6525P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.78$ e Å⁻³

$\Delta\rho_{\min} = -0.34$ e Å⁻³

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. A colorless crystal of approximate dimensions 0.280 x 0.325 x 0.454 mm was mounted in a cryoloop and transferred to a Bruker SMART APEX II diffractometer. The APEX2 program package was used to determine the unit-cell parameters and for data collection (60 sec/frame scan time for a sphere of diffraction data). The raw frame data was processed using SAINT and SADABS to yield the reflection data file. Subsequent calculations were carried out using the SHELXTL program. The diffraction symmetry was mmm and the systematic absences were consistent with the orthorhombic space groups Pbcm and Pca21. It was later determined that space group Pca21 was correct.

The structure was solved by dual space methods and refined on F2 by full-matrix least-squares techniques. The analytical scattering factors for neutral atoms were used throughout the analysis. Hydrogen atoms were included using a riding model. There were two molecules of tetrahydrofuran solvent present. One solvent molecule was disordered and included using multiple components with partial site-occupancy-factors.

Least-squares analysis yielded $wR2 = 0.2089$ and $Goof = 1.064$ for 988 variables refined against 15958 data (0.82 Å), $R1 = 0.0742$ for those 14236 data with $I > 2.0\sigma(I)$. The structure was refined as a two component inversion twin.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	U_{iso}^*/U_{eq}	Occ. (<1)
B1	0.8783 (3)	0.5804 (5)	0.3189 (4)	0.0416 (15)	
B2	0.7983 (2)	0.5517 (5)	0.3727 (3)	0.0363 (14)	
B3	0.8194 (2)	0.7051 (5)	0.3472 (3)	0.0357 (14)	
O1	0.84187 (16)	0.5214 (3)	0.3531 (2)	0.0438 (10)	
O2	0.78518 (15)	0.6428 (3)	0.3691 (2)	0.0409 (10)	
O3	0.86284 (16)	0.6798 (3)	0.3227 (2)	0.0439 (10)	
C1	0.9323 (2)	0.5724 (4)	0.3521 (3)	0.0373 (13)	
C2	0.9683 (2)	0.6407 (4)	0.3493 (3)	0.0358 (13)	
H2A	0.9617	0.6944	0.3254	0.043*	
C3	1.0129 (2)	0.6337 (4)	0.3795 (3)	0.0371 (13)	
H3A	1.0363	0.6819	0.3765	0.044*	
C4	1.0229 (3)	0.5565 (6)	0.4139 (4)	0.059 (2)	
H4A	1.0536	0.5503	0.4348	0.071*	
C5	0.9875 (3)	0.4864 (6)	0.4181 (4)	0.069 (2)	
H5A	0.9938	0.4331	0.4427	0.083*	
C6	0.9438 (3)	0.4954 (5)	0.3866 (4)	0.0526 (17)	
H6A	0.9205	0.4467	0.3887	0.063*	
C7	0.8806 (2)	0.5476 (4)	0.2443 (3)	0.0407 (14)	
C8	0.8981 (2)	0.6029 (6)	0.1968 (4)	0.0556 (19)	
H8A	0.9101	0.6625	0.2067	0.067*	
C9	0.8986 (3)	0.5735 (7)	0.1358 (4)	0.071 (2)	
H9A	0.9106	0.6134	0.1039	0.085*	
C10	0.8822 (3)	0.4880 (7)	0.1195 (3)	0.067 (2)	
H10A	0.8833	0.4685	0.0768	0.081*	
C11	0.8641 (3)	0.4304 (6)	0.1651 (4)	0.060 (2)	
H11A	0.8521	0.3712	0.1541	0.072*	
C12	0.8633 (2)	0.4599 (5)	0.2283 (4)	0.0454 (16)	
H12A	0.8511	0.4202	0.2601	0.054*	
C13	0.7585 (2)	0.4816 (5)	0.3991 (3)	0.0424 (14)	

C14	0.7634 (3)	0.3885 (5)	0.3920 (3)	0.0503 (16)
H14A	0.7918	0.3640	0.3720	0.060*
C15	0.7271 (3)	0.3302 (5)	0.4140 (4)	0.0558 (19)
H15A	0.7321	0.2658	0.4104	0.067*
C16	0.6842 (3)	0.3609 (7)	0.4406 (4)	0.068 (2)
H16A	0.6594	0.3187	0.4533	0.082*
C17	0.6773 (3)	0.4565 (5)	0.4489 (4)	0.0547 (18)
H17A	0.6482	0.4809	0.4671	0.066*
C18	0.7150 (2)	0.5120 (5)	0.4290 (3)	0.0455 (15)
H18A	0.7117	0.5762	0.4359	0.055*
C19	0.8057 (2)	0.8116 (5)	0.3545 (3)	0.0435 (14)
C20	0.7603 (3)	0.8361 (4)	0.3792 (3)	0.0457 (15)
H20A	0.7384	0.7892	0.3928	0.055*
C21	0.7462 (3)	0.9267 (5)	0.3846 (4)	0.0552 (17)
H21A	0.7146	0.9416	0.4008	0.066*
C22	0.7791 (2)	0.9987 (5)	0.3657 (3)	0.0451 (15)
H22A	0.7707	1.0617	0.3702	0.054*
C23	0.8234 (3)	0.9716 (4)	0.3409 (3)	0.0442 (15)
H23A	0.8455	1.0174	0.3264	0.053*
C24	0.8372 (2)	0.8797 (4)	0.3362 (3)	0.0411 (14)
H24A	0.8688	0.8642	0.3201	0.049*
B4	0.6001 (3)	0.1179 (6)	0.5064 (5)	0.054 (2)
N1	0.6889 (3)	0.1126 (5)	0.4428 (3)	0.0676 (19)
C25	0.6501 (2)	0.1121 (5)	0.4657 (4)	0.0526 (19)
C26	0.5909 (2)	0.2136 (4)	0.5281 (3)	0.0405 (14)
H26A	0.6005	0.2195	0.5724	0.061*
H26B	0.5558	0.2279	0.5237	0.061*
H26C	0.6102	0.2564	0.5025	0.061*
C27	0.6088 (2)	0.0457 (5)	0.5642 (3)	0.0436 (15)
C28	0.5940 (2)	0.0708 (6)	0.6250 (4)	0.062 (2)
H28A	0.5778	0.1279	0.6315	0.074*
C29	0.6028 (3)	0.0116 (8)	0.6771 (4)	0.074 (3)
H29A	0.5939	0.0302	0.7185	0.089*
C30	0.6238 (3)	-0.0709 (8)	0.6675 (5)	0.070 (3)
H30A	0.6291	-0.1110	0.7022	0.084*
C31	0.6373 (3)	-0.0972 (6)	0.6097 (4)	0.062 (2)
H31A	0.6521	-0.1557	0.6036	0.074*
C32	0.6300 (2)	-0.0405 (5)	0.5593 (4)	0.0457 (16)
H32A	0.6400	-0.0612	0.5188	0.055*
C33	0.5551 (2)	0.0858 (4)	0.4593 (3)	0.0444 (15)
C34	0.5151 (2)	0.1423 (4)	0.4492 (3)	0.0413 (14)
H34A	0.5147	0.2013	0.4683	0.050*
C35	0.4758 (2)	0.1170 (5)	0.4127 (3)	0.0423 (15)
H35A	0.4493	0.1587	0.4074	0.051*
C36	0.4739 (2)	0.0349 (5)	0.3842 (4)	0.0493 (16)
H36A	0.4462	0.0179	0.3595	0.059*
C37	0.5126 (3)	-0.0236 (6)	0.3918 (5)	0.074 (3)
H37A	0.5117	-0.0827	0.3728	0.089*

C38	0.5538 (3)	0.0023 (6)	0.4272 (5)	0.067 (2)
H38A	0.5813	-0.0378	0.4295	0.081*
K1	0.63521 (4)	0.17698 (7)	0.13192 (5)	0.0225 (2)
O4	0.6908 (2)	0.1969 (3)	0.2438 (2)	0.0539 (13)
O5	0.59021 (16)	0.1512 (3)	0.2501 (2)	0.0440 (10)
O6	0.68052 (14)	0.3229 (2)	0.0665 (2)	0.0321 (9)
O7	0.57967 (15)	0.3352 (3)	0.0934 (2)	0.0370 (9)
O8	0.68719 (13)	0.0226 (3)	0.0800 (2)	0.0323 (9)
O9	0.58499 (13)	0.0382 (3)	0.06660 (19)	0.0316 (8)
N2	0.74407 (17)	0.1841 (3)	0.1252 (3)	0.0399 (12)
N3	0.52603 (17)	0.1712 (3)	0.1390 (3)	0.0367 (11)
C39	0.7649 (3)	0.2152 (5)	0.1857 (4)	0.0568 (19)
H39A	0.7612	0.2827	0.1886	0.068*
H39B	0.8006	0.2013	0.1860	0.068*
C40	0.7421 (3)	0.1730 (5)	0.2421 (4)	0.065 (2)
H40A	0.7459	0.1053	0.2404	0.077*
H40B	0.7587	0.1955	0.2806	0.077*
C41	0.6699 (3)	0.1606 (5)	0.2991 (3)	0.0543 (19)
H41A	0.6882	0.1831	0.3364	0.065*
H41B	0.6719	0.0927	0.2984	0.065*
C42	0.6184 (3)	0.1899 (5)	0.3028 (3)	0.0520 (18)
H42A	0.6166	0.2578	0.3014	0.062*
H42B	0.6040	0.1690	0.3431	0.062*
C43	0.5393 (3)	0.1761 (5)	0.2556 (3)	0.0495 (17)
H43A	0.5358	0.2438	0.2549	0.059*
H43B	0.5258	0.1531	0.2960	0.059*
C44	0.5117 (2)	0.1341 (5)	0.2009 (3)	0.0485 (16)
H44A	0.5173	0.0669	0.2009	0.058*
H44B	0.4760	0.1445	0.2071	0.058*
C45	0.7583 (2)	0.2471 (4)	0.0750 (3)	0.0430 (15)
H45A	0.7518	0.2173	0.0339	0.052*
H45B	0.7941	0.2585	0.0779	0.052*
C46	0.7319 (2)	0.3381 (4)	0.0766 (3)	0.0406 (15)
H46A	0.7371	0.3682	0.1179	0.049*
H46B	0.7451	0.3790	0.0434	0.049*
C47	0.6549 (2)	0.4078 (4)	0.0635 (3)	0.0367 (13)
H47A	0.6704	0.4481	0.0316	0.044*
H47B	0.6566	0.4391	0.1048	0.044*
C48	0.6024 (2)	0.3903 (4)	0.0461 (3)	0.0404 (14)
H48A	0.5847	0.4494	0.0422	0.049*
H48B	0.6008	0.3583	0.0050	0.049*
C49	0.5290 (2)	0.3193 (5)	0.0787 (4)	0.0483 (16)
H49A	0.5263	0.2853	0.0384	0.058*
H49B	0.5117	0.3787	0.0740	0.058*
C50	0.5058 (2)	0.2641 (5)	0.1312 (4)	0.0495 (16)
H50A	0.5100	0.2980	0.1712	0.059*
H50B	0.4701	0.2590	0.1229	0.059*
C51	0.7628 (2)	0.0919 (4)	0.1118 (4)	0.0478 (17)

H51A	0.7610	0.0549	0.1508	0.057*
H51B	0.7979	0.0969	0.0998	0.057*
C52	0.7361 (2)	0.0415 (4)	0.0605 (4)	0.0450 (16)
H52A	0.7356	0.0794	0.0218	0.054*
H52B	0.7534	-0.0167	0.0510	0.054*
C53	0.6610 (2)	-0.0244 (4)	0.0311 (3)	0.0382 (13)
H53A	0.6783	-0.0821	0.0201	0.046*
H53B	0.6596	0.0146	-0.0071	0.046*
C54	0.6099 (2)	-0.0454 (4)	0.0535 (3)	0.0356 (13)
H54A	0.5919	-0.0802	0.0207	0.043*
H54B	0.6114	-0.0837	0.0920	0.043*
C55	0.5348 (2)	0.0197 (4)	0.0827 (3)	0.0399 (14)
H55A	0.5332	-0.0146	0.1230	0.048*
H55B	0.5192	-0.0182	0.0495	0.048*
C56	0.5077 (2)	0.1108 (4)	0.0891 (3)	0.0388 (13)
H56A	0.5098	0.1439	0.0485	0.047*
H56B	0.4725	0.0978	0.0972	0.047*
K2	0.37144 (4)	0.32115 (7)	0.67152 (5)	0.0278 (3)
O10	0.4324 (2)	0.3508 (4)	0.5644 (2)	0.0635 (15)
O11	0.33319 (18)	0.2979 (4)	0.5500 (2)	0.0503 (12)
O12	0.42078 (19)	0.1605 (3)	0.7159 (3)	0.0540 (12)
O13	0.31818 (18)	0.1755 (3)	0.7270 (2)	0.0441 (11)
O14	0.41652 (16)	0.4584 (3)	0.7459 (2)	0.0422 (10)
O15	0.31684 (15)	0.4755 (3)	0.7123 (2)	0.0370 (9)
N4	0.4804 (2)	0.3234 (4)	0.6868 (4)	0.0608 (19)
N5	0.26389 (19)	0.3164 (3)	0.6549 (3)	0.0411 (13)
C57	0.5027 (3)	0.3645 (8)	0.6302 (5)	0.088 (3)
H57A	0.5386	0.3531	0.6311	0.106*
H57B	0.4976	0.4318	0.6313	0.106*
C58	0.4825 (4)	0.3279 (10)	0.5704 (5)	0.099 (4)
H58A	0.5011	0.3538	0.5344	0.119*
H58B	0.4863	0.2603	0.5695	0.119*
C59	0.4153 (3)	0.3079 (5)	0.5103 (3)	0.059 (2)
H59A	0.4170	0.2404	0.5153	0.071*
H59B	0.4358	0.3256	0.4737	0.071*
C60	0.3647 (3)	0.3362 (5)	0.4999 (3)	0.057 (2)
H60A	0.3534	0.3139	0.4583	0.068*
H60B	0.3626	0.4041	0.5001	0.068*
C61	0.2829 (3)	0.3242 (5)	0.5402 (4)	0.0575 (19)
H61A	0.2715	0.3015	0.4987	0.069*
H61B	0.2801	0.3920	0.5403	0.069*
C62	0.2513 (3)	0.2849 (5)	0.5910 (4)	0.0525 (17)
H62A	0.2166	0.3011	0.5823	0.063*
H62B	0.2539	0.2170	0.5897	0.063*
C63	0.4985 (3)	0.2297 (6)	0.6946 (5)	0.068 (2)
H63A	0.5330	0.2324	0.7097	0.081*
H63B	0.4987	0.1991	0.6530	0.081*
C64	0.4697 (3)	0.1739 (6)	0.7386 (5)	0.073 (3)

H64A	0.4859	0.1135	0.7441	0.088*	
H64B	0.4686	0.2047	0.7801	0.088*	
C65	0.3931 (3)	0.1040 (5)	0.7584 (4)	0.062 (2)	
H65A	0.3911	0.1339	0.8002	0.074*	
H65B	0.4095	0.0438	0.7635	0.074*	
C66	0.3428 (3)	0.0905 (4)	0.7324 (4)	0.0543 (19)	
H66A	0.3451	0.0610	0.6905	0.065*	
H66B	0.3240	0.0492	0.7605	0.065*	
C67	0.2685 (3)	0.1627 (4)	0.7081 (4)	0.0492 (18)	
H67A	0.2513	0.1233	0.7391	0.059*	
H67B	0.2675	0.1317	0.6666	0.059*	
C68	0.2434 (3)	0.2546 (4)	0.7039 (4)	0.0523 (18)	
H68A	0.2080	0.2446	0.6950	0.063*	
H68B	0.2459	0.2856	0.7453	0.063*	
C69	0.4936 (3)	0.3785 (5)	0.7419 (5)	0.065 (2)	
H69A	0.5296	0.3887	0.7420	0.078*	
H69B	0.4850	0.3440	0.7806	0.078*	
C70	0.4678 (3)	0.4699 (5)	0.7429 (5)	0.071 (3)	
H70A	0.4791	0.5057	0.7799	0.085*	
H70B	0.4764	0.5049	0.7045	0.085*	
C71	0.3906 (2)	0.5435 (4)	0.7514 (3)	0.0429 (15)	
H71A	0.3940	0.5790	0.7119	0.051*	
H71B	0.4047	0.5803	0.7862	0.051*	
C72	0.3381 (2)	0.5254 (4)	0.7641 (3)	0.0410 (14)	
H72A	0.3347	0.4892	0.8033	0.049*	
H72B	0.3205	0.5843	0.7698	0.049*	
C73	0.2657 (2)	0.4600 (4)	0.7209 (4)	0.0483 (17)	
H73A	0.2487	0.5194	0.7270	0.058*	
H73B	0.2604	0.4221	0.7591	0.058*	
C74	0.2446 (2)	0.4112 (4)	0.6642 (4)	0.0506 (17)	
H74A	0.2084	0.4082	0.6687	0.061*	
H74B	0.2519	0.4480	0.6261	0.061*	
O16	0.8367 (2)	0.2092 (4)	0.3526 (3)	0.0673 (15)	
C75	0.8791 (4)	0.2395 (8)	0.3223 (6)	0.091 (3)	
H75A	0.8781	0.3072	0.3176	0.110*	
H75B	0.8812	0.2119	0.2797	0.110*	
C76	0.9223 (3)	0.2129 (9)	0.3595 (7)	0.103 (4)	
H76A	0.9443	0.2662	0.3664	0.123*	
H76B	0.9410	0.1637	0.3381	0.123*	
C77	0.9007 (4)	0.1783 (6)	0.4226 (6)	0.080 (3)	
H77A	0.9038	0.1107	0.4263	0.096*	
H77B	0.9172	0.2076	0.4589	0.096*	
C78	0.8485 (3)	0.2063 (6)	0.4185 (4)	0.0595 (19)	
H78A	0.8273	0.1615	0.4406	0.071*	
H78B	0.8438	0.2676	0.4379	0.071*	
O17	0.4260 (5)	0.2513 (10)	0.8786 (7)	0.075 (4)*	0.5
C79	0.3802 (9)	0.3057 (16)	0.8880 (10)	0.072 (5)*	0.5
H79A	0.3533	0.2838	0.8604	0.087*	0.5

H79B	0.3859	0.3720	0.8802	0.087*	0.5
C80	0.3684 (7)	0.2852 (15)	0.9628 (9)	0.068 (5)*	0.5
H80A	0.3447	0.3291	0.9814	0.081*	0.5
H80B	0.3578	0.2211	0.9708	0.081*	0.5
C81	0.4237 (13)	0.304 (2)	0.9836 (18)	0.123 (11)*	0.5
H81A	0.4314	0.3706	0.9806	0.148*	0.5
H81B	0.4293	0.2839	1.0275	0.148*	0.5
C82	0.4539 (10)	0.2522 (19)	0.9406 (14)	0.103 (7)*	0.5
H82A	0.4588	0.1889	0.9564	0.123*	0.5
H82B	0.4864	0.2818	0.9353	0.123*	0.5
O17B	0.4414 (7)	0.2715 (13)	0.9034 (10)	0.100 (5)*	0.5
C79B	0.4032 (9)	0.3234 (16)	0.8637 (13)	0.089 (6)*	0.5
H79C	0.4046	0.3904	0.8718	0.106*	0.5
H79D	0.4081	0.3121	0.8181	0.106*	0.5
C80B	0.3583 (8)	0.2855 (14)	0.8849 (9)	0.066 (5)*	0.5
H80C	0.3326	0.3336	0.8838	0.079*	0.5
H80D	0.3482	0.2359	0.8557	0.079*	0.5
C81B	0.3612 (7)	0.2500 (15)	0.9457 (10)	0.070 (5)*	0.5
H81C	0.3536	0.1835	0.9449	0.084*	0.5
H81D	0.3365	0.2807	0.9729	0.084*	0.5
C82B	0.4160 (8)	0.2659 (16)	0.9746 (11)	0.074 (5)*	0.5
H82C	0.4279	0.2133	0.9998	0.089*	0.5
H82D	0.4190	0.3239	0.9987	0.089*	0.5

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
B1	0.043 (4)	0.041 (4)	0.040 (4)	0.000 (3)	0.008 (3)	0.000 (3)
B2	0.035 (3)	0.041 (4)	0.034 (3)	0.000 (3)	-0.001 (3)	0.000 (3)
B3	0.036 (3)	0.037 (3)	0.034 (3)	-0.001 (3)	0.001 (3)	0.005 (3)
O1	0.041 (2)	0.038 (2)	0.052 (3)	-0.0030 (18)	0.007 (2)	0.0022 (19)
O2	0.039 (2)	0.037 (2)	0.046 (2)	-0.0065 (18)	0.0003 (19)	0.0097 (19)
O3	0.045 (2)	0.038 (2)	0.049 (3)	-0.0006 (18)	0.007 (2)	0.0067 (19)
C1	0.040 (3)	0.036 (3)	0.036 (3)	0.005 (2)	0.003 (2)	-0.010 (3)
C2	0.049 (3)	0.031 (3)	0.028 (3)	0.005 (2)	0.010 (2)	-0.008 (2)
C3	0.042 (3)	0.034 (3)	0.035 (3)	-0.005 (2)	0.017 (3)	-0.010 (2)
C4	0.037 (4)	0.087 (6)	0.053 (4)	0.006 (4)	-0.007 (3)	-0.005 (4)
C5	0.067 (5)	0.061 (5)	0.080 (6)	-0.002 (4)	-0.009 (4)	0.036 (4)
C6	0.044 (4)	0.042 (3)	0.072 (5)	-0.007 (3)	0.002 (3)	0.015 (3)
C7	0.031 (3)	0.048 (4)	0.043 (3)	0.005 (3)	0.001 (3)	0.001 (3)
C8	0.034 (3)	0.079 (5)	0.054 (4)	-0.019 (3)	-0.005 (3)	0.012 (4)
C9	0.041 (4)	0.115 (8)	0.056 (5)	-0.019 (4)	0.005 (3)	0.021 (5)
C10	0.045 (4)	0.124 (8)	0.033 (4)	0.009 (5)	-0.013 (3)	-0.021 (4)
C11	0.046 (4)	0.066 (5)	0.069 (5)	0.004 (3)	-0.017 (4)	-0.019 (4)
C12	0.037 (3)	0.043 (4)	0.055 (4)	0.008 (3)	0.005 (3)	0.001 (3)
C13	0.045 (3)	0.055 (4)	0.028 (3)	-0.003 (3)	-0.006 (2)	0.002 (3)
C14	0.056 (4)	0.045 (4)	0.049 (4)	-0.014 (3)	-0.008 (3)	0.006 (3)
C15	0.074 (5)	0.037 (3)	0.057 (4)	-0.018 (3)	-0.002 (4)	0.009 (3)

C16	0.056 (5)	0.095 (7)	0.053 (4)	-0.030 (4)	-0.010 (4)	0.022 (4)
C17	0.040 (3)	0.063 (5)	0.061 (4)	-0.008 (3)	-0.007 (3)	0.026 (4)
C18	0.033 (3)	0.062 (4)	0.041 (3)	-0.003 (3)	0.000 (3)	0.003 (3)
C19	0.043 (3)	0.053 (4)	0.034 (3)	-0.001 (3)	-0.005 (3)	0.004 (3)
C20	0.049 (4)	0.039 (3)	0.049 (4)	-0.001 (3)	0.010 (3)	0.005 (3)
C21	0.052 (4)	0.057 (4)	0.056 (4)	0.014 (3)	-0.006 (3)	-0.013 (4)
C22	0.048 (4)	0.041 (3)	0.047 (4)	0.005 (3)	-0.010 (3)	0.007 (3)
C23	0.058 (4)	0.034 (3)	0.041 (3)	-0.004 (3)	-0.011 (3)	-0.001 (3)
C24	0.050 (4)	0.036 (3)	0.037 (3)	-0.002 (3)	-0.009 (3)	0.001 (3)
B4	0.041 (4)	0.044 (4)	0.076 (6)	-0.004 (3)	-0.001 (4)	0.013 (4)
N1	0.086 (5)	0.062 (4)	0.055 (4)	0.019 (4)	-0.008 (4)	0.002 (3)
C25	0.023 (3)	0.063 (4)	0.071 (5)	-0.004 (3)	0.015 (3)	0.032 (4)
C26	0.039 (3)	0.020 (3)	0.062 (4)	0.001 (2)	-0.012 (3)	-0.005 (3)
C27	0.028 (3)	0.049 (4)	0.054 (4)	-0.006 (3)	-0.003 (3)	0.007 (3)
C28	0.025 (3)	0.068 (5)	0.092 (6)	0.002 (3)	-0.001 (3)	-0.032 (5)
C29	0.039 (4)	0.144 (9)	0.038 (4)	-0.019 (5)	0.002 (3)	-0.013 (5)
C30	0.033 (4)	0.105 (7)	0.073 (6)	-0.016 (4)	-0.014 (4)	0.027 (6)
C31	0.040 (4)	0.065 (5)	0.080 (6)	-0.006 (3)	-0.008 (4)	0.030 (4)
C32	0.036 (3)	0.042 (4)	0.060 (4)	-0.006 (3)	0.007 (3)	0.005 (3)
C33	0.039 (3)	0.041 (3)	0.053 (4)	-0.009 (3)	0.008 (3)	0.011 (3)
C34	0.052 (4)	0.032 (3)	0.040 (3)	-0.001 (3)	0.008 (3)	0.010 (3)
C35	0.042 (3)	0.048 (4)	0.036 (3)	0.008 (3)	0.011 (3)	0.016 (3)
C36	0.035 (3)	0.048 (4)	0.065 (4)	-0.001 (3)	-0.002 (3)	0.002 (3)
C37	0.053 (5)	0.051 (4)	0.118 (8)	0.013 (3)	-0.023 (5)	-0.037 (5)
C38	0.033 (4)	0.065 (5)	0.103 (7)	0.012 (3)	-0.008 (4)	-0.006 (5)
K1	0.0211 (5)	0.0246 (5)	0.0218 (5)	-0.0034 (4)	0.0017 (4)	-0.0003 (4)
O4	0.073 (3)	0.059 (3)	0.031 (2)	-0.024 (3)	-0.012 (2)	0.008 (2)
O5	0.044 (2)	0.060 (3)	0.028 (2)	-0.008 (2)	0.0118 (18)	-0.0045 (19)
O6	0.0296 (19)	0.0231 (19)	0.044 (2)	-0.0051 (14)	0.0056 (17)	0.0003 (16)
O7	0.031 (2)	0.034 (2)	0.046 (2)	0.0045 (16)	0.0025 (17)	0.0027 (18)
O8	0.0252 (18)	0.0274 (19)	0.044 (2)	-0.0023 (14)	0.0005 (16)	-0.0018 (17)
O9	0.0266 (18)	0.033 (2)	0.035 (2)	-0.0052 (15)	-0.0020 (15)	-0.0078 (16)
N2	0.024 (2)	0.037 (3)	0.058 (3)	-0.0057 (19)	-0.013 (2)	-0.002 (2)
N3	0.027 (2)	0.037 (2)	0.046 (3)	-0.0024 (18)	0.008 (2)	-0.005 (2)
C39	0.049 (4)	0.052 (4)	0.070 (5)	-0.015 (3)	-0.025 (4)	-0.001 (4)
C40	0.066 (5)	0.062 (5)	0.066 (5)	-0.014 (4)	-0.043 (4)	0.015 (4)
C41	0.090 (6)	0.036 (3)	0.037 (3)	-0.008 (3)	-0.013 (4)	0.004 (3)
C42	0.081 (5)	0.055 (4)	0.020 (3)	0.008 (4)	0.003 (3)	-0.004 (3)
C43	0.050 (4)	0.062 (4)	0.037 (3)	-0.007 (3)	0.024 (3)	-0.005 (3)
C44	0.041 (3)	0.054 (4)	0.050 (4)	-0.018 (3)	0.019 (3)	-0.013 (3)
C45	0.020 (3)	0.044 (3)	0.065 (4)	-0.011 (2)	0.006 (3)	-0.002 (3)
C46	0.036 (3)	0.028 (3)	0.058 (4)	-0.012 (2)	0.013 (3)	0.001 (3)
C47	0.045 (3)	0.021 (3)	0.044 (3)	0.000 (2)	0.013 (3)	0.004 (2)
C48	0.045 (3)	0.030 (3)	0.046 (4)	0.010 (3)	-0.002 (3)	0.011 (3)
C49	0.035 (3)	0.044 (4)	0.067 (5)	0.009 (3)	0.001 (3)	-0.001 (3)
C50	0.027 (3)	0.048 (4)	0.074 (5)	0.006 (2)	0.013 (3)	-0.006 (4)
C51	0.024 (3)	0.038 (3)	0.082 (5)	0.000 (2)	-0.010 (3)	0.000 (3)
C52	0.022 (3)	0.037 (3)	0.076 (5)	0.006 (2)	0.004 (3)	-0.001 (3)

C53	0.039 (3)	0.039 (3)	0.037 (3)	0.004 (3)	0.001 (2)	-0.009 (3)
C54	0.038 (3)	0.030 (3)	0.038 (3)	-0.007 (2)	-0.006 (2)	-0.006 (2)
C55	0.028 (3)	0.040 (3)	0.051 (3)	-0.013 (2)	-0.003 (2)	-0.009 (3)
C56	0.021 (2)	0.052 (4)	0.044 (3)	-0.004 (2)	0.001 (2)	-0.001 (3)
K2	0.0351 (6)	0.0252 (6)	0.0231 (5)	-0.0068 (4)	0.0050 (4)	-0.0041 (4)
O10	0.072 (4)	0.078 (4)	0.041 (3)	-0.021 (3)	0.015 (3)	-0.002 (3)
O11	0.055 (3)	0.067 (3)	0.029 (2)	0.001 (2)	-0.006 (2)	-0.004 (2)
O12	0.058 (3)	0.045 (3)	0.059 (3)	0.008 (2)	0.000 (2)	-0.004 (2)
O13	0.062 (3)	0.024 (2)	0.046 (3)	-0.0058 (18)	0.020 (2)	0.0017 (17)
O14	0.043 (2)	0.031 (2)	0.053 (3)	-0.0010 (18)	-0.005 (2)	-0.0147 (19)
O15	0.033 (2)	0.036 (2)	0.042 (2)	-0.0053 (17)	0.0069 (17)	-0.0058 (18)
N4	0.045 (3)	0.056 (4)	0.081 (5)	-0.009 (3)	0.013 (3)	-0.031 (3)
N5	0.035 (3)	0.032 (3)	0.056 (3)	-0.008 (2)	0.006 (2)	-0.006 (2)
C57	0.055 (5)	0.107 (8)	0.102 (8)	-0.039 (5)	0.032 (5)	-0.022 (7)
C58	0.064 (6)	0.156 (11)	0.077 (7)	-0.011 (6)	0.046 (6)	0.011 (7)
C59	0.087 (6)	0.046 (4)	0.043 (4)	0.006 (4)	0.032 (4)	0.006 (3)
C60	0.089 (6)	0.059 (4)	0.023 (3)	0.009 (4)	0.008 (3)	-0.002 (3)
C61	0.064 (5)	0.051 (4)	0.057 (4)	-0.010 (3)	-0.018 (4)	-0.005 (3)
C62	0.044 (3)	0.042 (3)	0.072 (5)	-0.010 (3)	-0.013 (3)	-0.014 (3)
C63	0.041 (4)	0.075 (5)	0.087 (6)	0.011 (4)	0.006 (4)	-0.030 (5)
C64	0.068 (5)	0.059 (5)	0.093 (7)	0.029 (4)	-0.016 (5)	-0.020 (5)
C65	0.092 (6)	0.036 (4)	0.059 (5)	0.017 (4)	0.006 (4)	0.005 (3)
C66	0.077 (5)	0.030 (3)	0.055 (4)	0.001 (3)	0.018 (4)	0.004 (3)
C67	0.056 (4)	0.036 (3)	0.056 (4)	-0.022 (3)	0.031 (3)	-0.007 (3)
C68	0.043 (4)	0.038 (3)	0.076 (5)	-0.016 (3)	0.023 (3)	-0.005 (3)
C69	0.047 (4)	0.062 (5)	0.086 (6)	0.007 (3)	-0.019 (4)	-0.028 (4)
C70	0.040 (4)	0.052 (4)	0.120 (8)	-0.002 (3)	-0.022 (4)	-0.035 (5)
C71	0.050 (4)	0.033 (3)	0.046 (4)	0.002 (3)	-0.009 (3)	-0.014 (3)
C72	0.050 (4)	0.032 (3)	0.041 (3)	0.003 (3)	0.001 (3)	-0.010 (3)
C73	0.037 (3)	0.030 (3)	0.078 (5)	-0.003 (2)	0.018 (3)	-0.011 (3)
C74	0.030 (3)	0.035 (3)	0.086 (5)	-0.004 (2)	0.008 (3)	-0.004 (3)
O16	0.069 (4)	0.077 (4)	0.056 (3)	-0.006 (3)	-0.006 (3)	-0.009 (3)
C75	0.076 (6)	0.094 (7)	0.104 (8)	-0.011 (5)	-0.020 (6)	0.038 (7)
C76	0.044 (5)	0.120 (9)	0.144 (11)	0.006 (5)	0.012 (6)	0.041 (8)
C77	0.088 (7)	0.056 (5)	0.097 (7)	0.016 (4)	-0.022 (6)	0.009 (5)
C78	0.072 (5)	0.057 (4)	0.050 (4)	0.005 (4)	-0.005 (4)	0.005 (4)

Geometric parameters (Å, °)

B1—O1	1.499 (8)	O8—C53	1.433 (7)
B1—O3	1.508 (8)	O9—C54	1.420 (7)
B1—C1	1.634 (10)	O9—C55	1.432 (7)
B1—C7	1.655 (10)	N2—C45	1.457 (9)
B2—O1	1.331 (8)	N2—C51	1.463 (8)
B2—O2	1.374 (8)	N2—C39	1.475 (9)
B2—C13	1.589 (9)	N3—C56	1.464 (8)
B3—O3	1.342 (8)	N3—C50	1.469 (8)
B3—O2	1.380 (8)	N3—C44	1.472 (9)

B3—C19	1.601 (9)	C39—C40	1.481 (12)
C1—C6	1.374 (9)	C41—C42	1.466 (12)
C1—C2	1.394 (9)	C43—C44	1.513 (10)
C2—C3	1.375 (9)	C45—C46	1.507 (9)
C3—C4	1.368 (11)	C47—C48	1.497 (9)
C4—C5	1.405 (12)	C49—C50	1.511 (10)
C5—C6	1.370 (11)	C51—C52	1.499 (10)
C7—C8	1.375 (10)	C53—C54	1.500 (9)
C7—C12	1.401 (9)	C55—C56	1.522 (9)
C8—C9	1.364 (12)	K2—O11	2.802 (4)
C9—C10	1.366 (14)	K2—O13	2.824 (4)
C10—C11	1.371 (13)	K2—O14	2.826 (4)
C11—C12	1.408 (11)	K2—O15	2.828 (4)
C13—C14	1.368 (10)	K2—O10	2.848 (5)
C13—C18	1.413 (9)	K2—O12	2.855 (5)
C14—C15	1.382 (10)	K2—N5	2.947 (5)
C15—C16	1.372 (12)	K2—N4	2.980 (7)
C16—C17	1.414 (13)	O10—C59	1.387 (10)
C17—C18	1.373 (10)	O10—C58	1.407 (12)
C19—C24	1.367 (9)	O11—C61	1.436 (10)
C19—C20	1.387 (9)	O11—C60	1.475 (9)
C20—C21	1.379 (9)	O12—C64	1.429 (11)
C21—C22	1.435 (11)	O12—C65	1.433 (10)
C22—C23	1.371 (10)	O13—C66	1.412 (8)
C23—C24	1.394 (9)	O13—C67	1.421 (9)
B4—C26	1.488 (10)	O14—C70	1.406 (9)
B4—C25	1.614 (10)	O14—C71	1.430 (7)
B4—C27	1.633 (11)	O15—C73	1.421 (7)
B4—C33	1.648 (11)	O15—C72	1.438 (7)
N1—C25	1.163 (10)	N4—C63	1.459 (11)
C27—C32	1.384 (10)	N4—C69	1.462 (10)
C27—C28	1.399 (11)	N4—C57	1.474 (12)
C28—C29	1.423 (13)	N5—C62	1.471 (9)
C29—C30	1.345 (14)	N5—C68	1.483 (9)
C30—C31	1.336 (14)	N5—C74	1.490 (8)
C31—C32	1.365 (11)	C57—C58	1.482 (16)
C33—C34	1.380 (9)	C59—C60	1.452 (12)
C33—C38	1.394 (11)	C61—C62	1.493 (11)
C34—C35	1.369 (10)	C63—C64	1.464 (14)
C35—C36	1.339 (10)	C65—C66	1.486 (12)
C36—C37	1.365 (10)	C67—C68	1.506 (10)
C37—C38	1.400 (12)	C69—C70	1.504 (10)
K1—O9	2.805 (4)	C71—C72	1.475 (9)
K1—O5	2.815 (4)	C73—C74	1.510 (11)
K1—O6	2.819 (4)	O16—C75	1.393 (12)
K1—O4	2.830 (5)	O16—C78	1.435 (9)
K1—O7	2.872 (4)	C75—C76	1.467 (14)
K1—O8	2.873 (4)	C76—C77	1.546 (17)

K1—N2	2.965 (5)	C77—C78	1.478 (13)
K1—N3	2.974 (5)	O17—C79	1.49 (3)
O4—C41	1.405 (9)	O17—C82	1.52 (3)
O4—C40	1.440 (11)	C79—C80	1.65 (3)
O5—C43	1.435 (8)	C80—C81	1.59 (4)
O5—C42	1.468 (8)	C81—C82	1.44 (4)
O6—C47	1.420 (7)	O17B—C79B	1.54 (3)
O6—C46	1.431 (7)	O17B—C82B	1.66 (3)
O7—C48	1.425 (7)	C79B—C80B	1.41 (3)
O7—C49	1.431 (8)	C80B—C81B	1.39 (3)
O8—C52	1.420 (7)	C81B—C82B	1.63 (3)
O1—B1—O3	109.9 (5)	C45—N2—K1	108.8 (3)
O1—B1—C1	110.1 (5)	C51—N2—K1	109.0 (3)
O3—B1—C1	107.2 (5)	C39—N2—K1	110.6 (4)
O1—B1—C7	108.8 (5)	C56—N3—C50	110.1 (5)
O3—B1—C7	109.8 (5)	C56—N3—C44	109.6 (5)
C1—B1—C7	111.0 (5)	C50—N3—C44	109.8 (5)
O1—B2—O2	122.2 (6)	C56—N3—K1	108.7 (3)
O1—B2—C13	120.2 (6)	C50—N3—K1	110.0 (3)
O2—B2—C13	117.6 (5)	C44—N3—K1	108.7 (4)
O3—B3—O2	123.0 (6)	N2—C39—C40	114.6 (6)
O3—B3—C19	120.5 (5)	O4—C40—C39	109.0 (6)
O2—B3—C19	116.5 (5)	O4—C41—C42	108.7 (6)
B2—O1—B1	123.3 (5)	C41—C42—O5	110.3 (5)
B2—O2—B3	118.5 (5)	O5—C43—C44	108.3 (5)
B3—O3—B1	121.9 (5)	N3—C44—C43	113.9 (5)
C6—C1—C2	116.4 (6)	N2—C45—C46	114.2 (5)
C6—C1—B1	119.4 (6)	O6—C46—C45	109.0 (5)
C2—C1—B1	124.1 (6)	O6—C47—C48	109.3 (5)
C3—C2—C1	123.1 (6)	O7—C48—C47	109.6 (5)
C4—C3—C2	119.0 (6)	O7—C49—C50	109.1 (6)
C3—C4—C5	119.7 (7)	N3—C50—C49	114.6 (5)
C6—C5—C4	119.6 (7)	N2—C51—C52	114.9 (5)
C5—C6—C1	122.3 (7)	O8—C52—C51	109.7 (6)
C8—C7—C12	118.1 (7)	O8—C53—C54	109.1 (5)
C8—C7—B1	123.0 (6)	O9—C54—C53	109.3 (5)
C12—C7—B1	118.9 (6)	O9—C55—C56	108.6 (5)
C9—C8—C7	121.1 (8)	N3—C56—C55	115.0 (5)
C8—C9—C10	121.5 (8)	O11—K2—O13	95.89 (15)
C9—C10—C11	119.7 (7)	O11—K2—O14	139.54 (15)
C10—C11—C12	119.4 (7)	O13—K2—O14	121.34 (15)
C7—C12—C11	120.3 (7)	O11—K2—O15	100.52 (14)
C14—C13—C18	116.2 (6)	O13—K2—O15	101.51 (13)
C14—C13—B2	122.1 (6)	O14—K2—O15	59.70 (12)
C18—C13—B2	121.7 (6)	O11—K2—O10	59.95 (16)
C13—C14—C15	120.1 (7)	O13—K2—O10	138.17 (15)
C16—C15—C14	123.1 (7)	O14—K2—O10	94.97 (15)

C15—C16—C17	118.9 (7)	O15—K2—O10	115.46 (16)
C18—C17—C16	116.3 (7)	O11—K2—O12	112.31 (15)
C17—C18—C13	125.3 (7)	O13—K2—O12	59.27 (15)
C24—C19—C20	118.6 (6)	O14—K2—O12	100.96 (14)
C24—C19—B3	121.8 (6)	O15—K2—O12	142.74 (15)
C20—C19—B3	119.6 (6)	O10—K2—O12	96.54 (17)
C21—C20—C19	121.7 (6)	O11—K2—N5	61.21 (15)
C20—C21—C22	120.0 (7)	O13—K2—N5	61.52 (15)
C23—C22—C21	116.5 (6)	O14—K2—N5	120.95 (13)
C22—C23—C24	122.7 (6)	O15—K2—N5	62.25 (13)
C19—C24—C23	120.4 (7)	O10—K2—N5	119.07 (17)
C26—B4—C25	110.9 (6)	O12—K2—N5	119.12 (15)
C26—B4—C27	113.2 (7)	O11—K2—N4	118.08 (18)
C25—B4—C27	104.2 (6)	O13—K2—N4	118.20 (18)
C26—B4—C33	109.1 (6)	O14—K2—N4	60.00 (14)
C25—B4—C33	106.6 (7)	O15—K2—N4	118.67 (14)
C27—B4—C33	112.5 (6)	O10—K2—N4	60.4 (2)
N1—C25—B4	171.6 (9)	O12—K2—N4	60.39 (17)
C32—C27—C28	115.2 (7)	N5—K2—N4	179.04 (16)
C32—C27—B4	125.9 (7)	C59—O10—C58	107.1 (7)
C28—C27—B4	118.8 (7)	C59—O10—K2	113.4 (4)
C27—C28—C29	120.6 (7)	C58—O10—K2	117.1 (5)
C30—C29—C28	119.6 (8)	C61—O11—C60	110.4 (6)
C31—C30—C29	120.7 (8)	C61—O11—K2	117.1 (4)
C30—C31—C32	120.4 (9)	C60—O11—K2	113.7 (4)
C31—C32—C27	123.3 (8)	C64—O12—C65	110.9 (7)
C34—C33—C38	115.1 (6)	C64—O12—K2	115.9 (4)
C34—C33—B4	120.8 (6)	C65—O12—K2	115.5 (4)
C38—C33—B4	124.2 (6)	C66—O13—C67	111.1 (5)
C35—C34—C33	122.8 (6)	C66—O13—K2	116.6 (4)
C36—C35—C34	121.7 (6)	C67—O13—K2	117.9 (4)
C35—C36—C37	118.3 (7)	C70—O14—C71	112.9 (5)
C36—C37—C38	120.8 (7)	C70—O14—K2	119.3 (4)
C33—C38—C37	121.1 (7)	C71—O14—K2	116.3 (3)
O9—K1—O5	97.61 (12)	C73—O15—C72	112.0 (5)
O9—K1—O6	120.79 (13)	C73—O15—K2	115.3 (3)
O5—K1—O6	136.77 (13)	C72—O15—K2	115.1 (3)
O9—K1—O4	138.46 (13)	C63—N4—C69	109.8 (7)
O5—K1—O4	59.89 (14)	C63—N4—C57	109.4 (7)
O6—K1—O4	95.88 (13)	C69—N4—C57	109.2 (7)
O9—K1—O7	100.40 (12)	C63—N4—K2	109.8 (4)
O5—K1—O7	97.57 (13)	C69—N4—K2	109.7 (4)
O6—K1—O7	59.09 (12)	C57—N4—K2	109.0 (6)
O4—K1—O7	115.96 (14)	C62—N5—C68	111.7 (5)
O9—K1—O8	59.14 (11)	C62—N5—C74	109.2 (6)
O5—K1—O8	116.81 (13)	C68—N5—C74	109.6 (5)
O6—K1—O8	100.66 (11)	C62—N5—K2	110.4 (4)
O4—K1—O8	97.99 (15)	C68—N5—K2	107.7 (4)

O7—K1—O8	140.91 (13)	C74—N5—K2	108.1 (3)
O9—K1—N2	119.17 (13)	N4—C57—C58	113.6 (7)
O5—K1—N2	118.79 (15)	O10—C58—C57	110.6 (9)
O6—K1—N2	60.90 (13)	O10—C59—C60	108.5 (6)
O4—K1—N2	60.26 (16)	C59—C60—O11	109.5 (6)
O7—K1—N2	118.90 (13)	O11—C61—C62	109.9 (6)
O8—K1—N2	61.17 (12)	N5—C62—C61	114.4 (5)
O9—K1—N3	61.21 (12)	N4—C63—C64	114.2 (6)
O5—K1—N3	61.12 (15)	O12—C64—C63	111.1 (8)
O6—K1—N3	118.86 (13)	O12—C65—C66	109.1 (6)
O4—K1—N3	119.57 (16)	O13—C66—C65	110.6 (6)
O7—K1—N3	60.81 (13)	O13—C67—C68	109.4 (5)
O8—K1—N3	119.23 (12)	N5—C68—C67	114.1 (5)
N2—K1—N3	179.61 (15)	N4—C69—C70	112.5 (7)
C41—O4—C40	108.8 (6)	O14—C70—C69	110.9 (6)
C41—O4—K1	116.5 (4)	O14—C71—C72	109.8 (5)
C40—O4—K1	118.1 (4)	O15—C72—C71	109.9 (5)
C43—O5—C42	110.1 (5)	O15—C73—C74	110.1 (5)
C43—O5—K1	117.3 (4)	N5—C74—C73	114.0 (6)
C42—O5—K1	113.6 (4)	C75—O16—C78	105.8 (7)
C47—O6—C46	110.5 (4)	O16—C75—C76	109.3 (9)
C47—O6—K1	117.6 (3)	C75—C76—C77	104.4 (8)
C46—O6—K1	118.1 (3)	C78—C77—C76	102.9 (8)
C48—O7—C49	110.8 (5)	O16—C78—C77	106.3 (7)
C48—O7—K1	115.1 (3)	C79—O17—C82	107.3 (16)
C49—O7—K1	116.0 (3)	O17—C79—C80	101.2 (15)
C52—O8—C53	110.3 (5)	C81—C80—C79	93 (2)
C52—O8—K1	114.9 (3)	C82—C81—C80	106 (3)
C53—O8—K1	114.0 (3)	C81—C82—O17	106 (2)
C54—O9—C55	110.0 (4)	C79B—O17B—C82B	103.9 (17)
C54—O9—K1	118.8 (3)	C80B—C79B—O17B	102.6 (19)
C55—O9—K1	118.7 (3)	C81B—C80B—C79B	113 (2)
C45—N2—C51	110.0 (6)	C80B—C81B—C82B	110.4 (18)
C45—N2—C39	109.9 (5)	C81B—C82B—O17B	92.6 (15)
C51—N2—C39	108.5 (5)		
