

UC Davis

UC Davis Previously Published Works

Title

Environmental variability can prolong parasite co-occurrence within hosts and promote transmission among hosts

Permalink

<https://escholarship.org/uc/item/8mk605bt>

Journal

Theoretical Ecology, 18(1)

ISSN

1874-1738

Authors

Jarvis-Cross, Madeline

Krkošek, Martin

Publication Date

2025-12-01

DOI

10.1007/s12080-025-00613-y

Copyright Information

This work is made available under the terms of a Creative Commons Attribution-NonCommercial License, available at <https://creativecommons.org/licenses/by-nc/4.0/>

Peer reviewed



Environmental variability can prolong parasite co-occurrence within hosts and promote transmission among hosts

Madeline Jarvis-Cross¹ · Martin Krkošek¹

Received: 2 August 2024 / Accepted: 26 March 2025 / Published online: 7 April 2025
© The Author(s), under exclusive licence to Springer Nature B.V. 2025

Abstract

While the mechanisms that govern disease emergence and spread among hosts are increasingly well-described, the mechanisms that promote parasite diversity within-hosts, affecting host outcomes and spillover potential, have been comparatively understudied. Furthermore, while attention has been paid to the effects of increasing temperatures on disease systems, the effects of environmental variability have been left underexplored, despite rising climatic variability. To investigate the impacts of environmental variability on parasite co-occurrence within-hosts, we analysed a model of within-host population dynamics wherein two parasites indirectly compete through the host's immune response. We simulated the model under constant, demographically stochastic, environmentally stochastic, and demographically and environmentally stochastic conditions, and analysed the viability and longevity of non-equilibrium parasite co-occurrence. We found that environmental stochasticity increased the viability and longevity of parasite co-occurrence, suggesting that thermal variability arising from climatic change and as a physiological trait may promote parasite diversity within ectotherms and help explain bats' propensity to support diverse communities of parasites. Further, we found that under certain conditions, the transmissibility of co-occurring parasites can surpass the transmissibility of single parasites, suggesting that thermal variability may increase the transmissibility of co-occurring parasites.

Keywords Within-host · Co-infection · Environmental variability · Co-existence · Modelling

Introduction

It has become clear that climate and land-use change can facilitate novel species interactions and increase contact rates between wild populations and humans, resulting in devastating epidemics and enabling spillover and disease emergence events among wild and human hosts alike (Daszak et al. 2000, 2001; Plowright et al. 2017). There is strong theoretical and empirical evidence that within-host parasite diversity influences disease outcomes in individual hosts (Bromenshenk et al. 2010; Johnson & Hoverman 2012), transmissibility among hosts (Jolles et al. 2008; Susi et al. 2015), and community structure and functioning (Lafferty et al. 2006). However, the mechanisms that promote within-host parasite diversity have been understudied.

Ecological theory predicts a relationship between environmental variability and species diversity. For example, the Intermediate Disturbance Hypothesis (IDH) posits that environmental disturbances to competitive and successional processes will maximize species richness by preventing competitive exclusion (Chesson 1994; Connell 1978). In the case of apparent competition, analogized by immune suppression of two infecting parasites (Holt and Bonsall 2017; Holt and Lawton 1994), theory indicates that intermediate levels of disturbance will promote coexistence when a consumer has a larger impact on the stronger of two competitors (Wootton 1998). More recent work suggests that within-host parasite diversity may be maintained by stabilizing and equalizing mechanisms (Bashey 2015), including niche differentiation (Ramesh and Hall 2023), competition-colonization trade-offs (Dallas et al. 2019), and environmental heterogeneity (Greenrod et al. 2024). Song and Spaak (2024) determined that in lower trophic levels, diversity is primarily constrained by fitness differences, which can be minimized via equalizing mechanisms like environmental variation (Song and Spaak 2024; Spaak et al. 2021). We may extend this finding

✉ Madeline Jarvis-Cross
madeline.jarvis.cross@mail.utoronto.ca

¹ Department of Ecology and Evolutionary Biology,
University of Toronto, 25 Willcocks Street, Toronto,
ON M5S 3B2, Canada

to consider models of within-host population dynamics, wherein the interaction between the host's immune system and infecting parasites is analogous to the interaction between a predator and its prey (Anderson 1994; Antia et al. 1994, 1996; Bell 1973; Nowak and May 2000; Pilyugin and Antia 2000), creating a food web that contains two trophic levels. While coexistence theory concerns the maintenance of equilibrium coexistence, the same mechanisms may play a role in mediating nonequilibrium co-occurrence (Comins and Noble 1985). The impacts of these mechanisms on the maintenance of nonequilibrium co-occurrence may be of particular interest when we consider within-host disease systems, which are often characterised by multiparasite infections (Balmer and Tanner 2011; Petney and Andrews 1998) and seldom exhibit long-term coexistence.

We considered the relationship between environmental variation and the within-host co-occurrence of two parasites along two axes: magnitude and frequency of stochastic disturbance (Mackey and Currie 2001; Miller et al. 2011, 2012). First, we considered rising temperature variability as a feature of climate change (Lawson et al. 2015; Stocker et al. 2013; Vasseur et al. 2014). In ectothermic hosts, increasingly erratic external conditions will affect the internal conditions of the host, thereby also affecting any parasites the host may harbour (Cizauskas et al. 2017). A relationship between thermal variability and within-host parasite diversity may be of particular interest when considered alongside the long-debated relationship between climate change and disease-mediated amphibian declines (Daszak et al. 1999; Johnson 2006; Lips et al. 2008; Rohr et al. 2008). While it has been posited that increasing temperatures may drive these declines, Rohr and Raffel (2010) demonstrated that El Niño events and associated regional temperature variability increase host susceptibility, and thus contribute to host population declines (Cohen et al. 2019; Rohr et al. 2008; Rohr and Raffel 2010). Second, we considered environmental variability as a host trait. Bats (order *Chiroptera*) have long been acknowledged as prolific reservoirs for disease and are unique among mammals in their ability to control body temperature in accordance with activity level (e.g. torpor, flight) (Brook and Dobson 2015; Calisher et al. 2006; Irving et al. 2021; Luo et al. 2021). It has been theorized that thermal variation may contribute to the efficacy of bats' immune systems, allowing them to carry microparasites without suffering increased morbidity (Fumagalli et al. 2021; Luo et al. 2021). Similarly, thermal variability may affect parasite coexistence. In combination, the effects of thermal variability on immune responses and parasite coexistence may help explain bats' propensity to support large communities of parasites.

We used a model to investigate the impacts of environmental variability on the viability and maintenance of nonequilibrium parasite co-occurrence within hosts. We

began by adapting Antia et al.'s (1994) model of within-host population dynamics to include two parasites that indirectly compete via the host's immune system, and thus exhibit apparent competition (Holt 1977; Råberg et al. 2006; Williamson 1957). We analysed deterministic, demographically stochastic, environmentally stochastic, and demographically and environmentally stochastic implementations of the model. Within the environmentally stochastic implementations of the model, we draw on the metabolic theory of ecology to model the parasites' replication rates as temperature-dependent and vary the size and frequency of thermal disturbance within the system. Between and among the deterministic and stochastic implementations of the model, we compared the ability of the secondary parasite to invade a system containing the primary parasite, and the duration of time for which the primary and secondary parasites co-occurred. Further, we compared the total transmission of co-occurring parasites with the total transmission of a single parasite. We found that compared to a constant environment, a variable environment tended to increase the ability of the secondary parasite to invade the system, facilitated longer periods of co-occurrence between the primary and secondary parasites, and increased the transmissibility of co-occurring parasites. These results (1) suggest that climatic changes may have significant effects on the occurrence and longevity of multiparasite infections in ectothermic hosts, (2) provide insight on a host's ability to support large communities of parasites, and propose that variability within-hosts may influence transmission dynamics among-hosts.

Methods

Model description

Within-host population dynamics model

We began with a model of within host population dynamics based on Antia et al. (1994) that represents the interaction between an infecting parasite and its host's immune system as analogous to the interaction between a prey item and its predator. Per Antia et al. (1994), the density of the parasite population, P , is determined by its replication rate, r , and the rate at which it is cleared by the host's immune system, k . The model assumes that the immune system's replication rate is proportional to parasite density at low parasite densities and saturates at high parasite densities, allowing the host's immune system to either extirpate or succumb to the infection (Antia et al. 1994). We adapted the model to include a second parasite, P_j , that indirectly competes with the primary parasite, P_i , through the host's immune response (Eqs. 1, 2 and 3, Table 1). As such, the model

Table 1 Parameter definitions and values

Symbol	Parameter	Values stated in Antia et al. (1994)	Values used
P_i and P_j	Parasite abundance	1*	1*
I	Host immune cell abundance	1*	1*
r_i and r_j	Replication rates of parasites	0.1–10.0	0.1–2.0 [†]
k	Rate at which parasites are destroyed by host's immune system	10^{-3}	10^{-2}
ρ	Maximum replication rate of immune cells	1.0	1.0
ϕ	Parasite density at which replication rate of immune cells is half its maximum	10^3	10^3
D	Lethal within-host parasite density	10^9	10^9

*Initial conditions

[†]Dependent on the temperature of system in environmentally stochastic simulations

describes an immune-mediated apparent competition (Holt 1977; Råberg et al. 2006; Williamson 1957). Parameter values were not chosen to describe a specific system, but to produce the dynamical behaviour described above and shown in Fig. 1. Given that any number of parameter value combinations may produce this behaviour, we chose to maintain the initial conditions and select parameter values

proposed by Antia et al. (1994). However, we increased the rate at which parasites are destroyed by the host's immune system (k) and decreased the maximum replication rates of the primary and secondary parasites (r_i, r_j) to favour extirpation and eliminate host death as an infection outcome (Table 1). We also conducted a sensitivity analysis to determine how variations in (1) the value of k and (2)




Fig. 1 A schematic detailing simulation experiment. When simulating each of four models (deterministic, demographically stochastic, environmentally stochastic, and demographically and environmentally stochastic), we considered four thermal scenarios (identical responses, offset responses, P_i -advantage, and P_j -advantage). When simulating the environmentally stochastic models (in blue), we con-

sidered fifteen sizes and thirteen frequencies of disturbance, totalling 195 unique disturbance regimes. We additionally considered fifty days on which the secondary parasite could be introduced. We conducted one thousand simulations per stochastic model and treatment. Examples of such simulations are shown as time series

the shape of the relationship between parasite replication rate and the temperature of the system would affect our results and found the model to be robust (see pages 2, 4–8 of supplement for complete results).

$$P_i(t+1) = (r_i + e^{-kI(t)})P_i(t) \quad (1)$$

$$P_j(t+1) = (r_j + e^{-kI(t)})P_j(t) \quad (2)$$

$$I(t+1) = (1 + \rho \left(\frac{P_i(t)}{P_i(t) + \phi} + \frac{P_j(t)}{P_j(t) + \phi} \right))I(t) \quad (3)$$

To compare the impacts of different types of stochasticity on parasite diversity, we wrote deterministic (D), demographically stochastic (S_D), environmentally stochastic (S_E), and demographically and environmentally stochastic (S_{DE}) implementations of the model. Within the demographically stochastic implementation of the model, we wrote the model's replication terms ($r_i \cdot P_i(t)$, $r_j \cdot P_j(t)$, and $\rho \left(\frac{P_i(t)}{P_i(t) + \phi} + \frac{P_j(t)}{P_j(t) + \phi} \right) I(t)$) as Poisson processes, and the parasite mortality term ($1 - e^{-kI(t)}$) as a binomial process (Eqs. 4, 5 and 6).

$$P_i(t+1) = P_i(t) + B_{P_i}(t) - D_{P_i}(t) \quad (4)$$

$$P_j(t+1) = P_j(t) + B_{P_j}(t) - D_{P_j}(t) \quad (5)$$

$$I(t+1) = I(t) + B_I(t) \quad (6)$$

In Eqs. (4) and (5), $B_{P_i}(t)$ and $B_{P_j}(t)$ are Poisson random variables such that $B_{P_i}(t) \sim \text{Poisson}(\lambda_i)$, where $\lambda_i = r_i \cdot P_i(t)$, and $B_{P_j}(t) \sim \text{Poisson}(\lambda_j)$, where $\lambda_j = r_j \cdot P_j(t)$. In Eq. (6), $B_I(t)$ is a Poisson random variable such that $B_I(t) \sim \text{Poisson}(\lambda)$, where $\lambda = \rho \left(\frac{P_i(t)}{P_i(t) + \phi} + \frac{P_j(t)}{P_j(t) + \phi} \right) I(t)$. In Eqs. (4) and (5), $D_{P_i}(t)$ and $D_{P_j}(t)$ are binomial random variables such that $D_{P_i}(t) \sim \text{Binomial}(P_i(t), p)$, where $p = 1 - e^{-kI(t)}$, and $D_{P_j}(t) \sim \text{Binomial}(P_j(t), p)$. To write the environmentally stochastic implementation of the model, we used randomly generated temperature series to dictate the temperature of the system (T), and allowed the replication rates of each parasite (r_i , r_j) to vary in response (Eqs. 7, 8 and 9). We chose to write r_i and r_j , though not k , as temperature-dependent to isolate the effect of thermal variability on traits that can be metabolically scaled and constrain the scope of the work. We did not write ρ or ϕ as temperature-dependent to constrain the scope of the work.

$$P_i(t+1) = (r_i(T(t)) + e^{-kI(t)})P_i(t) \quad (7)$$

$$P_j(t+1) = (r_j(T(t)) + e^{-kI(t)})P_j(t) \quad (8)$$

$$I(t+1) = (1 + \rho \left(\frac{P_i(t)}{P_i(t) + \phi} + \frac{P_j(t)}{P_j(t) + \phi} \right))I(t) \quad (9)$$

We used first-order autocorrelated Gaussian noise to generate temperature series. We initiated the series with a value of zero, and generated the next value by multiplying the previous value by the coefficient of autocorrelation (0.25), and summing the resulting value with a value drawn from a normal random variable with a mean of zero. We varied the standard deviation of the normal random variable to control the size of disturbance, as described below. We then added the average temperature of the system to each value in the series. Resulting values outside the thermal range of the system were re-drawn. We then converted the temperature series into replication rates using various parameterizations of the Sharpe-Schoolfield sub-model, as described below (Eqs. S1.1, S1.2; Table S2).

Sharpe-Schoolfield sub-model

The metabolic theory of ecology (MTE) predicts how metabolic rates control ecological processes across levels of organization and explains how metabolic rates vary with body size and temperature (Brown et al. 2004). Building off the MTE, the Boltzmann-Arrhenius (BA) relation describes metabolism as a function of temperature. The Sharpe-Schoolfield model characterizes the performance of a trait as a function of temperature through a thermal performance curve (TPC) (Dell et al. 2011; Molnár et al. 2017; Schoolfield et al. 1981). When applied to disease systems, the Sharpe-Schoolfield model has been shown to describe relationships between temperature and species traits and has proven useful in investigating the relationship between rising temperatures and disease transmission (Kirk et al. 2018; Molnár et al. 2013).

We used various parameterizations of the Sharpe-Schoolfield model to generate paired thermal performance curves relating the temperature of the system to the replication rates of two parasites (r_i and r_j) (Fig. S9). We generated four paired thermal performance curves, representing four thermal scenarios: (1) the identical responses scenario, (2) the offset responses scenario, (3) the P_i -advantage scenario, and (4) the P_j -advantage scenario (Fig. 1, Table S2). The identical responses scenario describes r_i and r_j as responding identically to the temperature of the system and exhibits an average temperature of 19.9 °C (Fig. 1, Table S2, Fig. S9). The remaining scenarios describe a system within which P_i exhibits a higher replication rate at lower temperatures and P_j exhibits a higher replication rate at higher temperatures and are differentiated by the average temperature of the system. The offset responses scenario exhibits an average temperature of 19.9 °C. At this temperature, the replication

rate of P_i is equal to that of P_j (Fig. 1, Table S2, Fig. S9). The identical and offset responses scenarios share an environment and specify different thermal responses, enabling us to examine how species' responses drive competitive outcomes. The P_i -advantage scenario exhibits an average temperature of 18 °C, giving P_i a thermal advantage over P_j (Fig. 1, Table S2, Fig. S9). We considered two versions, (I) and (II), of the P_j -advantage scenario, each of which give P_j a thermal advantage over P_i . (I) exhibits an average temperature of 20.9 °C, such that P_j 's thermal advantage is comparable in size to that of P_i , as described above (Fig. 1, Table S2, Fig. S9). (II) replicates the P_i -advantage scenario but re-assigns the low-temperature curve to P_j and the high-temperature curve to P_i , to isolate the effects of asymmetry (Fig. S9). Given that (I) and (II) produced qualitatively comparable results, we will only discuss (I) in the main text (see supplement for complete results). The offset responses scenario, the P_i -advantage scenario, and the P_j -advantage scenario specify different environments and share thermal responses, enabling us to examine how competitive differences benefit or disadvantage one parasite relative to another in different environments.

Given that the Sharpe-Schoolfield model cannot generate negative values, it cannot be used to explore cases in which extreme temperatures result in parasite mortality and population decline. To compensate, we developed an alternative set of paired thermal performance curves, through which extreme temperatures produce “negative” replication rates, and spur population decline (see “Extended analysis” in supplement; Fig. S19).

Model simulations

Within each thermal scenario, we simulated each model (D , S_D , S_E , S_{DE}) for fifty days. Simulating for fifty days proved sufficient to capture the full course of parasite infection and clearance across all treatments. We began each set of simulations with the primary parasite and the host's immune system in the system, at initial abundances of one. We then considered all cases for the introduction of the secondary parasite from its introduction at the onset of infection, to its introduction on the forty-ninth day of infection.

For each environmentally stochastic model (S_E , S_{DE}) simulation, we generated a new temperature series and two corresponding series of replication rates. When determining how new temperatures would be selected, we considered two axes of thermal disturbance: size and frequency. We controlled the size of disturbance by changing the standard deviation of the normal random variable in the temperature selection function, starting from two and increasing by two until the effect was fully characterized at a value of thirty, resulting in fifteen sizes of disturbance (Fig. 1).

We considered standard deviations between two and ten to denote small disturbances, between twelve and twenty to denote intermediately sized disturbances, and between twenty-two and thirty to denote large disturbances (Fig. 1). To control the frequency of disturbance, we re-assigned the temperature of the system every ten, nine, eight, seven, six, five, four, three, and two days, once per day, and two, five, and ten times per day to fully characterise the effect, resulting in thirteen frequencies of disturbance (Fig. 1). We considered re-assignments every ten to seven days to denote infrequent disturbances, every six to two days to denote intermediately frequent disturbances, and every day to ten times per day to denote frequent disturbances (Fig. 1). Considering size and frequency together resulted in 195 unique disturbance regimes. We conducted one thousand simulations per stochastic model and treatment (Fig. 1). Two hundred such simulations are shown in Figs. 1 and S10.

Co-occurrence analysis

Given successful invasion by the secondary parasite ($P_j(t+1) > P_j(t)$), we defined longevity of co-occurrence as the amount of time for which both parasites exhibited population abundances equal to or greater than one.

We summarised these results as distributions for each stochastic model and treatment. Using the S_D model as a baseline, we investigated the effects of environmental variability on the viability and maintenance of parasite diversity by comparing the ability of the secondary parasite to invade a single-parasite system, and the duration of time for which the primary and secondary parasites co-occurred across stochastic treatments. Additionally, we analysed (1) the proportion of stochastic simulations per model and treatment that resulted in longer periods of co-occurrence relative to the deterministic model, and (2) the competitive outcomes of each treatment to gain insight on underlying mechanisms.

Transmissibility analysis

To explore the impacts of co-infection on transmission potential, we compared the lifetime replicative outputs of co-occurring parasites with those of a single parasite in constant and variable environments. As per Antia et al., (1994), we determined the total transmission (q_i , q_j , q , where q is the total transmission of a single parasite) of a parasite (P_i , P_j , P , where P is a single parasite) from an infected host by taking the integral of the rate of transmission (u) of each parasite from the host over the duration of infection (Eqs. 10, 11 and 12). For simplicity, we set the rate of transmission to one.

$$q_i = \int_0^{\infty} uP_i(t)dt \quad (10)$$

$$q_j = \int_0^{\infty} uP_j(t)dt \quad (11)$$

$$q = \int_0^{\infty} uP(t)dt \quad (12)$$

To compare constant and variable environments, we analysed multi-parasite and single-parasite versions of the S_D and S_{DE} models.

Results

Analytically, the secondary parasite could invade ($P_j(t+1) > P_j(t)$) when $I(t) < \frac{-\ln(1-r_j)}{k}$. Our parameterization of the S_D model allowed the secondary parasite to invade when introduced before the tenth to twelfth day of infection, resulting in 8.4 to 9.2 days of co-occurrence, depending on thermal scenario (Fig. S11, Supplementary Datafile 1–1:10). After this point, which we will refer to as the invasion threshold, and assuming the primary parasite had not yet been extirpated, the immune response stimulated by the primary parasite prevented invasion by the secondary parasite.

Co-occurrence analysis

Simulation outputs show that compared to a deterministic baseline, the addition of demographic stochasticity has little effect on co-occurrence (Supplementary Datafile 2–1:10), while the addition of thermal variability increases the ability of the secondary parasite to invade the system and facilitates longer periods of co-occurrence between parasites. While longevity of co-occurrence are maximised under the P_j -advantage scenario, the difference between deterministic and environmentally stochastic implementations is not consistently maximised by any one thermal scenario. Additionally, while larger and more frequent disturbances tend to promote longer periods of co-occurrence, increasingly large frequent disturbances produce a less steep effect than do increasingly frequent disturbances. These findings are common among thermal scenarios.

When both parasites were introduced simultaneously, S_D model simulations resulted in periods of co-occurrence that were, on average, 4.5% to 3.1% shorter (0.4 to 0.3 days) than deterministically simulated periods of co-occurrence (Figs. 2, S12). In comparison, simulations of the environmentally stochastic models (S_E , S_{DE}) resulted in periods of co-occurrence that were up to 46.2% (S_E) and 36.0% (S_{DE})




Fig. 2 The mean percent difference in longevity of co-occurrence relative to deterministically simulated periods of co-occurrence, when both parasites are introduced at the same time. Single points above “ S_D ” labels represent the mean percent difference in longevity of co-occurrence relative to deterministic thresholds across one thousand demographically stochastic simulations. Each point above “ S_E ”/“ S_{DE} ”

labels represents the mean percent difference in longevity of co-occurrence relative to deterministic thresholds across one thousand environmentally stochastic simulations. As such, there are 195 points above each “ S_E ”/“ S_{DE} ” label, corresponding to 195 environmental disturbance regimes. The small, horizontal black lines denote the 25th, 50th, and 75th percentiles

longer (3.9 (S_E) and 3.0 (S_{DE}) days) than deterministically simulated periods of co-occurrence (Figs. 2, S12). Across all 195 disturbance regimes, simulations of the S_E model resulted in mean periods of co-occurrence that were between 2.7% shorter (0.2 days) and 46.2% longer (3.9 days) than their deterministic counterparts (Figs. 2, S12). The identical responses scenario promoted the shortest and longest mean periods of co-occurrence (Figs. 2, 3, S12, S13). Simulations of the S_{DE} model resulted in mean periods of co-occurrence that were between 6.8% shorter (0.6 days) and 36.0% longer (3.0 days) than deterministically simulated periods of co-occurrence (Figs. 2, S12). Again, the identical responses scenario promoted the shortest and longest mean periods of co-occurrence (Figs. 2, 3, S12, S13). Across thermal scenarios, mean periods of co-occurrence were shortest when disturbances were large (standard deviation of the normal random variable in the temperature autocorrelation function = 22 to 30) and infrequent (once every seven to nine days) (Figs. 3, S13). Mean periods of co-occurrence were longest when disturbances were large ($SD = 24$ to 30) and frequent (five to ten times per day) (Figs. 3, S13).

More generally, as disturbances increased in size, periods of co-occurrence increased in length towards a saturation point (standard deviation of the normal random variable in the temperature autocorrelation function ≈ 24 –30, depending on thermal scenario) (Figs. S14, S18, Supplementary Datafiles 3–1:5, 4–1:5). However, an analysis of simulations within which an alternative set of TPCs allowed for extreme

temperatures to cause population decline revealed a parabolic relationship between disturbance size and longevity of co-occurrence, that peaked when disturbances were intermediately sized (standard deviation of the normal random variable in the temperature autocorrelation function ≈ 8 –16) (Fig. S20).

When disturbances were intermediately frequent (once every two to six days), periods of co-occurrence were similar in length to deterministically simulated periods of co-occurrence. Interestingly, when disturbances were infrequent (once every six to ten days), mean periods of co-occurrence were shorter than deterministically simulated periods of co-occurrence (Fig. S14, S21). Per the deterministic model, P_i and P_j can only co-occur for 8.4 to 9.2 days. Thus, a system subjected to infrequent disturbances is functionally similar to the deterministic system, making the outcome of infection reliant on initial conditions. When disturbances were frequent (ten times per day to once per day), mean periods of co-occurrence were longer than deterministically simulated periods of co-occurrence. Within a given disturbance regime, longevities of co-occurrence were normally distributed (Figs. S14–S15, S23).

Rarity and competitive outcomes

When the secondary parasite was introduced before or on the invasion threshold (tenth or twelfth day of infection, depending on thermal scenario), 29%–71% (S_E) and




Fig. 3 Fully factorial representations of the impacts of varying sized and timed disturbances on the longevity of co-occurrence when P_i and P_j are introduced at the same time. The plot shows the mean number of time steps (ten time steps per day) for which P_i and P_j have

population abundances greater than one as per the S_{DE} model. The circled numbers denote the number of time steps for which P_i and P_j can co-occur, per the deterministic model

46%–58% (S_{DE}) of environmentally stochastic simulations resulted in longer periods of co-occurrence relative to deterministic simulations (Fig. 4). When the secondary parasite was introduced after the invasion threshold, the likelihood that an environmentally stochastic simulation resulted in some period of co-occurrence sharply declined. When the secondary parasite was introduced one to twelve days after the invasion threshold, 1%–31% (S_E) and 1%–46% (S_{DE}) of environmentally stochastic simulations resulted in some period of co-occurrence (Fig. 4). Thus, environmental variability extended the invasion threshold by 50%–75% (S_E) and 100%–120% (S_{DE}), depending on thermal scenario (Fig. 4).

Analysing competitive outcomes at intermediate levels of disturbance ($SD = 10$, frequency of disturbance = once per day) revealed that under the offset responses scenario, when P_i and P_j were introduced at the same time, P_j outlasted P_i 65.5% (S_E) and 63.4% (S_{DE}) of the time (Fig. S22–S23). Additionally, under the P_i -advantage scenario, when P_i and P_j were introduced at the same time, P_i and P_j were equally likely to outlast one another (P_j outlasts P_i 49.9% (S_E) and 51.0% (S_{DE}) of the time), but under the P_j -advantage, P_j was much more likely to outlast P_i (P_j outlasts P_i 76.6% (S_E) and 70.4% (S_{DE}) of the time) (Fig. S21–S23). Across thermal scenarios, when the secondary parasite, P_j , was introduced in the middle of the infection period, P_i was more likely to outlast P_j . When P_j was introduced late in the infection period, P_j was more likely to outlast P_i (Fig. S22–S23). Regardless of when P_j was introduced, simulations resulting in especially long periods of co-occurrence are equally likely to show either parasite outlasting the other (Fig. S22–S23).

Transmissibility analysis


Across thermal scenarios, simulations of the multi-parasite S_D model produced lifetime replicative outputs (LROs) that were one to two orders of magnitude lower than those produced by simulations of the single-parasite S_D model

(Supplementary Datafiles 5–1:16). Similarly, under the first three thermal scenarios, simulations of the multi-parasite S_{DE} model produced LROs that were one to two orders of magnitude lower than those produced by simulations of the single-parasite S_{DE} model. However, under the P_j -advantage scenario and among 62 of 195 (32%) disturbance regimes, the average transmission potential of the co-occurring secondary parasite, P_j , exceeded that of a single parasite (Fig. 5). Among these disturbance regimes, the transmission potential of P_j exceeded that of a single parasite by an average of 2% to 135% (Supplementary Datafiles 5–8). The transmission potentials of co-occurring parasites peaked when disturbances were infrequent (once every ten days) to intermediately frequent (once every six days) (Fig. S24). While intermediately large and frequent disturbances maximally increased longevity of co-occurrence, such infections did not result in particularly high transmission potentials, indicating that especially long periods of co-occurrence may be facilitated by lower lifetime replicative outputs that do not trigger a large immune response, and thus allow for more persistent, but less transmissible, infections.

Discussion

We investigated the potential for environmental variability to affect the viability and maintenance of within-host parasite co-occurrence. Our work provides theoretical evidence that relative to a constant environment, a variable environment can increase the ability of a secondary parasite to invade a single-parasite system, promote longer periods of co-occurrence between parasites, and increase the transmissibility of co-occurring parasites. In more detail, thermal disturbances of various sizes and frequencies allowed a secondary parasite to invade a single-parasite system up to twelve days after the deterministic invasion threshold, extending the threshold by 100% to 120% (S_{DE} model, Fig. 4). Large and frequent

Fig. 4 Proportion of environmentally stochastic simulations ($SD = 10$, frequency of disturbance = once per day) resulting in longer periods of co-occurrence relative to deterministic simulations by the day on which the secondary parasite was introduced. Vertical lines show the first day on which the introduction of the secondary parasite does not result in any co-occurrence between the two parasites, as per the deterministic model (D)






Fig. 5 **A** The impacts of environmental variability on transmission potential under the P_j -advantage scenario. The plot shows a fully factorial representation of the impacts of thermal disturbance on the transmission potential of P_j , relative to the equivalent single-parasite treatment. The black line indicates disturbance regimes under which the average transmission potential of P_j was equal to that of a single parasite. Colours warmer than denoted by the black line indicate disturbance regimes under which the average transmission potential of P_j exceeded that of a single parasite; colours cooler than denoted by the black line indicate the opposite. **B**, **C** Simulated time series

thermal disturbances enabled parasites to co-occur for up to 37.8% longer, relative to a constant environment (S_{DE} model, Fig. 2, S12). Under the P_j -advantage scenario and among 32% disturbance regimes, the average transmission potential of the co-occurring secondary parasite exceeded that of a single parasite (S_{DE} model, Fig. 5). Finally, the disturbance regimes that enabled longer periods of co-occurrence differed from the disturbance regimes that promoted increased transmission potential (Fig. 5).

Across thermal scenarios, the relationship between disturbance size and longevity of co-occurrence was initially

under large/frequent ($SD = 30$, frequency = ten times per day), and large/infrequent ($SD = 30$, frequency = every nine days) disturbance regimes. Abundances of the primary and secondary parasites are shown in yellow and green, respectively. Notably, large/frequent disturbance regimes resulted in prolonged co-occurrence and decreased transmission potential relative to a single parasite, while large/infrequent disturbance regimes resulted in limited co-occurrence and increased transmission potential relative to a single parasite. In **B**, parasite populations are shown to just exceed $8 \cdot 10^5$. In **C**, this value is denoted by a dotted red line

positive and eventually saturating (Figs. S14, S18). Large disturbances produced a saturating response rather than the expected parabolic response due to the structure of the Sharpe-Schoolfield sub-model. Allowing for extreme temperatures to result in parasite mortality and population decline produced a parabolic response, implying that intermediate sizes of disturbance maximised the viability and longevity of parasite co-occurrence (Figs. S19–S20). When thermal disturbances were small, the parasite with the higher initial replication rate was more likely to grow quickly and trigger a commensurate immune response that

would competitively exclude the other. As thermal disturbances increased in size, it became increasingly likely that when the temperature of the system changed, the competitive advantage would shift from one parasite to the other, minimizing fitness differences over the course of an infection and promoting co-occurrence. However, per the extended analysis, large thermal disturbances caused parasite populations to alternatively explode and crash, triggering a sizable immune response, increasing the likelihood of extirpation, and precluding co-occurrence.

The relationship between climate change and disease-mediated amphibian declines has long been debated. While some argue that climatic changes may impede immune function or result in performance gaps between parasites and their hosts, making amphibians more susceptible to infection, others argue that insufficient evidence should preclude a causal link (Cohen et al. 2017, 2019, 2020; Rohr et al. 2008; Rohr and Raffel 2010). While much of the literature proposes mean-climate hypotheses, some evidence suggests that intensifying El Niño events are increasing thermal variability, which may in turn leave amphibians more susceptible to or less able to defend against infection (Rohr and Raffel 2010). Independently, our findings add another piece to the puzzle by providing theoretical evidence that thermal variability may also promote the viability and maintenance of within-host parasite diversity. As such, attention should be paid to shifts in microparasite communities and their impacts on ectothermic hosts. In particular, well-documented climate-driven increases in disease incidence among a wide variety of marine organisms may constitute an important starting point for future work (Bates et al. 2009; Bruno et al. 2007; Tracy et al. 2019).

Across thermal scenarios, low and high frequencies of disturbance impeded and promoted the viability and longevity of parasite co-occurrence, respectively (Figs. 3, S13–S14, S21). Given the transient nature of infection specified by the underlying model, infrequent disturbances (once every 7 to 10 days) resulted in the temperature of the system being disturbed towards the end of the infection period, when parasite populations were small and especially susceptible to unfavourable conditions. Conversely, frequent disturbances (ten times per day to once per day) resulted in the temperature of the system being disturbed throughout the duration of the infection period, and as such, neither favourable nor unfavourable conditions lasted long enough for one parasite to gain a decisive advantage over and exclude the other, and thus promoted extended periods of co-occurrence. While it has been theorized that internal temperature variation may aid bats' immune responses, allowing them to carry microparasites without suffering outsized morbidity (Fumagalli et al. 2021), these findings suggest that internal temperature variation may also facilitate parasite co-occurrence within

bats, and thus may help explain bats' propensity to support diverse communities of parasites.

Under the offset responses scenario, P_j often outlasted P_i , while under the P_i -advantage scenario, P_i and P_j were equally likely to outlast each other (Figs. S22, S23). Under the P_j -advantage scenario, P_j was much more likely to outlast P_i (Figs. S22, S23). These outcomes are consequences of the asymmetrical structure of the thermal performance curves, which made unfavourable conditions more disadvantageous for P_i than P_j (Sinclair et al. 2016). In more detail, Sharpe-Schoolfield thermal performance curves are characterised by a shallow slope to the left of the optimum, and a steep slope to the right of optimum. Given that P_i exhibits a lower thermal optimum than P_j , temperatures to the right of its thermal optimum corresponded to lower growth rates, than did temperatures to the left of P_j 's thermal optimum. We explored a scenario in which P_j exhibits a lower thermal optimum than P_i (P_j -advantage scenario (II)), and identified qualitatively comparable results (see supplement for complete results). Further, regardless of when P_j was introduced, simulations resulting in especially long periods of co-occurrence were equally likely to show either parasite outlasting the other. Taken together, these results suggest that to predict the competitive outcome of a multiparasite infection, one may need to consider the interplay between how well adapted a parasite is to its host's internal environment, priority effects, and the relative responses of biological rates to environmental factors. While one might expect a specialist or well-adapted parasite (e.g. P_i under the P_i -advantage scenario) to exploit its host more efficiently than a generalist, poorly-adapted, or novel parasite (e.g. P_j under the P_i -advantage scenario), the results of this work suggest that environmental variability can even the scales, and allow an underdog to establish and persist, as predicted by the intermediate disturbance hypothesis (Chesson 1994; Connell 1978; Roxburgh et al. 2004). Similarly, while one might additionally expect a colonizing parasite to benefit from an initial lack of competition, the results of this work again suggest that environmental variability and disturbance may contribute to the structuring of communities by allowing latecomers to establish and persist more often and for longer than they otherwise would (Jiang and Patel 2008; Symons and Arnott 2014).

While the MTE posits that certain biological rates (e.g. metabolic rate, replication/growth rate) will exhibit an asymmetrical and unimodal response to temperature, other vital rates may exhibit symmetrical or bimodal responses to temperature (Dell et al. 2014; Molnár et al. 2017; Sinclair et al. 2016). This work suggests that the shape of a thermal performance curve can meaningfully affect the nature and outcome of an infection. The shape of a thermal performance curve can also dictate the effects of stochasticity

on persistence. Here, thermal variation pushes the system away from a parasite's optimal temperature, and thus, away from its maximum replication rate. In accordance with Jensen's inequality, compared to a deterministic system, the introduction of stochasticity results in a lower average population growth rate (Denny 2017; Jensen 1906; Vasseur et al. 2014). In the absence of thermal dependency, the introduction of stochasticity will also result in a lower average population growth rate over time (Tuljapurkar 1990). In this system, lower average parasite replication rates slow down the infection and subsequently, the immune response, allowing parasites to co-occur for longer. As anthropogenic changes alter host-parasite interaction networks, understanding how evolutionary and environmental factors combine to affect interactions between parasites will be essential in predicting and managing spillover and emergence in wildlife and human populations (Paull et al. 2012; Pedersen and Fenton 2007).

Developing an understanding of how the impacts of environmental variability on parasite co-occurrence within hosts may scale to affect among-host processes and events, including transmission, spillover, and emergence will be essential to the maintenance of wildlife, human, and ecosystem health (Allocati et al. 2016; Halpin et al. 2000; Johnson 2006). While increased within-host parasite diversity provides more strains or species of a parasite or parasites an opportunity to transmit, within-host competition may result in decreased parasite loads, thus decreasing the transmission probability of any one strain or species, and altering expected transmission patterns (Johnson et al. 2013; Pedersen and Fenton 2007). Similarly, our findings provide theoretical evidence that when environmental variations are large and frequent, within-host competition decreases the total transmission of co-occurring parasites. As such, while large and frequent disturbances may maximally increase the amount of time for which two parasites co-occur, such infections do not result in particularly high transmission potentials, indicating that especially long periods of co-occurrence may be facilitated by lower parasite replication rates that do not trigger a large immune response, and thus allow for more persistent infections. However, we found that approximately a third of tested disturbance regimes allowed the replicative outputs of co-occurring parasites to exceed those of single parasites, indicating that the effects of environmental variability may not be limited to the prolonging of within-host parasite co-occurrence, but may also promote transmission between hosts (Fig. 5, S24). Disturbances of intermediate size and low to intermediate frequency may constitute a sizable region within which environmental variation can decrease fitness differences between parasites and promote within-host parasite diversity, while allowing parasites to achieve sufficiently high parasite loads, resulting in more diverse and transmissible infections (Fig. 5, S24).

While both Antia et al.'s (1994) model and our modified model should be broadly applicable to an array of within-host multi-parasite systems, including multi-strain or cross-species, both models are relatively simple as to remain generalizable, and thus cannot represent chronic infections, systems within which parasites exhibit more complex interactions with their host or their host's immune system, nor systems within which parasites directly interact with one another. As such, it may be interesting to validate the theoretical insights of this work in an empirical system. Similarly, this work examines the viability and longevity of two-parasite co-occurrence as a proxy for the maintenance of within-host parasite diversity. While our findings suggest that environmental variability may, in this case, promote parasite diversity, it is interesting to consider the impacts of environmental variability on the viability and longevity of parasite co-occurrence between more than two strains or species. The co-occurrence of more than two strains, may, for example, trigger a more substantial immune response, making it harder for additional strains to invade (Cox 2001; Rigaud et al. 2010). In such a case, we would only expect environmental variability to promote co-occurrence when invading parasites arrive early and experience relatively substantial competitive (here, thermal) advantages.

This work is also limited by the scope with which thermal dependency was considered. Given the thermal sensitivity of immune function in ectotherms and endotherms, it may be worthwhile to consider the thermal dependency of immune processes like response time and replication rate in tandem with the thermal dependency of parasite replication rates (Butler et al. 2013; Wright and Cooper 1981). For example, if the replication rate of the immune system (ρ) peaks at some optimal temperature (Molnár et al. 2017), nonlinear averaging would predict lower values of ρ , or, immune suppression (Rohr and Raffel 2010), in a variable environment. Relatedly, using nonlinear averaging to assess the response of a nonlinear process to variability is contentious. While many theoretical works and some empirical studies support the use of nonlinear averaging and find evidence for Jensen's inequality, borne out as disproportionate effects of thermal variability on performance, many empirical studies have contested an implicit assumption of nonlinear averaging: that traits should respond instantaneously to change, without acclimation (Bernhardt et al. 2018; Estay et al. 2011; Krichel et al. 2023; Sinclair et al. 2016). Thermal acclimation through phenotypic plasticity alters thermal performance curves, and as such, produces a more complex relationship between temperature and biological rates that may not be well-characterized by nonlinear averaging (Schulte et al. 2011; Sckrabulis et al. 2022). With regard to host-parasite interactions, the temperature variability hypothesis posits that owing to their small relative size, parasites should acclimate more quickly than their hosts, though the speed

at which various parasites may acclimate remains unclear (Raffel et al. 2013; Rohr et al. 2013). Interestingly, some empirical studies of the impacts of environmental variability on host-parasite interactions have considered thermal acclimation, yielding a range of conclusions including decreased pathogen prevalence, reduced infection prevalence, and decreased resistance in hosts (Dallas and Drake 2016; Duncan et al. 2011; Raffel et al. 2013).

Ecological theory has long predicted that environmental variability supports species diversity, from the IDH (Chesson 1994; Connell 1978; Huston 2014; Roxburgh et al. 2004) to modern coexistence theory (Barabás et al. 2018; Chesson 1994, 2000, 2003; Chesson & Huntly 1997). Work on trophic interactions, wherein apparent competition is considered analogous to interactions between the host's immune system and infecting parasites (Holt and Bonsall 2017; Holt and Lawton 1994), indicates that intermediate levels of disturbance can promote coexistence (Wootton 1998) and that within lower trophic levels (e.g. among competing parasites), diversity is primarily constrained by fitness differences, which can be minimized via equalizing mechanisms like environmental variation (Song and Spaak 2024). Here, we provide theoretical evidence that the predicted relationship between environmental variability and species diversity is borne out in within-host disease systems. Camus wrote that plagues, despite their commonality, find us unprepared (Camus 1991). Improving our understanding of how environmental variability affects parasite diversity within- and between-hosts will expand our understanding of the mechanisms that can facilitate disease spread and emergence, and allow us to make robust predictions, and develop and implement mitigation strategies as we move into an increasingly uncertain future. In brief, we will be found *less* unprepared.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s12080-025-00613-y>.

Acknowledgements We thank a number of anonymous reviewers, who, through their generous comments, greatly improved the quality of the work. This work was supported by The Emerging and Pandemic Infections Consortium Doctoral Award (MJ-C), and Ontario Graduate Scholarship (MJ-C), a Natural Sciences and Engineering Research Council of Canada Discovery Grant (Grant Number RGPIN- 2024-05054) (MK), and a Canada Research Chair (MK).

Author contributions MJC and MK conceived the ideas and designed the methodology; MJC collected the data; MJC analysed the data; MJC led the writing of the manuscript. All authors contributed critically to drafts and gave final approval for publication.

Funding This work was supported by The Emerging and Pandemic Infections Consortium Doctoral Award (MJ-C), an Ontario Graduate Scholarship (MJ-C), a Natural Sciences and Engineering Research Council of Canada Discovery Grant (Grant Number RGPIN- 2024-05054) (MK), and a Canada Research Chair (MK).

Data availability Time series data available from the Federated Research Data Repository at <https://doi.org/https://doi.org/10.20383/103.0764>.

Scripts and replication materials available from GitHub at https://github.com/MadelineJC/EnvironmentalVariability_Repo.

Declarations

Competing interests The authors declare no competing interests.

References

- Allocati N, Petrucci AG, Di Giovanni P, Masulli M, Di Ilio C, De Laurenzi V (2016) Bat–man disease transmission: zoonotic pathogens from wildlife reservoirs to human populations. *Cell Death Discov* 2(1):16048. <https://doi.org/10.1038/cddiscovery.2016.48>
- Anderson RM (1994) Mathematical studies of parasitic infection and immunity. *Science* 264(5167):1884–1886. <https://doi.org/10.1126/science.8009218>
- Antia R, Levin BR, May RM (1994) Within-host population dynamics and the evolution and maintenance of microparasite virulence. *Am Nat* 144(3):457–472. <https://doi.org/10.1086/285686>
- Antia R, Koella JC, Perrot V (1996) Models of the within-host dynamics of persistent mycobacterial infections. *Proc R Soc Lond. Ser B: Biol Sci* 263(1368):257–263. <https://doi.org/10.1098/rspb.1996.0040>
- Balmer O, Tanner M (2011) Prevalence and implications of multiple-strain infections. *Lancet Infect Dis* 11(11):868–878. [https://doi.org/10.1016/S1473-3099\(11\)70241-9](https://doi.org/10.1016/S1473-3099(11)70241-9)
- Barabás G, D'Andrea R, Stump SM (2018) Chesson's coexistence theory. *Ecol Monogr* 88(3):277–303. <https://doi.org/10.1002/ecm.1302>
- Bashey F (2015) Within-host competitive interactions as a mechanism for the maintenance of parasite diversity. *Philos Trans R Soc B: Biol Sci* 370(1675):20140301. <https://doi.org/10.1098/rstb.2014.0301>
- Bates A, Hilton B, Harley C (2009) Effects of temperature, season and locality on wasting disease in the keystone predatory sea star *Pisaster ochraceus*. *Dis Aquat Org* 86:245–251. <https://doi.org/10.3354/dao02125>
- Bell GI (1973) Predator-prey equations simulating an immune response. *Math Biosci* 16(3–4):291–314. [https://doi.org/10.1016/0025-5564\(73\)90036-9](https://doi.org/10.1016/0025-5564(73)90036-9)
- Bernhardt JR, Sunday JM, Thompson PL, O'Connor MI (2018) Non-linear averaging of thermal experience predicts population growth rates in a thermally variable environment. *Proc R Soc B: Biol Sci* 285(1886):20181076. <https://doi.org/10.1098/rspb.2018.1076>
- Bromenshenk JJ, Henderson CB, Wick CH, Stanford MF, Zulich AW, Jabbour RE, Deshpande SV, McCubbin PE, Seccomb RA, Welch PM, Williams T, Firth DR, Skowronski E, Lehmann MM, Bili-moria SL, Gress J, Wanner KW, Cramer RA (2010) Iridovirus and microsporidian linked to honey bee colony decline. *PLoS ONE* 5(10):e13181. <https://doi.org/10.1371/journal.pone.0013181>
- Brook CE, Dobson AP (2015) Bats as 'special' reservoirs for emerging zoonotic pathogens. *Trends Microbiol* 23(3):172–180. <https://doi.org/10.1016/j.tim.2014.12.004>
- Brown JH, Gillooly JF, Allen AP, Savage VM, West GB (2004) Toward a metabolic theory of ecology. *Ecology* 85(7):1771–1789. <https://doi.org/10.1890/03-9000>
- Bruno JF, Selig ER, Casey KS, Page CA, Willis BL, Harvell CD, Sweatman H, Melendy AM (2007) Thermal stress and coral cover as drivers of coral disease outbreaks. *PLoS Biol* 5(6):e124. <https://doi.org/10.1371/journal.pbio.0050124>
- Butler MW, Stahlschmidt ZR, Ardia DR, Davies S, Davis J, Guillet-Lette LJ, Johnson N, McCormick SD, McGraw KJ, DeNardo DF (2013) Thermal sensitivity of immune function: evidence against

- a generalist-specialist trade-off among endothermic and ectothermic vertebrates. *Am Nat* 181(6):761–774. <https://doi.org/10.1086/670191>
- Calisher CH, Childs JE, Field HE, Holmes KV, Schountz T (2006) Bats: important reservoir hosts of emerging viruses. *Clin Microbiol Rev* 19(3):531–545. <https://doi.org/10.1128/CMR.00017-06>
- Camus A (1991) The plague. (trans: Stuart G) 1st edn. Vintage international. Vintage Books
- Chesson P (2000) Mechanisms of maintenance of species diversity. *Annu Rev Ecol Syst* 31(1):343–366. <https://doi.org/10.1146/annurev.ecolsys.31.1.343>
- Chesson P (2003) Quantifying and testing coexistence mechanisms arising from recruitment fluctuations. *Theor Popul Biol* 64(3):345–357. [https://doi.org/10.1016/S0040-5809\(03\)00095-9](https://doi.org/10.1016/S0040-5809(03)00095-9)
- Chesson P (1994) Multispecies competition in variable environments. *Theor Popul Biol* 45(3):227–276. <https://doi.org/10.1006/tpbi.1994.1013>
- Chesson P, Huntly N (1997) The roles of harsh and fluctuating conditions in the dynamics of ecological communities. *Am Nat* 150(5):519–553. <https://doi.org/10.1086/286080>
- Cizauskas CA, Carlson CJ, Burgio KR, Clements CF, Dougherty ER, Harris NC, Phillips AJ (2017) Parasite vulnerability to climate change: an evidence-based functional trait approach. *R Soc Open Sci* 4(1):160535. <https://doi.org/10.1098/rsos.160535>
- Cohen JM, Venesky MD, Sauer EL, Civitello DJ, McMahon TA, Roznik EA, Rohr JR (2017) The thermal mismatch hypothesis explains host susceptibility to an emerging infectious disease. *Ecol Lett* 20(2):184–193. <https://doi.org/10.1111/ele.12720>
- Cohen JM, Civitello DJ, Venesky MD, McMahon TA, Rohr JR (2019) An interaction between climate change and infectious disease drove widespread amphibian declines. *Glob Chang Biol* 25(3):927–937. <https://doi.org/10.1111/gcb.14489>
- Cohen JM, Sauer EL, Santiago O, Spencer S, Rohr JR (2020) Divergent impacts of warming weather on wildlife disease risk across climates. *Science* 370(6519):eabb1702. <https://doi.org/10.1126/science.abb1702>
- Comins HN, Noble IR (1985) Dispersal, variability, and transient niches: species coexistence in a uniformly variable environment. *Am Nat* 126(5):706–723. <https://doi.org/10.1086/284448>
- Connell JH (1978) Diversity in tropical rain forests and coral reefs: high diversity of trees and corals is maintained only in a nonequilibrium state. *Science* 199(4335):1302–1310. <https://doi.org/10.1126/science.199.4335.1302>
- Cox FEG (2001) Concomitant infections, parasites and immune responses. *Parasitology* 122(S1):S23–S38. <https://doi.org/10.1017/S003118200001698X>
- Dallas T, Drake JM (2016) Fluctuating temperatures alter environmental pathogen transmission in a *Daphnia*–pathogen system. *Ecol Evol* 6(21):7931–7938. <https://doi.org/10.1002/ece3.2539>
- Dallas TA, Laine A-L, Ovaskainen O (2019) Detecting parasite associations within multi-species host and parasite communities. *Proc R Soc B: Biol Sci* 286(1912):20191109. <https://doi.org/10.1098/rspb.2019.1109>
- Daszak P, Berger L, Cunningham AA, Hyatt AD, Green DE, Speare R (1999) Emerging infectious diseases and amphibian population declines. *Emerg Infect Dis* 5(6):735–748. <https://doi.org/10.3201/eid0506.990601>
- Daszak P, Cunningham AA, Hyatt AD (2000) Emerging infectious diseases of wildlife—threats to biodiversity and human health. *Science* 287(5452):443–449. <https://doi.org/10.1126/science.287.5452.443>
- Daszak P, Cunningham AA, Hyatt AD (2001) Anthropogenic environmental change and the emergence of infectious diseases in wildlife. *Acta Trop* 78(2):103–116. [https://doi.org/10.1016/S0001-706X\(00\)00179-0](https://doi.org/10.1016/S0001-706X(00)00179-0)
- Dell AI, Pawar S, Savage VM (2011) Systematic variation in the temperature dependence of physiological and ecological traits. *Proc Natl Acad Sci* 108(26):10591–10596. <https://doi.org/10.1073/pnas.1015178108>
- Dell AI, Pawar S, Savage VM (2014) Temperature dependence of trophic interactions are driven by asymmetry of species responses and foraging strategy. *J Anim Ecol* 83(1):70–84. <https://doi.org/10.1111/1365-2656.12081>
- Denny M (2017) The fallacy of the average: on the ubiquity, utility and continuing novelty of Jensen’s inequality. *J Exp Biol* 220(2):139–146. <https://doi.org/10.1242/jeb.140368>
- Duncan AB, Fellous S, Kaltz O (2011) Temporal variation in temperature determines disease spread and maintenance in *Paramecium* microcosm populations. *Proc R Soc B: Biol Sci* 278(1723):3412–3420. <https://doi.org/10.1098/rspb.2011.0287>
- Estay SA, Clavijo-Baquet S, Lima M, Bozinovic F (2011) Beyond average: an experimental test of temperature variability on the population dynamics of *Tribolium confusum*. *Popul Ecol* 53(1):53–58. <https://doi.org/10.1007/s10144-010-0216-7>
- Fumagalli MR, Zapperi S, La Porta CAM (2021) Role of body temperature variations in bat immune response to viral infections. *J R Soc Interface* 18(180):20210211. <https://doi.org/10.1098/rsif.2021.0211>
- Greenrod STE, Cazares D, Johnson S, Hector TE, Stevens EJ, MacLean RC, King KC (2024) Warming alters life-history traits and competition in a phage community. *Appl Environ Microbiol* 90(5):e00286–e324. <https://doi.org/10.1128/aem.00286-24>
- Halpin K, Young PL, Field HE, Mackenzie JS (2000) Isolation of Hendra virus from pteropid bats: a natural reservoir of Hendra virus. *J Gen Virol* 81(8):1927–1932. <https://doi.org/10.1099/0022-1317-81-8-1927>
- Holt RD (1977) Predation, apparent competition, and the structure of prey communities. *Theor Popul Biol* 12(2):197–229. [https://doi.org/10.1016/0040-5809\(77\)90042-9](https://doi.org/10.1016/0040-5809(77)90042-9)
- Holt RD, Bonsall MB (2017) Apparent competition. *Annu Rev Ecol Evol Syst* 48(1):447–471. <https://doi.org/10.1146/annurev-ecolsys-110316-022628>
- Holt RD, Lawton JH (1994) The ecological consequences of shared natural enemies. *Annu Rev Ecol Syst* 25(1):495–520. <https://doi.org/10.1146/annurev.es.25.110194.002431>
- Huston MA (2014) Disturbance, productivity, and species diversity: empiricism vs. logic in ecological theory. *Ecology* 95(9):2382–2396. <https://doi.org/10.1890/13-1397.1>
- Irving AT, Ahn M, Goh G, Anderson DE, Wang L-F (2021) Lessons from the host defenses of bats, a unique viral reservoir. *Nature* 589(7842):363–370. <https://doi.org/10.1038/s41586-020-03128-0>
- Jensen JLWV (1906) Sur les fonctions convexes et les inégalités entre les valeurs moyennes. *Acta Math* 30:175–193. <https://doi.org/10.1007/BF02418571>
- Jiang L, Patel SN (2008) Community assembly in the presence of disturbance: a microcosm experiment. *Ecology* 89(7):1931–1940. <https://doi.org/10.1890/07-1263.1>
- Johnson PTJ (2006) Amphibian diversity: decimation by disease. *Proc Natl Acad Sci* 103(9):3011–3012. <https://doi.org/10.1073/pnas.0600293103>
- Johnson PTJ, Hoverman JT (2012) Parasite diversity and coinfection determine pathogen infection success and host fitness. *Proc Natl Acad Sci* 109(23):9006–9011. <https://doi.org/10.1073/pnas.1201790109>
- Johnson PTJ, Preston DL, Hoverman JT, LaFonte BE (2013) Host and parasite diversity jointly control disease risk in complex communities. *Proc Natl Acad Sci* 110(42):16916–16921. <https://doi.org/10.1073/pnas.1310557110>
- Jolles AE, Ezenwa VO, Etienne RS, Turner WC, Olff H (2008) Interactions between macroparasites and microparasites drive infection

- patterns in free-ranging African buffalo. *Ecology* 89(8):2239–2250. <https://doi.org/10.1890/07-0995.1>
- Kirk D, Jones N, Peacock S, Phillips J, Molnár PK, Krkošek M, Luijckx P (2018) Empirical evidence that metabolic theory describes the temperature dependency of within-host parasite dynamics. *PLoS Biol* 16(2):e2004608. <https://doi.org/10.1371/journal.pbio.2004608>
- Krichel L, Kirk D, Pencer C, Hönig M, Wadhawan K, Krkošek M (2023) Short-term temperature fluctuations increase disease in a *Daphnia*-parasite infectious disease system. *PLoS Biol* 21(9):e3002260. <https://doi.org/10.1371/journal.pbio.3002260>
- Lafferty KD, Dobson AP, Kuris AM (2006) Parasites dominate food web links. *Proc Natl Acad Sci* 103(30):11211–11216. <https://doi.org/10.1073/pnas.0604755103>
- Lawson CR, Vindenes Y, Bailey L, van de Pol M (2015) Environmental variation and population responses to global change. *Ecol Lett* 18(7):724–736. <https://doi.org/10.1111/ele.12437>
- Lips KR, Diffendorfer J, Mendelson JR, Sears MW (2008) Riding the wave: reconciling the roles of disease and climate change in amphibian declines. *PLoS Biol* 6(3):e72. <https://doi.org/10.1371/journal.pbio.0060072>
- Luo J, Greif S, Ye H, Bumrungsri S, Eitan O, Yovel Y (2021) Flight rapidly modulates body temperature in freely behaving bats. *Anim Biotelemetry* 9(1):45. <https://doi.org/10.1186/s40317-021-00268-6>
- Mackey RL, Currie DJ (2001) The diversity-disturbance relationship: is it generally strong and peaked? *Ecology* 82(12):3479–3492. [https://doi.org/10.1890/0012-9658\(2001\)082\[3479:TDDRII\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2001)082[3479:TDDRII]2.0.CO;2)
- Miller AD, Roxburgh SH, Shea K (2011) How frequency and intensity shape diversity–disturbance relationships. *Proc Natl Acad Sci* 108(14):5643–5648. <https://doi.org/10.1073/pnas.1018594108>
- Miller AD, Roxburgh SH, Shea K (2012) Timing of disturbance alters competitive outcomes and mechanisms of coexistence in an annual plant model. *Thyroid Res* 5(3):419–432. <https://doi.org/10.1007/s12080-011-0133-1>
- Molnár PK, Kutz SJ, Hoar BM, Dobson AP (2013) Metabolic approaches to understanding climate change impacts on seasonal host-macroparasite dynamics. *Ecol Lett* 16(1):9–21. <https://doi.org/10.1111/ele.12022>
- Molnár PK, Skrabulis JP, Altman KA, Raffel TR (2017) Thermal performance curves and the metabolic theory of ecology—a practical guide to models and experiments for parasitologists. *J Parasitol* 103(5):423. <https://doi.org/10.1645/16-148>
- Nowak MA, May RM (2000) *Virus dynamics: mathematical principles of immunology and virology*. Oxford University Press. <https://doi.org/10.1093/oso/9780198504184.001.0001>
- Paull SH, LaFonte BE, Johnson PTJ (2012) Temperature-driven shifts in a host-parasite interaction drive nonlinear changes in disease risk. *Glob Chang Biol* 18(12):3558–3567. <https://doi.org/10.1111/gcb.12018>
- Pedersen AB, Fenton A (2007) Emphasizing the ecology in parasite community ecology. *Trends Ecol Evol* 22(3):133–139. <https://doi.org/10.1016/j.tree.2006.11.005>
- Petney TN, Andrews RH (1998) Multiparasite communities in animals and humans: frequency, structure and pathogenic significance. *Int J Parasitol* 28(3):377–393. [https://doi.org/10.1016/S0020-7519\(97\)00189-6](https://doi.org/10.1016/S0020-7519(97)00189-6)
- Pilyugin S, Antia R (2000) Modeling immune responses with handling time. *Bull Math Biol* 62(5):869–890. <https://doi.org/10.1006/bulm.2000.0181>
- Plowright RK, Parrish CR, McCallum H, Hudson PJ, Ko AI, Graham AL, Lloyd-Smith JO (2017) Pathways to zoonotic spillover. *Nat Rev Microbiol* 15(8):502–510. <https://doi.org/10.1038/nrmicro.2017.45>
- Råberg L, De Roode JC, Bell AS, Stamou P, Gray D, Read AF (2006) the role of immune-mediated apparent competition in genetically diverse malaria infections. *Am Nat* 168(1):41–53. <https://doi.org/10.1086/505160>
- Raffel TR, Romansic JM, Halstead NT, McMahon TA, Venesky MD, Rohr JR (2013) Disease and thermal acclimation in a more variable and unpredictable climate. *Nat Clim Chang* 3(2):146–151. <https://doi.org/10.1038/nclimate1659>
- Ramesh A, Hall SR (2023) Niche theory for within-host parasite dynamics: analogies to food web modules via feedback loops. *Ecol Lett* 26(3):351–368. <https://doi.org/10.1111/ele.14142>
- Rigaud T, Perrot-Minnot M-J, Brown MJF (2010) Parasite and host assemblages: embracing the reality will improve our knowledge of parasite transmission and virulence. *Proc R Soc B: Biol Sci* 277(1701):3693–3702. <https://doi.org/10.1098/rspb.2010.1163>
- Rohr JR, Raffel TR (2010) Linking global climate and temperature variability to widespread amphibian declines putatively caused by disease. *Proc Natl Acad Sci* 107(18):8269–8274. <https://doi.org/10.1073/pnas.0912883107>
- Rohr JR, Raffel TR, Romansic JM, McCallum H, Hudson PJ (2008) Evaluating the links between climate, disease spread, and amphibian declines. *Proc Natl Acad Sci* 105(45):17436–17441. <https://doi.org/10.1073/pnas.0806368105>
- Rohr JR, Raffel TR, Blaustein AR, Johnson PTJ, Paull SH, Young S (2013) Using physiology to understand climate-driven changes in disease and their implications for conservation. *Conserv Physiol* 1(1):cot022–cot022. <https://doi.org/10.1093/conphys/cot022>
- Roxburgh SH, Shea K, Wilson JB (2004) The intermediate disturbance hypothesis: patch dynamics and mechanisms of species coexistence. *Ecology* 85(2):359–371. <https://doi.org/10.1890/03-0266>
- Schoolfield RM, Sharpe PJH, Magnuson CE (1981) Non-linear regression of biological temperature-dependent rate models based on absolute reaction-rate theory. *J Theor Biol* 88(4):719–731. [https://doi.org/10.1016/0022-5193\(81\)90246-0](https://doi.org/10.1016/0022-5193(81)90246-0)
- Schulte PM, Healy TM, Fanguie NA (2011) Thermal performance curves, phenotypic plasticity, and the time scales of temperature exposure. *Integr Comp Biol* 51(5):691–702. <https://doi.org/10.1093/icb/ucr097>
- Skrabulis JP, Altman KA, Raffel TR (2022) Using metabolic theory to describe temperature and thermal acclimation effects on parasitic infection. *Am Nat* 199(6):789–803. <https://doi.org/10.1086/719409>
- Sinclair BJ, Marshall KE, Sewell MA, Levesque DL, Willett CS, Slotsbo S, Dong Y, Harley CDG, Marshall DJ, Helmuth BS, Huey RB (2016) Can we predict ectotherm responses to climate change using thermal performance curves and body temperatures? *Ecol Lett* 19(11):1372–1385. <https://doi.org/10.1111/ele.12686>
- Song C, Spaak JW (2024) Trophic tug-of-war: coexistence mechanisms within and across trophic levels. *Ecol Lett* 27(4):e14409. <https://doi.org/10.1111/ele.14409>
- Spaak JW, Godoy O, De Laender F (2021) Mapping species niche and fitness differences for communities with multiple interaction types. *Oikos* 130(12):2065–2077. <https://doi.org/10.1111/oik.08362>
- Stocker TF, Qin D, Plattner G-K, Tignor M, Allen SK, Boschung J, Nauels A, Xia Y, Bex V, Midgley PM (2013) *Climate Change 2013: The physical science basis - IPCC. Contribution of working group I to the fifth assessment report of the intergovernmental panel on climate change*. Cambridge University Press, Cambridge, pp 1535
- Susi H, Barrès B, Vale PF, Laine A-L (2015) Co-infection alters population dynamics of infectious disease. *Nat Commun* 6(1):5975. <https://doi.org/10.1038/ncomms6975>
- Symons CC, Arnott SE (2014) Timing is everything: priority effects alter community invasibility after disturbance. *Ecol Evol* 4(4):397–407. <https://doi.org/10.1002/ece3.940>

- Tracy AM, Pielmeier ML, Yoshioka RM, Heron SF, Harvell CD (2019) Increases and decreases in marine disease reports in an era of global change. *Proceedings of the Royal Society b: Biological Sciences* 286(1912):20191718. <https://doi.org/10.1098/rspb.2019.1718>
- Tuljapurkar S (1990) *Population dynamics in variable environments* (vol 85). Springer Berlin Heidelberg. <https://doi.org/10.1007/978-3-642-51652-8>
- Vasseur DA, DeLong JP, Gilbert B, Greig HS, Harley CDG, McCann KS, Savage V, Tunney TD, O'Connor MI (2014) Increased temperature variation poses a greater risk to species than climate warming. *Proc R Soc B: Biol Sci* 281(1779):20132612. <https://doi.org/10.1098/rspb.2013.2612>
- Williamson MH (1957) An elementary theory of interspecific competition. *Nature* 180(4583):422–425. <https://doi.org/10.1038/180422a0>
- Wootton JT (1998) Effects of disturbance on species diversity: a multitrophic perspective. *Am Nat* 152(6):803–825. <https://doi.org/10.1086/286210>
- Wright RK, Cooper EL (1981) Temperature effects on ectotherm immune responses. *Dev Comp Immunol* 5:117–122
- Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.
- Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.