Lawrence Berkeley National Laboratory

Recent Work

Title

SYNTHESIS, STRUCTURAL ELUCIDATION AND STEREOCHEMISTRY OF FIVE-COORDINATE
ORGANOARSENIC CATECHOLATES

Permalink

<https://escholarship.org/uc/item/8mw9p81t>

Authors

Fish, R.H.
Tannous, R.S.

Publication Date

1982-04-01

Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

ENERGY & ENVIRONMENT DIVISION

RECEIVED
LAWRENCE
BERKELEY LABORATORY

APR 16 1982

LIBRARY AND
DOCUMENTS SECTION

Submitted to Organometallics

SYNTHESIS, STRUCTURAL ELUCIDATION AND STEREOCHEMISTRY
OF FIVE-COORDINATE ORGANOARSENIC CATECHOLATES

Richard H. Fish and Raja S. Tannous

April 1982

TWO-WEEK LOAN COPY

This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 6782

LBL-14199
e-2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Synthesis, Structural Elucidation and Stereochemistry
of Five-Coordinate Organoarsenic Catecholates

Richard H. Fish and Raja S. Tannous
Energy and Environment Division
Lawrence Berkeley Laboratory
University of California
Berkeley, CA 94720

This work was supported by the Assistant Secretaries for Fossil Energy and Oil, Gas and Shale Technology, and the Bartlesville Energy Technology Center Of the U.S. Department of Energy under Contract number DE-AC03-76SF00096.

Abstract

The reactions of methyl and phenylarsonic acids with substituted catechols was studied. These reactions produced five-coordinate organoarsenic catecholates, which were characterized by 250MHz ^1H NMR spectroscopy, IR spectroscopy, EI mass spectrometry and elemental analyses. The possibility of cis and trans isomers for several of the substituted catecholate organoarsenic compounds was verified by NMR spectroscopy, and in one case, single crystal x-ray crystallography.

This latter x-ray study provided, for the first time, unequivocal evidence for cis stereochemistry of methyl groups attached to the catecholate rings, a rectangular pyramidal configuration (95%) of the oxygen atoms surrounding the arsenic atom with a phenyl group in the axial position.

Synthesis, Structural Elucidation and Stereochemistry
of Five-Coordinate Organoarsenic Catecholates

Richard H. Fish* and Raja S. Tannous
Energy and Environment Division
Lawrence Berkeley Laboratory
University of California
Berkeley, CA 94720

Summary: The synthesis of five-coordinate organoarsenic catecholates, with substitution of 3-CH₃-; 4-CH₃OC(=O)-; 4,5- , and 3,3'-C(=O)-N(CH₂)₄NH- groups on the catechol ring and CH₃ or Ph on arsenic, was performed. These reactions provided new insights into the structures and stereochemistry of these organoarsenic compounds. In one case, 3-methylcatecholate of phenylarsonic acid, a single crystal x-ray analysis provided unequivocal evidence for cis stereochemistry of the methyl groups on the catechol rings, an essentially rectangular pyramidal configuration around arsenic, and a preferred conformation of the phenyl group in relation to the oxygens surrounding the arsenic atom.

Recently, we have been investigating the speciation or molecular characterization of organoarsenic compounds thought to present in biogeochemical materials such as oil shale kerogen and the products of the pyrolysis of oil shale kerogen, those of the shale oils and the retort waters^{1a,b}. These studies led to the identification, for the first time, of methyl and phenylarsonic acids in these precursors and products.

In view of these discoveries, we have initiated studies to find innovative methods for the removal of these compounds and other organometallics from fossil fuel products. In this regard, we have been experimenting with a method that utilizes substituted catechols as

potential ligands that could be placed in a polymeric matrix for the future removal of organoarsenic compounds from the above mentioned products.

Surprisingly, we found very few references on the reactions of catechols with alkyl or arylarsonic acids^{2a-c} and none on similar reactions with substituted catechols³. Thus, in this communication, we present our initial results on the synthesis, structural elucidation and stereochemistry of the five-coordinate organoarsenic catecholates we prepared as model compounds for the above mentioned purposes.

Chart 1 shows the catechols, 1-4, we utilized in the reactions with methyl or phenylarsonic acid, 5 and 6. Compound 1 reacts with either 5 or 6 to provide a mixture of cis and trans five-coordinate organoarsenic catecholates, 7 - 10 (Eq. 1).

Compounds 7-10 were characterized by a combination of nuclear magnetic resonance spectroscopy (NMR), mass spectrometry (MS), infrared spectroscopy (IR) and elemental analysis.⁴ The 250 MHz NMR spectrum provided definitive evidence that compound 1 reacted with either 5 or 6 to give a mixture of cis and trans isomers (7, 8 R = CH₃ and 9, 10 R = Ph). Thus, compounds 7 and 8 showed two methyl resonances (catecholate ring) at 2.21 and 2.19 ppm (benzene-d₆, TMS) and two methyl resonances for

groups bonded to arsenic at 1.33 and 1.32 ppm in the ratio of 53:47. The corresponding cis and trans compounds, 9 and 10, where R=Ph, had methyl resonances at 2.26 and 2.12 ppm (benzene-d₆, TMS) in the ratio of 90:10. The complexity of the phenyl region at 250 MHz did not allow a separation of catecholate protons and phenylarsenic protons and thus the ¹H NMR spectrum was obtained at 400 MHz. The ¹H 400 MHz NMR spectrum of 9, 10 (benzene-d₆, TMS) showed resonances at 7.81 (doublet, J=8.3Hz); 6.83 (multiplet) and 6.74 ppm (overlapping triplets J=8.3Hz) for the phenylarsenic protons in the ratio of 2:1:2. The catecholate protons were found at 6.56 (doublet of doublets, J_{ortho}=7.8 Hz, J_{meta}=1.4 Hz); 6.65 (overlapping triplets, J=7.8 Hz), 6.93 (triplet, J=7.8 Hz) and 2.24 ppm (singlet) in the ratio of 2:2:2:3. The complexity of the phenyl region, where protons on the phenyl group attached to arsenic appeared to be all non-equivalent, provided tentative evidence for the cis isomer, 9, rather than the trans isomer, 10, as the major product in this reaction.

In order to unequivocally ascertain the stereochemistry of the major isomer, either 9 (cis) or 10 (trans), we obtained a single-crystal x-ray analysis.⁵ Figure 1 shows the ORTEP drawing of the major isomer, 9, with the methyl groups clearly cis to each other and the geometry around the arsenic, essentially rectangular pyramidal(95%), while the axial phenyl group is twisted so that it lies in the same plane as oxygen 2 and oxygen 3. The angle between the carbons 1 and 6 on the phenyl group attached to arsenic and the oxygen-arsenic-oxygen plane is 4.7°. This represents the first rectangular pyramidal and stereochemical assignment that has been made for a five-coordinate organoarsenic

catecholate with substitution on the catecholate rings (see Table 1 for pertinent bond lengths and angles).⁶

Compound 2 reacted with 5 to provide a compound, 11, with a single methylarsenic resonance at 1.92 ppm (¹H, 250MHz, DMSO-d₆, TMS) and a single methoxyl resonance at 3.78 ppm and is indicative of one isomer, which we presume to have cis stereochemistry as in 9. Compound 2 also reacts with 6 to provide a mixture of compounds, 12 and 13, with two methoxyl signals (¹H, 250 MHz, DMSO-d₆, TMS) at 3.76 and 3.74 ppm in the ratio of 95:5. Again, as with 11, we presume cis stereochemistry for the major isomer. Compound 3, a benzo-substituted catechol, reacted with 5 or with 6 to give the five-coordinate organoarsenic compounds 14 and 15 respectively. While no stereochemistry is involved in the formation of either 14 or 15, it is important to note that substitution on the naphthyl ring is certainly possible for future attachment of this type of compound to a polymeric backbone. The pertinent NMR data (¹H 250 MHz, DMSO-d₆, TMS) provided an upfield shift, as with the NMR spectra of compounds 7-13, for the catechol ring protons of 14 and 15 when compared to 3. Thus, 14 had the catechol protons (singlet) at 7.09 ppm, 15 at 7.10 and 3 at 7.12 ppm indicative of the arsenic atoms influence on shifting, to higher fields, protons on catechol rings. A similar NMR result was obtained by Raymond et al.⁷ for some gallium and rhodium catecholate complexes.

Our final ligand of interest, 4⁸, was important to study, since it represented a model for a recently reported polymer of potential use for our future applications⁹. We chose 4 (4-LICAM) after making Dreiding models that clearly showed the central cavity being able to accomodate

an arsenic atom (~3.58 to 3.63 Å, see Figure 1). Reaction of 4, with either 5 or 6, provided the intramolecular five-coordinate organoarsenic derivatives, 16 and 17 (Eq. 2).

The ^1H 250 MHz NMR and 70 ev MS (solid probe) data were consistent with the structures assigned. Notably, the mass spectra provided the parent ion and an ion resulting from a loss of the catechol group with a carbonyl attached. This was followed by a fragmentation of the $-\text{CH}_2\text{CH}_2-\text{NH}$ groupings. For example, with 17 the MS ions of interest were: m/e 508 ($\text{M}^{+\bullet}$); 373 ($\text{M}-\text{C}_7\text{H}_{13}\text{N}_2\text{O}_3$); 331 ($\text{M}-\text{C}_9\text{H}_8\text{NO}_3$); and 287 ($\text{M}-\text{C}_{11}\text{H}_{13}\text{N}_2\text{O}_3$).

A typical procedure for the preparation of a five-coordinate organoarsenic catecholate derivative is described as follows for 9.

In a 50 ml flask, equipped with condenser and drying tube, was placed 1.29g (10.42 mmole) of phenylarsonic acid and 1.05 g (5.21 mmole) 3-methylcatechol (freshly sublimed) in 30 ml of benzene. The reaction mixture was refluxed for 5h. The benzene was removed on a rotary evaporator and the compound recrystallized from carbon tetrachloride/methanol and dried under vacuum to give 1.88g (91% yield) of 9, mp, 134-135°C. Calculated for $\text{C}_{20}\text{H}_{17}\text{O}_4\text{As}$; C, 60.6; H, 4.3, Found: C, 60.39; H, 4.46; EIMS (70 ev, solid probe) m/e 396 ($\text{M}^{+\bullet}$), 274, 197,

151, 106.¹⁰

In future experiments, we hope to place several of our catechol derivatives in polymeric backbones to see if their reactivity remains in reactions with organoarsonic acids.

Acknowledgements

We wish to thank Drs. F. L. Weitzl and K. N. Raymond for several samples of LICAM derivatives for evaluation and helpful suggestions during the initial stages of this project. Helpful discussions with Drs. K. Irgolic and R. Zingaro are also acknowledged. We wish to thank Drs. F.E. Brinckman and R.B. Johannesen, of the National Bureau of Standards, for obtaining the 400 MHz ¹H NMR spectrum of 9. The crystal structure analysis for 9 was performed by Dr. F.J. Hollander at U.C. Berkeley x-ray Crystallographic Facility (CHEXRAY). The work was supported by the Assistant Secretaries for Fossil Energy and Oil, Gas and Shale Technology, and the Bartlesville Energy Technology Center (project manager, Dexter Sutterfield) of the U.S. Department of Energy under contract No. DE-AC03-76SF00098.

Supplementary Material Available

A listing of observed and calculated structure factor amplitudes (18 pages). Ordering information is given on any current masthead page.

References

- (1) a) Fish R.H., Brinckman F.E. and Jewett K.L., Environ. Sci. Technol., 1982, 16, 174.
b) Weiss C.S., Brinckman F.E. and Fish R.H., Proceedings on

Environmental Speciation of Trace Metal-Containing Compounds in Energy-Related Processes, May 18-20, 1981, Gaithersburg, MD. National Bureau of Standards Special Publication 1981 618, 197.

- (2) a) Salmi E.J., Merivuori K. and Laaksonen E., *Kemistilehti Suomen* 1946, 19B, 102, (CA, 41, 5440a).
b) Bacher J.H. and Van Oosten R.P., *Rec. Trav. Chem.* 1940, 59, 41.
c) Englund B., *Chem. Ber.*, 1926, 59, 2669.
- (3) Fish R.H. and Weitz F.L., Xth International Conference on Organometallic Chemistry, Toronto, Canada, August 9-14, 1981, Abstract 1D12, p. 62.
- (4) All new compounds we report had elemental analyses (C, H, N) within $\pm 0.5\%$ of theory. The EIMS, IR and NMR spectra for all compounds supported their structures.
- (5) Crystal data for 9: space group $P\bar{1}$, $a=9.3734(8)\text{\AA}$, $b=9.3823(12)\text{\AA}$, $c=11.3718(15)\text{\AA}$, $\alpha=69.433(10)^\circ$, $\beta=70.809(9)^\circ$, $\gamma=72.631(8)^\circ$, $V=865.7(2)\text{\AA}^3$, formula weight: 396.28 amu, $Z = 2$, $d = 1.52\text{g/cm}^3$, $\mu=19.76\text{ cm}^{-1}$, size: 0.21 x 0.33 x 0.37 mm. The structure was solved by Patterson and Fourier Techniques and refined via normal least squares procedures: $R=3.47\%$ and $R_w=5.49\%$ using 2242 reflections with $F^2 > 3 F^2$. The crystals of 9 were grown in a solvent mixture of carbon disulfide and pentane (1:1) by slow evaporation.
- (6) Several other x-ray studies on five-coordinate organoarsenic glycolate and catecholate derivatives have been done. However, in both cases a distorted rectangular pyramidal configuration around arsenic (60-75% RP) was found and no stereochemical characteriza-

tion was attempted, i.e., substituted catechols were not used.
(See Goldwhite H. and Teller R.G. , J. Am. Chem. Soc. 1978, 100,
5357. Wunderlich, H. Acta Cryst. 1978, B34, 1000.)

See Holmes, R.R. and Deiters, J. J. Am. Chem. Soc. 1977, 99, 3318
for idealized angles for square pyramidal, rectangular pyramidal and
trigonal bipyramidal structures and methods of analysis. Using the
angle deviation analysis method of Holmes and Deiter, we find a
minimum deviation sum for the rectangular pyramidal structure of
 11.2° (~95% distortion from trigonal bipyramidal toward rectangular
pyramidal) for compound 9.

- (7) Mcardle J.V., Sofen S.R., Cooper S.R., and Raymond K.N., Inorg.
Chem. 1978, 17, 3075.
- (8) The series of LICAM derivatives of increasing methylene chain, 2-6,
were prepared according to the procedures of Dr. F.L. Weitl (cf.
F.L. Weitl and K.N. Raymond, J.Amer. Chem. Soc. 1980, 102, 2289).
- (9) Dawson M. I., Chan R.L.-S., Cloudsdale I.S. and Harris W.R.,
Tetrahedron Letters, 1981, 22, 2739.
- (10) We have used this procedure to derivatize phenylarsonic acid, 6,
that had been isolated from a Green River oil shale kerogen by
extraction with methanol. Since 9 can be chromatographed on a 30 m
fused silica capillary column (OV101), the use of GC-MS will
enhance the utility of these organoarsonic acid derivatives for
other applications. Fish, R.H., Tannous, R.S., Weiss, C.S., and
Brinckman, F.E., in preparation.

Table I

Selected Bond Lengths and Angles for

cis (3-CH₃-C₆H₃O₂)₂AsC₆H₅, 9^a

Atoms	Bond length Å	Atoms	Angle, Deg
As-O ₁	1.806 (.001)	O ₁ -As-O ₂	87.93 (.07)
As-O ₂	1.799 (.002)	O ₁ -As-O ₃	85.43 (.07)
As-O ₃	1.784 (.002)	O ₁ -As-O ₄	150.84 (.09)
As-O ₄	1.825 (.002)	O ₂ -As-O ₃	149.85 (.09)
As-C ₁	1.899 (.002)	O ₂ -As-O ₄	82.71 (.08)
C ₇ -O ₁	1.365 (.003)	O ₃ -As-O ₄	88.95 (.09)
C ₈ -O ₂	1.370 (.003)	O ₁ -As-C ₁	105.5 (.08)
C ₁₄ -O ₃	1.413 (.003)	O ₂ -As-C ₁	104.99 (.09)
C ₁₅ -O ₄	1.345 (.003)	O ₃ -As-C ₁	105.12 (.09)
C ₁₂ -C ₁₃	1.474 (.004)	O ₄ -As-C ₁	103.59 (.09)

^aEstimated standard deviations in parenthesis

Figure 1. ORTEP diagram of cis (3-CH₃-C₆H₃O₂)₂ AsC₆H₅, 9, showing 20% probability ellipsoids. ✓

Chart 1. Catechols used in the synthesis of
five coordinate organoarsenic catecholates

1

2

3

4

XBL 823-8458

XBL 823-8156

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

TECHNICAL INFORMATION DEPARTMENT
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720