Lawrence Berkeley National Laboratory

Recent Work

Title

REFORMULATION OF QUASILINEAR THEORY

Permalink

<https://escholarship.org/uc/item/8nh016wn>

Author

Kaufman, Allan N.

Publication Date

1970-06-24

c.2

RECEIVED
LAWRENCE
RADIATION LABORATORY
JUL 29 1970
LIBRARY AND
DOCUMENTS SECTION

REFORMULATION OF QUASILINEAR THEORY

Allan N. Kaufman

June 24, 1970

AEC Contract No. W-7405-eng-48

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

20 LAWRENCE RADIATION LABORATORY
UNIVERSITY of CALIFORNIA BERKELEY

UCRL-19869
cy

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

REFORMULATION OF QUASILINEAR THEORY*

Allan N. Kaufman

Department of Physics and Lawrence Radiation Laboratory
University of California, Berkeley, California 94720

June 24, 1970

ABSTRACT

A new formulation of quasilinear theory is presented, which allows for only resonant diffusion, caused by both growing and damped waves. Nonresonant terms do not appear in the diffusion equation, but contribute to wave momentum and energy, and ensure conservation of total momentum and energy. Analyticity is not required, since the complex plane is not used.

* This work was supported by the U. S. Atomic Energy Commission and by NASA.

I. INTRODUCTION

Users of quasilinear theory¹ have often expressed uneasiness about its applicability to damped waves,² because the standard expression of the diffusivity is then negative. Attempts^{3,4} to establish a quasilinear theory uniformly valid for growing and decaying waves have been based on the Klimontovich⁵ picture, rather than on a Vlasov analysis, and have further required analytic continuation into the complex wave-number and velocity planes. As pointed out by Weitzner,⁶ analytic continuation may impose unrealistic continuity requirements on physical distributions.

The present paper is devoted to a fresh look at quasilinear theory, with a treatment of growing and decaying waves on an equal footing. The complex plane is completely avoided, even for frequency, and Landau's analysis of the linear Vlasov problem is replaced by a different technique. The nonlinear Vlasov equation is manipulated to a form exhibiting only resonant diffusion, without the need for the limit $\gamma \rightarrow 0+$. The term previously denoted non-resonant, adiabatic, or "fake" diffusion¹ no longer appears explicitly; it contributes to wave momentum and energy, and ensures the conservation of total momentum and energy.

No attempt is made here to maintain mathematical rigor. The ordering of terms is made heuristically, without the explicit introduction of multiple-time techniques.

To keep things as simple as possible, only the uniform one-dimensional Coulomb plasma is treated here. The generalization to more dimensions,⁷ and to a nonuniform⁸ Maxwell⁹ system is straightforward.

II. LINEAR ANALYSIS

When the Vlasov equation

$$\frac{\partial f(x, v; t)}{\partial t} + v \frac{\partial f}{\partial x} + \frac{e}{m} E(x, t) \frac{\partial f}{\partial v} = 0 \quad (1)$$

is linearized about a uniform state $f_0(v; t)$ which is slowly varying in time, the first-order solution, obtained by integration along characteristics, is

$$f_1(x, v; t) = -(e/m) f'_0(v; t) \int_0^\infty d\tau E(x - v\tau, t - \tau). \quad (2)$$

Here the prime denotes velocity-differentiation. The time-integration is on the fast time-scale; accordingly $f'_0(v; t - \tau)$ has been replaced by $f'_0(v; t)$ and taken outside the integral.

We look for solutions of the form:

$$E(x, t) = E_k(t) \exp(ikx - i\omega_k t), \quad (3)$$

$$f_1(x, v; t) = f_k(v; t) \exp(ikx - i\omega_k t),$$

where k and ω_k are both real, and the amplitudes have a slow time dependence representing wave growth or decay. But we note that the variation of $E_k(t)$, which is of order γ_k , is not as slow as that of $f_0(v; t)$, which is of order E_k^2 . (Our ordering thus breaks down when E_k^2 and γ_k are of comparable size, in appropriate units.)

Insertion of (3) into (2) yields

$$f_k(v; t) = -(e/m) f'_0(v; t) \int_0^\infty d\tau E_k(t - \tau) \exp i(\omega_k - kv)\tau. \quad (4)$$

The Poisson equation then reads

$$ikE_k(t) = -(4\pi e^2/m) \int dv f'_0(v; t) \int_0^\infty d\tau E_k(t - \tau) \exp i(\omega_k - kv)\tau, \quad (5)$$

where a sum over species, if needed, is implicit.

Appealing to the slow variation of $E_k(t)$, we expand

$$E_k(t - \tau) = E_k(t) - \tau \dot{E}_k(t) + \dots \quad (6)$$

and keep only the two terms shown. We then use

$$\begin{aligned} \int_0^\infty d\tau \exp i(\omega_k - kv)\tau &= \pi \delta(\omega_k - kv) + i P(\omega_k - kv)^{-1} \\ &\equiv \pi \delta_+(\omega_k - kv) \end{aligned} \quad (7)$$

and

$$\int_0^\infty d\tau \tau \exp i(\omega_k - kv)\tau = -i \partial[\pi \delta_+(\omega_k - kv)]/\partial\omega_k, \quad (8)$$

to obtain

$$\begin{aligned} ikE_k(t) &= -(4\pi e^2/m)[E_k(t) \int dv f'_0(v; t) \pi \delta_+(\omega_k - kv) \\ &\quad + i \dot{E}_k(t)(\partial/\partial\omega_k) \int dv f'_0(v; t) \pi \delta_+(\omega_k - kv)]. \end{aligned} \quad (9)$$

We now multiply (9) by $E_k^*(t)$, and multiply the complex conjugate of (9) by $E_k(t)$. Adding and subtracting the resulting equations, we find equations for the growth rate and for the eigenfrequency ω_k . The subtraction yields

$$1 = -(4\pi e^2/mk) \int dv f'_0(v; t) P(\omega_k - kv)^{-1} \quad (10)$$

(plus terms of relative order $\dot{E}_k/E_k$), which is the standard dispersion equation for the real eigenfrequency $\omega_k(t)$. The addition yields

$$\frac{d \ln |E_k(t)|^2}{dt} = \frac{2 \int dv f'_0(v; t) \pi \delta(\omega_k - kv)}{(\partial/\partial \omega_k) \int dv f'_0(v; t) P(\omega_k - kv)^{-1}} \quad (11)$$

(plus terms of relative order $\dot{E}_k/E_k$), which is the standard equation for the growth (or decay) rate $2 \gamma_k(t)$. The slow time-dependence of $\omega_k(t)$, implied by (10), is of order E^2 , and therefore is not included in (11), which is zero-order in E^2 .

III. QUASILINEAR ANALYSIS

In expanding the nonlinear Vlasov equation (1) to second order in the wave amplitudes, we shall distinguish between $f_2(v; t)$, the second-order perturbation in f (averaged over x and over oscillation period), whose size is proportional to E^2 , and $f_0(v; t)$, the part of f independent of E , whose evolution is proportional to E^2 .

Inserting (2) into the space-time average of (1), we thus obtain

$$\frac{\partial f_0}{\partial t} + \frac{\partial f_2}{\partial t} = \left(\frac{e}{m}\right)^2 \frac{\partial}{\partial v} \left[I(v, t) \frac{\partial f_0(v; t)}{\partial v} \right], \quad (12)$$

where

$$I(v, t) \equiv \int_0^\infty d\tau \langle E(x, t) E(x - v\tau, t - \tau) \rangle \quad (13)$$

(the angular bracket denotes spatial and fast-time averaging), and where $\partial f_0(v; t)/\partial t$ represents the slow evolution of f_0 due to diffusion, while $\partial f_2(v; t)/\partial t$ represents the slow variation of f_2 due to wave growth and decay. On the right side of (12), we have dropped $\partial f_2/\partial v$, but have retained $\partial f_0/\partial v$; we justify this below.

In expression (13), we generalize (3) to include all k , and several branches ℓ :

$$E(x, t) = \sum_{k, \ell} E_k^\ell(t) \exp(ikx - i\omega_k^\ell t), \quad (14)$$

with $E_{-k}^\ell(t) = E_k^{\ell*}(t)$, and $\omega_{-k}^\ell = -\omega_k^\ell$; we then obtain

$$I(v, t) = \sum_{k, \ell} \int_0^\infty d\tau E_k^{\ell*}(t) E_k^\ell(t - \tau) \exp i(\omega_k^\ell - kv)\tau. \quad (15)$$

Again we use Eqs. (6), (7), and (8), and find

$$I(v, t) = \sum_{k>0, \ell} \{ |E_k^\ell(t)|^2 2\pi \delta(\omega_k^\ell - kv) - (d|E_k^\ell|^2/dt) \partial P(\omega_k^\ell - kv)^{-1} / \partial \omega_k^\ell \}. \quad (16)$$

We insert (16) in (12), associating the first term of (16) with $\partial f_0/\partial t$, and the second term of (16) with $\partial f_2/\partial t$. We then obtain for f_0 the resonant diffusion equation:

$$\frac{\partial f_0(v; t)}{\partial t} = \frac{\partial}{\partial v} \left[D(v, t) \frac{\partial f_0}{\partial v} \right], \quad (17)$$

with

$$D(v, t) \equiv (e/m)^2 \sum_{k>0, \ell} |E_k^\ell(t)|^2 2\pi \delta(\omega_k^\ell - kv), \quad (18)$$

while the equation for f_2 can be explicitly integrated to yield

$$f_2(v; t) = -\left(\frac{e}{m}\right)^2 \frac{\partial}{\partial v} \left[f_0'(v; t) \sum_{k>0, \ell} |E_k^\ell(t)|^2 \partial P(\omega_k^\ell - kv)^{-1} / \partial \omega_k^\ell \right]. \quad (19)$$

The diffusion equation (17) includes the effects of all waves in the diffusivity (18), whether growing or damped. The second-order perturbation f_2 appears singular, but is actually quite smooth for a continuous spectrum of waves. Replacing $\sum_k$ by $V \int dk/2\pi$, and defining the spectral density

$$\mathcal{E}^{\ell}(k; t) \equiv \lim_{V \rightarrow \infty} V |E_k^{\ell}(t)|^2, \quad (20)$$

we have

$$f_2(v; t) = \left(\frac{e}{m}\right)^2 \frac{\partial}{\partial v} \left\{ f'_0(v; t) \times \int_0^{\infty} (dk/2\pi) \sum_{\ell} P(\omega_k^{\ell} - kv)^{-1} \frac{\partial}{\partial k} \left[\mathcal{E}^{\ell}(k; t) \left(\frac{\partial \omega_k^{\ell}}{\partial k} \right)^{-1} \right] \right\}. \quad (21)$$

Estimating the ratio of f'_2 to f'_0 , we find it to be of order $(\omega_k \tau_D)^{-1}$ (where τ_D is the diffusion time for f_0), and therefore very small. Hence we are justified in keeping only f'_0 on the right side of (12) and (17).

The moments of f_2 yield the particle contribution to the wave energy and momentum. The second moment, which is the wave kinetic energy density, is

$$\begin{aligned} K_2(t) &\equiv \int dv \frac{1}{2} m v^2 f_2(v; t) \\ &= \left(\frac{e}{m}\right)^2 \sum_{k>0, \ell} |E_k^{\ell}(t)|^2 \frac{\partial}{\partial \omega_k^{\ell}} \int dv m v f'_0(v; t) P(\omega_k^{\ell} - kv)^{-1} \\ &= (e^2/m) \sum_{k>0, \ell} |E_k^{\ell}(t)|^2 \frac{\partial}{\partial \omega_k^{\ell}} \left[(\omega_k^{\ell}/k) \int dv f'_0(v; t) P(\omega_k^{\ell} - kv)^{-1} \right]. \end{aligned} \quad (22)$$

When we add this to the wave electric energy density

$$\begin{aligned} W_E(t) &= \langle |E(x, t)|^2 / 8\pi \rangle \\ &= (4\pi)^{-1} \sum_{k>0, \ell} |E_k^{\ell}(t)|^2, \end{aligned} \quad (23)$$

we obtain the total wave energy density:

$$\begin{aligned} W(t) &\equiv W_E(t) + K_2(t) \\ &= (4\pi)^{-1} \sum_{k>0, \ell} |E_k^\ell(t)|^2 \frac{\partial}{\partial \omega_k^\ell} [\omega_k^\ell \epsilon(k, \omega_k^\ell; t)], \end{aligned} \quad (24a)$$

where

$$\epsilon(k, \omega_k^\ell; t) \equiv 1 + (4\pi e^2/mk) \int dv f'_0(v; t) P(\omega_k^\ell - kv)^{-1} \quad (25)$$

is the real part of the dielectric function. We note that Eq. (10) is equivalent to $\epsilon(k, \omega_k^\ell; t) = 0$, whence (24a) may be written

$$W(t) = (4\pi)^{-1} \sum_{k>0, \ell} |E_k^\ell(t)|^2 \omega_k^\ell \partial \epsilon / \partial \omega_k^\ell. \quad (24b)$$

The first moment of f_2 is the wave momentum density:

$$\begin{aligned} P_2(t) &\equiv \int dv m v f_2(v; t) \\ &= (e^2/m) \sum_{k>0, \ell} |E_k^\ell(t)|^2 \frac{\partial}{\partial \omega_k^\ell} \int dv f'_0(v; t) P(\omega_k^\ell - kv)^{-1} \\ &= (4\pi)^{-1} \sum_{k>0, \ell} |E_k^\ell(t)|^2 k \partial \epsilon(k, \omega_k^\ell) / \partial \omega_k^\ell. \end{aligned} \quad (26)$$

Comparing (26) with (24b), we see that wave momentum and wave energy satisfy the standard relation

$$\frac{(P_2)_k^\ell}{W_k^\ell} = \frac{k}{\omega_k^\ell}. \quad (27)$$

The standard conservation laws follow from these equations, if we ignore the time-dependence of ω_k^ℓ and ϵ . From (24b) and (11), we have

$$dW/dt = (e^2/m) \sum_{k>0, \ell} |E_k^\ell|^2 (\omega_k^\ell/k) \int dv f'_0(v; t) 2\pi \delta(\omega_k^\ell - kv). \quad (28)$$

The zero order kinetic energy density

$$K_0(t) \equiv \int dv \frac{1}{2} m v^2 f_0(v; t) \quad (29)$$

evolves according to (17) and (18):

$$\begin{aligned} dK_0/dt &= - \int dv m v D(v, t) f'_0 \\ &= -(e^2/m) \sum_{k>0, \ell} |E_k^\ell|^2 (\omega_k^\ell/k) \int dv f'_0(v; t) 2\pi \delta(\omega_k^\ell - kv). \end{aligned} \quad (30)$$

Comparing (28) and (30), we see that the total energy density

$$U(t) \equiv W(t) + K_0(t) \quad (31)$$

is indeed conserved.

The total momentum density

$$P(t) \equiv P_2(t) + P_0(t) \quad (32)$$

is proved constant in the same way, using

$$P_0(t) \equiv \int dv m v f_0(v; t) \quad (33)$$

and Eq. (26), with Eqs. (17), (18), and (11) for their evolution.

ACKNOWLEDGMENT

I have benefited from discussing these ideas with Blake
Patterson and David Montgomery.

REFERENCES

1. For a survey, see R. Sagdeev and A. Galeev, Nonlinear Plasma Theory, T. O'Neil and D. Book, Eds. (W. A. Benjamin, N. Y., 1969).
2. For a recent critique, see G. Vahala and D. Montgomery, Discrete Spectra and Damped Waves in Quasilinear Theory (submitted to J. Plasma Phys.).
3. J. C. Price, Phys. Fluids 10, 1623 (1967).
4. A. Rogister and C. Oberman, J. Plasma Phys. 2, 33 (1968).
5. Y. L. Klimontovich, The Statistical Theory of Non-Equilibrium Processes in a Plasma (Massachusetts Institute of Technology Press, 1967).
6. H. Weitzner, Comm. Pure and App. Math. 18, 307 (1965).
7. I. Bernstein and F. Engelmann, Phys. Fluids 9, 937 (1966).
8. A. N. Kaufman and T. Nakayama, Phys. Fluids 13, 956 (1970).
9. A. N. Kaufman, Resonant Interactions Between Particles and Normal Modes in a Cylindrical Plasma, Lawrence Radiation Laboratory Report UCRL-19839, May 1970 (submitted to Phys. Fluids).

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or*
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.*

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

TECHNICAL INFORMATION DIVISION
LAWRENCE RADIATION LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720