

UC San Diego

UC San Diego Previously Published Works

Title

Systematic light exposure in the treatment of cancer-related fatigue: a preliminary study.

Permalink

<https://escholarship.org/uc/item/8nh9d096>

Journal

Psycho-oncology, 23(12)

ISSN

1057-9249

Authors

Redd, William H
Valdimarsdottir, Heiddis
Wu, Lisa M
et al.

Publication Date

2014-12-01

DOI

10.1002/pon.3553

Peer reviewed

Clinical Correspondence

Systematic light exposure in the treatment of cancer-related fatigue: a preliminary study

William H. Redd^{1*}, Heiddis Valdimarsdottir^{1,2}, Lisa M. Wu¹, Gary Winkel¹, Emily E. Byrne¹, Melba A. Beltre¹, Elizabeth S. Liebman¹, Tanya Erazo¹, Judy A. Hayes¹, Luis Isola³, Eileen Scigliano³, Yeraz Meschian¹, Susan Lutgendorf⁴ and Sonia Ancoli-Israel⁵

¹Department of Oncological Sciences, Icahn School of Medicine at Mount Sinai, New York, NY, USA

²Department of Psychology, Reykjavik University, Iceland

³Department of Medicine, Icahn School of Medicine at Mount Sinai, New York, NY, USA

⁴Department of Psychology, University of Iowa, Iowa, IA, USA

⁵Departments of Psychiatry and Medicine, University of California, San Diego, CA, USA

*Correspondence to:

Department of Oncological Sciences, Icahn School of Medicine at Mount Sinai, New York, NY, USA. E-mail: william.redd@mssm.edu

Received: 1 October 2013

Revised: 13 March 2014

Accepted: 25 March 2014

Dear Editor,

Introduction

Cancer-related fatigue (CRF) is the most commonly reported side effect of cancer treatment [1]. It is defined by the National Comprehensive Cancer Network as ‘a distressing, persistent, subjective sense of physical, emotional, and cognitive tiredness or exhaustion related to cancer or cancer treatment that is not proportional to recent activity’. Patients feel tired even after resting, have reduced capacity to carry out normal activities, experience slow physical recovery from tasks, and report diminished concentration. Most patients experience CRF during their treatment, and many continue to experience CRF long after all treatment has ended. In our research with survivors of hematopoietic stem cell transplantation (HSCT) 1–3 years off treatment, 40% of those interviewed reported clinically significant CRF.

Various pharmacological and nonpharmacological interventions have been studied to treat CRF with generally mild to moderate effect sizes [1]. An alternate intervention may be systematic bright white light (BWL) exposure, which is far less costly and involves less patient burden than other interventions. Reviews and meta-analyses [2] have reported that BWL is effective in reducing sleep and circadian rhythm problems associated with other disorders (e.g., depression and jet lag) but has not been previously tested for the effect on fatigue. Recent research from our group has shown that, when given during chemotherapy, BWL compared with

dim red light (DRL) keeps CRF from getting worse [3]. However, to date, the impact of BWL on fatigue among survivors of cancer has not been examined.

This preliminary efficacy trial sought to determine the impact of BWL on CRF among survivors of breast and gynecologic cancers who had completed all cancer treatment and survivors of hematological malignancy who had completed HSCT. We tested the hypothesis that the BWL condition would result in a significant reduction in CRF compared with the DRL condition.

Methods

Participants

Thirty-six survivors of cancer from the Mount Sinai Hospital in New York, NY, participated in the study (Figure 1) across all four seasons. Inclusion criteria were the following: up to 3.5 years post-HSCT, up to 3 years postcompletion of chemotherapy or chemotherapy and radiation for breast cancer, or any time postcompletion of treatment for gynecologic cancer; 18 years of age or older; and reporting clinically significant fatigue as measured by the Functional Assessment of Chronic Illness Therapy-Fatigue (FACIT-Fatigue) scale. Exclusion criteria were the following: pregnancy, confounding underlying medical illnesses such as significant preexisting anemia, history of mania (which is a contraindication for light treatment) or current clinical depression, and any other physical or psychological impairment including a sleep disorder that would limit participation. The study was approved by the Program for the Protection of Human Subjects at Mount Sinai Hospital.




Figure 1. Mean FACIT-Fatigue scores for bright white light and dim red light conditions over four assessment time points. Higher FACIT-Fatigue score corresponds to less fatigue. FACIT-Fatigue score equal to or less than 30 constitutes clinically significant fatigue. Standard errors are given

Procedure

Survivors of cancer were approached by study personnel during their regular follow-up clinical visits at Mount Sinai. The study was described, signed consent obtained, and the screening/eligibility measures administered. Participants who qualified were then randomized to either the BWL ($n = 18$) or the DRL ($n = 18$) group. All participants completed questionnaires on fatigue at four time points: baseline, during the second week of the intervention, at the end of the 4-week intervention, and 3 weeks after the completion of the intervention.

Light treatment

The BWL condition was compared with the DRL condition. Traditionally, DRL has been used as a comparison for BWL as the biological clock is less sensitive to wavelengths in the red light area [4]. Both BWL and DRL treatments were administered via a Litebook 1.2 (Litebook®, Ltd., Medicine Hat, Canada). For the BWL condition, the Litebook® used 60 premium white light emitting diode (LED) lights that mimic the visible spectrum of sunlight (full spectrum white light) for minimum glare and maximum eye comfort. An identical-appearing device utilizing red LEDs emitting <50 lux was used for the DRL comparison group. Participants were instructed to self-administer the light treatment in their homes by placing the light box at a 45° angle, 18 inches from their face, for 30 min every morning within 30 min of waking throughout the 4-week intervention period.

Measures

Cancer-related fatigue was measured with the FACIT-Fatigue scale. A FACIT-Fatigue score ≤ 30 constitutes clinically significant fatigue [5]. Because depression is a known

correlate of fatigue, the Brief Symptom Inventory depression scale was also used [6].

Results

Participant characteristics

Table 1 contains descriptive medical and demographic information for the sample. The general linear model analysis

Table 1. Patient demographics and medical variables

	<i>n</i> = 36
Gender	
Male	7 (19) ^a
Female	29 (81)
Race/ethnicity	
Black	11 (31)
White	12 (33)
Other	13 (36)
Living arrangement	
Live alone	4 (13)
Live with spouse/partner	12 (40)
Live with parents	2 (7)
Other	12 (40)
Marital status	
Single	4 (13)
Married/partnered	14 (47)
Separated	5 (17)
Divorced	5 (17)
Other	2 (6)
Children	
Yes	22 (76)
No	7 (24)
Education	
High school graduate	4 (14)
Some college/postcollege	7 (24)
College graduate	10 (34)
Graduate degree	8 (28)
Employed	
No	16 (62)
Yes	10 (38)
Annual income (in dollars)	
0–9999	4 (15)
10,000–19,999	3 (12)
30,000–39,999	1 (4)
40,000–49,999	3 (11)
50,000–59,999	1 (4)
60,000–69,999	1 (4)
70,000–79,999	1 (4)
Above 80,000	12 (46)
Diagnosis	
Acute myeloid leukemia	4 (10)
Follicular non-Hodgkin's lymphoma	2 (6)
Hodgkin's lymphoma	1 (3)
Multiple myeloma	10 (28)
Myelofibrosis	1 (3)
Breast cancer	15 (41)
Endometrial cancer	2 (6)
Ovarian cancer	1 (3)
Months postdiagnosis	17.22 (10.11) ^b

^a*n* (percentage)

^bMean (SD)

of baseline fatigue indicated no significant ($p=0.4840$) difference between the DRL and BWL groups at baseline. There was also no significant ($F(1, 23)=0.95$; $p=0.3389$) difference between the two light conditions at baseline in participants' belief that the treatment to which they had been assigned would successfully reduce their fatigue.

Effect of light on CRF

Using a linear mixed model repeated measures analysis (SAS Proc MIXED), there was a significant time effect ($F(3, 68)=5.93$; $p=0.0012$) suggesting that fatigue levels changed over time. There was a significant group effect ($F(1, 28)=7.12$; $p=0.0125$) suggesting that across all time points, the BWL group had less fatigue than the DRL group. In addition, there was a significant time times treatment effect ($F(3, 68)=5.27$; $p=0.0025$) indicating that the groups differed with respect to change in fatigue over time. As can be seen in Figure 1, there was a significant improvement in fatigue over time for those in the BWL condition, while in the DRL condition, fatigue improved at 2 weeks but became worse at 4 weeks and at follow-up. There was a large effect size (d) at 0.98 [7]. When depression was added to the model as a control variable with assessment time and the treatment variable, depression was not a statistically significant ($p=0.3322$) predictor of CRF, and BWL remained a statistically significant predictor of fatigue.

At the end of the 4-week intervention, no patients in the BWL condition were still clinically fatigued (FACIT-Fatigue score >30), whereas 55% of patients in the DRL condition were still clinically fatigued.

Discussion

This preliminary efficacy trial showed that CRF among clinically fatigued survivors of cancer was significantly reduced among those who were exposed to BWL but not among those exposed to DRL. To our knowledge, this is the first study to show beneficial effects of BWL exposure on fatigue in survivors of cancer.

Although other studies have examined nonpharmacologic treatments of fatigue, the effect size ($d=0.98$) in this study was larger than the small to moderate effects seen in other interventions [1]. Three weeks following the completion of the intervention, none of the participants in the BWL condition reported clinical levels of fatigue, whereas most of the participants in the DRL condition continued to report clinical levels of fatigue. It is important to note that the BWL intervention is a low-cost and low-burden intervention compared with alternate approaches (most of which require travel to meet with a clinician and intensive clinician involvement).

Although these results are encouraging, this study is preliminary. Because the follow-up assessment was only 3 weeks, the long-term effect of BWL on CRF is not yet known. In addition, potential mechanisms underlying the

effects of BWL on CRF were not examined. BWL may affect CRF through its effects on circadian activity rhythms (CARs)—a chronobiological mechanism theorized to underlie CRF. CARs serve the function of preparing the body for restful sleep at some times of the day and for active wakefulness at others. CARs are biological cycles that are slightly longer than 24 h and are entrained to the 24 h day by environmental zeitgebers such as bright light. Several studies have observed desynchronized CARs among patients with cancer and survivors of cancer [8], and CARs have been found to be associated with fatigue in patients with cancer [9]. A recent study from our group found that BWL prevented CAR deterioration among patients with breast cancer undergoing chemotherapy [10]. Other mechanisms involving circadian neuroendocrine regulation and inflammatory dynamics may also be involved. Future studies with larger and more diverse samples are needed to determine the long-term effects of BWL on CRF and to examine the potential mechanisms underlying the effects of BWL on CRF among survivors of cancer.

Despite these limitations, the present study suggests that BWL, a safe, inexpensive, and easily delivered treatment, may be a clinically effective intervention to treat CRF among survivors of cancer who continue to experience CRF long after medical treatment has ended.

Acknowledgements

Research reported in this publication was supported by the National Cancer Institute of the National Institutes of Health under Award Numbers R21CA158954 and K05CA108955. Additional support was provided by grant UL1TR000067 from the National Center for Advancing Translational Sciences, National Institutes of Health.

Key points

- CRF lasts well beyond the end of cancer treatment.
- Light has a strong effect on sleep and circadian rhythms in other medical conditions, but the effect of light on CRF has only just begun to be tested.
- Preliminary studies in breast cancer suggest that BWL, compared with DRL, kept CRF from worsening during chemotherapy.
- The current preliminary study tested BWL versus DRL on 36 patients post-HSCT, postchemotherapy or chemotherapy and radiation for breast cancer, or postcompletion of treatment for gynecologic cancer.
- Results suggest that BWL reduced fatigue in these patients, while DRL did not. In addition, postlight treatment, no patients in the BWL group reported any residual fatigue, while 55% of those in the DRL condition still reported fatigue.

References

1. Kangas M, Bovbjerg DH, Montgomery GH. Cancer-related fatigue: a systematic and meta-analytic review of non-pharmacological therapies for cancer patients. *Psychol Bull* 2008;**134**:700–741. doi: 10.1037/a0012825.
2. Golden RN, Gaynes BN, Ekstrom RD et al. The efficacy of light therapy in the treatment of mood disorders: a review and meta-analysis of the evidence. *Am J Psychiatry* 2005;**162**(4):656–662.
3. Ancoli-Israel S, Rissling M, Neikrug A, et al. Light treatment prevents fatigue in women undergoing chemotherapy for breast cancer. *Support Care Cancer* 2012;**20**(6):1211–1219. doi: 10.1007/s00520-011-1203-z.
4. Glickman G, Byrne B, Pineda C, Hauck WW, Brainard GC. Light therapy for seasonal affective disorder with blue narrow-band light-emitting diodes (LEDs). *Biol Psychiatry* 2006;**59**(6):502–507.
5. Cella D. The Functional Assessment of Chronic Illness Therapy-Fatigue (FACIT-F) scale: Summary of development and validation. September 3 2013 Update. 2013.
6. Derogatis L. *Brief Symptom Inventory 18: administration, scoring, and procedures manual*. Minneapolis: NCS Assessments, 2001.
7. Cohen J. *Statistical Power Analysis for the Behavioral Sciences* (2nd edn), Lawrence Erlbaum Associates: Hillsdale, NJ, 1988.
8. Pati AK, Parganiha A, Kar A, Soni R, Roy S, Choudhary V. Alterations of the characteristics of the circadian rest-activity rhythm of cancer in-patients. *Chronobiol Int* 2007;**24**(6):1179–1197. doi: 10.1080/07420520701800868.
9. Payne JK. Altered circadian rhythms and cancer-related fatigue outcomes. *Integr Cancer Ther* 2011;**10**(3):221–233. doi: 10.1177/1534735410392581.
10. Neikrug AB, Rissling M, Trofimenko V, et al. Bright light therapy protects women from circadian rhythm desynchronization during chemotherapy for breast cancer. *Behav Sleep Med* 2012;**10**(3):202–216. doi: 10.1080/15402002.2011.634940.