UC Davis

UC Davis Previously Published Works

Title

Kinesin family deregulation coordinated by bromodomain protein ANCCA and histone methyltransferase MLL for breast cancer cell growth, survival, and tamoxifen resistance.

Permalink

<https://escholarship.org/uc/item/8nh9r11f>

Journal

Molecular cancer research : MCR, 12(4)

ISSN

1541-7786

Authors

Zou, June X
Duan, Zhijian
Wang, Junjian
et al.

Publication Date

2014-04-01

DOI

10.1158/1541-7786.mcr-13-0459

Peer reviewed

Molecular Cancer Research

Kinesin Family Deregulation Coordinated by Bromodomain Protein ANCCA and Histone Methyltransferase MLL for Breast Cancer Cell Growth, Survival, and Tamoxifen Resistance

June X. Zou, Zhijian Duan, Junjian Wang, et al.

Mol Cancer Res 2014;12:539-549. Published OnlineFirst January 3, 2014.

Updated version Access the most recent version of this article at:
doi:[10.1158/1541-7786.MCR-13-0459](https://doi.org/10.1158/1541-7786.MCR-13-0459)

Supplementary Material Access the most recent supplemental material at:
<http://mcr.aacrjournals.org/molcanres/suppl/2014/01/09/1541-7786.MCR-13-0459.DC1.html>

Cited Articles This article cites by 49 articles, 28 of which you can access for free at:
<http://mcr.aacrjournals.org/content/12/4/539.full.html#ref-list-1>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, contact the AACR Publications Department at permissions@aacr.org.

Kinesin Family Deregulation Coordinated by Bromodomain Protein ANCCA and Histone Methyltransferase MLL for Breast Cancer Cell Growth, Survival, and Tamoxifen Resistance

June X. Zou^{1,4}, Zhijian Duan^{2,4}, Junjian Wang^{2,4}, Alex Sokolov^{2,4}, Jianzhen Xu⁵, Christopher Z. Chen^{2,4}, Jian Jian Li³, and Hong-Wu Chen^{2,4}

Abstract

Kinesins are a superfamily of motor proteins and often deregulated in different cancers. However, the mechanism of their deregulation has been poorly understood. Through examining kinesin gene family expression in estrogen receptor (ER)-positive breast cancer cells, we found that estrogen stimulation of cancer cell proliferation involves a concerted regulation of specific kinesins. Estrogen strongly induces expression of 19 kinesin genes such as *Kif4A/4B*, *Kif5A/5B*, *Kif10*, *Kif11*, *Kif15*, *Kif18A/18B*, *Kif20A/20B*, *Kif21*, *Kif23*, *Kif24*, *Kif25*, and *KifC1*, whereas suppresses the expression of seven others, including *Kif1A*, *Kif1C*, *Kif7*, and *KifC3*. Interestingly, the bromodomain protein ANCCA/ATAD2, previously shown to be an estrogen-induced chromatin regulator, plays a crucial role in the up- and downregulation of kinesins by estrogen. Its overexpression drives estrogen-independent upregulation of specific kinesins. Mechanistically, ANCCA (AAA nuclear coregulator cancer associated) mediates E2-dependent recruitment of E2F and MLL1 histone methyltransferase at kinesin gene promoters for gene activation-associated H3K4me3 methylation. Importantly, elevated levels of *Kif4A*, *Kif15*, *Kif20A*, and *Kif23* correlate with that of ANCCA in the tumors and with poor relapse-free survival of patients with ER-positive breast cancer. Their knockdown strongly impeded proliferation and induced apoptosis of both tamoxifen-sensitive and resistant cancer cells. Together, the study reveals ANCCA as a key mediator of kinesin family deregulation in breast cancer and the crucial role of multiple kinesins in growth and survival of the tumor cells.

Implications: These findings support the development of novel inhibitors of cancer-associated kinesins and their regulator ANCCA for effective treatment of cancers including tamoxifen-resistant breast cancers. *Mol Cancer Res*; 12(4): 539–49. ©2014 AACR.

Introduction

The kinesin superfamily of motor proteins is encoded by more than 40 different genes in humans. They function by traveling unidirectionally along the microtubules for intracellular transport of molecules or organelles. Many of them such as *Kif4A*, *Kif10/CENPE*, and *Kif11/Eg5* play important roles in cell division, particularly in different stages of

mitosis and cytokinesis (1). Several kinesins including *Kif1A*, *Kif7*, and *KifC3* function primarily in nonmitotic processes such as protein complex or organelle movement and cellular signaling (2–6). Overexpression of mitotic kinesins including *KifC1*, *Kif10/CENPE*, and *Kif18A* was found in several human cancers including breast cancer (7–11). However, it is unclear whether other kinesins are aberrantly expressed in breast cancer and whether they play roles in the cancer cell proliferation and survival. Moreover, little is known about the mechanisms how the kinesin gene expression is deregulated in cancer cells.

The mitogenic effect of estrogen 17 β -estradiol (E2) is primarily through its receptor estrogen receptor- α (ER α) to control gene expression. E2, through binding to ERs, plays pivotal roles in breast cancer development and progression. In cells, E2 treatment elicits changes in expression of hundreds of genes with the ones involved in cell-cycle progression, DNA synthesis and replication being the most robustly upregulated. Interestingly, estrogen also strongly represses many antiproliferative and proapoptotic genes. Although the majority of them are likely direct targets of

Authors' Affiliations: Departments of ¹Internal Medicine, ²Biochemistry and Molecular Medicine, and ³Radiation Oncology, School of Medicine, University of California, Davis; ⁴UC Davis Cancer Center/Basic Sciences, Sacramento, California; and ⁵College of Bioengineering, Henan University of Technology, Zhengzhou, China

Note: Supplementary data for this article are available at Molecular Cancer Research Online (<http://mcr.aacrjournals.org/>).

Corresponding Author: Hong-Wu Chen, UC Davis Cancer Center/Basic Sciences, University of California, Davis, UCDMC Research III, Room 1400B, 4645 2nd Avenue, Sacramento, CA 95817. Phone: 916-734-7743; Fax: 916-734-0190; E-mail: hwzchen@ucdavis.edu

doi: 10.1158/1541-7786.MCR-13-0459

©2014 American Association for Cancer Research.

ER α , many are regulated indirectly through estrogen control of other transcriptional regulators (12–18).

ER-mediated transcriptional regulation involves assembly at target gene loci of chromatin regulators such as the p160/SRC coactivators, histone modifying or demodifying enzymes, and the ATP-dependent chromatin-remodeling proteins (15, 16, 19–21). Different ER target genes may use different sets of specific chromatin regulators, which often act through transient and dynamic protein–protein interactions (19, 22). ANCCA (AAA nuclear coregulator cancer associated), a bromodomain containing, ATPase protein, was identified by us as a novel ER α coregulator and shown to play a critical function in mediating ER activation of CCND1, MYC, and E2F1 as well as E2-stimulated breast cancer cell proliferation (23–28). Importantly, ANCCA expression is strongly induced by E2 in ER-positive breast cancer cells. Its overexpression in tumors is highly associated with poor outcomes of patients with breast cancer (26). Here, we examined the function of ANCCA in estrogen control of kinesin family gene expression. Our results revealed a pivotal role played by ANCCA in mediating both estrogen induction and repression of specific kinesins and demonstrated a critical function of ANCCA-activated kinesins in breast cancer cell proliferation and survival.

Materials and Methods

Cell culture, hormone treatment, and siRNA transfection

MCF-7 and T-47D cells were obtained from the American Type Culture Collection (ATCC) and cultured in Dulbecco's Modified Eagle Medium (DMEM) or RPMI-1640 (Invitrogen), respectively, with 10% FBS (Gemini). For ligand treatment, cells were grown in phenol red–free medium supplemented with 10% charcoal–dextran–stripped (cds) FBS (J R Scientific) for 3 days before being treated with E2 (at 10^{-8} mol/L), 4-hydroxytamoxifen (Tam, at 10^{-6} mol/L), or fulvestrant (at 10^{-7} mol/L; all from Sigma-Aldrich) for indicated times. MCF-7 cells that ectopically express ANCCA were generated by transfection with pcD-HCMV-ANCCA (23, 28), and stable transfectant cells were clonally isolated and expanded in the presence of G418 (300 μ g/mL). Tamoxifen-resistant MCF-7 subline LCC2 cells (a kind gift from Dr. Robert Clarke at Georgetown University, Washington, DC) were cultured in DMEM + 10% cds–FBS containing 10^{-6} mol/L 4-hydroxytamoxifen. For siRNA transfection, cells were seeded at 2×10^5 cells per well in 6-well plates in phenol red–free DMEM supplemented with 5% cds–FBS and transfected using DharmaFECT with siRNAs (Dharmacon) targeting ANCCA (29), MLL1 (30) or Kif4A, Kif15, Kif20A, and Kif23 (Supplementary Materials and Methods).

Quantitative real-time PCR and immunoblot analysis

Total RNA was isolated and the cDNA was prepared, amplified, and detected in the presence of SYBR as previously (29). The fluorescent values were collected and a melting curve analysis was performed. Fold difference was

calculated as described previously (29). Cell lysates were analyzed by immunoblotting with antibodies against specific kinesins, ANCCA, and other proteins. The PCR primers, details of cell lysates, and immunoblotting and antibodies are listed in the Supplementary Materials and methods.

Apoptosis and cell growth assays

For apoptosis, cells were transfected with siRNA on glass chamber slides and processed for TUNEL assay using the In Situ Cell Death Detection Kit (Roche) as previously described (26). Random fields (10 fields per condition) of TUNEL-positive cells were counted and averaged. For cell growth, cells were seeded in 6-well plates at 1×10^5 per well and treated as indicated. Total cell numbers were counted using a Coulter cell counter. The assays were performed in triplicates and the entire experiments were repeated three times.

Chromatin immunoprecipitation (ChIP) assay

ChIP was performed essentially as described previously (27, 28) with the following modifications. Cells were lysed and sonicated using Sonic Dismembrator 550 (Fisher Scientific). The crude chromatin solutions were first cleared with protein A beads (Invitrogen) that had been precoated with preimmune serum or normal immunoglobulin G (IgG) for 2 hours at 4°C. Then, the precleared solutions were incubated at 4°C overnight with antibodies against ANCCA (26), E2F1 (1:1 mixture of C-20 and KH-95; Santa Cruz Biotechnology), MLL (clone 9–12; Millipore), H3K4me3 and H3K27ac (ab8580 and ab4729, ChIP grade; both from Abcam), before precipitation with protein A beads that had been preblocked with bovine serum albumin (BSA) and sonicated salmon sperm DNA. The precipitated DNA was analyzed by real-time PCR with SYBR green on an iCycler instrument. Enrichment of genomic DNA was presented as the percentage recovery relative to the input. The primers are listed in the Supplementary Materials and Methods.

Analysis of kinesin expression in tumors for clinical outcome and ANCCA correlation and statistics

Microarray data from GSE9195 (31) were downloaded from the National Center for Biotechnology Information (NCBI) Gene Expression Omnibus (GEO) website (<http://www.ncbi.nlm.nih.gov/gds>). The dataset consists of gene expression profiles of early stage, ER-positive breast cancer tumors that had received tamoxifen only as adjuvant treatment. Normalized probe set expression intensities were obtained using robust multi-array average for probe summarization and normalization as before (29). Correlations of ANCCA/atad2 and kinesin expression were assessed by computing the Pearson correlation coefficient (r) and a two-tailed t test for significance. The Kaplan–Meier estimates were used to compute the survival curves. All above computations were conducted in R statistical package (<http://www.r-project.org/>). Kinesin expression and patient survival were also analyzed using

an online survival analysis tool (32). For the assays, the paired *t* test was performed as previously described (26).

Results

Estrogen stimulation of breast cancer cell proliferation involves a concerted up- and downregulation of specific kinesin expression

We previously demonstrated that ANCCA is an estrogen responsive gene and controls the expression of cyclins and other genes important for cell proliferation and survival (23, 26). This prompted us to investigate whether ANCCA plays any function in control of mitotic kinesins in ER-positive breast cancer cells. Thus, we first identified kinesins with expression regulated by E2 in estrogen-sensitive MCF7 cells. Remarkably, among the 38 kinesin genes with mRNA expression detected in MCF7 cells, E2 strongly stimulated the expression of a large number (19 out of 38, more than 2-fold in 12 hours and/or 24 hours of E2 treatment) of the detected kinesins, which include Kif2A, Kif3A, Kif3B, Kif4A, Kif4B, Kif5B, Kif10/CENPE, Kif11/EG5, Kif15, Kif16A, Kif18A, Kif18B, Kif20A, Kif20B, Kif21A, Kif23, Kif24, Kif25, and KifC1 (Fig. 1A). In most cases, the induction can be observed

by 12 hours of E2 stimulation. Except Kif2A, Kif5B, and Kif21A, most of them play important roles in mitosis and/or cytokinesis (1). Intriguingly, E2 also significantly repressed several kinesins, including Kif1A, Kif1C, Kif3C, Kif7, Kif13B, Kif16B, and KifC3. Notably, most of the repressed kinesins have primary functions in nonmitotic processes such as synaptic vesicle transport in neurons (Kif1A), integrin transport for cell migration (Kif1C), control of the Hedgehog (Hh)-Gli signaling (Kif7) and golgi positioning and integration with dynein (KifC3; refs. 2, 5, 33).

Given the prominent function of E2-ER in promoting breast cancer cell proliferation, we focused our further analysis on mitotic kinesins. As shown before, ANCCA and its targets cyclin D1 and CDC6 are induced by E2 in MCF7 cells (Fig. 1B). Western blotting with available antibodies confirmed the E2 induction of mitotic kinesin proteins of Kif4A, Kif11, Kif15, Kif20A, and Kif23 (Fig. 1B). To examine whether the E2 regulation is through ER α , cells were treated with ER α pure antagonist fulvestrant. Indeed, when cells were treated simultaneously with E2 and fulvestrant, the kinesin induction by E2 was mostly suppressed (Fig. 1B, top), indicating that E2 induction of Kif4A, Kif11,

Figure 1. Coordinated regulation of kinesin family expression in breast cancer cells by estrogen-ER α . A, MCF-7 cells were hormone depleted for 3 days and then treated with E2 at 10^{-8} mol/L for the indicated hours before harvesting for real-time RT-PCR analysis. Fold change was obtained by comparing the normalized quantitative PCR value from E2-treated cells with the value from cells without E2 harvested at the same time point. The data are expressed as the mean $\pm$ SD from three independent experiments. B, MCF-7 and T-47D cells were treated with E2 as in A, or fulvestran/ICI182,780 (ICI) at 10^{-7} mol/L for the indicated hours before harvesting for immunoblotting with the indicated antibodies.

Figure 2. ANCCA plays a crucial role in mediating E2-ER regulation of kinesin expression. A, MCF-7 cells were in hormone-depleted medium for 24 hours, transfected with control or ANCCA siRNA, and 48 hours posttransfection treated with E2 at 10^{-8} mol/L for 12 or 24 hours before harvesting for real-time RT-PCR analysis. Fold change was obtained by comparing the E2-induced quantitative PCR value change from ANCCA siRNA-transfected cells with the value change from control siRNA-transfected cells harvested at the same time point of E2 treatment. The data are expressed as the mean $\pm$ SD of three independent experiments. B, MCF-7 cells were siRNA transfected and treated with E2 as in A for the indicated hours before harvesting for immunoblotting with indicated antibodies.

Kif15, Kif20A, and Kif23 is through ER α . Similar results were obtained from another estrogen-sensitive cell T-47D (Fig. 1B, bottom). Together, the results suggest that estrogen

via ER α coordinately regulates kinesin family gene expression with upregulation of mitotic kinesins and downregulation of nonmitotic kinesins.

Figure 3. ANCCA overexpression enhances estrogen-dependent and independent expression of kinesins. A and B, MCF-7 cells without (vector) or with (ANCCA) ectopic expression were hormone depleted and then treated with E2 at 10^{-8} mol/L for 24 hours before harvesting for real-time RT-PCR analysis (A) or immunoblotting (B). Data represent the mean $\pm$ SD of three independent experiments (*, $P < 0.01$ for groups without E2 for each *Kif* gene; **, $P < 0.05$ for groups with E2 for each *Kif* gene; paired *t* test).

ANCCA plays a crucial role in mediating E2 regulation of kinesins

To determine whether ANCCA mediates E2 control of the kinesins, we measured their expression in MCF7 cells with ANCCA suppressed by siRNA. As shown in Fig. 2A, ANCCA suppression markedly diminished E2 induction of most (18 out of 19) of the kinesins, which include Kif2A, Kif3A, Kif3B, Kif4A, Kif4B, Kif5B, Kif10/CENPE, Kif11/EG5, Kif15, Kif16A, Kif18A, Kif18B, Kif20A,

Kif20B, Kif21A, Kif23, Kif25, and KifC1. More strikingly, for the seven kinesins that are repressed by E2, upon ANCCA silencing by siRNA, their repression by E2 was largely lost (except for Kif16B). Kif14 is the only one that was not significantly regulated by E2 but decreased by ANCCA suppression. Suppression of ANCCA by a different siRNA yielded essentially the same effects on E2-regulated kinesin mRNAs (data not shown). As expected, ANCCA suppression by the two siRNAs resulted in strong reduction

Figure 4. Estrogen induction of kinesins involves assembly and function of ANCCA, E2F, and MLL at the kinesin promoters. A and B, MCF-7 cells were treated with E2 for 1 or 3 hours and harvested for ChIP with indicated antibodies. ChIP and input DNA was analyzed by real-time PCR with primers amplifying a region in the promoter (presented by the long line for about 0.8 kb). ChIP data represent the mean $\pm$ SD of three independent experiments (*, $P < 0.006$ for groups between 0 and 3 hours of E2). C and D, MCF-7 cells were transfected with indicated siRNAs and treated with E2 for 3 hours before harvesting for ChIP with antibodies for E2F1, MLL1, and H3K4me3. ChIP data were analyzed as above and the normalized quantitative PCR values of ChIP DNA from cells treated with control siRNA and without E2 were set as 1. ChIP data represent the mean $\pm$ SD of three independent experiments (**, $P < 0.01$ for groups between siCont and siANCCA for each antibody). E and F, MCF-7 cells were treated with E2 and siRNA and analyzed, as in Fig. 2 for protein (E) and mRNA (F), except that mRNA analysis was performed with 24-hour E2 treatment (***, $P < 0.01$ for groups between siCont and siMLL1 for each Kif expression).

of protein levels of mitotic kinesin Kif4A, Kif11, Kif15, Kif20A, and Kif23 in the cancer cells (Fig. 2B).

To further examine the function of ANCCA in control of kinesin expression, we analyzed whether elevated ANCCA promotes the kinesin expression. Strikingly, in the absence of E2, ectopic ANCCA was able to stimulate the mRNA expression of Kif4A, Kif11, Kif15, Kif20A, and Kif23 to the level almost equivalent to that induced by E2 in the vector control cells. Except Kif4A, E2 further stimulated their expression in the ANCCA-overexpressing cells (Fig. 3A), suggesting that part of the ANCCA enhancement of kinesin expression is through ER α . Moreover, the ANCCA stimulatory effects can be observed on the kinesin proteins (Fig. 3B). Thus, these results strongly suggest that chromatin-associated protein ANCCA is a key mediator of estrogen-coordinated regulation of kinesin gene family in breast cancer cells.

E2 induction of kinesins involves the assembly and function of ANCCA, E2F, and their associated histone methyltransferase MLL

To examine the mechanism of ANCCA function in estrogen control of kinesin expression, we first performed ChIP assays for potential direct involvement of ANCCA. Indeed, marked ANCCA occupancy was observed at promoters of Kif4A, Kif15, Kif20A, and Kif23 upon E2 treatment for 1 or 3 hours (Fig. 4A and B and Supplementary Fig. S1). Inspection of the local sequences with ANCCA occupancy suggests binding sites for transcription factors E2Fs and c-Myc at the kinesin promoters. ChIP with anti-E2F1 antibody demonstrated a significant E2 induction of recruitment at the same locations, in support of our previous finding that ANCCA interacts with E2Fs and acts as a potent coactivator of E2Fs (27). However, no consistent, E2-induced c-Myc binding was

observed with a c-Myc antibody (data not shown). Notably, the ANCCA recruitment was associated with a significant increase of gene activation-linked histone marks such as H3K4me3 and H3K27ac, with H3K4me3 being highly elevated by E2. Because H3K4me3 is primarily deposited by histone methyltransferases such as MLL1, we next examined whether MLL1 is involved. Indeed, strong E2-induced recruitment of MLL1 at the kinesin gene promoters was observed. To determine the role of ANCCA in the E2-induced chromatin events, we performed ChIP with si-ANCCA knockdown cells. Remarkably, suppression of ANCCA strongly diminished E2-dependent recruitment of E2F1 and MLL1 at the kinesin promoters. Expectedly, MLL1-associated H3K4me3 level was also largely reduced (Fig. 4C and D and Supplementary Fig. S1). Knockdown of MLL1 strongly mitigated E2 induction of the kinesin expression (Fig. 4E and F). Together, these results suggest that ANCCA plays an important role in mediating the assembly of E2F and MLL complexes, and H3K4me3 mark elevation at the kinesin gene promoters for E2 induction of their expression.

Kif4A, Kif15, Kif20A, and Kif23 are important for proliferation and survival of tamoxifen-sensitive and resistant breast cancer cells

Several kinesins such as Kif11, Kif14, and Kif18A have recently been shown to be important for breast cancer cell proliferation. We, thus, focused on our analysis on the role of ANCCA-controlled Kif4A, Kif15, Kif20A, and Kif23 in cell growth and survival. Each of the four kinesins was knocked down by two different siRNAs. Each knockdown of Kif4A, Kif15, Kif20A, and Kif23 resulted in strong inhibition of E2-stimulated proliferation of tamoxifen-sensitive MCF7 cells (Fig. 5A). Consistent with the marked reduction of

Figure 5. Knockdown of kinesins decreases proliferation and survival of tamoxifen-sensitive breast cancer cells. A, MCF-7 cells were hormone depleted, siRNA transfected, and treated with E2 at 10^{-8} mol/L for indicated days before harvesting, for counting of viable cells. B and C, MCF-7 cells were treated with siRNA and E2 as in A for 3 days before harvesting for TUNEL assay and for immunoblotting with indicated antibodies. The experiments were repeated three times. The *P* values for differences between cells treated with control siRNA and kinesin siRNA were <0.001 .

Figure 6. Kinesins are important for proliferation and survival of tamoxifen-resistant breast cancer cells. A, MCF-7 and its tamoxifen-resistant subline LCC2 cells were treated with 4-hydroxytamoxifen (Tam) at

viable cells, depletion of each of the kinesins caused a pronounced apoptotic cell death as indicated by the number of TUNEL-positive cells and the induction of cleaved caspase-7 and PARP1 proteins (Fig. 5B and C).

To further examine the function of ANCCA in control of kinesins, we performed the knockdowns in a tamoxifen-resistant subline (LCC2) of MCF-7 cells (34–37). Comparing with the MCF-7 cells in which the kinesin and ANCCA expression was strongly inhibited by tamoxifen, LCC2 cells were essentially unresponsive to the inhibitory effect of tamoxifen on Kif4A, Kif15, Kif20A, and Kif23 (Fig. 6A). Interestingly, ANCCA level was significantly higher in LCC2 cells than MCF-7 cells, especially when the cells were treated with tamoxifen. As expected, tamoxifen treatment alone did not alter LCC2 cell growth. Silencing each kinesin in LCC2 cells, however, markedly inhibited their hormone-independent proliferation (Fig. 6B and Supplementary Fig. S2, compare siCont-Tam and siKif-Tam). Importantly, the kinesin silencing also dramatically increased LCC2 cell sensitivity to tamoxifen inhibition (compare si-Cont+Tam and siKif+Tam). Altogether, the results suggest that ANCCA-regulated kinesins play crucial roles in proliferation and survival of both tamoxifen-sensitive and resistant breast cancer cells.

High level of kinesins correlates with that of ANCCA and with poor outcome of ER-positive tumors

The above results that ANCCA-regulated kinesins play an important role in breast cancer cell growth and survival prompted us to examine the clinical significance of our experimental findings. In a dataset of gene expression profiles obtained from early stage, ER-positive breast cancer tumors that had received tamoxifen only as adjuvant treatment, high expression of Kif4A, Kif15, Kif20A, and Kif23 strongly correlated with poor relapse-free survival. Importantly, their expression also correlated significantly with the expression of ANCCA in the tumors (Fig. 7A and B and Supplementary Fig. S3). To test further, the impact of high specific kinesin expression, we used an online survival analysis tool with a large database of breast cancer (1,413 ER-positive tumors) and found that high levels of each of the four kinesins tend to associate with poor prognosis (relapse-free survival; Fig. 7C and Supplementary Fig. S3). Together, these clinical data support the role of ANCCA as a key regulator of specific kinesins and indicate that their overexpression may drive breast cancer progression such as resistance to tamoxifen therapy.

Discussion

Little is known about the mechanism of kinesin family deregulation in human cancers. In this regard, this study made several unique and unexpected findings. First, a large

10^{-6} mol/L for the indicated hours before harvesting for immunoblotting with the indicated antibodies. B, LCC2 cells were siRNA transfected and treated with 4-hydroxytamoxifen at 10^{-6} mol/L for the indicated days before harvesting for counting of viable cells. Cell numbers are presented as the mean $\pm$ SD of three independent experiments.

Figure 7. Expression of ANCCA-controlled kinesins is associated with poor outcomes in ER-positive breast tumors. A, Kaplan-Meier plots of relapse-free survival (RFS) of tamoxifen-treated patients stratified by high or low (above or below median) expression of indicated kinesins in a microarray data set as described in Materials and Methods. B, scatter plot showing a positive correlation between ANCCA and indicated kinesin expression in an ER-positive tumor data set. C, correlation of high expression of indicated kinesins with poor relapse-free survival of patients with ER-positive tumors was determined using an online survival analysis tool as described in Materials and Methods. The P value for A and C was calculated by a log-rank test and the P value for B was obtained by a two-tailed t test.

number of kinesins are regulated by estrogen in breast cancer cells. Mammalian kinesin family has 45 members and many of them are expressed in tissue or cell-specific manner (1). Remarkably, among the 38 kinesins with expression

detected in breast cancer MCF-7 cells, 26 of them are regulated by E2, therefore, likely making it one of the most closely regulated gene families in estrogen signaling. Second, estrogen regulation of kinesins is bidirectional. Although

many of the kinesins are upregulated by E2, at least seven are strongly downregulated by E2. Third, estrogen-regulated kinesins seem to play distinct and nonredundant function. Although many are mitotic regulators (Kif4A, Kif10, Kif11, Kif15, Kif18A, Kif20A, and Kif23), Kif3B is primarily involved in cell migration, and Kif24 is a centriolar kinesin and functions to remodel the local microtubules for cilia assembly (38). Knockdown of individual mitotic kinesins such as Kif4A, Kif15, Kif20A, and Kif23 demonstrated that each of them is critical for cell proliferation or survival. Interestingly, kinesins repressed by E2, which include Kif1C, Kif3C, Kif7, Kif16B, and KifC3, are mostly non-mitotic kinesins (1) and they likely play different functions too. For instance, Kif1C is involved in stabilization of cellular trailing adhesions (33). Kif1C gene is found mutated in a subset of metastatic, ER α -positive breast cancer (39). Interestingly, Kif7 plays an important role in regulation of Hedgehog (Hh)-Gli signaling and its deletion contributes to the development of skin basal cell carcinoma (40). One unexpected finding is that both E2-dependent induction and repression of kinesins involves the function of ANCCA. We reported previously that ANCCA acts as a novel coactivator of ER α (23). The fact that ANCCA depletion results in derepression of specific kinesins strongly suggests that ANCCA also plays an important role in E2-dependent transcriptional repression. Although future study is needed to understand the underlying mechanism, it is conceivable that ANCCA, through its direct interaction with ER α , is recruited to the ER-repressed target genes and that recruited ANCCA, via its AAA type ATPase activity, may facilitate the assembly of repressive protein complexes such as the corepressor-HDAC, NRIP1/RIP140, or p300-CtBP (41, 42).

We focused our mechanistic analysis on the mitotic kinesins Kif4A, Kif15, Kif20A, and Kif23. Our ChIP analysis clearly showed that ANCCA directly occupies their promoter in the absence of hormone and that its occupancy is strongly increased upon E2 treatment. Thus, ANCCA may be involved in both estrogen-dependent and independent activation of the kinesins. This is consistent with recent findings that these mitotic kinesins are overexpressed in many types of cancers such as lung cancer, pancreatic cancer, and glioma that generally do not involve strong estrogen stimulation but also show ANCCA overexpression (41–45). Interestingly, we showed here that ANCCA is required for E2F1 recruitment to the kinesin genes. We also found that ANCCA is required for E2-induced recruitment of MLL1 histone H3K4 methylase and the E2 induction of H3K4me3 mark at the kinesin gene promoter, as we demonstrated for androgen induction of EZH2 (46). H3K4me3 mark is necessary for the RNA polymerase II preinitiation complex assembly (47). Thus, together, our results suggest that upon estrogen stimulation, ANCCA occupancy at the kinesin genes is enhanced to facilitate the recruitment of E2F1 and MLL-mediated H3K4 methylation, which results in the assembly of Pol-II complex for transcriptional activation. How E2 stimulates the function of ANCCA at kinesin promoters is currently unknown. Although we were able to detect E2-induced

robust recruitment of ER α to cyclin D1 enhancer and promoter, we were unable to detect any significant ER α recruitment to the ANCCA-occupied promoter regions of the kinesin genes examined (data not shown). It is possible that ER α acts through tethering to other transcription factor(s) (14, 21) or is recruited to an uncharacterized distal enhancer of the kinesin to facilitate ANCCA loading to the promoter through chromosomal looping. Future studies will be needed to determine the functional mechanism of ER α in the induction of specific kinesins.

Importantly, this study revealed that multiple kinesins including Kif4A, Kif15, Kif20A, and Kif23 are crucial for growth and survival of both tamoxifen-sensitive and resistant breast cancer cells. We also found that among the patients treated with tamoxifen, high levels of the four kinesins are strongly associated with poor recurrence-free survival. Thus, the results strongly suggest that kinesins such as Kif4A, Kif15, Kif20A, and Kif23 can have important values as both prognostic factors and new therapeutic targets for endocrine therapy-resistant breast cancer. However, kinesins may play similar cellular functions (1), which would necessitate simultaneous targeting of multiple kinesins for effective therapy. One alternative can be blocking MLL-mediated histone methylation, which is required for the kinesin gene activation. However, members of the MLL family are found mutated or deleted in multiple human cancers (48, 49). Given ANCCA being a common activator of these kinesins, targeting ANCCA can be a more attractive option. With the small molecules targeting bromodomain proteins such as BRD4 are near clinical trial, development of inhibitors specifically targeting ANCCA bromodomain or its ATPase is highly desirable for effective treatment of aggressive breast cancer.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Authors' Contributions

Conception and design: J.J. Li, H.-W. Chen

Development of methodology: J.X. Zou, J. Wang

Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): J.X. Zou, Z. Duan, A. Sokolov, C.Z. Chen

Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis): J.X. Zou, J. Xu, H.-W. Chen

Writing, review, and/or revision of the manuscript: J.J. Li, H.-W. Chen

Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases): J. Wang

Study supervision: H.-W. Chen

Acknowledgments

The authors thank Dr. Robert Clarke for kindly providing the LCC2 cells.

Grant Support

This study was supported by a grant from National Natural Science Foundation of China (U1304311) to J.Z. Xu, a NIH grant R01CA133402 to J.J. Li, and R01DK060019 to H.-W. Chen.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received August 29, 2013; revised December 13, 2013; accepted December 13, 2013; published OnlineFirst January 3, 2014.

References

- Rath O, Kozielski F. Kinesins and cancer. *Nat Rev Cancer* 2012;12:527–39.
- Xu Y, Takeda S, Nakata T, Noda Y, Tanaka Y, Hirokawa N. Role of KIFC3 motor protein in Golgi positioning and integration. *J Cell Biol* 2002;158:293–303.
- Semiz S, Park JG, Nicoloso SM, Furcinitti P, Zhang C, Chawla A, et al. Conventional kinesin KIF5B mediates insulin-stimulated GLUT4 movements on microtubules. *EMBO J* 2003;22:2387–99.
- Wiesner C, Faix J, Himmel M, Bentzien F, Linder S. KIF5B and KIF3A/KIF3B kinesins drive MT1-MMP surface exposure, CD44 shedding, and extracellular matrix degradation in primary macrophages. *Blood* 2010;116:1559–69.
- Cheung HO, Zhang X, Ribeiro A, Mo R, Makino S, Puvion-Rodan V, et al. The kinesin protein Kif7 is a critical regulator of Gli transcription factors in mammalian hedgehog signaling. *Sci Signal* 2009;2:ra29.
- Scholey JM. Kinesin-2: a family of heterotrimeric and homodimeric motors with diverse intracellular transport functions. *Annu Rev Cell Dev Biol* 2013;29:443–69.
- Agarwal R, Gonzalez-Angulo AM, Myhre S, Carey M, Lee JS, Overgaard J, et al. Integrative analysis of cyclin protein levels identifies cyclin b1 as a classifier and predictor of outcomes in breast cancer. *Clin Cancer Res* 2009;15:3654–62.
- De S, Cipriano R, Jackson MW, Stark GR. Overexpression of kinesins mediates docetaxel resistance in breast cancer cells. *Cancer Res* 2009;69:8035–42.
- Tan MH, De S, Bebek G, Orloff MS, Wesolowski R, Downs-Kelly E, et al. Specific kinesin expression profiles associated with taxane resistance in basal-like breast cancer. *Breast Cancer Res Treat* 2012;131:849–58.
- Zhang C, Zhu C, Chen H, Li L, Guo L, Jiang W, et al. Kif18A is involved in human breast carcinogenesis. *Carcinogenesis* 2010;31:1676–84.
- Singel SM, Corneliussen C, Batten K, Fasciani G, Wright WE, Lum L, et al. A targeted RNAi screen of the breast cancer genome identifies KIF14 and TLN1 as genes that modulate docetaxel chemosensitivity in triple-negative breast cancer. *Clin Cancer Res* 2013;19:2061–70.
- Frasor J, Danes JM, Komm B, Chang KC, Lytle CR, Katzenellenbogen BS. Profiling of estrogen up- and down-regulated gene expression in human breast cancer cells: insights into gene networks and pathways underlying estrogenic control of proliferation and cell phenotype. *Endocrinology* 2003;144:4562–74.
- Bourdeau V, Deschenes J, Laperriere D, Aid M, White JH, Mader S. Mechanisms of primary and secondary estrogen target gene regulation in breast cancer cells. *Nucleic Acids Res* 2008;36:76–93.
- Kininis M, Chen BS, Diehl AG, Isaacs GD, Zhang T, Siepel AC, et al. Genomic analyses of transcription factor binding, histone acetylation, and gene expression reveal mechanistically distinct classes of estrogen-regulated promoters. *Mol Cell Biol* 2007;27:5090–104.
- Carroll JS, Meyer CA, Song J, Li W, Geistlinger TR, Eeckhoutte J, et al. Genome-wide analysis of estrogen receptor binding sites. *Nat Genet* 2006;38:1289–97.
- Magnani L, Stoeck A, Zhang X, Lanczky A, Mirabella AC, Wang TL, et al. Genome-wide reprogramming of the chromatin landscape underlies endocrine therapy resistance in breast cancer. *Proc Natl Acad Sci U S A* 2013;110:E1490–9.
- Casa AJ, Potter AS, Malik S, Lazard Z, Kuatse I, Kim HT, et al. Estrogen and insulin-like growth factor-I (IGF-I) independently down-regulate critical repressors of breast cancer growth. *Breast Cancer Res Treat* 2012;132:61–73.
- Miller TW, Balko JM, Ghazoui Z, Dunbier A, Anderson H, Dowsett M, et al. A gene expression signature from human breast cancer cells with acquired hormone independence identifies MYC as a mediator of antiestrogen resistance. *Clin Cancer Res* 2011;17:2024–34.
- Won Jeong K, Chodankar R, Purcell DJ, Bittencourt D, Stallcup MR. Gene-specific patterns of coregulator requirements by estrogen receptor- α in breast cancer cells. *Mol Endocrinol* 2012;26:955–66.
- Chen H, Lin RJ, Xie W, Wilpitz D, Evans RM. Regulation of hormone-induced histone hyperacetylation and gene activation via acetylation of an acetylase. *Cell* 1999;98:675–86.
- Stender JD, Kim K, Charn TH, Komm B, Chang KC, Kraus WL, et al. Genome-wide analysis of estrogen receptor α DNA binding and tethering mechanisms identifies Runx1 as a novel tethering factor in receptor-mediated transcriptional activation. *Mol Cell Biol* 2010;30:3943–55.
- Hah N, Danko CG, Core L, Waterfall JJ, Siepel A, Lis JT, et al. A rapid, extensive, and transient transcriptional response to estrogen signaling in breast cancer cells. *Cell* 2011;145:622–34.
- Zou JX, Revenko AS, Li LB, Gemo AT, Chen HW. ANCCA, an estrogen-regulated AAA+ ATPase coactivator for ER α , is required for coregulator occupancy and chromatin modification. *Proc Natl Acad Sci U S A* 2007;104:18067–72.
- Caron C, Lestrat C, Marsal S, Escoffier E, Curtet S, Virolle V, et al. Functional characterization of ATAD2 as a new cancer/testis factor and a predictor of poor prognosis in breast and lung cancers. *Oncogene* 2010;29:5171–81.
- Ciro M, Prosperini E, Quarto M, Grazini U, Walfridsson J, McBlane F, et al. ATAD2 is a novel cofactor for MYC, overexpressed and amplified in aggressive tumors. *Cancer Res* 2009;69:8491–8.
- Kalashnikova EV, Revenko AS, Gemo AT, Andrews NP, Tepper CG, Zou JX, et al. ANCCA/ATAD2 overexpression identifies breast cancer patients with poor prognosis, acting to drive proliferation and survival of triple-negative cells through control of B-Myb and EZH2. *Cancer Res* 2010;70:9402–12.
- Revenko AS, Kalashnikova EV, Gemo AT, Zou JX, Chen HW. Chromatin loading of E2F-MLL complex by cancer-associated coregulator ANCCA via reading a specific histone mark. *Mol Cell Biol* 2010;30:5260–72.
- Zou JX, Guo L, Revenko AS, Tepper CG, Gemo AT, Kung HJ, et al. Androgen-induced coactivator ANCCA mediates specific androgen receptor signaling in prostate cancer. *Cancer Res* 2009;69:3339–46.
- Yang P, Guo L, Duan ZJ, Tepper CG, Xue L, Chen X, et al. Histone methyltransferase NSD2/MMSET mediates constitutive NF- κ B signaling for cancer cell proliferation, survival, and tumor growth via a feed-forward loop. *Mol Cell Biol* 2012;32:3121–31.
- Caslini C, Connelly JA, Serna A, Broccoli D, Hess JL. MLL associates with telomeres and regulates telomeric repeat-containing RNA transcription. *Mol Cell Biol* 2009;29:4519–26.
- Loi S, Haibe-Kains B, Desmedt C, Wirapati P, Lallemand F, Tutt AM, et al. Predicting prognosis using molecular profiling in estrogen receptor-positive breast cancer treated with tamoxifen. *BMC Genomics* 2008;9:239.
- Gyorffy B, Lanczky A, Eklund AC, Denkert C, Budczies J, Li Q, et al. An online survival analysis tool to rapidly assess the effect of 22,277 genes on breast cancer prognosis using microarray data of 1,809 patients. *Breast Cancer Res Treat* 2010;123:725–31.
- Theisen U, Straube A, Straube A. Directional persistence of migrating cells requires Kif1C-mediated stabilization of trailing adhesions. *Dev Cell* 2012;23:1153–66.
- Arteaga CL, Koli KM, Dugger TC, Clarke R. Reversal of tamoxifen resistance of human breast carcinomas *in vivo* by neutralizing antibodies to transforming growth factor- β . *J Natl Cancer Inst* 1999;91:46–53.
- Brunner N, Frandsen TL, Holst-Hansen C, Bei M, Thompson EW, Wakeling AE, et al. MCF7/LCC2: a 4-hydroxytamoxifen resistant human breast cancer variant that retains sensitivity to the steroidal antiestrogen ICI 182,780. *Cancer Res* 1993;53:3229–32.
- Karmakar S, Foster EA, Blackmore JK, Smith CL. Distinctive functions of p160 steroid receptor coactivators in proliferation of an estrogen-independent, tamoxifen-resistant breast cancer cell line. *Endocr Relat Cancer* 2011;18:113–27.
- Riggs KA, Wickramasinghe NS, Cochrum RK, Watts MB, Klinge CM. Decreased chicken ovalbumin upstream promoter transcription factor II expression in tamoxifen-resistant breast cancer cells. *Cancer Res* 2006;66:10188–98.
- Kobayashi T, Tsang WY, Li J, Lane W, Dynlacht BD. Centriolar kinesin Kif24 interacts with CP110 to remodel microtubules and regulate ciliogenesis. *Cell* 2011;145:914–25.

39. Shah SP, Morin RD, Khattri J, Prentice L, Pugh T, Burleigh A, et al. Mutational evolution in a lobular breast tumour profiled at single nucleotide resolution. *Nature* 2009;461:809–13.
40. Li ZJ, Nieuwenhuis E, Nien W, Zhang X, Zhang J, Puvion-Randall V, et al. Kif7 regulates Gli2 through Sufu-dependent and -independent functions during skin development and tumorigenesis. *Development* 2012;139:4152–61.
41. Hao H, d'Alincourt-Salazar M, Kelley KM, Shatnawi A, Mukherjee S, Shah YM, et al. Estrogen-induced and TAFII30-mediated gene repression by direct recruitment of the estrogen receptor and co-repressors to the core promoter and its reversal by tamoxifen. *Oncogene* 2007;26:7872–84.
42. Stossi F, Madak-Erdogan Z, Katzenellenbogen BS. Estrogen receptor alpha represses transcription of early target genes via p300 and CtBP1. *Mol Cell Biol* 2009;29:1749–59.
43. Imai K, Hirata S, Irie A, Senju S, Ikuta Y, Yokomine K, et al. Identification of HLA-A2-restricted CTL epitopes of a novel tumour-associated antigen, KIF20A, overexpressed in pancreatic cancer. *Br J Cancer* 2011;104:300–7.
44. Takahashi S, Fusaki N, Ohta S, Iwahori Y, Iizuka Y, Inagawa K, et al. Downregulation of KIF23 suppresses glioma proliferation. *J Neurooncol* 2012;106:519–29.
45. Taniwaki M, Takano A, Ishikawa N, Yasui W, Inai K, Nishimura H, et al. Activation of KIF4A as a prognostic biomarker and therapeutic target for lung cancer. *Clin Cancer Res* 2007;13:6624–31.
46. Duan Z, Zou JX, Yang P, Wang Y, Borowsky AD, Gao AC, et al. Developmental and androgenic regulation of chromatin regulators EZH2 and ANCCA/ATAD2 in the prostate via MLL histone methylase complex. *Prostate* 2013;73:455–66.
47. Lin JJ, Lehmann LW, Bonora G, Sridharan R, Vashisht AA, Tran N, et al. Mediator coordinates PIC assembly with recruitment of CHD1. *Genes Dev* 2011;25:2198–209.
48. Ellis MJ, Ding L, Shen D, Luo J, Suman VJ, Wallis JW, et al. Whole-genome analysis informs breast cancer response to aromatase inhibition. *Nature* 2012;486:353–60.
49. Grasso CS, Wu YM, Robinson DR, Cao X, Dhanasekaran SM, Khan AP, et al. The mutational landscape of lethal castration-resistant prostate cancer. *Nature* 2012;487:239–43.