

UC Berkeley

UC Berkeley Previously Published Works

Title

Trifluoromethylation of Arylsilanes with [(phen)CuCF₃]

Permalink

<https://escholarship.org/uc/item/8p93j3df>

Journal

Angewandte Chemie International Edition, 55(28)

ISSN

1433-7851

Authors

Morstein, Johannes

Hou, Haiyun

Cheng, Chen

et al.

Publication Date

2016-07-04

DOI

10.1002/anie.201601163

Peer reviewed



Published in final edited form as:

Angew Chem Int Ed Engl. 2016 July 4; 55(28): 8054–8057. doi:10.1002/anie.201601163.

Trifluoromethylation of Arylsilanes with (phen)CuCF₃


Johannes Morstein[†], Haiyun Hou[†], Chen Cheng, and Prof. Dr. John F. Hartwig^{*}

Department of Chemistry, University of California, Berkeley Berkeley, CA 94720 (USA)

Abstract

A method for the trifluoromethylation of arylsilanes is reported. The reaction occurs with (phen)CuCF₃ as the CF₃-source under mild, oxidative conditions with high functional group compatibility. This transformation complements prior trifluoromethylation of arenes in several ways. Most important, this method converts arylsilanes formed by the silylation of aryl C-H bonds to trifluoromethylarenes, thereby allowing the conversion of arenes to trifluoromethylarenes. The unique capabilities of the reported method are demonstrated by the conversion of a C-H bond to a C-CF₃ bond in active pharmaceutical ingredients that do not undergo this overall transformation by alternative functionalization processes, including a combination of borylation and trifluoromethylation.

Graphical abstract



Keywords

Arylsilanes; Cross Coupling; C-Si Functionalization; Late Stage Functionalization; Trifluoromethylation

Fluoroalkyl substituents modulate physicochemical properties, metabolic stability, and protein-ligand interactions.^[1] For this reason, approximately 30% of the most recently approved drugs contain fluorine, and nearly half of these fluorinated molecules contain a trifluoromethyl group.^[2] Yet, most commercial syntheses of trifluoromethylarenes still rely on classical, harsh Swarts reactions^[3] of benzotrichlorides with HF and SF₅ that are unsuitable for transformations of molecules containing most functional groups. Most commercial syntheses of trifluoromethyl heteroarenes are conducted by condensation reactions of trifluoroacetic acid derivatives.^[4] Although milder than the Swarts reactions,

^{*} jhartwig@berkeley.edu.

[†] J. Morstein and H. Hou contributed equally to this work.

these methods limit the position on the heteroarene at which a trifluoromethyl group can be installed. In both cases, these methods require that the trifluoromethyl group be installed early in a synthetic sequence.

More recently, arylboron reagents have been shown to undergo copper-mediated, oxidative trifluoromethylation reactions.^[5–11] The ability to form trifluoromethylarenes from arylboronates is important because the arylboronates can be formed by metal-catalyzed borylation of aryl C-H bonds, which occurs with regioselectivities that are controlled by the steric environment at the aryl C-H bonds.^[12–13]

Yet, organosilanes would be attractive alternatives to arylboronates as precursors to trifluoromethylarenes because silanes are inexpensive^[14] and because arylsilanes and heteroarylsilanes are more stable than arylboronates.^[15] Recent advances in the regioselective, catalytic silylation of C-H bonds in unactivated arenes and heteroarenes have greatly expanded the accessibility of arylsilanes.^[16–19] The silylation of C-H bonds forms arylsilanes with regioselectivities that are enhanced or distinct from those of C-H borylation.^[17, 20] The greater selectivity and stability enables the late stage functionalization of arenes and heteroarenes in complex molecules to form arylsilane intermediates that would be unstable as the analogous arylboronate. Thus, if a method to convert arylsilanes to trifluoromethyl arenes could be developed, a method to prepare trifluoromethylarenes from a wide range of arenes and heteroarenes, via the products of C-H bond silylation, would result.

We report the trifluoromethylation of readily accessible, stable arylsilanes. These reactions occur with (phen)CuCF₃ as the CF₃-source and air as the oxidant. The reactions occur with a broad scope of arenes, heteroarenes and active pharmaceutical ingredients and, when combined with C-H bond silylation, provide access to trifluoromethylarenes directly from both simple and complex arenes.

To initiate our search for reagents and conditions that lead to the trifluoromethylation of arylsilanes, we investigated conditions reported for the trifluoromethylation of arylboranes.^[5–11] We tested this reaction with PhSiMe(OSiMe₃)₂ (Ph-HMTS for phenyl heptamethyltrisiloxy) because Ar-HMTS compounds can be formed by the silylation of aryl and heteroaryl C-H bonds. A range of common trifluoromethylation reagents, including both electrophilic CF₃-sources (Togni's reagent^[21] and Umemoto's reagent^[22]) and nucleophilic CF₃-sources (Ruppert's reagent^[23], Langlois' reagent^[24], Chen's reagent^[25], and (phen)CuCF₃^[26]) (Scheme 1) were tested as the source of the CF₃ group.

We conducted reactions with a series of fluoride sources to activate the arylsilane by forming a hypervalent silicon species. The identity of the source of fluoride is likely to be crucial for the activation of the silane; the fluoride must be sufficiently nucleophilic to add to the silane, but it must not trigger rapid cleavage of the carbon-silicon bond. Most combinations of the sources of trifluoromethyl groups and fluoride we tested yielded only trace-amounts or no desired product (see the Supporting Information for a list of these conditions).

However, reactions of Ph-HMTS with (phen)CuCF₃ (**1**) and added fluoride in air formed the trifluoromethylarene product in a measurable yield (20%).^[27] (phen)CuCF₃ is commercially available and is a free-flowing solid that can be stored indefinitely under nitrogen and

weighed in air. Our group previously reported the trifluoromethylation of aryl iodides^[26], aryl boronates^[27], and heteroaryl bromides^[28] with **1** (Scheme 2). Thus, a convenient method to convert arylsilanes to trifluoromethylarenes would be created if the yields of this initial reaction could be improved.

Further studies showed that the trifluoromethylation of Ph-HMTS (Table 1) can occur in high yields with the proper fluoride source and solvent. The reactions with AgF under air occurred in higher yields than those with KF, KHF₂, or TBAF. Interestingly, reactions in the presence of benzoquinone formed trifluoromethylbenzene in a higher yield (71%) than with air as oxidant alone (62% yield). The same reaction with benzoquinone under nitrogen gave the product in 42% yield, suggesting that oxygen serves as the oxidant in this system.^[29] Finally, an evaluation of these conditions with a series of arylsilanes, including phenyl trimethylsilane, triphenyl silanol, and both dimethoxymethyl and trimethoxy phenylsilane (see the Supporting Information) showed that the yield of trifluoromethylbenzene was highest from the reaction of phenyl heptamethyltrisiloxane.

With these conditions in hand, we explored the scope of the trifluoromethylation of arylsilanes bearing the HMTS group (Table 2). The reaction of arylsilanes with **1** tolerates a wide range of functional groups. Substrates bearing chloride, ether, ketone, ester, amide, nitrile, or trifluoromethyl functionality (**3a–f**) gave the trifluoromethylarene in good yields. In general, the reactions of more electron-deficient arenes (**3a–e**) formed the trifluoromethylarene products in higher yields than did reaction with electron-rich arenes (**3g**). More specifically, the 3,4-dimethylphenylsilane gave the corresponding trifluoromethylxylene in a yield (34%) that was lower than that of most other examples. Reaction of a substrate containing a bromide substituent (**3h**) also gave the product in a moderate yield (39%), most likely because of competing trifluoromethylation at the aryl bromide.^[30]

We subsequently explored the scope of the trifluoromethylation of heteroarylsilanes (Table 3). The reaction occurs with five-membered ring heteroarylsilanes, including benzo-fused examples, such as silylindoles. The reaction also occurs with six-membered ring heteroarylsilanes, including those derived from pyridines, pyrazines and quinolines. The ability to conduct the reaction with both electron-rich and electron-poor heteroarenes demonstrates the exceptionally broad scope of the conditions we developed.

The trifluoromethylation of heteroarylsilanes is particularly valuable because heteroarylsilanes are less prone to oxidation and protodemetalation than are heteroarylboronates. For example, trifluoromethyl pyrazines (**6c**) cannot be prepared from an isolated heteroarylboronate because of rapid protodeborylation.^[31] Moreover, the combination of C-H bond silylation and trifluoromethylation of the heteroarylsilane provides isomers that complement those generated from direct trifluoromethylation. For example, the reaction of **5a** gave 3-trifluoromethylpyrrole in an excellent yield (96%), whereas direct trifluoromethylation with common trifluoromethylation reagents, such as Togni's reagent,^[32] trifluoromethyl iodide,^[33] and Umemoto's reagent,^[34] form the product from reaction at the 2'-position of pyrrole. Although the borylation of Boc-protected pyrrole

occurs at the 3'-position, the yields of trifluoromethyl pyrroles from the corresponding heteroarylboronates are below 50%.^[35]

Encouraged by the high functional group tolerance and broad substrate scope of the trifluoromethylation of arylsilanes, we evaluated the applicability of our conditions to the late-stage trifluoromethylation of pharmaceutically active molecules (Table 4). We prepared analogs of clopidogrel (antiaggregant) and aripiprazole (antipsychotic) containing HMDS groups under conditions we reported previously.^[18] Most striking, the trifluoromethylation of the silylated clopidogrel gave the corresponding trifluoromethylthiophene in good yield (89%), tolerating all four functional groups present with only minor epimerization of the stereocenter (3.8 % of **7** epimerized). The overall yield of **8** for the two-step sequence comprising silylation and trifluoromethylation was 77% (68% isolated, 95.1% ee). Likewise, reaction of the silylated aripiprazole gave the corresponding trifluoromethylarene in 53% yield (41% isolated). Compounds **8** and **10** are not accessible by an analogous sequence comprising borylation and trifluoromethylation. The borylation of clopidogrel gave a mixture of constitutional isomers, as determined by GC and ¹¹B-NMR spectroscopy, and the borylated aripiprazole is unstable; this arylboronate decomposed during purification on silica gel.

In conclusion, we report the first method for the trifluoromethylation of arylsilanes. Trifluoromethylation of arylsilanes with **1** proceeds with high functional group compatibility, broad scope of arenes and heteroarenes, and is suitable for the functionalization of pharmaceutically active molecules. Furthermore, the trifluoromethylation of arylsilanes gives access to complex, trifluoromethylated molecules that are inaccessible by the sequence of borylation and trifluoromethylation.

Experimental Section

General procedure for the synthesis of trifluoromethyl arenes from arylsilanes with **1**: To an oven-dried 20 mL vial was added **1** (0.36 mmol, 1.2 equiv), dry DMF (3.0 mL, stored under nitrogen), AgF (0.60 mmol, 2.0 equiv), *p*-benzoquinone (0.90 mmol, 3.0 equiv), and arylsilane (0.30 mmol, 1.0 equiv) under air. The suspension was purged with dry air for 30 seconds, and the vial was sealed with a Teflon-lined cap and heated at 50 °C for 16 h under vigorous stirring. The suspension was allowed to cool to room temperature. Water was added to the reaction mixture (5 mL) and extracted with diethylether (3 × 5 mL). The organic layer was washed with a saturated NaCl-solution (1 × 2 mL), dried over anhydrous MgSO₄, concentrated, and the residue purified by flash column chromatography or preparative thin layer chromatography. See Supporting Information for detailed procedures.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank the NIH (GM-11581201) for support of this work. J. M. thanks the German National Academic Foundation (Studienstiftung) for supporting his research in the Hartwig lab. We thank Michael Mormino for a

sample of (phen)CuCF₃, Jeff Pelton (UC Berkeley NMR facility) for expert technical assistance, and Johnson-Matthey for a gift of [Ir(COD)Cl]₂.

References

1. Purser S, Moore PR, Swallow S, Gouverneur V. *Chem. Soc. Rev.* 2008; 37:320–330. [PubMed: 18197348]
2. Zhou Y, Wang J, Gu Z, Wang S, Zhu W, Aceña JL, Soloshonok VA, Izawa K, Liu H. *Chem. Rev.* 2016
3. Swarts F. *Bull. Acad. Roy. Belg.* 1892; 24:309.
4. Petrov, VA. *Flourinated Heterocyclic Compounds: Synthesis, Chemistry, and Applications*. Vol. 1. Weinheim: John Wiley & Sons; 2009.
5. Chu L, Qing F-L. *Org. Lett.* 2010; 12:5060–5063. [PubMed: 20923196]
6. Khan BA, Buba AE, Gooßen LJ. *Chem. Eur. J.* 2012; 18:1577–1581. [PubMed: 22238176]
7. Li Y, Wu L, Neumann H, Beller M. *Chem. Comm.* 2013; 49:2628–2630. [PubMed: 23431556]
8. Liu T, Shao X, Wu Y, Shen Q. *Angew. Chem. Int. Ed.* 2012; 51:540–543.
9. Senecal TD, Parsons AT, Buchwald SL. *J. Org. Chem.* 2011; 76:1174–1176. [PubMed: 21235259]
10. Xu J, Luo D-F, Xiao B, Liu Z-J, Gong T-J, Fu Y, Liu L. *Chem. Comm.* 2011; 47:4300–4302. [PubMed: 21380454]
11. Ye Y, Künzi SA, Sanford MS. *Org. Lett.* 2012; 14:4979–4981. [PubMed: 22984900]
12. Hartwig JF. *Acc. Chem. Res.* 2012; 45:864–873. [PubMed: 22075137]
13. Ishiyama T, Takagi J, Ishida K, Miyaura N, Anastasi NR, Hartwig JF. *J. Am. Chem. Soc.* 2002; 124:390–391. [PubMed: 11792205]
14. Yaroshevsky AA. *Geochem. Int.* 2006; 44:48–55.
15. Denmark SE, Ambrosi A. *Org. Process Res. Dev.* 2015; 19:982–994. [PubMed: 26478695]
16. Cheng C, Hartwig JF. *J. Am. Chem. Soc.* 2014; 136:12064–12072. [PubMed: 25082802]
17. Cheng C, Hartwig JF. *Science.* 2014; 343:853–857. [PubMed: 24558154]
18. Cheng C, Hartwig JF. *J. Am. Chem. Soc.* 2015; 137:592–595. [PubMed: 25514197]
19. Toutov AA, Liu W-B, Betz KN, Fedorov A, Stoltz BM, Grubbs RH. *Nature.* 2015; 518:80–84. [PubMed: 25652999]
20. Cheng C, Hartwig JF. *Chem. Rev.* 2015
21. Eisenberger P, Gischig S, Togni A. *Chem. Eur. J.* 2006; 12:2579–2586. [PubMed: 16402401]
22. Teruo U, Sumi I. *Tetrahedron Lett.* 1990; 31:3579–3582.
23. Ruppert I, Schlich K, Volbach W. *Tetrahedron Lett.* 1984; 25:2195–2198.
24. Langlois BR, Laurent E, Roidot N. *Tetrahedron Lett.* 1991; 32:7525–7528.
25. Chen Q-Y, Wu S-W. *J. Chem. Soc., Chem. Commun.* 1989:705–706.
26. Morimoto H, Tsubogo T, Litvinas ND, Hartwig JF. *Angew. Chem. Int. Ed.* 2011; 50:3793–3798.
27. Litvinas ND, Fier PS, Hartwig JF. *Angew. Chem. Int. Ed.* 2012; 51:536–539.
28. Mormino MG, Fier PS, Hartwig JF. *Org. Lett.* 2014; 16:1744–1747. [PubMed: 24621138]
29. Although some reactions without added benzoquinone form the trifluoromethylarene in good yield, most reactions of arenes that are not sensitive to oxidation occurred in about 20% higher yield in the presence of benzoquinone than in the absence of this oxidant. See the Supporting Information for a series of comparisons in the presence and absence of benzoquinone.
30. This assertion is supported by the observation of additional Aryl-CF₃ compounds by 19F-NMR spectroscopy.
31. Sadler SA, Tajuddin H, Mkhaldid IAI, Batsanov AS, Albesa-Jove D, Cheung MS, Maxwell AC, Shukla L, Roberts B, Blakemore DC, Lin Z, Marder TB, Steel PG. *Organic & Biomolecular Chemistry.* 2014; 12:7318–7327. [PubMed: 25116330]
32. Wiehn MS, Vinogradova EV, Togni A. *J. Fluorine Chem.* 2010; 131:951–957.
33. Kino T, Nagase Y, Ohtsuka Y, Yamamoto K, Uraguchi D, Tokuhisa K, Yamakawa T. *J. Fluorine Chem.* 2010; 131:98–105.


34. Yang J-J, Kirchmeier RL, Shreeve JnM. *J. Org. Chem.* 1998; 63:2656–2660. [PubMed: 11672133]
35. Reaxys Search

Author Manuscript


Author Manuscript

Author Manuscript

Author Manuscript




X = Hal, B(OR)₂, NH₂



Togni's reagent

TMSCF₃


Rupperts' reagent



Umemoto's reagent

CF₃CO₂K

Langlois' reagent



(phen)CuCF₃


FO₂SCF₂CO₂Me

Chen's reagent


Scheme 1.

Common reagents for the trifluoromethylation of prefunctionalized arenes.



previous work




X = I, B(OR)₂



this work



Scheme 2.
Copper-mediated trifluoromethylation of arylsilanes.



Het = Heteroarene

Table 1

Optimization of trifluoromethylation of $\text{PhSiMe(OSiMe}_3)_2$.^[a]

Entry	Oxidant	Additive	Fluoride (2.0 eq)	Solvent	Yield ^[b]
1	Air	-	KF	DMF	20%
2	Air	-	KHF ₂	DMF	38%
3	Air	-	AgF	DMF	62%
4	Air	-	TBAF	DMF	17%
5	Air	KOTMS (2.0 eq)	-	DMF	-
6	-	-	KHF ₂	DMF	8%
7	Air	-	AgF	DMSO	58%
8	Air	BQ (3.0 eq) ^[c]	AgF	DMF	71%
9	-	BQ (3.0 eq) ^[c]	AgF	DMF	42%

^[a]Reactions run on a 0.1 mmol scale.

^[b]¹⁹F-NMR yields are determined with 4-trifluoromethoxyanisole as an internal standard.

^[c]BQ = *p*-Benzoquinone.





Table 2

Trifluoromethylation of Arylsilanes.^[a,b]

^[a]Reactions run on a 0.1 mmol scale to determine ¹⁹F-NMR yields and run on a 0.3 mmol scale to obtain yields of the isolated products. ¹⁹F-NMR yields are listed first followed by yields of isolated products in parenthesis.

^[b]¹⁹F-NMR yields are determined with 4-trifluoromethoxyanisole as an internal standard.

[c] Reaction run without benzoquinone.

Author Manuscript







Author Manuscript

Author Manuscript

Author Manuscript

Table 3

Trifluoromethylation of Heteroarylsilanes.^[a,b]

Heteroaryl—[Si]	1 (1.2 eq), AgF (2.0 eq) Benzoquinone (3.0 eq)	Heteroaryl—CF ₃
5	Air, DMF, 50°C, 16h [Si] = SiMe(OSiMe ₃) ₂	6
		
6a 96 %		6c 60% ^[c]
		
6d 63% (51%)		6f 89% (76%)
		
6g 62%		6i 51% (39%)

^[a]Reactions run on a 0.1 mmol scale to determine ¹⁹F-NMR yields and run on a 0.3 mmol scale to obtain yields of the isolated products. ¹⁹F-NMR yields are listed first followed by yields of isolated products in parenthesis.


^[b]¹⁹F-NMR yields are determined with 4-trifluoromethoxyanisole as an internal standard.

^[c]Reaction run at 80°C.

Table 4

Late-stage trifluoromethylation of pharmaceutically active molecules.^[a,b]

Yield of Trifluoromethylation: 89% (77%)



Yield of Trifluoromethylation: 53% (41%)

^[a]¹⁹F-NMR yields are listed first followed by yields of isolated products in parenthesis.^[b]¹⁹F-NMR yields are determined with 4-trifluoromethoxyanisole as an internal standard.