Lawrence Berkeley National Laboratory

Recent Work

Title

REGGE TRAJECTORIES FOR A SINGLE ATTRACTIVE YUKAWA POTENTIAL

Permalink

<https://escholarship.org/uc/item/8pr0v64d>

Authors

Ahmadzadeh, Akabar
Burde, Phillip G.
Tate, Cecil.

Publication Date

1962-03-27

RECEIVED
LIBRARY
RADIATION LABORATORY

UCRL 10140

c. 2

DOCUMENTS SECTION

University of California

Ernest O. Lawrence Radiation Laboratory

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

REGGE TRAJECTORIES FOR A SINGLE ATTRACTIVE YUKAWA POTENTIAL

Berkeley, California

UCRL-10140C

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Research and Development

UCRL-10140
UC-34 Physics
TID-4500 (17th Ed.)

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

Contract No. W-7405-eng-48

REGGE TRAJECTORIES FOR
A SINGLE ATTRACTIVE YUKAWA POTENTIAL

Akbar Ahmadzadeh, Phillip G. Burke, and Cecil Tate

March 27, 1962

REGGE TRAJECTORIES FOR
A SINGLE ATTRACTIVE YUKAWA POTENTIAL

Akbar Ahmadzadeh, Philip G. Burke, and Cecil Tate

Lawrence Radiation Laboratory
University of California
Berkeley, California

March 27, 1962

ABSTRACT

The Regge trajectories of the poles in the S matrix for nonrelativistic scattering by a single attractive Yukawa potential have been investigated. They are found to be closed loops in the complex angular-momentum plane starting from, and returning to, the negative integers as the energy goes from $-\infty$ to $+\infty$. Detailed results are graphically presented.

REGGE TRAJECTORIES FOR
A SINGLE ATTRACTIVE YUKAWA POTENTIAL*

Akbar Ahmadzadeh, Philip G. Burke, and Cecil Tate

Lawrence Radiation Laboratory
University of California
Berkeley, California

March 27, 1962

Recent conclusions by Regge and coworkers¹ concerning poles of the scattering amplitude in the complex angular-momentum plane for nonrelativistic scattering have been shown by Chew et al.² to have an important bearing on the interpretation of relativistic dispersion relations. Many features of low-energy resonances and high-energy cross sections can also be understood in terms of these poles.² It is of interest to investigate in detail the behavior of these pole "trajectories" in the nonrelativistic case to gain further insight into this and also the more general relativistic problem.

The Schrödinger equation for a single Yukawa potential has been solved numerically on IBM 709/7090 computers, and the poles of the S matrix found. Both positive and negative energies were investigated, and the Regge trajectories of the poles as a function of energy were traced in the complex angular-momentum plane. Several attractive potential strengths were investigated.

The Schrödinger equation can be written

$$\left[\frac{d^2}{dr^2} + k^2 - \frac{\ell(\ell+1)}{r^2} + \frac{A \exp(-r)}{r} \right] u(r) = 0. \quad (1)$$

The boundary condition imposed at the origin for all ℓ was

$$u(r) \sim r^{\ell+1}, \quad (2)$$

and Eq. (1) was then integrated numerically, by using standard techniques, to a region where the asymptotic form

-2-

$$u(r) \sim f_+(\ell, k)e^{ikr} - f_-(\ell, k)e^{-ikr} \quad (3)$$

is applicable.

The S matrix is defined by

$$S(\ell, k) = \frac{f_+(\ell, k)}{f_-(\ell, k)} e^{i\pi\ell} \quad (4)$$

For a fixed energy, k^2 , we vary ℓ to search for the zeros of $[S(\ell, k)]^{-1}$ corresponding to the poles of $S(\ell, k)$. A standard Newton's method is used in this search. Once a pole is located, k^2 is varied slightly and the search repeated starting in the neighborhood of the original zero. By varying k^2 from large negative to large positive values, we trace a complete trajectory of the Regge poles, which is found to be a closed loop in the complex ℓ plane.

It should be noted that, at least as far as the first Regge trajectory (that trajectory which lies furthest to the right in the complex ℓ plane) is concerned, no matter how small the potential strength, $\text{Re } \ell$ is greater than -0.5 near $k^2 = 0$. It is an interesting hypothesis that this may be true for all Regge curves (see Fig. 1). However, we have not yet been able to obtain conclusive evidence on this point.

For small k^2 , the standard threshold behavior for the curves is

$$\text{Im } \ell \sim k^{2\ell_0+1} \quad (5a)$$

and

$$\text{Re } \ell \sim A k^2 + B k^{2\ell_0+1} + \ell_0, \quad (5b)$$

where ℓ_0 is the threshold value of the position of the pole. This gives confidence in the correctness of the numerical procedure. Another check is provided in the large-energy behavior ($k^2 \rightarrow \pm\infty$) of Eq. (1). In these limits,

only that region of r near the origin, where the potential is comparable to k^2 , is important. The potential can be replaced in Eq. (1) by A/r in this region, and Singh³ has shown analytically that for this Coulomb potential, the position of the poles is given by

$$\ell = -n + \frac{1}{2} \frac{A}{\sqrt{k^2}} \quad (6)$$

where $n = 1, 2, 3, \dots$ gives the first, second, third, etc. poles respectively. It is found that the Regge curves for large $|k^2|$ go over smoothly to the analytic form in Eq. (6), and, in particular, the curves go to $-n$ as $|k^2| \rightarrow \infty$. It is only necessary to carry out the numerical investigation in a region where Eq. (6) is not applicable in order to trace the full curve. The final results are presented in Figs. 1 and 2.

We are greatly indebted to Professor G. F. Chew and other members of the Physics Department of the Lawrence Radiation Laboratory for advice and encouragement in this work.

FOOTNOTES AND REFERENCES

- * This work was performed under the auspices of the U. S. Atomic Energy Commission.
1. T. Regge, Nuovo Cimento 14, 951 (1959); 18, 947 (1960). See also A. Bottino, A. M. Longini, and T. Regge, Potential Scattering for Complex Energy and Angular Momentum (University of Turin preprint, 1961).
 2. G. F. Chew, S. C. Frautschi, and S. Mandelstam, Regge Poles in $\pi\pi$ Scattering, UCRL-9925, November 9, 1961 and Phys. Rev. (to be published). See also Geoffrey F. Chew, S-Matrix Theory of Strong Interactions without Elementary Particles, UCRL-10058, February 1962.
 3. Virendra Singh, Analyticity in the Complex Angular-Momentum Plane of the Coulomb-Scattering Amplitude, UCRL-9972, December 1961.

FIGURE LEGENDS

Fig. 1. The first few Regge trajectories for the potential strength $A = 5$.

The k^2 values are written adjacent to the curves. The small positive k^2 values for the third and fourth curves have not yet been obtained owing to numerical difficulties.

Fig. 2. The first Regge trajectory for a number of potential strengths A .

The k^2 values are written adjacent to the curves (When k^2 is negative, some of these values are shown displaced from the real axis for the sake of clarity.)

MU-26482

Fig. 1. The first few Regge trajectories for the potential strength $A = 5$. The k^2 values are written adjacent to the curves. The small positive k^2 values for the third and fourth curves have not yet been obtained owing to numerical difficulties.

MU-26481

Fig. 2. The first Regge trajectory for a number of potential strengths A . The k^2 values are written adjacent to the curves (When k^2 is negative, some of these values are shown displaced from the real axis for the sake of clarity.)

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.