



Published in final edited form as:

Curr Opin Microbiol. 2021 June ; 61: 58–66. doi:10.1016/j.mib.2021.03.002.

New discoveries expand possibilities for carboxysome engineering

Julia Borden, David Savage

Department of Molecular & Cellular Biology, UC Berkeley, Berkeley, CA 94720 USA

Abstract

Carboxysomes are CO₂-fixing protein compartments present in all cyanobacteria and some proteobacteria. These structures are attractive candidates for carbon assimilation bioengineering because they concentrate carbon, allowing the fixation reaction to occur near its maximum rate, and because they self-assemble in diverse organisms with a set of standard biological parts. Recent discoveries have expanded our understanding of how the carboxysome assembles, distributes itself, and sustains its metabolism. These studies have already led to substantial advances in engineering the carboxysome and carbon concentrating mechanism into recombinant organisms, with an eye towards establishing the system in industrial microbes and plants. Future studies may also consider the potential of in vitro carboxysomes for both discovery and applied science.

Keywords

carboxysome; carbon fixation; Self-assembly; bioengineering; Plant engineering; Biotechnology; bacterial microcompartment

Introduction

All cyanobacteria and many chemoautotrophic proteobacteria use specialized proteinaceous organelles called carboxysomes to facilitate CO₂ fixation. Carboxysomes have fascinated researchers and biotechnologists for both their icosahedral structure and ability to enable efficient carbon fixation kinetics. It's estimated that ~10–25% of CO₂ fixed globally passes through these compartments annually [1,2]. Since they were first purified in 1973 [3], researchers have sought to both understand and engineer carboxysomes. Although the presence of Rubisco signaled a critical role in CO₂ fixation, studies on carboxysomes continue to reveal new and unexpected components, structures, and potential applications.

Corresponding Author: Savage, David (dsavage@berkeley.edu).

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Competing interests

D.F.S. is a co-founder of Scribe Therapeutics and a scientific advisory board member of Scribe Therapeutics and Mammoth Biosciences. All other authors declare no competing interests.

Carboxysomes are icosahedral protein assemblies ranging from 100–500 nm in diameter, depending on the species [4]. They have a proteinaceous shell composed primarily of hexameric proteins and are capped with pentameric proteins at the icosahedral vertices. In general, they enclose Rubisco, carbonic anhydrase (CA), a Rubisco nucleating protein, and, likely, Rubisco activase. There are two lineages of carboxysomes, α and β , which evolved convergently in freshwater and coastal cyanobacteria (β lineage) and marine cyanobacteria and proteobacteria (α lineage) [5]. Remarkably, both lineages arrived at the same general carboxysome structure and function, though they differ in gene organization and protein sequences.

Carboxysomes function within a broader metabolic network called the Carbon Concentrating Mechanism, or CCM (Figure 1a). Inorganic carbon transporters in the cell membrane pump HCO_3^- into the lumen, raising its concentration to about 30x the equilibrium concentration in water [2,6]. The disequilibrium between HCO_3^- and CO_2 is advantageous because it stockpiles a charged, and therefore membrane-impermeable, form of carbon in the cell. This preferences the dehydration reaction in the carboxysome, concentrating CO_2 near Rubisco. Rubisco is thus poised to operate near its V_{max} when carboxylating ribulose-1,5-bisphosphate (RuBP) to produce two molecules of 3-phosphoglycerate (3-PG) (Figure 1b). In addition to carboxylation, Rubisco can also oxygenate RuBP; the product of this off-target reaction must be recycled via wasteful photorespiration pathways. The high CO_2 environment of the carboxysome therefore competitively inhibits oxygenation, and it remains an open question as to whether exclusion of O_2 by the shell is necessary for CCM function [2]. Finally, this unique environment enabled Rubisco evolution to maximize for carboxylation activity, and carboxysomal Rubiscos are among some of the fastest known Rubiscos, despite having low specificity for CO_2 over O_2 [7,8].

Knocking out various components of the CCM renders cells incapable of growing at the atmospheric CO_2 concentration ($\sim 0.04\%$), and they must be grown in high CO_2 ($\sim 1\text{--}10\%$) [9,10]. In particular, carbonic anhydrase must be active only inside of the carboxysome; knocking it out or expressing it in the cytosol destroys the CCM [11]. Carboxysomes must also limit CO_2 permeability so that it doesn't diffuse away from Rubisco. Pentamer deletion strains, which produce carboxysomes with pores at the icosahedral vertices, only grow in high CO_2 [12]. A precise understanding of how shell proteins limit CO_2 diffusion while allowing entry and exit of other intermediates such as 3-PG and RuBP is still not well understood. Despite speculation that the carboxysome shell is selectively permeable, promoting uptake of HCO_3^- while blocking O_2 , no direct evidence has been experimentally measured. Mathematical models show that the CCM does not require O_2 impermeability to function [2,13], though both O_2 and CO_2 may encounter an increased resistance at the hexamer pore compared to HCO_3^- and 3-PG [13]. In total, these results show that concentrating CO_2 near Rubisco by limiting CO_2 leakage from the carboxysome is essential to the function of the CCM and is an important principle in the development of biotechnological tools to concentrate CO_2 .

From carboxysome structure to functional CCM reconstitution

Reconstituting functional carboxysomes, i.e. those that can concentrate carbon, into an alternative host organism has been a major academic and bioengineering goal. However, it has been difficult to do because structure alone cannot recapitulate the CCM. The first recombinantly produced carboxysomes were α -carboxysomes from the model proteobacterium *Halothiobacillus neapolitanus*, expressed in *E. coli*. Expressing the native 10-gene operon was sufficient to produce wild-type looking carboxysomes [14]. Similarly, a synthetic operon of 12 genes from cyanobacterium *Synechococcus elongatus* PCC7942 produced wild-type looking β -carboxysomes in *E. coli* [15]. The engineered heterologous systems in both studies possessed active Rubisco, but it remained unclear whether recombinant carboxysomes could concentrate carbon, arguably the carboxysome's essential feature. A transposon mutagenesis screen of *H. neapolitanus* under high CO₂ vs. low CO₂ conditions revealed dozens of new genes responsible for the functioning of the α -carboxysome CCM, including several uncharacterized proteins in a secondary operon [16]. Characterizing unknown hits, as well as undertaking a systematic study in β -carboxysomes beyond the previous screens [9,10], will be crucial to uncovering what components are necessary to fully reconstitute the CCM. The following sections highlight recently discovered proteins and protein activities and how they may translate into using CCMs for enhancing metabolism.

New discoveries in carboxysome assembly and CCM function

Researchers made early progress in identifying and characterizing major players in the carboxysome and CCM such as Rubisco, CA, shell proteins, and carbon transporters, reviewed in greater detail in references [5,17]. This section reviews proteins discovered or characterized in recent years that have greatly increased our understanding of the carboxysome and CCM, and which are important new entries in the carboxysome biotechnological toolbox.

CsoS2 & CcmM –

CsoS2, from α -carboxysomes, and CcmM, from β -carboxysomes, are essential for carboxysome assembly and structure. Though they have no sequence or domain homology (Figure 2a & b), they share many striking similarities. Both are conserved, essential proteins located in the core carboxysome locus of their respective lineages [17]. Both are highly abundant in the carboxysome, with numbers roughly equal to Rubisco holoenzyme [5].

Notably, both CsoS2 and CcmM bind Rubisco and facilitate carboxysome nucleation. Both α - and β -carboxysome Rubiscos evolved binding sites that bridge two large subunits while making contacts with the small subunit (Figure 2c) [18,19]. This likely ensures that only the 16-subunit Rubisco holoenzyme is encapsulated during carboxysome assembly. Both CsoS2 and CcmM Rubisco-binding domains contain 3–5 repeat motifs separated by predicted disordered sequences (Figure 2a & b). Despite these shared features, the binding domains differ in their secondary structure. In CsoS2, the Rubisco-binding N-terminal domain (NTD) repeats are alpha helical, while the C-terminal CcmM repeats have structural similarity to the Rubisco small subunit (termed small subunit-like or SSUL) [18–20]. In CsoS2, a single

repeat binds with low affinity, but multivalent interactions could promote high affinity binding across multiple Rubiscos, thus nucleating carboxysome assembly (Figure 2d) [19]. Likewise, the three CcmM repeats together bind Rubisco with micromolar affinity [18,20]. Following a common theme for repetitive, multivalent proteins, both Rubisco-binding domains of CsoS2 and CcmM were shown to undergo liquid-liquid phase separation (LLPS) with Rubisco, though it should be noted both studies required salt concentrations below that of physiological 150 mM [18,19].

CsoS2 and CcmM both have a short and long isoform, and the significance of this is not fully understood (Figure 2a & b). The isoforms are produced in CsoS2 by ribosomal frameshifting and by an internal ribosome entry site (IRES) in CcmM [21,22]. In wild-type α -carboxysomes, both the short (CsoS2A) and long (CsoS2B) forms are present at a roughly equimolar ratio [21]. When the frameshifting site is mutated, CsoS2B is sufficient to reconstitute carboxysomes on its own, but CsoS2A cannot [21]. Relatedly, CcmM has a short (M35) and long (M58) form. Both M35 and M58 are needed for functional β -carboxysomes [22].

Both CsoS2 and CcmM may possess redox-regulated intrinsic flexibility. Suggestively, in *Thermosynechococcus elongatus* BP-1, the CcmM C-terminal γ -CA is only active under disulfide-forming oxidizing conditions [23]. Most repeat segments of both CsoS2 and CcmM contain 1–2 cysteines. Cells with β -carboxysomes with mutated CcmM cysteines grew 2–3 times slower than wild-type, and many carboxysomes were irregularly shaped [18]. The effect in α -carboxysomes is not yet known. In biochemical studies, the reduced form of CcmM repeats bound Rubisco with higher affinity but showed less mobility under LLPS conditions [18]. This suggests a model in which the carboxysome nucleates under reducing cytosolic conditions and, upon complete assembly, matures into a liquid-like oxidizing environment (Figure 2d). Microscopy of developing β -carboxysomes using a redox-sensitive GFP suggested that this model may be true in vivo [24]. The effect of redox regulation in these compartments remains an understudied, yet potentially highly significant, aspect of their assembly and function.

McdA & McdB –

Cells with β -carboxysomes arrange them linearly along a central longitudinal axis throughout growth and equally distribute carboxysomes to daughter cells during division [25]. This organization is driven by a pair of proteins, McdA and McdB [26]. McdA is a ParA-type ATPase that binds the nucleoid and shows a characteristic oscillatory behavior between cell poles. McdB, by analogy to plasmid partitioning systems [27], is thus thought to engage both the carboxysome and McdA. This facilitates an even distribution of carboxysomes, and those that lack either or both proteins show carboxysome clumping at a polar end. Expression of carboxysomes in a strain that lacks the proper positioning and partitioning machinery results in carboxysome aggregation, and loss of carbon fixation function in descendants without carboxysomes [25,28]. However, cells with McdA/B knockouts do not require high CO₂ to grow, likely because carboxysome-less cells can simply produce new ones, though their doubling time is significantly longer [25].

DabA & DabB –

Inorganic carbon (C_i) transporters are essential to the CCM as active C_i accumulation powers the downstream action of the carboxysome [2,6]. A review by Price et al. summarizes five of the C_i uptake systems [6]. Recently, the DAB complex joined this list. DabA and DabB form a membrane-bound complex that appears to couple CO_2 transport into the cell to a cation gradient [16,29,30].

Rubisco activases –

Rubisco is prone to inhibition by its substrate, RuBP, and other sugar derivatives. Rubisco activases catalyze release of this inhibitor. These enzymes are essential in plants and algae, but do not appear to be essential in carboxysome-containing bacteria [31,32]. The activases are divergent in the two carboxysomal lineages: α -lineages contain activase CbbQ and associated protein CbbO, while β -lineages contain β -Rca. Through convergent but different mechanisms, both activases bind Rubisco and are likely targeted to the carboxysome [32–35]. Though the biochemistry of these activases is increasingly understood, more research needs to be done to understand their role in carboxysomal carbon fixation.

Bioengineering the carboxysome and CCM

Optimizing the bacterial CCM

CCM-enhanced microbes could serve many bioindustrial applications seeking to take advantage of CO_2 -dependent metabolism. New discoveries suggest optimization could start with Rubisco (Figure 3c). Fixation flux could, in theory, be improved via encapsulation of a faster Rubisco, many of which were recently discovered and characterized [8]. In contrast to carboxysomal Form I Rubiscos, most of the fastest Rubiscos are Form II, and would need to be engineered for carboxysome targeting likely using CsoS2, CcmM, or other encapsulation peptides [36,37]. The carboxysome appears to be sensitive to the type of Rubisco it encapsulates - cells with an orthologous Form Ia Rubisco expressed in an α -carboxysome did not grow well in air, and replacement with a Form II lacked carboxysomes and required high CO_2 for growth [38,39]. A recombinant Rubisco may also require its cognate Rubisco activase to be expressed in the carboxysome.

To engineer a heterologous bacterial host to utilize a carboxysomal CCM, more genes are needed than just those in the major carboxysomal operon, which typically contains Rubisco, a carboxysomal nucleating protein, CA, and shells. A complete reconstitution of the *H. neapolitanus* CCM in *E. coli* required expression of a secondary operon alongside the major operon, thus enabling Rubisco-dependent *E. coli* to grow at atmospheric CO_2 [40]. This secondary operon included the DAB inorganic carbon transporter, the CbbO and CbbQ Rubisco activase complex, and acRAF, a proposed Rubisco chaperone [41], along with several other unknown ORFs. Of these, both the DAB and acRAF were shown to be essential CCM components in the native organism *H. neapolitanus* [16]. The DAB complex and bicarbonate transporter SbtA are the only transporters that have been demonstrated to be active when expressed in a heterologous system [16,42], making them useful candidates for CCM engineering. This successful reconstitution marks a substantial progression in our knowledge from carboxysome structure to CCM function.

Now that a first-principles study in *E. coli* has shown heterologous reconstitution of a carboxysomal CCM to be possible, expansion into industrial hosts is a logical next step (Figure 3a). Baumgart et al. expressed the *H. neapolitanus* carboxysome operon in the biotechnologically relevant bacterium *Corynebacterium glutamicum*, but carboxysomes were small and malformed [43]. Despite this, there is precedent for recombinant bacterial microcompartments to express in many diverse bacterial species, as was demonstrated with expression of the Pdu compartment in over 6 different hosts [44]. Introducing the CCM into eukaryotic hosts such as yeast may enable efforts to engineer bioindustrial strains that are better equipped to utilize carboxylation as part of a metabolic engineering strategy [45]. In other cases, the CO₂-dependent growth behavior of autotrophic strains, such as *C. necator*, could be improved through introduction of a CCM [46]. Finally, lower DNA payloads are advantageous for recombinant CCM engineering, and researchers are testing the limits of minimal carboxysome systems by eliminating unnecessary proteins or creating fusions (Figure 3d). Many of these minimal systems show structural integrity and Rubisco activity [47–50].

Optimizing the plant CCM

Much effort has gone towards creating bacterial CCMs in plants. Many agriculturally important C3 plants such as wheat and rice lack CCMs, and instead devote ~5% of leaf biomass to Rubisco [51], consuming large amounts of nitrogen in the process. Plants with engineered carboxysomal CCMs could theoretically increase yield while consuming far less nitrogen [52]. Many groups have proposed how to engineer carboxysomal CCMs into plants, and readers are directed to cited papers for more in-depth details [53,54]. In general, the engineering milestones are as follows: (1) Insert bicarbonate transporters into the chloroplast inner membrane to raise the concentration of bicarbonate in the stroma, (2) Express carboxysomes in the chloroplast, and (3) Knock out stromal carbonic anhydrases in order to maintain a high ratio of HCO₃⁻ to CO₂ (Figure 3b).

Initial efforts towards this ultimate goal are already underway. Minimal α - and β -carboxysomes have been expressed in chloroplasts, and studies showed the formation of carboxysome-like structures [49,55,56]. Cyanobacterial Rubiscos expressed in plants maintained kinetic properties equivalent to those of their native host [49,56,57]. As expected due to lack of bicarbonate transporters, plants only grew under high CO₂ conditions, though with severe growth deficiencies compared to wild-type. Single-gene bicarbonate transporters BicA and SbtA have been expressed in the chloroplast inner envelope membrane, though it is unclear if they had activity [58,59]. Future efforts will thus need to focus on identifying, characterizing, and testing transporters that are capable of functional heterologous expression. A recent survey of dissolved inorganic carbon transporters in bacteria may provide useful candidates [60]. It is also possible that additional components such as the partitioning proteins McdA and McdB will improve growth by ensuring even carboxysome distribution among dividing chloroplasts in leaf cells.

Future Directions

The relative simplicity of carboxysomes opens up the possibility of creating in vitro structures capable of performing carbon concentration, fixation, and other activities in order

to understand and engineer function (Figure 3e). Rubisco and CsoS2 or CcmM readily form liquid-separated droplets in vitro, achieving the first step of cargo nucleation [18,19]. A logical next step is to show partitioning of other cargo proteins such as a CA and Rubisco activase into the droplets, followed by shell encapsulation. Experiments to test preferential partitioning of metabolites such as RuBP, HCO_3^- , or CO_2 into the light or dense phase could probe whether or not LLPS plays a role in metabolite transfer and CO_2 concentration.

Recent insights on carboxysome structure and assembly are now enabling them to be re-engineered for alternative metabolisms, a domain which has mostly been limited to other types of bacterial microcompartments (Figure 3f) [61]. Recently, Li et al. expressed an [FeFe]-hydrogenase and ferredoxin in the α -carboxysome shell in *E. coli* to enhance H_2 production while shielding the hydrogenase from inactivating O_2 . They observed an increase in H_2 in an aerobic environment compared to unencapsulated enzyme [37]. This kind of study opens doors for exciting new biotechnological applications of carboxysomes and structures engineered from them, while continuing to shed light on basic carboxysome biology. In particular, this study suggests that the carboxysome is an O_2 -excluding environment, a theory which has generated significant discussion [2,13]. In addition, shell protein pore engineering (Figure 3g) may continue to further enable novel metabolism, including even redox-based reactions, while also providing exciting new insights into how carboxysomes permit entry of substrates, exit of products, and restrict loss of intermediates.

Conclusion

Carboxysomes are unique among protein microcompartments for their ability to concentrate CO_2 and turn it into a useful cellular product. They are a biotechnologist's dream: they self-assemble in diverse organisms with a set of standard biological parts. They are, however, deceptively simple structures. Recent studies reveal that we are still discovering many of the proteins necessary to build functional carboxysome-based CO_2 -concentrating systems. These and future discoveries will prove crucial to making meaningful engineering advances.

Acknowledgements

The authors thank Cecilia Blikstad and Luke Oltrogge for thoughtful comments. The work was supported by a grant from the U.S. Department of Energy (DE-SC00016240).

References

1. Flombaum P, Gallegos JL, Gordillo RA, Rincón J, Zabala LL, Jiao N, Karl DM, Li WKW, Lomas MW, Veneziano D, et al.: Present and future global distributions of the marine Cyanobacteria *Prochlorococcus* and *Synechococcus*. *Proc Natl Acad Sci USA* 2013, 110:9824–9829. [PubMed: 23703908]
2. Mangan NM, Flamholz A, Hood RD, Milo R, Savage DF: pH determines the energetic efficiency of the cyanobacterial CO_2 concentrating mechanism. *Proc Natl Acad Sci USA* 2016, 113:E5354–62. [PubMed: 27551079]
3. Shively JM, Ball F, Brown DH, Saunders RE: Functional organelles in prokaryotes: polyhedral inclusions (carboxysomes) of *Thiobacillus neapolitanus*. *Science* 1973, 182:584–586. [PubMed: 4355679]
4. Espie GS, Kimber MS: Carboxysomes: cyanobacterial RubisCO comes in small packages. *Photosyn. Res* 2011, 109:7–20.

5. Rae BD, Long BM, Badger MR, Price GD: Functions, compositions, and evolution of the two types of carboxysomes: polyhedral microcompartments that facilitate CO₂ fixation in cyanobacteria and some proteobacteria. *Microbiol. Mol. Biol. Rev* 2013, 77:357–379. [PubMed: 24006469]
6. Price GD: Inorganic carbon transporters of the cyanobacterial CO₂ concentrating mechanism. *Photosyn. Res* 2011, 109:47–57.
7. Flamholz AI, Prywes N, Moran U, Davidi D, Bar-On YM, Oltrogge LM, Alves R, Savage D, Milo R: Revisiting Trade-offs between Rubisco Kinetic Parameters. *Biochemistry* 2019, 58:3365–3376. [PubMed: 31259528]
8. Davidi D, Shamshoum M, Guo Z, Bar-On YM, Prywes N, Oz A, Jablonska J, Flamholz A, Wernick DG, Antonovsky N, et al.: Highly active rubiscos discovered by systematic interrogation of natural sequence diversity. *EMBO J* 2020, doi:10.15252/embj.2019104081.
9. Marcus Y, Schwarz R, Friedberg D, Kaplan A: High CO₂ Requiring Mutant of *Anacystis nidulans* R(2). *Plant Physiol* 1986, 82:610–612. [PubMed: 16665079]
10. Price GD, Badger MR: Isolation and Characterization of High CO₂-Requiring-Mutants of the Cyanobacterium *Synechococcus* PCC7942 : Two Phenotypes that Accumulate Inorganic Carbon but Are Apparently Unable to Generate CO₂ within the Carboxysome. *Plant Physiol* 1989, 91:514–525. [PubMed: 16667063]
11. Price GD, Badger MR: Expression of Human Carbonic Anhydrase in the Cyanobacterium *Synechococcus* PCC7942 Creates a High CO₂-Requiring Phenotype : Evidence for a Central Role for Carboxysomes in the CO₂ Concentrating Mechanism. *Plant Physiol* 1989, 91:505–513. [PubMed: 16667062]
12. Cai F, Menon BB, Cannon GC, Curry KJ, Shively JM, Heinhorst S: The pentameric vertex proteins are necessary for the icosahedral carboxysome shell to function as a CO₂ leakage barrier. *PLoS ONE* 2009, 4:e7521. [PubMed: 19844578]
13. Faulkner M, Szabó I, Weetman SL, Sicard F, Huber RG, Bond PJ, Rosta E, Liu L-N: Molecular simulations unravel the molecular principles that mediate selective permeability of carboxysome shell protein. *Sci. Rep* 2020, 10:17501. [PubMed: 33060756]
14. Bonacci W, Teng PK, Afonso B, Niederholtmeyer H, Grob P, Silver PA, Savage DF: Modularity of a carbon-fixing protein organelle. *Proc Natl Acad Sci USA* 2012, 109:478–483. [PubMed: 22184212]
15. Fang Y, Huang F, Faulkner M, Jiang Q, Dykes GF, Yang M, Liu L-N: Engineering and Modulating Functional Cyanobacterial CO₂-Fixing Organelles. *Front. Plant Sci* 2018, 9:739. [PubMed: 29922315]
16. Desmarais JJ, Flamholz AI, Blikstad C, Dugan EJ, Laughlin TG, Oltrogge LM, Chen AW, Wetmore K, Diamond S, Wang JY, et al.: DABs are inorganic carbon pumps found throughout prokaryotic phyla. *Nat. Microbiol* 2019, 4:2204–2215. [PubMed: 31406332]
17. Kerfeld CA, Melnicki MR: Assembly, function and evolution of cyanobacterial carboxysomes. *Curr. Opin. Plant Biol* 2016, 31:66–75. [PubMed: 27060669]
- 18**. Wang H, Yan X, Aigner H, Bracher A, Nguyen ND, Hee WY, Long BM, Price GD, Hartl FU, Hayer-Hartl M: Rubisco condensate formation by CcmM in β -carboxysome biogenesis. *Nature* 2019, 566:131–135. [PubMed: 30675061] Characterized the Rubisco-binding domain of CcmM, and implicated LLPS and redox as important factors in carboxysome biogenesis.
- 19**. Oltrogge LM, Chaijarasphong T, Chen AW, Bolin ER, Marqusee S, Savage DF: Multivalent interactions between CsoS2 and Rubisco mediate α -carboxysome formation. *Nat. Struct. Mol. Biol* 2020, 27:281–287. [PubMed: 32123388] Characterized the long enigmatic CsoS2 as a multivalent Rubisco-nucleating factor in α -carboxysomes.
20. Ryan P, Forrester TJB, Wroblewski C, Kenney TMG, Kitova EN, Klassen JS, Kimber MS: The small RbcS-like domains of the β -carboxysome structural protein CcmM bind RubisCO at a site distinct from that binding the RbcS subunit. *J. Biol. Chem* 2019, 294:2593–2603. [PubMed: 30591587]
21. Chaijarasphong T, Nichols RJ, Kortright KE, Nixon CF, Teng PK, Oltrogge LM, Savage DF: Programmed Ribosomal Frameshifting Mediates Expression of the α -Carboxysome. *J. Mol. Biol* 2016, 428:153–164. [PubMed: 26608811]

22. Long BM, Tucker L, Badger MR, Price GD: Functional cyanobacterial beta-carboxysomes have an absolute requirement for both long and short forms of the CcmM protein. *Plant Physiol* 2010, 153:285–293. [PubMed: 20304968]
23. Peña KL, Castel SE, de Araujo C, Espie GS, Kimber MS: Structural basis of the oxidative activation of the carboxysomal gamma-carbonic anhydrase, CcmM. *Proc Natl Acad Sci USA* 2010, 107:2455–2460. [PubMed: 20133749]
24. Chen AH, Robinson-Mosher A, Savage DF, Silver PA, Polka JK: The bacterial carbon-fixing organelle is formed by shell envelopment of preassembled cargo. *PLoS ONE* 2013, 8:e76127. [PubMed: 24023971]
25. Savage DF, Afonso B, Chen AH, Silver PA: Spatially ordered dynamics of the bacterial carbon fixation machinery. *Science* 2010, 327:1258–1261. [PubMed: 20203050]
- 26*. MacCready JS, Hakim P, Young EJ, Hu L, Liu J, Osteryoung KW, Vecchiarelli AG, Ducat DC: Protein gradients on the nucleoid position the carbon-fixing organelles of cyanobacteria. *elife* 2018, 7.Characterized the proteins responsible for β -carboxysome distribution and partitioning.
27. Garner EC, Campbell CS, Mullins RD: Dynamic instability in a DNA-segregating prokaryotic actin homolog. *Science* 2004, 306:1021–1025. [PubMed: 15528442]
28. MacCready JS, Tran L, Basalla JL, Hakim P, Vecchiarelli AG: The McdAB system positions α -carboxysomes in proteobacteria. *Mol. Microbiol* 2021, doi:10.1111/mmi.14708.
29. Scott KM, Leonard JM, Boden R, Chaput D, Dennison C, Haller E, Harmer TL, Anderson A, Arnold T, Budenstein S, et al.: Diversity in CO₂-Concentrating Mechanisms among Chemolithoautotrophs from the Genera *Hydrogenovibrio*, *Thiomicrothabdis*, and *Thiomicrospira*, Ubiquitous in Sulfidic Habitats Worldwide. *Appl. Environ. Microbiol* 2019, 85.
30. Mangiapia M, USF MCB4404L, Brown T-RW, Chaput D, Haller E, Harmer TL, Hashemy Z, Keeley R, Leonard J, Mancera P, et al.: Proteomic and Mutant Analysis of the CO₂ Concentrating Mechanism of Hydrothermal Vent Chemolithoautotroph *Thiomicrospira crunogena*. *J. Bacteriol* 2017, 199.
31. Sutter M, Roberts EW, Gonzalez RC, Bates C, Dawoud S, Landry K, Cannon GC, Heinhorst S, Kerfeld CA: Structural Characterization of a Newly Identified Component of α -Carboxysomes: The AAA+ Domain Protein CsoCbbQ. *Sci. Rep* 2015, 5:16243. [PubMed: 26538283]
32. Lechno-Yossef S, Rohnke BA, Belza ACO, Melnicki MR, Montgomery BL, Kerfeld CA: Cyanobacterial carboxysomes contain an unique rubisco-activase-like protein. *New Phytol* 2020, 225:793–806. [PubMed: 31518434]
33. Tsai Y-CC, Lapina MC, Bhushan S, Mueller-Cajar O: Identification and characterization of multiple rubisco activases in chemoautotrophic bacteria. *Nat. Commun* 2015, 6:8883. [PubMed: 26567524]
34. Tsai Y-CC, Ye F, Liew L, Liu D, Bhushan S, Gao Y-G, Mueller-Cajar O: Insights into the mechanism and regulation of the CbbQO-type Rubisco activase, a MoxR AAA+ ATPase. *Proc Natl Acad Sci USA* 2020, 117:381–387. [PubMed: 31848241]
35. Flecken M, Wang H, Popilka L, Hartl FU, Bracher A, Hayer-Hartl M: Dual functions of a rubisco activase in metabolic repair and recruitment to carboxysomes. *Cell* 2020, doi:10.1016/j.cell.2020.09.010.
36. Kinney JN, Salmeen A, Cai F, Kerfeld CA: Elucidating essential role of conserved carboxysomal protein CcmN reveals common feature of bacterial microcompartment assembly. *J. Biol. Chem* 2012, 287:17729–17736. [PubMed: 22461622]
- 37*. Li T, Jiang Q, Huang J, Aitchison CM, Huang F, Yang M, Dykes GF, He H-L, Wang Q, Sprick RS, et al.: Reprogramming bacterial protein organelles as a nanoreactor for hydrogen production. *Nat. Commun* 2020, 11:5448. [PubMed: 33116131] Demonstration of a metabolically repurposed carboxysome.
38. Menon BB, Dou Z, Heinhorst S, Shively JM, Cannon GC: *Halothiobacillus neapolitanus* carboxysomes sequester heterologous and chimeric RubisCO species. *PLoS ONE* 2008, 3:e3570. [PubMed: 18974784]
39. Pierce J, Carlson TJ, Williams JG: A cyanobacterial mutant requiring the expression of ribulose biphosphate carboxylase from a photosynthetic anaerobe. *Proc Natl Acad Sci USA* 1989, 86:5753–5757. [PubMed: 2503824]

- 40**. Flamholz AI, Dugan E, Blikstad C, Gleizer S, Ben-Nissan R, Amram S, Antonovsky N, Ravishankar S, Noor E, Bar-Even A, et al.: Functional reconstitution of a bacterial CO₂ concentrating mechanism in *E. coli*. *elife* 2020, 9. The first demonstration of a functional carboxysomal CCM in a recombinant host.
41. Wheatley NM, Sundberg CD, Gidaniyan SD, Cascio D, Yeates TO: Structure and identification of a pterin dehydratase-like protein as a ribulose-bisphosphate carboxylase/oxygenase (RuBisCO) assembly factor in the α -carboxysome. *J. Biol. Chem* 2014, 289:7973–7981. [PubMed: 24459150]
42. Du J, Förster B, Rourke L, Howitt SM, Price GD: Characterisation of cyanobacterial bicarbonate transporters in *E. coli* shows that SbtA homologs are functional in this heterologous expression system. *PLoS ONE* 2014, 9:e115905. [PubMed: 25536191]
43. Baumgart M, Huber I, Abdollahzadeh I, Gensch T, Frunzke J: Heterologous expression of the *Halothiobacillus neapolitanus* carboxysomal gene cluster in *Corynebacterium glutamicum*. *J. Biotechnol* 2017, 258:126–135. [PubMed: 28359868]
44. Graf L, Wu K, Wilson JW: Transfer and analysis of *Salmonella pdu* genes in a range of Gram-negative bacteria demonstrate exogenous microcompartment expression across a variety of species. *Microb. Biotechnol* 2018, 11:199–210. [PubMed: 28967207]
45. Xia P-F, Zhang G-C, Walker B, Seo S-O, Kwak S, Liu J-J, Kim H, Ort DR, Wang S-G, Jin Y-S: Recycling Carbon Dioxide during Xylose Fermentation by Engineered *Saccharomyces cerevisiae*. *ACS Synth. Biol* 2017, 6:276–283. [PubMed: 27744692]
46. Ahrens W, Schlegel HG: Carbon dioxide requiring mutants of *Hydrogenomonas eutropha* strain H 16. I. Growth and CO₂-fixation. *Arch. Mikrobiol* 1972, 85:142–152. [PubMed: 4627399]
47. Gonzalez-Esquer CR, Shubitowski TB, Kerfeld CA: Streamlined Construction of the Cyanobacterial CO₂-Fixing Organelle via Protein Domain Fusions for Use in Plant Synthetic Biology. *Plant Cell* 2015, 27:2637–2644. [PubMed: 26320224]
48. Frey R, Mantri S, Rocca M, Hilvert D: Bottom-up Construction of a Primordial Carboxysome Mimic. *J. Am. Chem. Soc* 2016, 138:10072–10075. [PubMed: 27479274]
- 49**. Long BM, Hee WY, Sharwood RE, Rae BD, Kaines S, Lim Y-L, Nguyen ND, Massey B, Bala S, von Caemmerer S, et al.: Carboxysome encapsulation of the CO₂-fixing enzyme Rubisco in tobacco chloroplasts. *Nat. Commun* 2018, 9:3570. [PubMed: 30177711] Demonstrated α -carboxysome-like structures in plants using only three protein components.
50. Cai F, Bernstein SL, Wilson SC, Kerfeld CA: Production and Characterization of Synthetic Carboxysome Shells with Incorporated Luminal Proteins. *Plant Physiol* 2016, 170:1868–1877. [PubMed: 26792123]
51. Bar-On YM, Milo R: The global mass and average rate of rubisco. *Proc Natl Acad Sci USA* 2019, 116:4738–4743. [PubMed: 30782794]
52. McGrath JM, Long SP: Can the cyanobacterial carbon-concentrating mechanism increase photosynthesis in crop species? A theoretical analysis. *Plant Physiol* 2014, 164:2247–2261. [PubMed: 24550242]
53. Price GD, Pengelly JLL, Forster B, Du J, Whitney SM, von Caemmerer S, Badger MR, Howitt SM, Evans JR: The cyanobacterial CCM as a source of genes for improving photosynthetic CO₂ fixation in crop species. *J. Exp. Bot* 2013, 64:753–768. [PubMed: 23028015]
54. Hanson MR, Lin MT, Carmo-Silva AE, Parry MAJ: Towards engineering carboxysomes into C3 plants. *Plant J* 2016, 87:38–50. [PubMed: 26867858]
55. Lin MT, Occhialini A, Andralojc PJ, Devonshire J, Hines KM, Parry MAJ, Hanson MR: β -Carboxysomal proteins assemble into highly organized structures in *Nicotiana* chloroplasts. *Plant J* 2014, 79:1–12. [PubMed: 24810513]
56. Lin MT, Occhialini A, Andralojc PJ, Parry MAJ, Hanson MR: A faster Rubisco with potential to increase photosynthesis in crops. *Nature* 2014, 513:547–550. [PubMed: 25231869]
57. Occhialini A, Lin MT, Andralojc PJ, Hanson MR, Parry MAJ: Transgenic tobacco plants with improved cyanobacterial Rubisco expression but no extra assembly factors grow at near wild-type rates if provided with elevated CO₂. *Plant J* 2016, 85:148–160. [PubMed: 26662726]
58. Pengelly JLL, Förster B, von Caemmerer S, Badger MR, Price GD, Whitney SM: Transplastomic integration of a cyanobacterial bicarbonate transporter into tobacco chloroplasts. *J. Exp. Bot* 2014, 65:3071–3080. [PubMed: 24965541]

59. Uehara S, Adachi F, Ito-Inaba Y, Inaba T: Specific and efficient targeting of cyanobacterial bicarbonate transporters to the inner envelope membrane of chloroplasts in arabidopsis. *Front. Plant Sci* 2016, 7:16. [PubMed: 26870048]
60. Scott KM, Harmer TL, Gemmell BJ, Kramer AM, Sutter M, Kerfeld CA, Barber KS, Bari S, Boling JW, Campbell CP, et al.: Ubiquity and functional uniformity in CO₂ concentrating mechanisms in multiple phyla of Bacteria is suggested by a diversity and prevalence of genes encoding candidate dissolved inorganic carbon transporters. *FEMS Microbiol. Lett* 2020, doi:10.1093/femsle/fnaa106.
61. Kirst H, Kerfeld CA: Bacterial microcompartments: catalysis-enhancing metabolic modules for next generation metabolic and biomedical engineering. *BMC Biol* 2019, 17:79. [PubMed: 31601225]






Figure 1.

a) The CCM in a cyanobacterial cell. Bicarbonate transporters and facilitated CO_2 uptake proteins raise the intracellular HCO_3^- concentration while CO_2 flows freely across the plasma membrane. b) Carboxysome metabolism. HCO_3^- enters the carboxysome along its concentration gradient, where it is converted to CO_2 via a carbonic anhydrase. CO_2 and RuBP serve as substrates for Rubisco, which produces two molecules of 3-PG. O_2 may occasionally serve as a Rubisco substrate, though at a minimal level.

**Figure 2.**

a) Domain structure of CsoS2 from *H. neapolitanus* (uniprot ID: O85041; CSOS2_HALNC), with marked short (CsoS2A) and long (CsoS2B) forms. b) Domain structure of CcmM from *S. elongatus* PCC7942 (uniprot ID: Q03513; CCMM_SYNE7), with marked short (M35) and long (M58) forms. SSUL stands for “small subunit-like” domain. c) Structures of the CsoS2 NTD bound to Form 1A Rubisco (PDB: 6UEW) and CcmM SSUL bound to Form 1B Rubisco (PDB: 6HBC). Structures were rendered in ChimeraX. d) Hypothesized model of carboxysome nucleation. The carboxysome nucleating protein binds Rubisco with high avidity and affinity in the reducing cytosol. Maturation may involve oxidation (or exclusion of reducing agents) and disulfide-bond induced conformational changes.

**Figure 3.**

a) Engineering a carboxysomal CCM into industrial microbes could convert atmospheric CO_2 into high value products. b) Engineering a carboxysomal CCM into plants could increase plant CO_2 efficiency and promote growth and biomass yield. c) Faster Rubiscos could enable more efficient carbon fixation. d) A minimal gene set lowers the DNA payload when engineering the CCM into new host organisms. e) In vitro carboxysomes are a novel platform to study carboxysome assembly, and could act as in vitro catalytic reactors. f) Carboxysomes can be repurposed for alternative metabolisms. Enzymatic activity may depend on whether or not the carboxysome is an oxygen privileged environment, which remains unknown. g) Pore engineering, such as changing the charge or size of the pore, may aid development of alternative metabolisms.