

Lawrence Berkeley National Laboratory

Recent Work

Title

EXCHANGE DEGENERATE REGGE TRAJECTORIES AND MESON π NUCLEON CHARGE EXCHANGE SCATTERING

Permalink

<https://escholarship.org/uc/item/8q82s1pz>

Authors

Ahmadzadeh, Akbar
Kaufmann, William B.

Publication Date

1969-08-12

Submitted to Physical Review

UCRL-19315
Preprint

RECEIVED
LAWRENCE
RADIATION LABORATORY

OCT 16 1969

LIBRARY AND
DOCUMENTS SECTION

EXCHANGE DEGENERATE REGGE TRAJECTORIES AND
MESON NUCLEON CHARGE EXCHANGE SCATTERING

Akbar Ahmadzadeh and William B. Kaufmann

August 12, 1969

AEC Contract No. W-7405-eng-48

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

LAWRENCE RADIATION LABORATORY
UNIVERSITY of CALIFORNIA BERKELEY

UCRL-19315

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

EXCHANGE DEGENERATE REGGE TRAJECTORIES AND
MESON NUCLEON CHARGE EXCHANGE SCATTERING*

Akbar Ahmadzadeh and William B. Kaufmann

Lawrence Radiation Laboratory
University of California
Berkeley, California

and

Department of Physics
Arizona State University†
Tempe, Arizona 85281

August 12, 1969

ABSTRACT

Based on exchange degeneracy and $SU(3)$ symmetry, a Regge pole model fitting the differential cross sections for pion-nucleon and kaon-nucleon charge exchange and η production as well as polarization in $\pi^- p \rightarrow \pi^0 n$ is given. Besides the $\rho-A_2$ trajectory, a conspiring and exchange degenerate $\rho'-A_2$ trajectory with the same slope as $\rho-A_2$ but with intercept near zero is taken into account. The residues are assumed to have $1/\Gamma(\alpha)$ behavior. A four parameter fit gives good agreement with the experimental data.

*Research supported by the Air Force Office of Scientific Research, Office of Aerospace Research, United States Air Force, under Grant No. AF-AFOSR-1294-67. This document has been approved for public release and sale; its distribution is unlimited.

†Permanent address.

INTRODUCTION

A possible classification of Regge trajectories according to $SU(3)$ symmetry, exchange degeneracy and the Lorentz pole quantum number has been given previously.^{1,2} Our aim here is to combine these considerations with the Veneziano-type representation³ in a phenomenological study of pion-nucleon and kaon-nucleon charge exchange reactions and η production. The model presented here has numerous other phenomenological implications which are subject to future investigations.

The basic ideas of Refs. 1 and 2 utilized here are the following: The vector and tensor octets of trajectories form an octet of exchange degenerate trajectories with $SU(3)$ splittings of trajectories determined by the masses of resonances. These trajectories are coupled to mesons and baryons $SU(3)$ symmetrically. Similarly, the octet of pseudoscalar trajectories (to which the pion belongs) together with the octet of axial vector trajectories (to which the B meson belongs) form an exchange degenerate octet of conspiring trajectories. The co-conspirator of this octet is an exchange degenerate octet of vector-tensor trajectories. For example, the exchange degenerate π -B trajectory conspires with an exchange degenerate $\rho'-A'_2$ trajectory of the same intercept.

In addition, we shall assume that the trajectories have a universal slope. In the spirit of the Veneziano formula this simplifying assumption makes extra baryon trajectories in the s and u channels unnecessary. Furthermore, the possible resonances

belonging to the co-conspirators would be hidden underneath the other (known) resonances. This would explain why they have not been observed experimentally.

For the reactions we consider in this article only the ρ and A_2 quantum numbers can be exchanged in the t channel. Our model consists of exchange degenerate ρ - A_2 plus exchange degenerate and conspiring ρ' - A'_2 trajectories. In the next section the formalism and the results of fitting the parameters are given.

2. FORMULATION AND RESULTS

A phenomenological study of pion nucleon charge exchange differential cross section and polarization based on ρ and ρ' trajectories has been given in a separate article.⁴ Here we include the η production and kaon-nucleon charge exchange as well. Because we assume exchange degeneracy and SU(3) symmetry of the couplings, no new parameter is needed.

Let us define the helicity nonflip and helicity flip parts of the ρ trajectory contribution in meson nucleon scattering to be of the form

$$A'_{\rho} = \frac{\beta_{\rho}^n}{\Gamma(\alpha_1)} \xi_{\rho} (as)^{\alpha_1} \quad (1)$$

and

$$B_{\rho} = \frac{\beta_{\rho}^f}{\Gamma(\alpha_1)} \xi_{\rho} (as)^{\alpha_1-1} \quad (2)$$

respectively; where $\xi_{\rho} = (1 - e^{-i\pi\alpha_1}) / \sin\pi\alpha_1$ and $\beta^n(\beta^f)$ is the helicity nonflip (flip) residue constant. Eqs. (1) and (2) can be considered as the leading term (in energy) of a suitable Veneziano-type representation. In fact, Igi⁵ has given such a representation for pion nucleon scattering and his formula reduce to Eqs. (1) and (2) in the high energy limit. Furthermore, even if one has to take an infinite number of Veneziano terms (corresponding, perhaps, to a single Lorentz pole), still the leading term in energy will be in the form of Eqs. (1) and (2). The exchange degenerate ρ - A_2 trajectory passing through $\rho(760)$ and $A_2(1310)$ is given by

$$\alpha_1 \approx 0.5 + at; a \approx 0.9 (\text{GeV})^{-2} \quad (3)$$

Similarly, the amplitudes for the A_2 trajectory contribution (denoted by the subscript R) are given by

$$A'_R = \frac{\beta_R^n}{\Gamma(\alpha_1)} \xi_R (as)^{\alpha_1} \quad (4)$$

$$B_R = \frac{\beta_R^f}{\Gamma(\alpha_1)} \xi_R (as)^{\alpha_1-1} \quad (5)$$

where $\xi_R = (1 + e^{-i\pi\alpha_1})/\sin \pi\alpha_1$

The exchange degenerate ρ - A_2 trajectory is assumed to have the same slope as the ρ' - A'_2 trajectory. This simplifying assumption is made in order to avoid introducing new baryon trajectories in the s and u channels. Namely, one can imagine a Veneziano-type formula in which the same baryon trajectories coexist with ρ - A_2 as with ρ' - A'_2 . Furthermore, as pointed out earlier⁴, this model would predict existence of a particle with the same mass as the B(1220) meson and with the ρ quantum numbers. This is because in our scheme ρ' - A'_2 trajectory coincides with the π -B trajectory. This coincidence would be a realization of chiral symmetry. The same coincidence should occur for higher recurrences. On the other hand, we assume that the A'_2 trajectory chooses nonsense⁶ at $\alpha = 0$.

Therefore we do not expect a scalar particle at the pion mass. This scheme is consistent with the quark model; namely, if as in Ref. 2 we assume that all trajectories are coupled to a quark-antiquark system, then the point $\alpha = 0$ is a nonsense point for the A'_2 trajectory. Note that the A'_2 trajectory couples to the spin triplet of $q\bar{q}$ system. The $\rho' - A'_2$ trajectory, based on the masses of the pion and the B meson is given by

$$\alpha_2 \approx -0.02 + 0.9 t \quad (6)$$

The ρ' contribution is given by

$$A'_{\rho'} = \frac{t \beta_{\rho'}^n}{\Gamma(\alpha_2)} \xi_{\rho'} (as)^{\alpha_2} \quad (7)$$

and

$$B_{\rho'} = \frac{\beta_{\rho'}^f}{\alpha_2 \Gamma(\alpha_2)} \xi_{\rho'} (as)^{\alpha_2 - 1} \quad (8)$$

Similarly, for the A'_2 trajectory we have

$$A'_{R'} = \frac{t \beta_{R'}^n}{\Gamma(\alpha_2)} \xi_{R'} (as)^{\alpha_2} \quad (9)$$

$$B_{R'} = \frac{\beta_{R'}^f}{\alpha_2 \Gamma(\alpha_2)} \xi_{R'} (as)^{\alpha_2 - 1} \quad (10)$$

The factor of t in Eqs. (7) and (9) is included because ρ' and A'_2 are assumed to be conspiring trajectories (see Ref. 7, for example).

Note the factor α in the denominators of the right hand sides of Eqs. (8) and (10). Due to this factor there is a pole in the flip amplitude of the A_2' contribution at $\alpha_2 = 0$. This is a nonsense pole and thus we are assuming the existence of a compensating trajectory⁶. In Ref. 4, this factor of α was not included in the flip amplitude of the ρ' . It turns out that this extra factor improves our fit for polarization in $\pi^- p \rightarrow \pi^0 n$.

Now using the exchange degeneracy of the residues we have

$$\beta_1^n \equiv \beta_{\rho}^n = \beta_{R'}^n; \quad \beta_1^f \equiv \beta_{\rho'}^f = \beta_{R'}^f \quad (11)$$

and similarly

$$\beta_2^n \equiv \beta_{\rho'}^n = \beta_{R'}^n; \quad \beta_2^f \equiv \beta_{\rho}^f = \beta_{R'}^f \quad (12)$$

Note that, since the trajectories are fully determined by Eqs. (3) and (6), we are left with only four parameters β_1^n , β_1^f , β_2^n and β_2^f .

Defining the new amplitudes $\mathcal{A}$ and $\mathcal{B}$ as

$$\mathcal{A}_{\rho} = A_{\rho}' + A_{\rho'}'; \quad \mathcal{A}_R = A_{R'}' + A_{R'}' \quad (13)$$

$$\mathcal{B}_{\rho} = B_{\rho} + B_{\rho'}; \quad \mathcal{B}_R = B_R + B_{R'}$$

from SU(3) symmetry of the couplings we have⁸

$$\begin{aligned} A'(\pi^- p \rightarrow \pi^0 n) &= \sqrt{2} \mathcal{A}_{\rho} \\ A'(K^- p \rightarrow \bar{K}^0 n) &= \mathcal{A}_{\rho} - \mathcal{A}_R \\ A'(\pi^- p \rightarrow \eta n) &= -(2/3)^{1/2} \mathcal{A}_R \\ A'(K^+ n \rightarrow K^0 p) &= -\mathcal{A}_{\rho} - \mathcal{A}_R \end{aligned} \quad (14)$$

with similar expressions for the B amplitudes. In what follows we

shall consider the first three reactions in Eqs. (14). We are not including the K^+n charge exchange case because of the meager high energy experimental data and because of the complications due to the Glauber corrections for deuteron target. Note however that with the four parameter determined below our model makes unambiguous prediction of the K^+n charge exchange reaction. The differentiated cross section is given by⁹

$$\frac{d\sigma}{dt} = \frac{1}{64 \pi q^2 s} \left\{ (4m^2 - t) |A'|^2 + \frac{t}{4m^2 - t} [4\mu^2 m^2 - ts - (s - m^2 - \mu^2)^2] |B|^2 \right\} \quad (15)$$

where q is the center of mass momentum, m is the nucleon mass, μ is the meson mass and s and t are the usual invariant energy and momentum transfer. Note that in η production we should use unequal mass kinematics. However, at the energies we are considering this extra complication is not necessary. The polarization is given⁹ by

$$P = \frac{-\sin \theta \operatorname{Im}(A' B^*)}{16 \pi \sqrt{s} \frac{d\sigma}{dt}} \quad (16)$$

where θ is the scattering angle in the center of mass system.

The experimental data¹⁰⁻¹³ are fitted using the four free parameters. The minimum Chi square is obtained for

$$\beta_1^n = 9.81 \pm 1, \beta_1^f = 119. \pm 9, \beta_2^n = -36 \pm 15, \beta_2^f = 38 \pm 15 \quad (17)$$

For a total of 223 data points we have obtained $\chi^2 = 369$ which we consider to be rather reasonable especially in view of ignoring the other (non-leading) terms. One could of course break SU(3) and/or exchange degeneracy and obtain a smaller χ^2 . But as long as we do not know how to take all other possible terms into account, there is very little information to be gained.

The errors given in Eqs. (17) indicate the change in each parameter necessary to increase χ^2 by 10 percent. The large uncertainty in β_2^n and β_2^f reflects the lack of extensive polarization data, to which these parameters are sensitive. We are looking forward to more polarization data in these reactions especially at higher momentum transfers to determine these parameters with better confidence. One feature of our model worth mentioning is that it predicts zero polarization in π^-p charge-exchange at $t \approx -0.6(\text{GeV}/c)^2$ where $\alpha_p = 0$, and also predicts a zero in the $\pi^-p \rightarrow \eta n$ polarization at $t \approx -1.6(\text{GeV}/c)^2$ where $\alpha_R = -1$.

Note also that our formalism can be used to calculate the differential cross section and polarization in $\pi^+p \rightarrow K^+\Sigma^+$ reaction. In this case only the d/f ratios remain as free parameters; for the $K^* - K^{**}$ and $K^{*'} - K^{**'}$ trajectories are determined from the masses in the same way as our $\rho - A_2$ and $\rho' - A_2'$ trajectories. The residues in this case are of course related to those of Eqs. (17).

Figures 1. through 8. show the experimental data as well as our theoretical curves. The error bars shown on these curves are statistical only. In calculating the χ^2 , systematic errors in the

data are also taken into account when given in the experimental articles. In the η production data we have assumed a branching ratio¹⁴ of 0.381.

Based on our model and the parameters obtained here predictions are made for future data up to 100 GeV/c. Figures 9. through 11. show the results of our prediction.

ACKNOWLEDGEMENT

We would like to thank Professor Geoffrey F. Chew for his hospitality at the Lawrence Radiation Laboratory, where this work was done.


REFERENCES

1. Akbar Ahmadzadeh, Phys. Rev. Letters 20, 1125 (1968).
2. Akbar Ahmadzadeh and Richard J. Jacob, Phys. Rev. 176, 1719 (1968).
3. G. Veneziano, Nuovo Cimento 57A, 190 (1968).
4. Akbar Ahmadzadeh and Jane C. Jackson, Arizona State University preprint (1969).
5. K. Igi, Physics Letters 28B, 330 (1968).
6. M. Gell-Mann, M. Goldberger, F. Low, E. Marx and F. Zachariasen, Phys. Rev. 133, B145 (1964).
7. L. Sertorio and M. Toller, Phys. Rev. Letters 19, 1146 (1967).
8. Akbar Ahmadzadeh and C. H. Chan, Physics Letters 22, 692 (1966).
9. V. Singh, Phys. Rev. 129, 1889 (1963).
10. P. Sonderegger, J. Kirz, O. Guisan, P. Falk-Vairant, C. Bruneton, P. Borgeaud, A. W. Stirling, C. Caverzasio, J. P. Guillard, M. Yvert and B. Amblard, Physics Letters 20, 75 (1966); M. A. Wahlig and I. Mannelli, Phys. Rev. 168, 1515 (1968).
11. P. Bonamy, P. Borgeaud, C. Bruneton, P. Falk-Vairant, O. Guisan, P. Sonderegger, C. Caverzasio, J+ P+ Guillard, J. Schneider, M. Yvert, I. Mannelli, F. Sergiampietri and L. Vincelli, Physics Letters 23, 501 (1966); J. Schneider, private communication.

12. P. Astbury, G. Brautti, G. Finocchiaro, A. Michelini,
K. Terwilliger, D. Websdale, C. H. West, P. Zanella,
W. Beusch, W. Fischer, B. Gobbi, M. Pepin and E. Polgar,
Physics Letters 23, 396 (1966).
13. O. Guisan, J. Kirz, P. Sonderegger, A. V. Stirling, P. Gorgeaud,
C. Bruneton, P. Falk-Vairant, B. Amblard, C. Caversasio,
J+ P+ Guillaud and M. Yvert, Physics Letters 18, 200 (1965);
M+ A+ Wahlig and I. Mannelli, Phys. Rev. 168, 1515 (1968).
14. Review of Particle Properties, N. Barash-Schmidt et al,
Reviews of Modern Physics, January 1969.


FIGURE CAPTIONS

- Fig. 1. $\frac{d\sigma}{dt}$ for $\pi^- p \rightarrow \pi^0 n$. Upper curve: $100 \times \frac{d\sigma}{dt}$ at 4.83 GeV/c laboratory momentum; middle curve: $10 \times \frac{d\sigma}{dt}$ at 5.85 GeV/c; lower curve $\frac{d\sigma}{dt}$ at 6.0 GeV/c.
- Fig. 2. $\frac{d\sigma}{dt}$ for $\pi^- p \rightarrow \pi^0 n$: Upper curve: $100 \times \frac{d\sigma}{dt}$ at 10 GeV/c; middle curve $10 \times \frac{d\sigma}{dt}$ at 13.3 GeV/c; lower curve $\frac{d\sigma}{dt}$ at 18.2 GeV/c.
- Fig. 3. Polarization in $\pi^- p \rightarrow \pi^0 n$ at 5.9 GeV/c.
- Fig. 4. Polarization in $\pi^- p \rightarrow \pi^0 n$ at 11.2 GeV/c.
- Fig. 5. $\frac{d\sigma}{dt}$ for $K^- p \rightarrow \bar{K}^0 n$: Upper curve $10 \times \frac{d\sigma}{dt}$ at 5 GeV/c; Lower curve $\frac{d\sigma}{dt}$ at 7.1 GeV/c.
- Fig. 6. $\frac{d\sigma}{dt}$ for $K^- p \rightarrow \bar{K}^0 n$: Upper curve $10 \times \frac{d\sigma}{dt}$ at 9.5 GeV/c; Lower curve $\frac{d\sigma}{dt}$ at 12.3 GeV/c.
- Fig. 7. $\frac{d\sigma}{dt}$ for $\pi^- p \rightarrow \eta n$. Upper curve $100 \times \frac{d\sigma}{dt}$ at 5.9 GeV/c; middle curve $10 \times \frac{d\sigma}{dt}$ at 10. GeV/c; lower curve $\frac{d\sigma}{dt}$ at 9.8 GeV/c.
- Fig. 8. $\frac{d\sigma}{dt}$ for $\pi^- p \rightarrow \eta n$: Upper curve $10 \times \frac{d\sigma}{dt}$ at 13.3 GeV/c; lower curve $\frac{d\sigma}{dt}$ at 18.2 GeV/c.
- Fig. 9. Predicted $\frac{d\sigma}{dt}$ for $\pi^- p \rightarrow \pi^0 n$. Laboratory momentum from top to bottom is 25, 50, 75, 100 GeV/c respectively.
- Fig. 10. Predicted $\frac{d\sigma}{dt}$ for $K^- p \rightarrow \bar{K}^0 n$. At same momenta as Fig. 8.
- Fig. 11. Predicted $\frac{d\sigma}{dt}$ for $\pi^- p \rightarrow \eta n$. At same momenta as Fig. 8.




XBL698-3624

Fig. 1




XBL698-3625

Fig. 2




XBL698-3626

Fig. 3




XBL698-3627

Fig. 4




XBL698-3628

Fig. 5




XBL698-3629

Fig. 6




XBL698 - 3630

Fig. 7




XBL698-3631

Fig. 8




XBL698-3632

Fig. 9



XBL698-3633

Fig. 10



XBL698-3634

Fig. 11

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or*
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.*

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

TECHNICAL INFORMATION DIVISION
LAWRENCE RADIATION LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720