

UC Irvine

UC Irvine Previously Published Works

Title

Molecular Mechanisms Associated with Nicotine Pharmacology and Dependence

Permalink

<https://escholarship.org/uc/item/8rw7h1ss>

Authors

Fowler, Christie D
Turner, Jill R
Imad Damaj, M

Publication Date

2019

DOI

10.1007/164_2019_252

Peer reviewed



Molecular Mechanisms Associated with Nicotine Pharmacology and Dependence

Christie D. Fowler, Jill R. Turner, and M. Imad Damaj

Contents

- 1 Introduction
 - 2 Basic Neurocircuitry of Nicotine Addiction
 - 3 Role of Nicotinic Receptors in Nicotine Dependence and Brain Function
 - 4 Modulatory Factors That Influence nAChR Expression and Signaling
 - 5 Genomics and Genetics of Nicotine Dependence
 - 5.1 Overview
 - 5.2 Human and Animal Genetic Studies
 - 5.3 Transcriptionally Adaptive Changes
 - 6 Other Constituents in Nicotine and Tobacco Products Mediating Dependence
 - 7 Therapeutic Approaches for Tobacco and Nicotine Dependence
 - 7.1 Nicotine Replacement Therapies
 - 7.2 Varenicline and Bupropion
 - 7.3 Novel Approaches
 - 8 Conclusion
- References

Abstract

Tobacco dependence is a leading cause of preventable disease and death worldwide. Nicotine, the main psychoactive component in tobacco cigarettes, has also

C. D. Fowler

Department of Neurobiology and Behavior, University of California Irvine, Irvine, CA, USA

J. R. Turner

Department of Pharmaceutical Sciences, College of Pharmacy, University of Kentucky, Lexington, KY, USA

M. Imad Damaj (✉)

Department of Pharmacology and Toxicology, Virginia Commonwealth University, Richmond, VA, USA

Translational Research Initiative for Pain and Neuropathy at VCU, Richmond, VA, USA

e-mail: m.damaj@vcuhealth.org

© Springer Nature Switzerland AG 2019

Handbook of Experimental Pharmacology, https://doi.org/10.1007/164_2019_252

been garnering increased popularity in its vaporized form, as derived from e-cigarette devices. Thus, an understanding of the molecular mechanisms underlying nicotine pharmacology and dependence is required to ascertain novel approaches to treat drug dependence. In this chapter, we review the field's current understanding of nicotine's actions in the brain, the neurocircuitry underlying drug dependence, factors that modulate the function of nicotinic acetylcholine receptors, and the role of specific genes in mitigating the vulnerability to develop nicotine dependence. In addition to nicotine's direct actions in the brain, other constituents in nicotine and tobacco products have also been found to alter drug use, and thus, evidence is provided to highlight this issue. Finally, currently available pharmacotherapeutic strategies are discussed, along with an outlook for future therapeutic directions to achieve to the goal of long-term nicotine cessation.

Keywords

Neurobiology nicotine dependence · Nicotine · Nicotinic receptors · Smoking cessation

1 Introduction

Cigarette smoking is the principal cause of premature death and disability in the United States. In 2014, about 480,000 deaths in the United States were caused by cigarette smoking. Globally, smoking-related illnesses result in over four million deaths annually. However, despite enormous educational efforts about the health hazards of smoking and other tobacco control efforts, many smokers continue to encounter extreme difficulty quitting and staying tobacco-free in the long-term. The 2017 CDC report estimated that 15.1% of the US population was “current smokers,” (11.2% (75%) of them are daily smokers).

Addiction to tobacco smoking depends not only on the positive reinforcing and hedonic actions of nicotine but also on escape from the aversive consequences of nicotine withdrawal. Many studies suggest that avoidance of the negative emotional state produced by nicotine withdrawal represents a motivational component that promotes continued tobacco use and relapse after smoking cessation. The difficulty in overcoming nicotine dependence is illustrated by the poor success rates among smokers who try to quit. While the majority of smokers (~70%) report an interest in quitting, and around 55% have attempted to quit in the previous year, ~7% of smokers are abstinent at 1 month after their quit date, and fewer than 2% are abstinent 1 year after quitting when they do not receive assistance in smoking cessation (CDC 2015).

While several smoking cessation therapies are available, the success rate of these therapies after 1 year remains only about 20–25% (Gonzales et al. 2006). Therefore, understanding the various mechanisms and factors involved in the different aspects of nicotine dependence is crucial to develop successful prevention and intervention approaches, including newer and more effective pharmacotherapies.

2 Basic Neurocircuitry of Nicotine Addiction

Tobacco smoke contains about 9,000 chemicals, among which about 70 are known carcinogens. However, nicotine is the major psychoactive ingredient in tobacco smoke and the component most associated with tobacco dependence. The development and persistence of dependence on tobacco is due to the actions of nicotine, acting at neuronal nicotinic acetylcholine receptors (nAChRs). nAChRs belong to the Cys-loop receptor family, which are ligand-gated ion channels that form pentamers arranged around a water-filled pore and allow for the influx of both Na⁺ and Ca²⁺ (Changeux et al. 1998). The subunits of mammalian neuronal nAChRs range from $\alpha 2$ – $\alpha 7$, $\alpha 9$, $\alpha 10$, to $\beta 2$ – $\beta 4$, which form multiple combinations of homomeric and heteromeric receptor subtypes having varying function (Changeux et al. 1998). These receptors have three broad conformational states: resting closed states, open states, and desensitized states (Changeux et al. 1998). The typical resting closed state is induced when the orthosteric site (traditional ligand binding site) is unoccupied and the cation channel is closed. Upon binding of an orthosteric agonist, the cation channel is opened, allowing for cation influx into the cell. Following the open state, the receptor is then desensitized; despite agonist binding, the cation channel is closed, rendering the receptor inactive (Changeux et al. 1998). Due to their predominant presynaptic location, nAChRs in the CNS primarily function via modulation of neurotransmitter release (Mansvelder and McGehee 2000). This modulation, in turn, results in long-term synaptic plasticity, which is a prominent neuronal signature of exposure to nicotine (Ji et al. 2001). The most abundant nAChRs found in the mammalian brain are the low-affinity homomeric $\alpha 7$ and the high-affinity heteromeric $\alpha 4\beta 2$ containing ($\alpha 4\beta 2^*$), which have diverse characteristics (Hill et al. 1993). The $\alpha 7$ nAChR has high calcium permeability, low probability of opening, and rapid desensitization (in milliseconds) (Williams et al. 2011). In contrast, the $\alpha 4\beta 2^*$ nAChR has a high probability of opening and desensitizes at a slower rate (in seconds) (Li and Steinbach 2010). These differing characteristics, however, do not necessarily drive divergent effects on neuronal plasticity. For example, previous studies have shown that both $\alpha 4\beta 2^*$ and $\alpha 7$ nAChR activation can either elicit (Lagostena et al. 2008; Tang and Dani 2009; Welsby et al. 2009) or prevent (Alkondon and Albuquerque 2001; Alkondon et al. 1997; Ji et al. 2001) long-term potentiation (LTP) in the hippocampus, with these variable effects attributed to activation of differing subtypes on specific interneuron populations. Further, accessory nAChR subunits, such as $\alpha 5$ and $\beta 3$, can integrate into the $\alpha 4\beta 2$, $\alpha 3\beta 4$, or $\alpha 3\beta 2$ nAChR subtypes to alter receptor function. For instance, insertion of the $\alpha 5$ subunit into the $\alpha 4\beta 2$ or $\alpha 3\beta 2$ nAChR subtypes results in increased ligand-mediated receptor activation, rate of desensitization, and conductance (Gerzanich et al. 1998; Ramirez-Latorre et al. 1996).

Nicotine initiates its rewarding effects by activating nAChRs in the natural reward system of the brain, the mesolimbic pathway. This pathway is comprised of dopaminergic neurons originating in the ventral tegmental area (VTA) that project to regions such as the nucleus accumbens (NAc), prefrontal cortex (PFC), amygdala, and hippocampus (De Biasi and Dani 2011; Lisman and Grace 2005).

Dopamine release, especially in the NAc, is associated with the rewarding and reinforcing effects of all drugs of abuse. nAChRs are localized throughout the mesolimbic circuitry and when activated, increase dopaminergic firing and release (De Biasi and Dani 2011; Di Chiara 2000). Further, infusion of nAChR antagonists directly into the VTA attenuates nicotine self-administration (Corrigall et al. 1994). This pathway has a complex circuitry that also involves other neurotransmitters; for instance, glutamatergic, GABAergic, and cholinergic inputs converge on dopamine neurons to modulate dopamine release (Dani and Bertrand 2007). Cholinergic neurons in the laterodorsal tegmentum and the pedunculo pontine tegmentum initiate excitation of dopamine neurons in VTA that project to the NAc (Maskos 2010; Omelchenko and Sesack 2005), and these cells in the pedunculo pontine tegmentum have been shown to regulate nicotine self-administration (Lanca et al. 2000). In opposition to reward-related signaling, dense nAChR expression is also found in the projection from the medial habenula (MHb) to the interpeduncular nucleus (IPN), a circuit involved in aversive processing and nicotine withdrawal (Fowler et al. 2011; Salas et al. 2009). The major neurotransmitters of this pathway are acetylcholine, glutamate, and substance P, and it is thought that presynaptic nAChRs on MHb axons facilitate glutamate release from cholinergic and glutamatergic coexpressing axons in the IPN to mediate the aversive signal to high doses of nicotine (Fowler et al. 2011; Girod and Role 2001), which serves to limit drug intake.

3 Role of Nicotinic Receptors in Nicotine Dependence and Brain Function

The utilization of genetically mutant mice, pharmacological interventions, and viral reexpression approaches have implicated particular brain areas and specific nAChR subtypes in nicotine dependence. For instance, in a $\beta 2$ knockout mouse model, the $\beta 2^*$ nAChRs have been shown to be required for nicotine reward and reinforcement, as revealed in nicotine conditioned place preference (CPP) and intravenous self-administration studies (Orejarena et al. 2012; Picciotto et al. 1998; Walters et al. 2006). The $\beta 2$ subunit co-assembles with the $\alpha 6$ and $\alpha 4$ subunits to form several $\alpha 6\beta 2^*$, $\alpha 4\beta 2^*$, and $\alpha 4\alpha 6\beta 2^*$ nAChR subtypes, which are notably expressed in the VTA-NAc circuit (Champtiaux et al. 2003; Klink et al. 2001; Salminen et al. 2004). These findings are consistent with the fact that stimulation of $\alpha 4\beta 2^*$ high-affinity nAChRs located on the dopaminergic cells in the VTA shifts firing from tonic to phasic modes, resulting in increased DA release in both the NAc and the PFC (Dani et al. 2011). Nicotine CPP revealed a critical role of the $\alpha 4$, $\alpha 6$, and $\beta 2$ subunits in the NAc via genetic mutant mice and site-specific infusions (Sanjakdar et al. 2015). In addition, genetic ablation of the $\beta 2$, $\alpha 6$, and $\alpha 4$ nAChR subunits attenuated nicotine self-administration in mice, an effect which could be rescued by reexpression of these subunits in the VTA via a lentiviral vector (Picciotto et al. 1998; Pons et al. 2008). Furthermore, $\alpha 4$ “knock-in” mice (Leu9’ Ala mutation renders animals hypersensitive to nicotine) demonstrated a preference for nicotine at a dose 50-fold

lower than the typical nicotine dose that induces a preference in wild-type (WT) mice in the CPP test (Tapper et al. 2004).

Reward systems in the brain undergo neuroadaptations after chronic exposure to nicotine in tobacco products, which likely underlie nicotine dependence. Cessation from cigarette smoking induces a withdrawal syndrome comprised of physical, affective, and cognitive symptoms. The severity of these symptoms is a risk factor for relapse (Le Foll and Goldberg 2005; Markou and Kenny 2002), and nAChRs are important mediators of nicotine withdrawal symptoms. The nonselective nAChR antagonist mecamylamine is known to precipitate nicotine withdrawal signs in nicotine-dependent rodents (Damaj et al. 2003). Pharmacological interventions and mouse knockout studies have revealed that nAChR subunits modulate different aspects of the nicotine withdrawal syndrome. For example, some affective signs of withdrawal such as aversion-, anxiety-, and anhedonia-like measures are mediated by the $\beta 2$, $\alpha 6$, $\beta 4$, and $\alpha 7$ nAChR subunits (Jackson et al. 2008, 2009). The physical signs of the nicotine withdrawal syndrome are mediated by $\alpha 3$, $\alpha 5$, $\alpha 2$, and $\beta 4$ (Jackson et al. 2008, 2013; Salas et al. 2009), and a subset are mediated by $\alpha 7$ subunits (Stoker et al. 2012). One interesting feature of chronic nicotine exposure is the upregulation of nAChRs, most notably $\alpha 4\beta 2^*$ (Flores et al. 1992). This phenomenon has been observed both *in vitro* and *in vivo* and in human imaging studies (Kassiou et al. 2001; Marks et al. 1983; Perry et al. 1999). Interestingly, rodent and human studies suggest a positive correlation of nicotine withdrawal signs with upregulation of $\alpha 4\beta 2^*$ nAChRs (Cosgrove et al. 2010; Turner et al. 2011). Furthermore, the MHB-IPN pathway has been selectively implicated in withdrawal-induced somatic signs with $\alpha 5^*$ and $\beta 4^*$ nAChRs (Salas et al. 2009). In addition, infusion of the $\alpha 6^*$ nAChR-selective antagonist α -conotoxin MII in the MHB attenuated anxiety-like behavior in nicotine-withdrawn mice (Pang et al. 2016). Aberrant synaptic and circuitry function is also thought to underlie abnormal behavioral phenotypes, including nicotine withdrawal phenotypes like cognitive impairments and affective dysfunction (Ashare et al. 2014; Turner et al. 2013). For example, the hippocampus and the orbitofrontal cortex (OFC) are two well-described circuits impinging upon these nicotine withdrawal symptoms (Schoenbaum et al. 2016; Turner et al. 2011; Zhou et al. 2018), including impulsivity, altered affect, and cognition in humans. Supporting data in human (Dani and Harris 2005) and animal (Jackson et al. 2008) models link hippocampal function with nicotine withdrawal-induced symptoms, which are reliable determinants for smoking cessation outcomes. Functional imaging studies in smokers have shown that activation of the hippocampus can be correlated with both cognitive and affective withdrawal symptoms (Froeliger et al. 2010; McClernon and Gilbert 2004). Additionally, human studies report a correlation between hippocampal volume and successful quit attempts (Froeliger et al. 2010). This link may be due to nAChRs present at both excitatory and inhibitory terminals (Alkondon and Albuquerque 2001; Jones and Yakel 1997; Wada et al. 1989), well-positioning nicotinic signaling to influence the balance of excitatory and inhibitory transmission within the hippocampus (John and Berg 2015). The OFC regulates impulsivity, affective value of reinforcers, and emotion-attention interactions (Schoenbaum et al. 2016). Previous studies reported that

nicotine self-administration in rodents alters synaptic morphology in the OFC (Vazquez-Sanroman et al. 2016), while tobacco smokers display both morphological and functional connectivity changes within this region (Claus et al. 2013; Li et al. 2015). For example, smoking has been consistently shown to reduce the thickness of gray matter volume in the OFC (Kuhn et al. 2010; Li et al. 2015), and acute nicotine increases blood oxygen level-dependent fMRI signal in the striato-thalamo-orbitofrontal circuit (Ashare et al. 2014). However, the neuronal mechanisms underlying these effects are not easily examined, given that nicotine modulates the release of a number of neurotransmitters, including glutamate, GABA, and dopamine, and can lead to both facilitation and suppression of neuronal firing. For example, electrophysiological experiments have shown that nicotine impacts long-term potentiation (LTP) generation in the orbitofrontal cortex (Couey et al. 2007; Zhou et al. 2018). Classical LTP is based on the observation that a neuron's excitability to a particular synaptic input is increased following high-frequency stimulation, representing the molecular basis for Hebb's postulate, which states that when two connected cells fire simultaneously, the connection between them is strengthened. Previous studies examining nicotine's effect on this phenomenon have reported enhancement of LTP in a number of brain regions, such as the hippocampus (Nakauchi and Sumikawa 2012), amygdala (Huang et al. 2008), and VTA (Mansvelder and McGehee 2000). However, these effects diverge in the OFC. Zhou and colleagues (Zhou et al. 2018) demonstrated that acute nicotine application to the OFC during LTP induction resulted in nicotine-mediated conversion of LTP to LTD, a form of "metaplasticity," due to enhanced GABAergic transmission. These effects were in agreement with studies in nearby frontal cortical regions, where nicotine was observed to raise the threshold for LTP induction via enhancing GABAergic transmission (Couey et al. 2007). As appreciation grows for the importance of frontocortical excitatory/inhibitory balance in nicotine dependence (Pittaras et al. 2016), understanding nicotine's effects in this region may not only lead to better understanding of circuit-level mechanisms of nicotine dependence but also to potential therapeutic interventions.

4 Modulatory Factors That Influence nAChR Expression and Signaling

Several mechanisms that regulate nAChR expression, assembly, and trafficking were reported in the last two decades. Recent studies have shown that nicotine can act as a "chaperone" which expedites the transport of nAChR subunits, including $\alpha 4$ and $\beta 2$ nAChRs, to the endoplasmic reticulum and facilitates the passage and insertion of assembled nAChRs to the plasma membrane (Henderson et al. 2014; Srinivasan et al. 2011). In this context, this pharmacological chaperone mechanism may represent an important molecular mechanism of the first step in neuroadaptation to chronic nicotine and possibility of the emergence of neuronal adaptations underlying nicotine dependence. Another class of nAChR signaling modulators is represented by the Ly-6/neurotoxin gene superfamily of proteins that exhibit cellular

specific expression patterns in the brain and include Lynx1, Lynx2, and Lypd6. These proteins are negative modulators of nAChR signaling and feature a three-looped fold, a structural characteristic shared with the snake venom toxin α -bungarotoxin. Thus, as endogenous prototoxins, these proteins can bind directly to the extracellular face of nAChRs (Arvaniti et al. 2016; Miwa et al. 1999). The presence of Lynx1 and Lynx2 increases the desensitization rate and decreases ligand binding efficiency for multiple nAChR subtypes (George et al. 2017; Ibanez-Tallon et al. 2002; Lyukmanova et al. 2011; Tekinay et al. 2009). In cortex, Lynx1 is expressed in both glutamatergic and γ -aminobutyric acid-ergic (GABAergic) neurons, whereas Lynx2 has been mainly localized in glutamatergic neurons (Demars and Morishita 2014). Results suggest that lynx proteins can modulate nAChR function in the brain with important consequences for cholinergic-dependent synaptic plasticity (reviewed in Miwa et al. 2011; Miwa and Walz 2012; Thomsen and Mikkelsen 2012). Recently, Nissen and colleagues reported that the antinociceptive effect of nicotine and epibatidine in acute thermal pain tests is enhanced in Lynx1 knockout mice (Nissen et al. 2018). Further, computer simulations predict preferential binding affinity of Lynx1 to the α : α interface that exists in the stoichiometry of the low sensitivity (α 4) β 2 nAChRs.

5 Genomics and Genetics of Nicotine Dependence

5.1 Overview

Nicotine addiction is a complex disorder with multiple factors contributing to its dependence. Though a large host of factors contribute to nicotine dependence, reward, withdrawal effects, and relapse, twin studies have shown that genetics play a pivotal role (Li et al. 2003; Sullivan and Kendler 1999). Approximately 70% of the variability in nicotine dependence and smoking persistence has been attributed to genetic influences (Broms et al. 2006; Carmelli et al. 1992; Kendler et al. 2000; Li et al. 2003). Furthermore, twin studies have shown that ~50% of the individual differences that contribute to smoking relapse can be attributed to heritability (Xian et al. 2003). Ongoing studies examining not only genetics, but genomics and epigenetics, are increasing our understanding of how individual differences drive vulnerability or resilience to nicotine dependence.

5.2 Human and Animal Genetic Studies

In recent years, genome-wide association studies in humans revealed that a variant in the CHR5A3/B4 gene cluster (encodes α 3, α 5, β 4 nAChR subunits), located in chromosome region 15q25, serves as a risk factor for lung cancer and nicotine dependence (Berrettini et al. 2008; Liu et al. 2010; Saccone et al. 2009). More specifically, a single nucleotide polymorphism (SNP) in the CHR5A5 gene (rs16969968) (D398N), which encodes the α 5 nAChR subunit, has been repeatedly

linked to increased risk for tobacco dependence (Bierut et al. 2008; Kuryatov et al. 2011). The mechanisms behind this increased risk have been investigated in *in vitro* and *in vivo* functional studies. The $\alpha 5$ SNP was shown to reduce the function of the $\alpha 3\beta 4$ and $\alpha 4\beta 2$ nAChR subtypes that incorporate the mutant subunit (Bierut et al. 2008), a loss of function that subsequently was shown to influence addiction-like behaviors *in vivo*. Initial studies were conducted in $\alpha 5$ nAChR subunit gene knockout mice (Fowler et al. 2011). The $\alpha 5$ knockout mice were found to exhibit far greater motivation to consume large quantities of nicotine, and reexpression of $\alpha 5$ subunits within this pathway attenuated nicotine intake to wild-type levels (Fowler et al. 2011). Further, decreased expression of $\alpha 5$ subunits in rats similarly increased nicotine intake while decreasing the inhibitory effects of higher nicotine doses on brain reward circuitries (Fowler et al. 2011, 2013). Similar observations occurred in the nicotine CPP paradigm where $\alpha 5$ knockout mice exhibited a maintained nicotine preference at higher doses not maintained by $\alpha 5$ wild-type mice (Jackson et al. 2010). In addition, in mice expressing the $\alpha 5$ human mutation, an increase in nicotine self-administration was reported (Wilking and Stitzel 2015). Furthermore, using rats carrying the $\alpha 5$ human mutation, Forget et al. (2018) found greater nicotine intake in the SNP-expressing mutant rats compared with wild-type rats, as well as an increase in nicotine motivation mutant rats. In addition, the SNP-expressing rats exhibited a higher reinstatement of nicotine-seeking lever-pressing responses than the wild-type rats (Forget et al. 2018). Collectively, these studies suggest that the $\alpha 5$ subunit acts as an inhibitory signal that limits nicotine consumption and rewarding effects in smokers.

5.3 Transcriptionally Adaptive Changes

A potential way smoking and genetics may interact is through transcriptionally driven adaptive changes. It is now clear that continued drug use induces adaptive changes in the central nervous system that lead to drug dependence. Long-term adaptations in cellular signaling mechanisms are likely part of the maintenance of drug dependence, which may be necessary for their development and persistence. One well-characterized protein responsible for regulating gene expression is the transcription factor cAMP response element binding protein (CREB). Both human and animal studies have shown that CREB-dependent transcription is an important molecular mechanism underlying dependence on multiple drugs of abuse, including nicotine (Nestler 2005). In human studies, CREB expression correlates with the number of cigarettes smoked per day (Lenz et al. 2010). In adult mice, CREB activation is necessary for nicotine reward (Walters et al. 2005). These studies and others suggest a role for CREB in mediating the neuroplasticity changes that characterize nicotine dependence (Kenney et al. 2012; Portugal et al. 2012; Turner et al. 2014). For example, Turner and colleagues (Fisher et al. 2017; Turner et al. 2014) showed that hippocampal CREB signaling and the associated changes in synaptic plasticity impacted nicotine withdrawal phenotypes in mice. Further studies (Fisher et al. 2017) then demonstrated that site-specific CREB deletion in the

hippocampus impacted both cognitive and affective nicotine withdrawal phenotypes due to reduced CREB-mediated transcription of neuroplasticity-related genes, such as Arc and TrkB. However, while CREB is an important regulator of transcription, its widespread function precludes its use for development of targeted therapeutics. Instead, current studies are examining genomic CREB targets as potential therapeutics. For example, CREB ChIP-Seq data show that CREB's activation by chronic nicotine and withdrawal differentially modulate its binding to the genome and network pathway analyses of these data highlight the importance of different families of neuroplasticity genes, such as neurotrophin, netrin, and neuregulin family members (Turner et al. 2014).

Genes encoding a member of the epidermal growth factor family, neuregulin 3 (NRG3), and its receptor, ErbB4, have been recently linked to smoking cessation outcomes (Loukola et al. 2014; Turner et al. 2014). NRG3 is present on excitatory cells and signals transsynaptically through the ErbB4 receptor, which is found on select inhibitory cell types (Vullhorst et al. 2017). Genetic variation in this pathway has been demonstrated to impact multiple dimensions of smoking behavior, including smoking initiation, amount smoked, and nicotine dependence (Loukola et al. 2008, 2014). In particular, single nucleotide polymorphisms in the gene for NRG3 result in impaired ability to quit smoking in the clinical population (Turner et al. 2014). Conserved and consistent association of variants in this pathway with nicotine dependence measures lends confidence to future mechanistic evaluation of these associations. Furthermore, these data suggest that while therapeutic interventions for molecules such as CREB are unlikely, evaluation of those gene families regulated by CREB has great potential for future therapeutic development. For example, compounds targeting downstream effectors of ErbB4, the receptor for the CREB target gene NRG3, are already being developed for clinical use in psychiatric conditions such as schizophrenia (Law et al. 2012), a condition highly comorbid with nicotine dependence.

6 Other Constituents in Nicotine and Tobacco Products Mediating Dependence

While the field has focused on nicotine as the main psychoactive constituent in cigarettes and e-cigarettes, it is important to consider other compounds in the products that may alter the pharmacokinetics of nicotine and/or exert independent reinforcing effects on the substance user. Accumulating research has provided evidence that some non-nicotine constituents have innate reinforcing properties, which may thereby increase product use. For instance, anatabine, anabasine, cotinine, and myosmine have all been shown to increase the reinforcing properties of nicotine (Clemens et al. 2009; Hall et al. 2014). Mesolimbic dopamine levels are also increased in the presence of cotinine, acetaldehyde, and nornicotine at a level similar to that found for other substances of abuse (Bardo et al. 1999; Dwoskin et al. 1993, 1999; Foddai et al. 2004). Acetaldehyde and several minor alkaloids have also been shown to act as reinforcers (Myers et al. 1982; Peana et al. 2010; Smith et al. 2015),

although it is debatable as to whether this potentiative effect occurs at the concentrations of product consumed by humans. Another potential candidate mediating the enhanced reinforcing effect of nicotine in tobacco cigarettes is MAO inhibition with chronic exposure (Fowler et al. 1996, 2000). Consistent with the findings in humans, pharmacological inhibition of MAO in rodents has been shown to increase low-dose nicotine self-administration (Smith et al. 2015). Furthermore, the β -carbolines, harman and norharman, appear to inhibit MAO and may partially explain the effects found with tobacco consumption (Truman et al. 2017). With specific regard to e-cigarettes, several factors may interact to affect nicotine absorption and bioavailability, including pH, concentration of propylene glycol to glycerine vehicle, alcohol, nicotine, temperature, concentration of nicotine, and user characteristics (e.g., puff topography, level of experience) (DeVito and Krishnan-Sarin 2018). In addition, propylene glycol has been shown to decrease the aversive effects of high-dose nicotine, which may subsequently promote higher levels of nicotine consumption (Harris et al. 2018).

Various flavorant additives are also found in tobacco and e-cigarette products, and this topic has garnered recent attention since product flavor has been reported to be a main reason for the initiation of e-cigarette use among adolescents (Kong et al. 2015). Interestingly, a fMRI study found that e-cigarette advertisements showing sweet- and fruit-flavored products elicited a greater increase in nucleus accumbens activity compared to tobacco e-cigarette advertisements or control images of sweets and fruits (Garrison et al. 2018), thus demonstrating the strong cue-associated effects of these flavorants on brain reward circuitry. In addition to enhancing the attractiveness and palatability of the cigarette, the additives may additionally interact with nicotine or other constituents at a biological level. For instance, menthol, a common flavoring additive to cigarettes and e-cigarettes, has garnered much attention recently given the preferential use of mentholated products among youth, adult women, and racial/ethnic minorities (FDA 2013; Villanti et al. 2017). In addition to focused marketing in targeted communities, the disproportional use by these populations has been proposed to be due to underlying genetic or biological factors, such as differences in nAChR expression or nicotine metabolism. Indeed, the presence of menthol in cigarettes has been demonstrated to alter nicotine's effects in smokers (Benowitz et al. 2004; Williams et al. 2007), which may be due to menthol-mediated inhibition of nicotine metabolism (Caraballo et al. 2011; Fagan et al. 2016) and potentiative effects on nicotine-mediated dopamine release in brain reward pathways (Zhang et al. 2018). Furthermore, menthol has also been shown to allosterically modulate $\alpha 7$ nAChRs (Ashoor et al. 2013) and can further upregulate nAChR expression (Alsharari et al. 2015). Thus, the pharmacological and addictive properties of nicotine may be enhanced and prolonged in the presence of menthol. This is further evidenced by the finding that mentholated cigarette smokers are less successful in maintaining abstinence following cessation (Caraballo et al. 2011; Fagan et al. 2016; Okuyemi et al. 2007).

7 Therapeutic Approaches for Tobacco and Nicotine Dependence

7.1 Nicotine Replacement Therapies

Nicotine replacement therapies (NRT) represent one of the first effective strategies to promote smoking cessation. In most formulations, nicotine is slowly administered over a prolonged period of time; this approach is thought to attenuate the negative somatic and cognitive effects found during drug withdrawal, while minimizing the reinforcing properties of the drug. A variety of available products include nicotine containing gums, lozenges, and patches. In controlled studies, NRT has been shown to be moderately efficacious in the short-term (days to weeks) (Hartmann-Boyce et al. 2018). However, over longer periods, relapse is often found in most individuals (Hartmann-Boyce et al. 2018), thus necessitating the development of alternate approaches. Along these lines, e-cigarette devices were developed as an NRT and harm reduction product. Compared to the traditional tobacco cigarette, e-cigarettes have been promoted as reducing exposure to carcinogens while providing reinforcing properties of nicotine via inhalation and quick delivery of the drug to the brain. Although e-cigarettes have been reported to assist some individuals in tobacco cessation, the emerging incidence of e-cigarette use among never smokers has represented a concerning trend for the promotion of nicotine dependence, especially among adolescents (Miech et al. 2019). Indeed, while e-cigarettes may be less harmful than tobacco cigarettes, they are by no means harmless, as evidenced by the multitude of chemicals and carcinogens emitted (Goniewicz et al. 2018). It is currently debatable as to whether electronic nicotine delivery devices should be employed by physicians for tobacco cessation since inconsistent findings have been reported with effectiveness and the potential harmful effects with short- and long-term use remain to be resolved (Livingston et al. 2019).

7.2 Varenicline and Bupropion

Given the direct action of nicotine on $\alpha 4\beta 2^*$ nAChRs to mediate the reinforcing properties of the drug, it is perhaps not surprising that the most efficacious pharmacotherapeutics available is varenicline, a partial agonist of $\alpha 4\beta 2^*$ nAChRs. Varenicline also has full agonist, but less potent, effects at $\alpha 7$ and $\alpha 3\beta 4^*$ nAChRs and serotonin 5-HT₃ receptors. Approved by the FDA in 2006, varenicline has been shown to have similar or greater effectiveness in promoting smoking cessation compared to NRT and other approved therapeutics, such as bupropion (Gonzales et al. 2006). Bupropion was first characterized as a dopamine and norepinephrine reuptake inhibitor with antidepressant actions but more recently became approved as a first-line treatment for tobacco cessation. In addition to its actions as a catecholamine reuptake inhibitor, bupropion has also been shown to result in noncompetitive antagonism of $\alpha 4\beta 2^*$ and $\alpha 3\beta 4^*$ nAChRs (Carroll et al. 2014) and negative allosteric modulation of serotonin 5HT_{3A} receptors (Pandhare et al. 2017), either of which

may underlie the beneficial effects found for smoking cessation. In addition to NRT, varenicline, and bupropion, the tricyclic antidepressant nortriptyline and the α -adrenergic agonist clonidine have also been prescribed for smoking cessation, although studies have generally found them to be less effective than the aforementioned therapeutics (Dodd et al. 2018).

7.3 Novel Approaches

With advances in our understanding of the biological mechanisms underlying nicotine's physiological, reinforcing, and aversive effects, novel approaches for therapeutic development hold the promise of achieving substantial long-term clinical outcomes. Since $\alpha 5^*$ nAChRs in the MHB-IPN pathway have been demonstrated to mediate the aversive properties of nicotine that limit intake (Fowler et al. 2011), drug development efforts are focused on generating positive allosteric modulators of these receptors, with the idea of enhancing aversive processing in the presence of nicotine to decrease further drug intake (Jin et al. 2014). Another compound, AT-1001, which is an $\alpha 3\beta 4$ partial agonist, has been shown to reduce nicotine relapse-related behaviors in rodents (Yuan et al. 2017), likely through action on the $\alpha 3\beta 4^*$ nAChRs expressed in the MHB. GLP-1 receptor signaling has also been implicated in MHB-IPN modulation of nicotine intake (Tuesta et al. 2017), and a GLP-1 receptor agonist, liraglutide, is currently being tested for smoking cessation in a clinical trial (Ashare 2019). Another potentially beneficial strategy is to inhibit the main enzyme responsible for metabolizing nicotine, CYP2A6. The foundation of this approach is based on the observation that individuals with allelic variation in the CYP2A6 enzyme exhibit lower levels of nicotine consumption and greater abstinence rates when attempting to quit (Strasser et al. 2007). With CYP2A6 inhibition, lower levels of drug consumption would result in higher levels of nicotine intake, which may thereby lead either to an aversive effect with moderate levels of nicotine consumption or a reinforcing effect at lower levels of nicotine. Methoxsalen, a CYP2A5/CYP2A6 inhibitor, was a promising candidate as it was shown to decrease nicotine dependence-associated behaviors in rodents (Alsharari et al. 2014; Bagdas et al. 2014), but this drug was not further advanced for smoking cessation due to carcinogenic side effects that were unrelated to the CYP2A6 inhibitor actions. As such, current drug development efforts are ongoing to derive alternative CYP2A6 inhibitors. In addition to pharmacotherapeutics, nicotine vaccines have been under development. Conceptually, vaccination results in the generation of antibodies that bind to nicotine in the blood, thereby reducing the amount of nicotine capable of entering the brain. However, double-blind randomized trials have failed to demonstrate sustained benefit in long-term cessation (Hartmann-Boyce et al. 2012; Tonstad et al. 2013), likely due to insufficiently sustained antibody levels. In another approach to minimize nicotine entry into the brain, NicA2-J1 has been developed as a reengineered nicotine-degrading enzyme (Kallupi et al. 2018). Interestingly, while NicA2-J1 did not induce significant differences from the control in nicotine

intake, an attenuation of withdrawal and relapse-related behaviors was found in rats (Kallupi et al. 2018).

8 Conclusion

Tobacco use disorder is the leading cause of preventable disease and death in the United States and worldwide. The health consequences of nicotine addiction resulting from prolonged drug use are tremendous and devastating. After more than three decades of research on the neurobiology of nicotine dependence, health professionals can now turn to several efficacious pharmacotherapies to treat smoking. These agents often double the odds for quitting over placebo and in some cases (i.e., varenicline) almost triple the odds of quitting over those of placebo. However, despite these advances, many smokers relapse, and unfortunately the long-term abstinence rates among smokers attempting to quit remain low. Therefore, a better understanding of the various genetic, behavioral, and biological mechanisms mediating the various aspects of nicotine dependence is paramount.

References

- Alkondon M, Albuquerque EX (2001) Nicotinic acetylcholine receptor alpha7 and alpha4beta2 subtypes differentially control GABAergic input to CA1 neurons in rat hippocampus. *J Neurophysiol* 86:3043–3055
- Alkondon M, Pereira EF, Barbosa CT, Albuquerque EX (1997) Neuronal nicotinic acetylcholine receptor activation modulates gamma-aminobutyric acid release from CA1 neurons of rat hippocampal slices. *J Pharmacol Exp Ther* 283:1396–1411
- Alsharari SD, Siu EC, Tyndale RF, Damaj MI (2014) Pharmacokinetic and pharmacodynamics studies of nicotine after oral administration in mice: effects of methoxsalen, a CYP2A5/6 inhibitor. *Nicotine Tob Res* 16:18–25
- Alsharari SD, King JR, Nordman JC, Muldoon PP, Jackson A, Zhu AZ, Tyndale RF, Kabbani N, Damaj MI (2015) Effects of menthol on nicotine pharmacokinetic, pharmacology and dependence in mice. *PLoS One* 10:e0137070
- Arvaniti M, Jensen MM, Soni N, Wang H, Klein AB, Thiriet N, Pinborg LH, Muldoon PP, Wienecke J, Imad Damaj M, Kohlmeier KA, Gondre-Lewis MC, Mikkelsen JD, Thomsen MS (2016) Functional interaction between Lypd6 and nicotinic acetylcholine receptors. *J Neurochem* 138:806–820
- Ashare R (2019) Daily liraglutide for nicotine dependence. USNLB, [ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT03712098). NCT03712098
- Ashare RL, Falcone M, Lerman C (2014) Cognitive function during nicotine withdrawal: implications for nicotine dependence treatment. *Neuropharmacology* 76(Pt B):581–591
- Ashoor A, Nordman JC, Veltri D, Yang KH, Al Kury L, Shuba Y, Mahgoub M, Howarth FC, Sadek B, Shehu A, Kabbani N, Oz M (2013) Menthol binding and inhibition of alpha7-nicotinic acetylcholine receptors. *PLoS One* 8:e67674
- Bagdas D, Muldoon PP, Zhu AZ, Tyndale RF, Damaj MI (2014) Effects of methoxsalen, a CYP2A5/6 inhibitor, on nicotine dependence behaviors in mice. *Neuropharmacology* 85:67–72
- Bardo MT, Green TA, Crooks PA, Dwoskin LP (1999) Nicotine is self-administered intravenously by rats. *Psychopharmacology* 146:290–296
- Benowitz NL, Herrera B, Jacob P (2004) Mentholated cigarette smoking inhibits nicotine metabolism. *J Pharmacol Exp Ther* 310:1208–1215

- Berrettini W, Yuan X, Tozzi F, Song K, Francks C, Chilcoat H, Waterworth D, Muglia P, Mooser V (2008) Alpha-5/alpha-3 nicotinic receptor subunit alleles increase risk for heavy smoking. *Mol Psychiatry* 13:368–373
- Bierut LJ, Stitzel JA, Wang JC, Hinrichs AL, Gruzza RA, Xuei X, Saccone NL, Saccone SF, Bertelsen S, Fox L, Horton WJ, Breslau N, Budde J, Cloninger CR, Dick DM, Foroud T, Hatsukami D, Hesselbrock V, Johnson EO, Kramer J, Kuperman S, Madden PA, Mayo K, Nurnberger J Jr, Pomerleau O, Porjesz B, Reyes O, Schuckit M, Swan G, Tischfield JA, Edenberg HJ, Rice JP, Goate AM (2008) Variants in nicotinic receptors and risk for nicotine dependence. *Am J Psychiatry* 165:1163–1171
- Broms U, Silventoinen K, Madden PAF, Heath AC, Kaprio J (2006) Genetic architecture of smoking behavior: a study of Finnish adult twins. *Twin Res Hum Genet* 9:64–72
- Caraballo RS, Holiday DB, Stellman SD, Mowery PD, Giovino GA, Muscat JE, Eriksen MP, Bernert JT, Richter PA, Kozlowski LT (2011) Comparison of serum cotinine concentration within and across smokers of menthol and nonmenthol cigarette brands among non-Hispanic black and non-Hispanic white U.S. adult smokers, 2001–2006. *Cancer Epidemiol Biomarkers Prev* 20:1329–1340
- Carmelli D, Swan GE, Robinette D, Fabsitz R (1992) Genetic influence on smoking – a study of male twins. *New Engl J Med* 327:829–833
- Carroll FI, Blough BE, Mascarella SW, Navarro HA, Lukas RJ, Damaj MI (2014) Bupropion and bupropion analogs as treatments for CNS disorders. *Adv Pharmacol* 69:177–216
- CDC (2015) Smoking-attributable mortality, years of potential life lost, and productivity losses – United States, 2000–2004. *MMWR Morb Mortal Wkly Rep* 57(45):1226–1228. Updated 4 Mar 2015
- Champtiaux N, Gotti C, Cordero-Erausquin M, David DJ, Przybylski C, Lena C, Clementi F, Moretti M, Rossi FM, Le Novere N, McIntosh JM, Gardier AM, Changeux JP (2003) Subunit composition of functional nicotinic receptors in dopaminergic neurons investigated with knock-out mice. *J Neurosci* 23:7820–7829
- Changeux JP, Bertrand D, Corringer PJ, Dehaene S, Edelstein S, Lena C, Le Novere N, Marubio L, Picciotto M, Zoli M (1998) Brain nicotinic receptors: structure and regulation, role in learning and reinforcement. *Brain Res Brain Res Rev* 26:198–216
- Claus ED, Blaine SK, Filbey FM, Mayer AR, Hutchison KE (2013) Association between nicotine dependence severity, BOLD response to smoking cues, and functional connectivity. *Neuropsychopharmacology* 38:2363–2372
- Clemens KJ, Caille S, Stinus L, Cador M (2009) The addition of five minor tobacco alkaloids increases nicotine-induced hyperactivity, sensitization and intravenous self-administration in rats. *Int J Neuropsychopharmacol* 12:1355–1366
- Corrigall WA, Coen KM, Adamson KL (1994) Self-administered nicotine activates the mesolimbic dopamine system through the ventral tegmental area. *Brain Res* 653:278–284
- Cosgrove KP, Esterlis I, McKee S, Bois F, Alagille D, Tamagnan GD, Seibyl JP, Krishnan-Sarin S, Staley JK (2010) Beta2* nicotinic acetylcholine receptors modulate pain sensitivity in acutely abstinent tobacco smokers. *Nicotine Tob Res* 12:535–539
- Couey JJ, Meredith RM, Spijker S, Poorthuis RB, Smit AB, Brussaard AB, Mansvelder HD (2007) Distributed network actions by nicotine increase the threshold for spike-timing-dependent plasticity in prefrontal cortex. *Neuron* 54:73–87
- Damaj MI, Kao W, Martin BR (2003) Characterization of spontaneous and precipitated nicotine withdrawal in the mouse. *J Pharmacol Exp Ther* 307:526–534
- Dani JA, Bertrand D (2007) Nicotinic acetylcholine receptors and nicotinic cholinergic mechanisms of the central nervous system. *Annu Rev Pharmacol Toxicol* 47:699–729
- Dani JA, Harris RA (2005) Nicotine addiction and comorbidity with alcohol abuse and mental illness. *Nat Neurosci* 8:1465–1470
- Dani JA, Jenson D, Broussard JI, De Biasi M (2011) Neurophysiology of nicotine addiction. *J Addict Res Ther* S1. pii: 001

- De Biasi M, Dani JA (2011) Reward, addiction, withdrawal to nicotine. *Annu Rev Neurosci* 34:105–130
- Demars MP, Morishita H (2014) Cortical parvalbumin and somatostatin GABA neurons express distinct endogenous modulators of nicotinic acetylcholine receptors. *Mol Brain* 7:75
- DeVito EE, Krishnan-Sarin S (2018) E-cigarettes: impact of e-liquid components and device characteristics on nicotine exposure. *Curr Neuropharmacol* 16:438–459
- Di Chiara G (2000) Role of dopamine in the behavioural actions of nicotine related to addiction. *Eur J Pharmacol* 393:295–314
- Dodd S, Arancini L, Gomez-Coronado N, Gasser R, Lubman DI, Dean OM, Berk M (2018) Considerations when selecting pharmacotherapy for nicotine dependence. *Expert Opin Pharmacother* 20:1–6
- Dvoskin LP, Buxton ST, Jewell AL, Crooks PA (1993) S(–)-Nornicotine increases dopamine release in a calcium-dependent manner from superfused rat striatal slices. *J Neurochem* 60:2167–2174
- Dvoskin LP, Teng L, Buxton ST, Crooks PA (1999) (S)-(–)-Cotinine, the major brain metabolite of nicotine, stimulates nicotinic receptors to evoke [3H]dopamine release from rat striatal slices in a calcium-dependent manner. *J Pharmacol Exp Ther* 288:905–911
- Fagan P, Pokhrel P, Herzog TA, Pagano IS, Franke AA, Clanton MS, Alexander LA, Trinidad DR, Sakuma KL, Johnson CA, Moolchan ET (2016) Nicotine metabolism in young adult daily menthol and nonmenthol smokers. *Nicotine Tob Res* 18:437–446
- FDA (2013) Preliminary scientific evaluation of the possible public health effects of menthol versus nonmenthol cigarettes. <http://www.fda.gov/downloads/UCM361598.pdf>. Accessed Apr 2017
- Fisher ML, LeMalfant RM, Zhou L, Huang G, Turner JR (2017) Distinct roles of CREB within the ventral and dorsal Hippocampus in mediating nicotine withdrawal phenotypes. *Neuropsychopharmacology* 42:1599–1609
- Flores CM, Rogers SW, Pabreza LA, Wolfe BB, Kellar KJ (1992) A subtype of nicotinic cholinergic receptor in rat brain is composed of alpha 4 and beta 2 subunits and is up-regulated by chronic nicotine treatment. *Mol Pharmacol* 41:31–37
- Foddai M, Dosia G, Spiga S, Diana M (2004) Acetaldehyde increases dopaminergic neuronal activity in the VTA. *Neuropsychopharmacology* 29:530–536
- Forget B, Scholze P, Langa F, Morel C, Pons S, Mondoloni S, Besson M, Durand-de Cuttoli R, Hay A, Tricoire L, Lambollez B, Mourot A, Faure P, Maskos U (2018) A human polymorphism in CHRNA5 is linked to relapse to nicotine seeking in transgenic rats. *Curr Biol* 28:3244–3253. e7
- Fowler JS, Volkow ND, Wang GJ, Pappas N, Logan J, Shea C, Alexoff D, MacGregor RR, Schlyer DJ, Zezulko I, Wolf AP (1996) Brain monoamine oxidase A inhibition in cigarette smokers. *Proc Natl Acad Sci U S A* 93:14065–14069
- Fowler JS, Wang GJ, Volkow ND, Franceschi D, Logan J, Pappas N, Shea C, MacGregor RR, Garza V (2000) Maintenance of brain monoamine oxidase B inhibition in smokers after overnight cigarette abstinence. *Am J Psychiatry* 157:1864–1866
- Fowler CD, Lu Q, Johnson PM, Marks MJ, Kenny PJ (2011) Habenular alpha5 nicotinic receptor subunit signalling controls nicotine intake. *Nature* 471:597–601
- Fowler CD, Tuesta L, Kenny PJ (2013) Role of alpha5* nicotinic acetylcholine receptors in the effects of acute and chronic nicotine treatment on brain reward function in mice. *Psychopharmacology* 229(3):503–513
- Froeliger B, Kozink RV, Rose JE, Behm FM, Salley AN, McClernon FJ (2010) Hippocampal and striatal gray matter volume are associated with a smoking cessation treatment outcome: results of an exploratory voxel-based morphometric analysis. *Psychopharmacology* 210:577–583
- Garrison KA, O'Malley SS, Gueorguieva R, Krishnan-Sarin S (2018) A fMRI study on the impact of advertising for flavored e-cigarettes on susceptible young adults. *Drug Alcohol Depend* 186:233–241

- George AA, Bloy A, Miwa JM, Lindstrom JM, Lukas RJ, Whiteaker P (2017) Isoform-specific mechanisms of $\alpha 3\beta 4^{*}$ -nicotinic acetylcholine receptor modulation by the protoxin lynx1. *FASEB J* 31:1398–1420
- Gerzanich V, Wang F, Kuryatov A, Lindstrom J (1998) Alpha 5 subunit alters desensitization, pharmacology, Ca^{++} permeability and Ca^{++} modulation of human neuronal alpha 3 nicotinic receptors. *J Pharmacol Exp Ther* 286:311–320
- Girod R, Role LW (2001) Long-lasting enhancement of glutamatergic synaptic transmission by acetylcholine contrasts with response adaptation after exposure to low-level nicotine. *J Neurosci* 21:5182–5190
- Goniewicz ML, Smith DM, Edwards KC, Blount BC, Caldwell KL, Feng J, Wang L, Christensen C, Ambrose B, Borek N, van Bommel D, Konkel K, Erives G, Stanton CA, Lambert E, Kimmel HL, Hatsukami D, Hecht SS, Niaura RS, Travers M, Lawrence C, Hyland AJ (2018) Comparison of nicotine and toxicant exposure in users of electronic cigarettes and combustible cigarettes. *JAMA Netw Open* 1:e185937
- Gonzales D, Rennard SI, Nides M, Oncken C, Azoulay S, Billing CB, Watsky EJ, Gong J, Williams KE, Reeves KR, Varenicline Phase 3 Study Group (2006) Varenicline, an $\alpha 4\beta 2$ nicotinic acetylcholine receptor partial agonist, vs sustained-release bupropion and placebo for smoking cessation: a randomized controlled trial. *JAMA* 296:47–55
- Hall BJ, Wells C, Allenby C, Lin MY, Hao I, Marshall L, Rose JE, Levin ED (2014) Differential effects of non-nicotine tobacco constituent compounds on nicotine self-administration in rats. *Pharmacol Biochem Behav* 120:103–108
- Harris AC, Muelken P, Haave Z, Swain Y, Smethells JR, LeSage MG (2018) Propylene glycol, a major electronic cigarette constituent, attenuates the adverse effects of high-dose nicotine as measured by intracranial self-stimulation in rats. *Drug Alcohol Depend* 193:162–168
- Hartmann-Boyce J, Cahill K, Hatsukami D, Cornuz J (2012) Nicotine vaccines for smoking cessation. *Cochrane Database Syst Rev* 8:CD007072
- Hartmann-Boyce J, Chepkin SC, Ye W, Bullen C, Lancaster T (2018) Nicotine replacement therapy versus control for smoking cessation. *Cochrane Database Syst Rev* 5:CD000146
- Henderson BJ, Srinivasan R, Nichols WA, Dilworth CN, Gutierrez DF, Mackey ED, McKinney S, Drenan RM, Richards CI, Lester HA (2014) Nicotine exploits a COPI-mediated process for chaperone-mediated up-regulation of its receptors. *J Gen Physiol* 143:51–66
- Hill JA Jr, Zoli M, Bourgeois JP, Changeux JP (1993) Immunocytochemical localization of a neuronal nicotinic receptor: the beta 2-subunit. *J Neurosci* 13:1551–1568
- Huang YY, Kandel ER, Levine A (2008) Chronic nicotine exposure induces a long-lasting and pathway-specific facilitation of LTP in the amygdala. *Learn Mem* 15:603–610
- Ibanez-Tallon I, Miwa JM, Wang HL, Adams NC, Crabtree GW, Sine SM, Heintz N (2002) Novel modulation of neuronal nicotinic acetylcholine receptors by association with the endogenous protoxin lynx1. *Neuron* 33:893–903
- Jackson KJ, Martin BR, Changeux JP, Damaj MI (2008) Differential role of nicotinic acetylcholine receptor subunits in physical and affective nicotine withdrawal signs. *J Pharmacol Exp Ther* 325:302–312
- Jackson KJ, Kota DH, Martin BR, Damaj MI (2009) The role of various nicotinic receptor subunits and factors influencing nicotine conditioned place aversion. *Neuropharmacology* 56:970–974
- Jackson KJ, Marks MJ, Vann RE, Chen X, Gamage TF, Warner JA, Damaj MI (2010) Role of $\alpha 5$ nicotinic acetylcholine receptors in pharmacological and behavioral effects of nicotine in mice. *J Pharmacol Exp Ther* 334:137–146
- Jackson KJ, Sanjakdar SS, Muldoon PP, McIntosh JM, Damaj MI (2013) The $\alpha 3\beta 4^{*}$ nicotinic acetylcholine receptor subtype mediates nicotine reward and physical nicotine withdrawal signs independently of the $\alpha 5$ subunit in the mouse. *Neuropharmacology* 70:228–235
- Ji D, Lape R, Dani JA (2001) Timing and location of nicotinic activity enhances or depresses hippocampal synaptic plasticity. *Neuron* 31:131–141

- Jin Z, Khan P, Shin Y, Wang J, Lin L, Cameron MD, Lindstrom JM, Kenny PJ, Kamenecka TM (2014) Synthesis and activity of substituted heteroaromatics as positive allosteric modulators for $\alpha 4\beta 2\alpha 5$ nicotinic acetylcholine receptors. *Bioorg Med Chem Lett* 24:674–678
- John D, Berg DK (2015) Long-lasting changes in neural networks to compensate for altered nicotinic input. *Biochem Pharmacol* 97:418–424
- Jones S, Yakel JL (1997) Functional nicotinic ACh receptors on interneurons in the rat hippocampus. *J Physiol* 504(Pt 3):603–610
- Kallupi M, Xue S, Zhou B, Janda KD, George O (2018) An enzymatic approach reverses nicotine dependence, decreases compulsive-like intake, and prevents relapse. *Sci Adv* 4:eaat4751
- Kassiou M, Eberl S, Meikle SR, Birrell A, Constable C, Fulham MJ, Wong DF, Musachio JL (2001) In vivo imaging of nicotinic receptor upregulation following chronic (–)-nicotine treatment in baboon using SPECT. *Nucl Med Biol* 28:165–175
- Kendler KS, Thornton LM, Pedersen NL (2000) Tobacco consumption in Swedish twins reared apart and reared together. *Arch Gen Psychiatry* 57:886–892
- Kenney JW, Poole RL, Adoff MD, Logue SF, Gould TJ (2012) Learning and nicotine interact to increase CREB phosphorylation at the *jnk1* promoter in the hippocampus. *PLoS One* 7:e39939
- Klink R, de Kerchove d'Exaerde A, Zoli M, Changeux JP (2001) Molecular and physiological diversity of nicotinic acetylcholine receptors in the midbrain dopaminergic nuclei. *J Neurosci* 21:1452–1463
- Kong G, Morean ME, Cavallo DA, Camenga DR, Krishnan-Sarin S (2015) Reasons for electronic cigarette experimentation and discontinuation among adolescents and young adults. *Nicotine Tob Res* 17:847–854
- Kuhn S, Schubert F, Gallinat J (2010) Reduced thickness of medial orbitofrontal cortex in smokers. *Biol Psychiatry* 68:1061–1065
- Kuryatov A, Berrettini W, Lindstrom J (2011) Acetylcholine receptor (AChR) $\alpha 5$ subunit variant associated with risk for nicotine dependence and lung cancer reduces ($\alpha 4\beta 2$) $\alpha 5$ AChR function. *Mol Pharmacol* 79:119–125
- Lagostena L, Trocme-Thibierge C, Morain P, Cherubini E (2008) The partial $\alpha 7$ nicotine acetylcholine receptor agonist S 24795 enhances long-term potentiation at CA3-CA1 synapses in the adult mouse hippocampus. *Neuropharmacology* 54:676–685
- Lanca AJ, Adamson KL, Coen KM, Chow BL, Corrigan WA (2000) The pedunculopontine tegmental nucleus and the role of cholinergic neurons in nicotine self-administration in the rat: a correlative neuroanatomical and behavioral study. *Neuroscience* 96:735–742
- Law AJ, Wang Y, Sei Y, O'Donnell P, Piantadosi P, Papaleo F, Straub RE, Huang W, Thomas CJ, Vakkalanka R, Besterman AD, Lipska BK, Hyde TM, Harrison PJ, Kleinman JE, Weinberger DR (2012) Neuregulin 1-ErbB4-PI3K signaling in schizophrenia and phosphoinositide 3-kinase-p110 δ inhibition as a potential therapeutic strategy. *Proc Natl Acad Sci U S A* 109:12165–12170
- Le Foll B, Goldberg SR (2005) Control of the reinforcing effects of nicotine by associated environmental stimuli in animals and humans. *Trends Pharmacol Sci* 26:287–293
- Lenz B, Klafki HW, Hillemecher T, Killisch N, Schaller G, Frieling H, Clepce M, Gossler A, Therauf N, Winterer G, Kornhuber J, Bleich S (2010) Smoking behaviour is associated with expression and phosphorylation of CREB in human buffy coat. *Int J Neuropsychopharmacol* 13:207–215
- Li P, Steinbach JH (2010) The neuronal nicotinic $\alpha 4\beta 2$ receptor has a high maximal probability of being open. *Br J Pharmacol* 160:1906–1915
- Li MD, Cheng R, Ma JZ, Swan GE (2003) A meta-analysis of estimated genetic and environmental effects on smoking behavior in male and female adult twins. *Addiction* 98:23–31
- Li Y, Yuan K, Cai C, Feng D, Yin J, Bi Y, Shi S, Yu D, Jin C, von Deneen KM, Qin W, Tian J (2015) Reduced frontal cortical thickness and increased caudate volume within fronto-striatal circuits in young adult smokers. *Drug Alcohol Depend* 151:211–219
- Lisman JE, Grace AA (2005) The hippocampal-VTA loop: controlling the entry of information into long-term memory. *Neuron* 46:703–713

- Liu JZ, Tozzi F, Waterworth DM, Pillai SG, Muglia P, Middleton L, Berrettini W, Knouff CW, Yuan X, Waeber G, Vollenweider P, Preisig M, Wareham NJ, Zhao JH, Loos RJ, Barroso I, Khaw KT, Grundy S, Barter P, Mahley R, Kesaniemi A, McPherson R, Vincent JB, Strauss J, Kennedy JL, Farmer A, McGuffin P, Day R, Matthews K, Bakke P, Gulsvik A, Lucae S, Ising M, Brueckl T, Horstmann S, Wichmann HE, Rawal R, Dahmen N, Lamina C, Polasek O, Zgaga L, Huffman J, Campbell S, Kooner J, Chambers JC, Burnett MS, Devaney JM, Pichard AD, Kent KM, Satler L, Lindsay JM, Waksman R, Epstein S, Wilson JF, Wild SH, Campbell H, Vitart V, Reilly MP, Li M, Qu L, Wilensky R, Matthai W, Hakonarson HH, Rader DJ, Franke A, Wittig M, Schafer A, Uda M, Terracciano A, Xiao X, Busonero F, Scheet P, Schlessinger D, St Clair D, Rujescu D, Abecasis GR, Grabe HJ, Teumer A, Volzke H, Petersmann A, John U, Rudan I, Hayward C, Wright AF, Kolcic I, Wright BJ, Thompson JR, Balmforth AJ, Hall AS, Samani NJ, Anderson CA, Ahmad T, Mathew CG, Parkes M, Satsangi J, Caulfield M, Munroe PB, Farrall M, Dominiczak A, Worthington J, Thomson W, Eyre S, Barton A, Mooser V, Francks C, Marchini J (2010) Meta-analysis and imputation refines the association of 15q25 with smoking quantity. *Nat Genet* 42:436–440
- Livingston CJ, Freeman RJ, Costales VC, Westhoff JL, Caplan LS, Sherin KM, Niebuhr DW (2019) Electronic nicotine delivery systems or e-cigarettes: American College of Preventive Medicine's practice statement. *Am J Prev Med* 56:167–178
- Loukola A, Broms U, Maunu H, Widen E, Heikkila K, Siivola M, Salo A, Pergadia ML, Nyman E, Sammalisto S, Perola M, Agrawal A, Heath AC, Martin NG, Madden PA, Peltonen L, Kaprio J (2008) Linkage of nicotine dependence and smoking behavior on 10q, 7q and 11p in twins with homogeneous genetic background. *Pharmacogenomics J* 8:209–219
- Loukola A, Wedenoja J, Keskitalo-Vuokko K, Broms U, Korhonen T, Ripatti S, Sarin AP, Pitkanieni J, He L, Happola A, Heikkila K, Chou YL, Pergadia ML, Heath AC, Montgomery GW, Martin NG, Madden PA, Kaprio J (2014) Genome-wide association study on detailed profiles of smoking behavior and nicotine dependence in a twin sample. *Mol Psychiatry* 19:615–624
- Lyukmanova EN, Shenkarev ZO, Shulepko MA, Mineev KS, D'Hoedt D, Kasheverov IE, Filkin SY, Krivolopova AP, Janickova H, Dolezal V, Dolgikh DA, Arseniev AS, Bertrand D, Tsetlin VI, Kirpichnikov MP (2011) NMR structure and action on nicotinic acetylcholine receptors of water-soluble domain of human LYNX1. *J Biol Chem* 286:10618–10627
- Mansvelder HD, McGehee DS (2000) Long-term potentiation of excitatory inputs to brain reward areas by nicotine. *Neuron* 27:349–357
- Markou A, Kenny PJ (2002) Neuroadaptations to chronic exposure to drugs of abuse: relevance to depressive symptomatology seen across psychiatric diagnostic categories. *Neurotox Res* 4:297–313
- Marks MJ, Burch JB, Collins AC (1983) Effects of chronic nicotine infusion on tolerance development and nicotinic receptors. *J Pharmacol Exp Ther* 226:817–825
- Maskos U (2010) Role of endogenous acetylcholine in the control of the dopaminergic system via nicotinic receptors. *J Neurochem* 114:641–646
- McClernon FJ, Gilbert DG (2004) Human functional neuroimaging in nicotine and tobacco research: basics, background, and beyond. *Nicotine Tob Res* 6:941–959
- Miech R, Johnston L, O'Malley PM, Bachman JG, Patrick ME (2019) Adolescent vaping and nicotine use in 2017–2018 – U.S. National Estimates. *N Engl J Med* 380:192–193
- Miwa JM, Walz A (2012) Enhancement in motor learning through genetic manipulation of the *Lynx1* gene. *PLoS One* 7:e43302
- Miwa JM, Ibanez-Tallon I, Crabtree GW, Sanchez R, Sali A, Role LW, Heintz N (1999) *Lynx1*, an endogenous toxin-like modulator of nicotinic acetylcholine receptors in the mammalian CNS. *Neuron* 23:105–114
- Miwa JM, Freedman R, Lester HA (2011) Neural systems governed by nicotinic acetylcholine receptors: emerging hypotheses. *Neuron* 70:20–33

- Myers WD, Ng KT, Singer G (1982) Intravenous self-administration of acetaldehyde in the rat as a function of schedule, food deprivation and photoperiod. *Pharmacol Biochem Behav* 17:807–811
- Nakauchi S, Sumikawa K (2012) Endogenously released ACh and exogenous nicotine differentially facilitate long-term potentiation induction in the hippocampal CA1 region of mice. *Eur J Neurosci* 35:1381–1395
- Nestler EJ (2005) Is there a common molecular pathway for addiction? *Nat Neurosci* 8:1445–1449
- Nissen NI, Anderson KR, Wang H, Lee HS, Garrison C, Eichelberger SA, Ackerman K, Im W, Miwa JM (2018) Augmenting the antinociceptive effects of nicotinic acetylcholine receptor activity through lynx1 modulation. *PLoS One* 13:e0199643
- Okuyemi KS, Faseru B, Sanderson Cox L, Bronars CA, Ahluwalia JS (2007) Relationship between menthol cigarettes and smoking cessation among African American light smokers. *Addiction* 102:1979–1986
- Omelchenko N, Sesack SR (2005) Laterodorsal tegmental projections to identified cell populations in the rat ventral tegmental area. *J Comp Neurol* 483:217–235
- Orejarena MJ, Herrera-Solis A, Pons S, Maskos U, Maldonado R, Robledo P (2012) Selective re-expression of beta2 nicotinic acetylcholine receptor subunits in the ventral tegmental area of the mouse restores intravenous nicotine self-administration. *Neuropharmacology* 63:235–241
- Pandhare A, Pappu AS, Wilms H, Blanton MP, Jansen M (2017) The antidepressant bupropion is a negative allosteric modulator of serotonin type 3A receptors. *Neuropharmacology* 113:89–99
- Pang X, Liu L, Ngolab J, Zhao-Shea R, McIntosh JM, Gardner PD, Tapper AR (2016) Habenula cholinergic neurons regulate anxiety during nicotine withdrawal via nicotinic acetylcholine receptors. *Neuropharmacology* 107:294–304
- Peana AT, Muggironi G, Diana M (2010) Acetaldehyde-reinforcing effects: a study on oral self-administration behavior. *Front Psych* 1:23
- Perry DC, Davila-Garcia MI, Stockmeier CA, Kellar KJ (1999) Increased nicotinic receptors in brains from smokers: membrane binding and autoradiography studies. *J Pharmacol Exp Ther* 289:1545–1552
- Picciozzo MR, Zoli M, Rimondini R, Lena C, Marubio LM, Pich EM, Fuxe K, Changeux JP (1998) Acetylcholine receptors containing the beta2 subunit are involved in the reinforcing properties of nicotine. *Nature* 391:173–177
- Pittaras EC, Faure A, Leray X, Moraitopoulou E, Cressant A, Rabat AA, Meunier C, Fossier P, Granon S (2016) Neuronal nicotinic receptors are crucial for tuning of E/I balance in prefrontal cortex and for decision-making processes. *Front Psych* 7:171
- Pons S, Fattore L, Cossu G, Tolu S, Porcu E, McIntosh JM, Changeux JP, Maskos U, Fratta W (2008) Crucial role of alpha4 and alpha6 nicotinic acetylcholine receptor subunits from ventral tegmental area in systemic nicotine self-administration. *J Neurosci* 28:12318–12327
- Portugal GS, Wilkinson DS, Turner JR, Blendy JA, Gould TJ (2012) Developmental effects of acute, chronic, and withdrawal from chronic nicotine on fear conditioning. *Neurobiol Learn Mem* 97:482–494
- Ramirez-Latorre J, Yu CR, Qu X, Perin F, Karlin A, Role L (1996) Functional contributions of alpha5 subunit to neuronal acetylcholine receptor channels. *Nature* 380:347–351
- Saccone NL, Wang JC, Breslau N, Johnson EO, Hatsukami D, Saccone SF, Grucza RA, Sun L, Duan W, Budde J, Culverhouse RC, Fox L, Hinrichs AL, Steinbach JH, Wu M, Rice JP, Goate AM, Bierut LJ (2009) The CHRNA5-CHRNA3-CHRNA4 nicotinic receptor subunit gene cluster affects risk for nicotine dependence in African-Americans and in European-Americans. *Cancer Res* 69:6848–6856
- Salas R, Sturm R, Boulter J, De Biasi M (2009) Nicotinic receptors in the habenulo-interpeduncular system are necessary for nicotine withdrawal in mice. *J Neurosci* 29:3014–3018
- Salminen O, Murphy KL, McIntosh JM, Drago J, Marks MJ, Collins AC, Grady SR (2004) Subunit composition and pharmacology of two classes of striatal presynaptic nicotinic acetylcholine receptors mediating dopamine release in mice. *Mol Pharmacol* 65:1526–1535

- Sanjakdar SS, Maldoon PP, Marks MJ, Brunzell DH, Maskos U, McIntosh JM, Bowers MS, Damaj MI (2015) Differential roles of alpha6beta2* and alpha4beta2* neuronal nicotinic receptors in nicotine- and cocaine-conditioned reward in mice. *Neuropsychopharmacology* 40:350–360
- Schoenbaum G, Chang CY, Lucantonio F, Takahashi YK (2016) Thinking outside the box: orbitofrontal cortex, imagination, and how we can treat addiction. *Neuropsychopharmacology* 41:2966–2976
- Smith TT, Schaff MB, Rupprecht LE, Schassburger RL, Buffalari DM, Murphy SE, Sved AF, Donny EC (2015) Effects of MAO inhibition and a combination of minor alkaloids, beta-carbolines, and acetaldehyde on nicotine self-administration in adult male rats. *Drug Alcohol Depend* 155:243–252
- Srinivasan R, Pantoja R, Moss FJ, Mackey ED, Son CD, Miwa J, Lester HA (2011) Nicotine up-regulates alpha4beta2 nicotinic receptors and ER exit sites via stoichiometry-dependent chaperoning. *J Gen Physiol* 137:59–79
- Stoker AK, Olivier B, Markou A (2012) Role of alpha7- and beta4-containing nicotinic acetylcholine receptors in the affective and somatic aspects of nicotine withdrawal: studies in knockout mice. *Behav Genet* 42:423–436
- Strasser AA, Malaiyandi V, Hoffmann E, Tyndale RF, Lerman C (2007) An association of CYP2A6 genotype and smoking topography. *Nicotine Tob Res* 9:511–518
- Sullivan PF, Kendler KS (1999) The genetic epidemiology of smoking. *Nicotine Tob Res* 1(Suppl 2):S51–S57; discussion S69–70
- Tang J, Dani JA (2009) Dopamine enables in vivo synaptic plasticity associated with the addictive drug nicotine. *Neuron* 63:673–682
- Tapper AR, McKinney SL, Nashmi R, Schwarz J, Deshpande P, Labarca C, Whiteaker P, Marks MJ, Collins AC, Lester HA (2004) Nicotine activation of alpha4* receptors: sufficient for reward, tolerance, and sensitization. *Science* 306:1029–1032
- Tekinay AB, Nong Y, Miwa JM, Lieberam I, Ibanez-Tallon I, Greengard P, Heintz N (2009) A role for LYNX2 in anxiety-related behavior. *Proc Natl Acad Sci U S A* 106:4477–4482
- Thomsen MS, Mikkelsen JD (2012) Type I and II positive allosteric modulators differentially modulate agonist-induced up-regulation of alpha7 nicotinic acetylcholine receptors. *J Neurochem* 123:73–83
- Tonstad S, Heggen E, Giljam H, Lagerback PA, Tonnesen P, Wikingson LD, Lindblom N, de Villiers S, Svensson TH, Fagerstrom KO (2013) Nicotine(R), a nicotine vaccine, for relapse prevention: a phase II, randomized, placebo-controlled, multicenter clinical trial. *Nicotine Tob Res* 15:1492–1501
- Truman P, Grounds P, Brennan KA (2017) Monoamine oxidase inhibitory activity in tobacco particulate matter: are harman and norharman the only physiologically relevant inhibitors? *Neurotoxicology* 59:22–26
- Tuesta LM, Chen Z, Duncan A, Fowler CD, Ishikawa M, Lee BR, Liu XA, Lu Q, Cameron M, Hayes MR, Kamenecka TM, Pletcher M, Kenny PJ (2017) GLP-1 acts on habenular avoidance circuits to control nicotine intake. *Nat Neurosci* 20:708–716
- Turner JR, Castellano LM, Blendy JA (2011) Parallel anxiolytic-like effects and upregulation of neuronal nicotinic acetylcholine receptors following chronic nicotine and varenicline. *Nicotine Tob Res* 13:41–46
- Turner JR, Wilkinson DS, Poole RL, Gould TJ, Carlson GC, Blendy JA (2013) Divergent functional effects of sazetidine-a and varenicline during nicotine withdrawal. *Neuropsychopharmacology* 38:2035–2047
- Turner JR, Ray R, Lee B, Everett L, Xiang J, Jepson C, Kaestner KH, Lerman C, Blendy JA (2014) Evidence from mouse and man for a role of neuregulin 3 in nicotine dependence. *Mol Psychiatry* 19:801–810
- Vazquez-Sanroman DB, Monje RD, Bardo MT (2016) Nicotine self-administration remodels perineuronal nets in ventral tegmental area and orbitofrontal cortex in adult male rats. *Addict Biol* 22(6):1743–1755

- Villanti AC, Johnson AL, Ambrose BK, Cummings KM, Stanton CA, Rose SW, Feirman SP, Tworek C, Glasser AM, Pearson JL, Cohn AM, Conway KP, Niaura RS, Bansal-Travers M, Hyland A (2017) Flavored tobacco product use in youth and adults: findings from the first wave of the PATH study (2013–2014). *Am J Prev Med* 53(2):139–151
- Vullhorst D, Ahmad T, Karavanova I, Keating C, Buonanno A (2017) Structural similarities between Neuregulin 1-3 isoforms determine their subcellular distribution and signaling mode in central neurons. *J Neurosci* 37:5232–5249
- Wada E, Wada K, Boulter J, Deneris E, Heinemann S, Patrick J, Swanson LW (1989) Distribution of alpha 2, alpha 3, alpha 4, and beta 2 neuronal nicotinic receptor subunit mRNAs in the central nervous system: a hybridization histochemical study in the rat. *J Comp Neurol* 284:314–335
- Walters CL, Cleck JN, Kuo YC, Blendy JA (2005) Mu-opioid receptor and CREB activation are required for nicotine reward. *Neuron* 46:933–943
- Walters CL, Brown S, Changeux JP, Martin B, Damaj MI (2006) The beta2 but not alpha7 subunit of the nicotinic acetylcholine receptor is required for nicotine-conditioned place preference in mice. *Psychopharmacology* 184:339–344
- Welsby PJ, Rowan MJ, Anwyl R (2009) Intracellular mechanisms underlying the nicotinic enhancement of LTP in the rat dentate gyrus. *Eur J Neurosci* 29:65–75
- Wilking JA, Stitzel JA (2015) Natural genetic variability of the neuronal nicotinic acetylcholine receptor subunit genes in mice: consequences and confounds. *Neuropharmacology* 96:205–212
- Williams JM, Gandhi KK, Steinberg ML, Foulds J, Ziedonis DM, Benowitz NL (2007) Higher nicotine and carbon monoxide levels in menthol cigarette smokers with and without schizophrenia. *Nicotine Tob Res* 9:873–881
- Williams DK, Stokes C, Horenstein NA, Papke RL (2011) The effective opening of nicotinic acetylcholine receptors with single agonist binding sites. *J Gen Physiol* 137:369–384
- Xian H, Scherrer JF, Madden PAF, Lyons MJ, Tsuang M, True WR, Eisen SA (2003) The heritability of failed smoking cessation and nicotine withdrawal in twins who smoked and attempted to quit. *Nicotine Tob Res* 5:245–254
- Yuan M, Malagon AM, Yasuda D, Belluzzi JD, Leslie FM, Zaveri NT (2017) The alpha3beta4 nAChR partial agonist AT-1001 attenuates stress-induced reinstatement of nicotine seeking in a rat model of relapse and induces minimal withdrawal in dependent rats. *Behav Brain Res* 333:251–257
- Zhang M, Harrison E, Biswas L, Tran T, Liu X (2018) Menthol facilitates dopamine-releasing effect of nicotine in rat nucleus accumbens. *Pharmacol Biochem Behav* 175:47–52
- Zhou L, Fisher ML, Cole RD, Gould TJ, Parikh V, Ortinski PI, Turner JR (2018) Neuregulin 3 signaling mediates nicotine-dependent synaptic plasticity in the orbitofrontal cortex and cognition. *Neuropsychopharmacology* 43:1343–1354