

UC Berkeley

UC Berkeley Previously Published Works

Title

The Utility of Outpatient Commitment: II. Mortality Risk and Protecting Health, Safety, and Quality of Life.

Permalink

<https://escholarship.org/uc/item/8rx8v6j7>

Journal

Psychiatric services (Washington, D.C.), 68(12)

ISSN

1075-2730

Authors

Segal, Steven P
Hayes, Stephania L
Rimes, Lachlan

Publication Date

2017-12-01

DOI

10.1176/appi.ps.201600164

Peer reviewed



HHS Public Access

Author manuscript

Psychiatr Serv. Author manuscript; available in PMC 2020 May 11.

Published in final edited form as:

Psychiatr Serv. 2017 December 01; 68(12): 1255–1261. doi:10.1176/appi.ps.201600164.

The Utility of Outpatient Commitment: II. Mortality Risk and Protecting Health, Safety, and Quality of Life

Steven P. Segal, M.S.W., Ph.D.,

School of Social Welfare, University of California, Berkeley.

Department of Social Work, Melbourne School of Health Sciences, Melbourne, Victoria, Australia.

Stephania L. Hayes, M.A., O.T.R.,

School of Social Welfare, University of California, Berkeley.

Lachlan Rimes, B.A.

Victoria Department of Health and Human Services, Melbourne.

Abstract

Objective: This study assessed the contribution of a form of outpatient commitment—community treatment orders (CTOs)—to mortality risk and quality of life of patients with severe mental illness.

Methods: Data (2000–2012) were obtained from the Australian National Death Index, Victoria Department of Health, Victoria police records, and National Outcomes and CaseMix Collection quality-of-life records for patients in the Victorian Psychiatric Case Register/RAPID with a history of psychiatric hospitalization: CTO cohort, N=11,424; non-CTO cohort, N=16,161. The contribution of CTOs to mortality risk associated with CTO facilitation of access to general medical care and prevention of criminal involvement was assessed with logistic regression models. Cohort differences in quality of life were also examined.

Results: A total of 2,727 patients (10%) in the overall sample died, and the sample had a higher mortality risk than the general population. Probability of death by any cause was 9% lower in the CTO cohort than in the non-CTO cohort. Facilitation of access to medical care accounted for a 20% reduction in risk of non-injury-related deaths in the CTO cohort, compared with the non-CTO cohort. Risk of death by self-harm was 32% higher, compared with the non-CTO cohort. CTO placement appeared to lead to a gain of 3.8 years of life among men and 2.4 years among women, compared with the non-CTO cohort. Quality-of-life scores were modestly less favorable for the non-CTO cohort.

Conclusions: CTO placement was associated with lower mortality risk via facilitated access to medical care and with modest enhancement of quality of life.

Population research has consistently shown increased mortality among persons with severe mental illness (1–8). After widespread deinstitutionalization (9), compulsory supervision outside a hospital has been developed internationally for the treatment of persons with

Send correspondence to Dr. Segal (spsegal@berkeley.edu).

The authors report no financial relationships with commercial interests.

mental illness. Outpatient commitment seeks to provide protection and care via compulsory supervision for people with severe mental illness (10,11). This study investigated the relationship between the use of outpatient commitment in Victoria, Australia, which operates by means of community treatment orders (CTOs), and risk of mortality experienced by individuals hospitalized for psychiatric reasons. [A description of CTO use in Victoria is included in an online supplement to this article.]

In 2000, Victoria closed its last psychiatric hospital and moved to provide integrated health care centered around its general hospitals. In this new era, some have pointed to an increase in the number of CTOs issued as an indication of overreliance on involuntary community treatment (12,13). An alternative view is that the increase in CTOs represents clinicians' assessments that briefer hospitalizations and early release from an inpatient episode place patients at increased risk of adverse outcomes (14), one of which is increased mortality without the care provided in a psychiatric hospital. In this view, CTOs are the less restrictive alternative to a psychiatric hospitalization. Indeed, supervision provided under CTOs between 1990 and 2000 in Victoria (15) and elsewhere (16) has been found to be associated with reduced mortality risk. However, the reasons for this association remain unclear, as do the effects of CTO placement on individuals' quality of life (17).

Given the changes in Victoria's health system, this study addressed the following questions: Did being placed on a CTO between 2000 and 2010 continue to be associated with reduced mortality risk? If so, can reduced mortality risk be attributed to CTO-facilitated access to acute medical care or to the potential of CTOs to reduce involvement in crimes against persons? If CTO placement continued to be associated with reduced mortality risk in 2000–2010 (15), are the years of life saved associated with a poorer quality of life (17,18)?

METHODS

Samples

Mental health records from the state of Victoria, Australia, for 2000 to 2010 were obtained from the Victorian Psychiatric Case Register/RAPID (VPCR/RAPID) system for 27,535 patients who experienced psychiatric hospitalization during the period. A total of 11,424 were first placed on a CTO during this period (2000–2010), and 16,161 were hospitalized but never placed on a CTO. The VPCR/RAPID system provides details on all clinical mental health contacts occurring in Victoria. The records were matched with the Australian National Death Index, a compilation of all deaths throughout Australia, during the index study period of July 1, 2000, to February 28, 2012. These combined records were linked to the Victoria Police Law Enforcement Assistance Program (LEAP) data (19), which document all police contacts in Victoria associated with perpetration of and victimization by major crimes against persons; the Victorian Emergency Minimum Data set, which consists of clinical episode data from emergency departments of Victorian public hospitals; the Victorian Admitted Episodes Data set, which consists of clinical episode data for admitted episodes of care in Victorian general medical hospitals; and the National Outcomes and Casemix Collection, which includes clinical quality-of-life assessments of patients that are conducted with the Health of the Nation Outcome Scales (HoNOS).

Measurement

The structuring of the data for analysis and the content of each of the data sets are reported in a companion article (14). Briefly, all treatment contacts were organized into inpatient and outpatient episodes of care. Occasions of community service are reported as community treatment days, and intensity of service provision was measured by treatment days per community care episode (14). Access to acute general medical care was measured by receipt of at least one *ICD-10* or AR-DRG (Australian Refined Diagnosis-Related Groups) diagnosis of a major general medical illness in a hospital or emergency department. Perpetration and victimization were documented by reports of major crimes against persons in the Victoria Police LEAP system reports.

In multivariate modeling, Socio-Economic Indexes for Areas (SEIFA) (20) postal code ranks for Victoria provided a measure of social disadvantage (lower scores indicate more disadvantage). The individual's most disadvantaged area of residence was taken into account to control for inherent social inequalities in life expectancy attributable to residence in a poverty area.

The problem of selection associated with comparing cohorts was addressed with preliminary matching on age, sex, and diagnosis in drawing the sample, propensity score control in the study models, and statistical adjustment for potential confounding variables in these models. The propensity score used in the models was based on service, social, and premorbid characteristics that distinguished patients placed on CTOs from other patients in the Victorian mental health system who were not placed on CTOs [see online supplement for additional details].

Quality of life was measured with profiles of the 12 HoNOS items (21): aggression, nonaccidental self-injury, drug or alcohol problems, cognitive problems, general medical illness or disability, hallucinations or delusions, depressed mood, other mental or behavioral problems, relationships, activities of daily living, living conditions, and occupation and activities (14). HoNOS items were independently scored (22) at various stages in a person's involvement with mental health services; scoring was by clinicians not associated with the patient's care. The average of all HoNOS assessments made for each patient during the periods associated with years of life saved for men and women separately was taken as the measure of each patient's quality of life.

Analyses

All analyses were completed with SPSS version 23 (23) and Excel 2016 spreadsheet software (24). Excel was used for computing standardized mortality ratios (SMRs), years of life lost (YLL), and relative risk (RR) statistics (25). SMR comparisons were based on deaths in the Victorian population and reported by age and gender (26). RR and YLL statistics among patients in the CTO and non-CTO cohorts were computed by using life table analyses. The analyses compared the expected age- and gender-adjusted mortality of those in the CTO cohort with their expected mortality, assuming that the CTO group had the same age and gender distribution as the non-CTO patients; SMRs and YLL were used to compare life expectancy at birth of the average-age sample member at midstudy. The YLL

was computed separately for men and women. The index year was 1971, when a man's life expectancy in Victoria was 68.7 years and a woman's was 75.2 years.

The contributions of CTOs to mortality risk and the role of access to general medical care and of crime and victimization involvement in interaction with CTO exposure were assessed with logistic regressions. Significant interactions were evaluated on the basis of the model statistics. The logistic models analyzed the contribution of a CTO to non-injury-related mortality risk and death from injury (that is, *ICD-10* codes including accidents, homicide, suicide, and other unexplained causes of death [27]). The models were adjusted for treatment days per community care episode (that is, how much supervision was actually provided); gender; age; age at entry into the mental health system; diagnoses (schizophrenia, major affective disorders, paranoia and other psychoses, and dementia); total number of inpatient days (a control for the protective character of hospitalization in a patient's treatment); time in the study (risk period); potential stereotype and communication effects (non-English speaker and Aboriginal or Torres Strait Islander status); propensity to be selected into the CTO cohort; the patient's psychosocial profile reflected in his or her 12 HoNOS scores at both inpatient admission and release; and, given findings of an association between community context and death (28), the lowest SEIFA postal code rank of a neighborhood in which the individual resided (20).

Implications for patient quality of life were determined by using analysis of variance to evaluate differences in the mean scores on the 12 HoNOS items over the periods of life saved, according to YLL statistics for males and females separately.

Ethics

Ethics committees of the Department of Health and Human Services of Victoria, the Australian Institute of Health and Welfare, the Australian National Death Index, the National Coronal Registry, the Victoria Police, and the Human Subjects Committee of the University of California, Berkeley, approved the human subjects procedures for the project. The project was compliant with data regulations set by all approving agencies and their ethics committees; no data breaches or other adverse events occurred during the study.

RESULTS

Sample Characteristics

The mean \pm SD age of the overall sample of hospitalized patients ($N=27,585$) was 34.0 ± 16.7 years at the study outset. In the overall sample, 15,480 (56%) were men and 12,103 (44%) were women, and two were of unknown gender. Compared with the non-CTO cohort, the CTO cohort had proportionally more men (58%, $N=6,646$, versus 55%, $N=8,834$) and was approximately three years younger (32.4 ± 15.3 versus 35.4 ± 17.5). The diagnostic, social, and service characteristics of the samples are reported in the companion article (14).

Mortality

The crude cumulative death rate per 1,000 for Victoria's population over the 11.8-year period (July 1, 2000, to February 28, 2012) was 77.85. For the overall sample of hospitalized

patients, it was 98.93—90.16 for the CTO cohort and 105.10 for the non-CTO cohort (Table 1). During the study period, the death rate in the CTO cohort was 1.27 times the rate in the general population, and the death rate in the non-CTO cohort was 1.35 times the general population rate. For the CTO cohort, the risk of death was lower than for the non-CTO cohort (RR=.86). On the basis of the age- and gender-specific death rates for the state of Victoria, the SMRs were 1.17 for patients in the CTO cohort (1.14 for men and 1.22 for women) and 1.36 for patients in the non-CTO cohort (1.36 for men and 1.37 for women).

During the 11.8-year study period, there were 2,727 deaths (10% of the overall sample). The CTO cohort accounted for 38% of these deaths (N=1,030), and the non-CTO cohort accounted for 62% (N=1,697, or N=1,695 where information on the gender of the decedent was available). The expected number of deaths in the CTO cohort was 879, indicating an excess of 151 deaths—or 12.8 excess deaths per year. In the CTO cohort, the number of excess deaths over the 11.8-year period was 72 among men and 79 among women. In the non-CTO cohort, the expected number of deaths was 1,243, indicating an excess of 452, or 38.3 excess deaths per year. In the non-CTO cohort, the number of excess deaths was 249 for men and 203 for women.

Figure 1 shows YLL for men and women in both cohorts. The non-CTO cohort lost 5,352 more years of expected life, compared with the CTO cohort. The 938 men who died in the non-CTO cohort lost 3,550 more years of life than the 590 men who died in the CTO cohort—or, on average, 3.8 more YLL per individual in the non-CTO cohort. The 757 women in the non-CTO cohort who died lost 1,802 more years of life than the 440 women who died in the CTO cohort—or, on average, 2.4 more YLL per individual in the non-CTO cohort.

CTO Exposure and Mortality Risk

The logistic regression models, which included the control factors noted above, were all significant ($p < .001$) (Table 2). Results indicated that the CTO cohort experienced a 9% reduction in overall mortality risk, compared with the non-CTO cohort. Overall risk has two components: one for external causes and one for other causes. The 9% is the net effect of the 17% ($\text{Exp}(b) = .83$) reduced mortality risk in deaths by other causes (that is, exclusive of the external causes attributable to assault, self-harm, and undetermined intent) and an elevated risk of 26% ($\text{Exp}(b) = 1.26$) associated with external causes, assault, self-harm, and undetermined intent. Furthermore, the elevated risk associated with external causes in the CTO cohort seemed to be primarily explained by a 32% ($\text{Exp}(b) = 1.32$) increased risk of death from self-harm.

The models were rerun to assess the impact of greater access to general medical care facilitated by CTO oversight and of crime and victimization prevention resulting from CTO involvement (data not shown). When death exclusive of assault, self-harm, and undetermined intent was the criterion variable, access to medical care had a significant interaction with CTO status. The model was significant ($\chi^2 = 5,049.24$, $df = 46$ and $27,129$, $p < .001$; correct classification = 93.3%). The interaction term indicated that among patients on CTOs, which facilitated access to medical care, the risk of death (exclusive of assault, self-harm, and undetermined intent) was 20% less ($\text{Exp}(b) = .80$, 95% confidence

interval=.68-.95). These results remained the same when the analysis took into account all the potential confounding factors listed above [see online supplement].

The model assessing the impact of CTO involvement on risk of death from prevention of crime and victimization was also significant ($\chi^2=306.88$, $df=46$ and $27,129$, $p=.001$; correct classification=98.3%), although the interaction term testing this hypothesis was not.

Quality-of-Life Outcomes

As noted above, CTO placement appeared to lead to a gain of 3.8 years of life among men and 2.4 years among women. Figure 2 presents mean scores on the 12 HoNOS items during the final 3.8 years of life for all men who died and the final 2.4 years for all women who died. Men in the CTO cohort who died and for whom HoNOS assessments were available for the final 3.8 years of life ($N=343$) had significantly higher scores than their non-CTO counterparts ($N=344$) on drug or alcohol problems ($p=.025$) and hallucinations or delusions ($p<.001$). In the non-CTO cohort, however, these men had higher scores on problems associated with self-injury ($p=.042$), cognitive functioning ($p<.001$), physical health ($p<.001$), depression ($p<.001$), and activities of daily living ($p=.011$). The two groups of men did not differ significantly on assessments of aggression, other mental disorders, relationships, living conditions, and occupation or activities.

Women in the CTO cohort who died and for whom HoNOS assessments were available for the final 2.4 years ($N=207$) had significantly higher scores than their non-CTO counterparts ($N=258$) on drug or alcohol problems ($p=.004$), relationships ($p=.003$), and living conditions ($p=.010$). Women in the non-CTO cohort scored higher on problems associated with aggression ($p=.003$), cognitive functioning ($p<.001$), physical health ($p=.043$), depression ($p=.002$), and activities of daily living ($p<.001$). The two groups of women did not differ significantly on problems associated with self-injury, hallucinations or delusions, other mental disorders, and occupation or activities.

DISCUSSION

The analyses examined mortality among patients in Victoria, Australia, with mental disorders severe enough to require psychiatric hospitalization. As a group, these individuals were at increased risk of mortality compared with the population of Victoria. Protective supervision, such as that afforded by CTOs, seems to be of particular importance in Victoria and in other jurisdictions in which a reduced commitment to providing community-based services is evident (14). The protective supervision offered to patients on CTOs, compared with patients not on CTOs, was found to be associated with reductions in excess deaths, with SMRs 14% lower in the CTO cohort. After the analysis controlled for multiple factors, the results indicated that the supervision afforded by CTOs contributed to a 9% reduction in overall risk of death and 17% reduction in risk of death exclusive of assault, self-harm, and undetermined intent. The reduction in overall risk occurred despite the CTO cohort's 32% increased risk of death due to self-harm. These findings replicate mortality risk reduction outcomes associated with CTO supervision that were found in a similar analysis of data from the previous decade in Victoria (15).

Findings demonstrated an association between CTO placement and facilitation of access to acute general medical care, which may partly explain how the mortality risk was reduced. Patients under CTO supervision, which facilitated access to acute medical care, were 20% less likely to experience a non-injury-related death (that is, exclusive of assault, self-harm, and undetermined intent).

Authors of some studies have concluded that CTOs are overused (9,12,17). Persons who want to restrict the use of CTOs have raised two philosophical concerns about the type of findings reported here. First, although death rates are elevated among psychiatric patients, these persons question whether the rates are elevated enough to justify involuntary supervision. During the study period, patients in the CTO cohort died at a rate 1.27 times that of the general population. Comparative rates in the United States indicate that among individuals with severe mental illness who die, SMRs range from 1.2 to 4.9 times higher than in the general population (29). Other studies have reported decreased death rates associated with CTO placement (15,16), and there also appears to be an association between reduced homicide rates and use of an outpatient commitment law with criteria similar to the law in Victoria (30). These results and the results of this study suggest that a possible reason for the lower comparative death rate for persons with severe mental illness in Victoria is the availability of the CTO mechanism.

The second concern raised by those who want to restrict CTO use is that CTOs are an assault on quality of life (17). However, our analysis of HoNOS quality-of-life ratings associated with years of life saved appeared to modestly favor the CTO cohort. Men in the CTO cohort scored significantly better than their non-CTO counterparts on five of the HoNOS dimensions, did not differ on five dimensions, and scored worse on two. Women in the CTO cohort also scored better than their non-CTO counterparts on five HoNOS dimensions, did not differ on four, and scored worse on three. It would seem that the years of life saved associated with CTO placement are not negatively affected by CTO placement.

The study had some limits. It was based on administrative data. Results did not reach the level of certainty of causal inference associated with randomized controlled trials. However, with the exception of one trial focused on victimization outcomes (31), no trials have been completed that have a criterion focused on the need for treatment to protect health and safety. In today's deinstitutionalized environment, the misplaced focus on reduction in hospitalization days may constitute a denial of needed treatment. Also, quality-of-life assessments were based on clinicians' perspectives, although patients' perspectives obtained via surveys evaluating the impact of outpatient commitment on quality of life have been mixed (32).

CONCLUSIONS

Patients with mental disorders severe enough to require hospitalization were found to be at increased risk of mortality compared with persons in the general population in Victoria, Australia. The results support the use of CTOs as an alternative to psychiatric hospitalization. They indicate that for patients who refuse needed treatment, involuntary care

delivered outside the hospital to protect health was associated with reduced mortality risk without a negative effect on quality of life.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

REFERENCES

1. Black DW, Warrack G, Winokur G: The Iowa Record-Linkage Study: II. excess mortality among patients with organic mental disorders. *Archives of General Psychiatry* 42:78–81, 1985 [PubMed: 3966855]
2. Black DW, Warrack G, Winokur G: The Iowa record-linkage study: III. excess mortality among patients with “functional” disorders. *Archives of General Psychiatry* 42:82–88, 1985 [PubMed: 3966856]
3. Corten P, Riboudouille M, Dramaix M: Premature death among outpatients at a community mental health center. *Psychiatric Services* 42:1248–1251, 1991
4. Felker B, Yazel JJ, Short D: Mortality and medical comorbidity among psychiatric patients: a review. *Psychiatric Services* 47: 1356–1363, 1996 [PubMed: 9117475]
5. Harris EC, Barraclough B: Excess mortality of mental disorder. *British Journal of Psychiatry* 173:11–53, 1998 [PubMed: 9850203]
6. Koranyi EK: Morbidity and rate of undiagnosed physical illnesses in a psychiatric clinic population. *Archives of General Psychiatry* 36:414–419, 1979 [PubMed: 426608]
7. Lawrence D, Jablensky AV, Holman CD, et al.: Mortality in Western Australian psychiatric patients. *Social Psychiatry and Psychiatric Epidemiology* 35:341–347, 2000 [PubMed: 11037302]
8. Parks J, Svendsen D, Singer P, et al.: Morbidity and Mortality in People With Serious Mental Illness. Alexandria, Va, National Association of State Mental Health Program Directors Medical Directors Council, 2006
9. Burns T, Rugkåsa J, Molodynski A, et al.: Community treatment orders for patients with psychosis (OCTET): a randomised controlled trial. *Lancet* 381:1627–1633, 2013 [PubMed: 23537605]
10. Civil commitment of the mentally ill: theories and procedures. *Harvard Law Review* 79:1288, 1966
11. Civil commitment of the mentally ill. *UCLA Law Review* 14:822, 1967
12. Brophy LM, Reece JE, McDermott F: A cluster analysis of people on Community Treatment Orders in Victoria, Australia. *International Journal of Law and Psychiatry* 29:469–481, 2006 [PubMed: 17084453]
13. Lacking Insight, Part 2: Introductory Material. Melbourne, Victoria, Mental Health Legal Centre, 2016
14. Segal SP, Hayes SL, Rimes L: The utility of outpatient commitment: I. a need for treatment and a least restrictive alternative to psychiatric hospitalization. *Psychiatric Services* (Epub ahead of print, August 1, 2017)
15. Segal SP, Burgess PM: Effect of conditional release from hospitalization on mortality risk. *Psychiatric Services* 57:1607–1613, 2006 [PubMed: 17085609]
16. Kisely S, Preston N, Xiao J, et al.: Reducing all-cause mortality among patients with psychiatric disorders: a population-based study. *Canadian Medical Association Journal*, 185:E50–E56, 2013 [PubMed: 23148054]
17. Light E: An Analysis of the Community Treatment Order System in New South Wales. Sydney, Australia, University of Sydney, School of Public Health, Centre for Values, Ethics and the Law in Medicine, 2015
18. Churchill R: International Experiences of Using Community Treatment Orders. London, Kings College London, Institute of Psychiatry, 2007
19. About Victoria Police: Crime Statistics. Melbourne, Victoria Police, 2015 http://www.police.vic.gov.au/content.asp?Document_ID=782

20. Socio-Economic Indexes for Areas (SEIFA) Commonwealth of Australia. Technical Paper Canberra, Australian Bureau of Statistics, 2011
21. Wing JK, Beevor AS, Curtis RH, et al.: Health of the Nation Outcome Scales (HoNOS): research and development. *British Journal of Psychiatry* 172:11–18, 1998 [PubMed: 9534825]
22. Eagar K, Buckingham B, Coombs T, et al.: Victorian Outcome Measurement Strategy Resource Manual: AMHOCN Health of the Nation Outcome Scales (HoNOS): Glossary. Parramata, New South Wales, Australian Mental Health Outcomes and Classification Network, 2000
23. IBM SPSS Statistics for Windows, Version 23.0. Armonk, NY, IBM, 2015
24. Microsoft Excel for Windows. Seattle, Microsoft, 2016
25. Selvin S: *Statistical Analysis of Epidemiologic Data*. Oxford, United Kingdom, Oxford University Press, 2004
26. Australian Historical Population Statistics, 2014. Canberra, Australian Bureau of Statistics, 2014
27. International Classification of Diseases (ICD). Geneva, World Health Organization, 2015
28. Smith GD, Hart C, Watt G, et al.: Individual social class, area-based deprivation, cardiovascular disease risk factors, and mortality: the Renfrew and Paisley Study. *Journal of Epidemiology and Community Health* 52:399–405, 1998 [PubMed: 9764262]
29. Parks J, Svendsen D, Singer P, et al.: *Morbidity and Mortality in People With Serious Mental Illness*. Alexandria, Va, National Association of State Mental Health Program Directors Medical Directors Council, 2006
30. Segal SP: Civil commitment law, mental health services, and US homicide rates. *Social Psychiatry and Psychiatric Epidemiology* 47: 1449–1458, 2012 [PubMed: 22072224]
31. Hiday VA, Swartz MS, Swanson JW, et al.: Impact of outpatient commitment on victimization of people with severe mental illness. *American Journal of Psychiatry* 159:1403–1411, 2002 [PubMed: 12153835]
32. Swartz M, Swanson J, Steadman H, et al.: *New York State Assisted Outpatient Treatment Program Evaluation*. Durham, NC, Duke University, 2009




FIGURE 1. Years of life lost (YLL) among psychiatric patients by whether they were placed on a community treatment order (CTO) in Victoria, Australia, from 2000 to 2010^a

^aYLL was based on the average life expectancy from birth in Victoria in 1971, the average birth year of patients in the sample. It was estimated separately for men and women.




FIGURE 2. Mean HoNOS quality-of-life scores during years of life saved, by whether patients were placed on a community treatment order (CTO) in Victoria, Australia, from 2000 to 2010^a
^a Problem areas on the Health of the Nation Outcome Scales (HoNOS) are rated on a scale from 0, no problem, to 4, extremely problematic. CTO placement appeared to lead to a gain of 3.8 years of life among men and 2.4 years among women. Scores were taken for the final 3.8 years of life for the 343 men in the CTO cohort and for the 344 men in the non-CTO cohort who died and for whom scores were available. Scores were taken for the final 2.4 years of life for the 207 women in the CTO cohort and for the 258 women in the non-CTO cohort who died and for whom scores were available.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Mortality during the 11.8-year study among psychiatric patients, by whether they were placed on a community treatment order (CTO) from 2000 to 2010 in Victoria, Australia

TABLE 1.

| Group | N | Deaths | Crude rate | Deaths per 1,000 | Crude RR of death for CTO cohort ^d | | | χ^2 ^{a,b} |
|---|--------|--------------------|------------|------------------|---|----------|------|-------------------------|
| | | | | | RR | 95% CI | OR | |
| Total sample | 27,565 | 2,727 | .099 | 98.93 | | | | |
| CTO cohort | 11,424 | 1,030 | .090 | 90.16 | .86 | .80-.92 | .85 | .78-.92 16.56 |
| Non-CTO cohort | 16,161 | 1,697 ^c | .105 | 105.10 | 1.00 | - | - | - |
| Characteristic of total sample | | | | | | | | |
| Men | 15,480 | 1,528 | .099 | 98.71 | .84 | .76-.92 | .82 | .74-.91 12.52 |
| Women | 12,103 | 1,199 | .099 | 99.07 | .38 | .15-.95 | .88 | .78-.99 4.26 |
| Schizophrenia | 18,260 | 1,241 | .068 | 67.96 | .98 | .88-1.09 | .98 | .88-1.03 .09 |
| Major affective disorder | 2,760 | 215 | .078 | 77.90 | 1.01 | .70-1.44 | 1.01 | .75-1.35 .00 |
| Paranoia or other psychosis | 2,157 | 200 | .093 | 92.72 | .92 | .36-2.36 | .91 | .67-1.25 .33 |
| Dementia or other nervous system disorder | 3,167 | 922 | .291 | 291.13 | .81 | .72-.91 | .07 | .63-.87 13.06 |
| Other psychiatric disorder ^d | 912 | 123 | .139 | 134.87 | 1.11 | .78-1.57 | 1.13 | .76-1.68 .35 |

^aThe reference group is the non-CTO cohort.

^bdf=1

^cThere were 1,697 deaths in the non-CTO cohort. There were two cases where information on the gender of decedent was not available. Given the need to adjust for gender in computing life expectancy and other statistics, the reported statistics other than the total deaths statistics are based on N=1,695.

^dOther affective and somatoform disorders, anxiety disorders, eating and obsessive-compulsive disorders, personality disorders, acute stress reactions, conduct disorders, sexual disorders, alcohol intoxication and withdrawal, drug intoxication and withdrawal, alcohol use disorder and dependence, other drug use disorder and dependence

TABLE 2. Logistic regression models of effects of placement on a community treatment order (CTO) in Victoria, Australia, from 2000 to 2010^a

| Variable | Model χ^2 ^b | B | SE | Exp(b) | 95% CI |
|--|-----------------------------|------|-----|--------|-----------|
| All deaths | 4,336.84 | | | | |
| CTO cohort | | -.09 | .06 | .91 | .79–.99 |
| Treatment days per community care episode | | .00 | .00 | 1.00 | 1.00–1.01 |
| Interaction CTO × treatment days per community care episode | | .00 | .00 | 1.00 | 1.00–1.01 |
| Deaths excluding assault, self-harm, and undetermined intent | 4,983.54 | | | | |
| CTO cohort | | -.18 | .07 | .83 | .75–.93 |
| Treatment days per community care episode | | .00 | .00 | .99 | .99–1.01 |
| Interaction CTO × treatment days per community care episode | | .00 | .00 | .99 | .99–1.01 |
| Deaths by external causes, assault, self-harm, and undetermined intent | 298.45 | | | | |
| CTO cohort | | .23 | .11 | 1.26 | 1.01–1.58 |
| Treatment days per community care episode | | .01 | .01 | .99 | .98–1.01 |
| Interaction CTO × treatment days per community care episode | | .01 | .01 | 1.01 | .99–1.02 |
| Deaths by self-harm | 298.45 | | | | |
| CTO cohort | | .28 | .12 | 1.32 | 1.04–1.69 |
| Treatment days per community care episode | | .01 | .01 | .99 | .98–1.01 |
| Interaction CTO × treatment days per community care episode | | .01 | .01 | 1.00 | .99–1.02 |
| Deaths by external causes, assault, and undetermined intent | 220.21 | | | | |
| CTO cohort | | -.04 | .13 | .96 | .74–1.24 |
| Treatment days per community care episode | | .00 | .00 | 1.00 | 1.00–1.01 |
| Interaction CTO × treatment days per community care episode | | .00 | .00 | 1.00 | 1.00–1.01 |

^a All models in addition to the primary explanatory variables regressed the following explanatory variables on the model criterion variable; propensity of a patient to be selected into the CTO sample from among hospitalized patients, age, gender, socioeconomic status or vocational challenge (that is, >11th grade education, unemployment, and the lowest Socio-Economic Indexes for Areas rank of a neighborhood in which the patient lived), risk period associated with study and institutional involvement (that is, age at first date known to the mental health system, total time known to the mental health system, and lifetime total inpatient days), diagnoses (major affective disorder, dementia, schizophrenia, and paranoia or other psychosis), imprisonment or police custody during the study, aboriginal or Torres Strait Islander status, language interpreter required at the mental health tribunal, and psychosocial profile reflected in the 12 Health of the Nation Outcome Scale scores at both inpatient admission and release [see online supplement].

^b N for regression=27,585; df=43 and 27,542. All models were significant (p<.001).