Lawrence Berkeley National Laboratory

Recent Work

Title

Cluster Cascading in the Geometrical Branching Model

Permalink

<https://escholarship.org/uc/item/8s47174z>

Journal

Physical Review D, 43(7)

Authors

Chen, W.R.

Hwa, R.C.

Wang, X.-N.

Publication Date

1990-09-01

Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

Submitted to Physical Review D

Cluster Cascading in the Geometrical Branching Model

W.R. Chen, R.C. Hwa, and X.-N. Wang

September 1990

LOAN COPY
Circulates
for 2 weeks

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Cluster Cascading in the Geometrical Branching Model[†]

Wei R. Chen

*Oklahoma School of Science and Mathematics
1515 North Lincoln Blvd., Oklahoma City, OK 73104 USA*

Rudolph C. Hwa

*Institute of Theoretical Science and Department of Physics
University of Oregon, Eugene, OR 97403 USA*

Xin-Nian Wang

*Nuclear Science Division, Mailstop 70A-3307
Lawrence Berkeley Laboratory
University of California, Berkeley, CA 94720 USA*

Abstract

The geometrical branching model is further developed in a direction that allows the branching part to be implemented by a Monte Carlo scheme of cascading cluster decays. A universal mass distribution is adopted for the determination of cluster masses. Good agreements with data on multiplicity distribution and average multiplicity are achieved.

[†]This work was supported by the Director, Office of Energy Research, Division of Nuclear Physics of the Office of High Energy and Nuclear Physics of the U.S. Department of Energy under Contract No. DE-AC03-76SF00098 and DE-FG06-85ER-4022

Multiparticle production at low p_T in high-energy hadronic collisions has been described by a large number of models, all of which can reproduce the gross features very well[1]. The geometrical branching model(GBM)[2] is one among them, which combines the geometrical properties of hadronic collisions[3],[4] with the stochastic properties of particle production[5]. After the suggestion by Białas and Peschanski[6] to investigate intermittency in multiplicity fluctuations in various scales of resolution, experimental data on normalized factorial moments have revealed power-law behaviors in leptonic, hadronic and nuclear processes[7]. Since random cascade processes can lead to intermittency, the GBM is well poised to account for the observed effect in small rapidity intervals. However, before a Monte Carlo code can be developed to demonstrate intermittency, there is one intermediate step that must be taken to render the GBM suitable for such considerations. That is, the branching process, which was summarized by the Furry distribution previously[2], must now be implemented by a specific scheme of successive branching. It is the aim of this paper to accomplish this limited objective.

Our basic input will be the massive cluster decays in cascading. This mechanism in itself is, of course, not new. Ochs and Wosiek[8] have already shown that it can lead to intermittency. Our concern at this point is not so much intermittency as the formulation of GBM in such a way that successive cluster decay can be amalgamated with the Glauber-Gribov approach[4],[9],[10] to high-energy collisions. Our immediate aim is to show that a sensible cluster cascading scheme can be used to reproduce the multiplicity distribution in the whole rapidity space. The subject of fluctuations in small rapidity intervals is deferred to a future investigation.

Let us first recall the eikonal description of σ_{in} , which is related by the Abramovski-Gribov-Kancheli(AGK) cutting rule[9] to the eikonal function $\Omega(b, s)$ for the elastic amplitude by[10]

$$\sigma_{in} = \int d^2b \sum_{\mu=1}^{\infty} \frac{1}{\mu!} [2\Omega(b, s)]^{\mu} e^{-2\Omega(b, s)}, \quad (1)$$

where the summand is the contribution from μ cut-Pomerons. To determine the multiplicity distribution P_n it is necessary to specify what a cut-Pomeron is, for which only model description have been given. For example, in the dual parton model(DPM)[11] which is formulated in the momentum space, the lowest order contribution to a cut-Pomeron consists of two chains stretched between quark-diquark pairs. In the GBM we have represented the multiplicity distribution of each cut-Pomeron by the Furry distribution[2] $F_n^{k_j}$, whose self-reproducing property gives rise to an effective $F_n^{k(\mu)}$ for a μ cut-Pomeron contribution, where $k(\mu)$ is a parameter denoting the number of initial branching clusters. Thus, if we use $\pi_\mu(b)$ to signify the summand in Eq. 1, we have[2]

$$P_n = \frac{1}{\sigma_{in}} \int d^2b \sum_{\mu=1}^{\infty} \pi_\mu(b) F_n^{k(\mu)}. \quad (2)$$

By adjusting $k(\mu)$ it has been possible to fit the data on P_n throughout the ISR energy range, and then with minijet contributions in the SPS collider range also.

As an initial attempt to combine geometrical and stochastic properties of multiparticle production, it was sensible to let the Furry distribution to represent approximately the effects of branching so that we could judge quickly whether we were aiming in the right direction. Now, we begin a second-level consideration in which the branching process is to be treated explicitly in a Monte Carlo simulation with energy-momentum conservation applied at each step. There are two issues to be addressed. First, how many initial clusters are there and what are their masses? Second, how does a cluster decay? Since these questions are central to our improved treatment, let us discuss them at some length separately.

Since the average number of cut-Pomerons at each impact parameter b is specified by the probability $\pi_\mu(b)$, we know that μ can vary on the average from 1 to a large value at small b . If a massive cluster is associated with each cut Pomeron,

then impact-parameter smearing results in an effective distribution of initial clusters for each collision. This picture is different from, and perhaps complementary to, the two currently fashionable models: DPM[11] and Fritiof model[12], which have no explicit impact-parameter-smearing, but either have two chains with varying lengths or have two excited strings with varying effective masses. In our model we use $\pi_\mu(b)$ and stochasticity to determine the number of initial clusters. More specifically, our procedure is to first convert Eq. 1 into an integral over the scaled impact parameter $R = b/b_0$, where $b_0 = (\sigma_{in}/\pi)^{1/2}$, as in Ref.[2]. Then we rewrite Eq. 2 in the form

$$P_n = \int_0^\infty dR^2 \sum_{\mu=1}^\infty \pi_\mu(R) B_n^\mu, \quad (3)$$

$$\pi_\mu(R) = \frac{1}{\mu!} [2\Omega(R)]^\mu e^{-2\Omega(R)}, \quad (4)$$

where B_n^μ is a new branching distribution that we now attempt to determine by Monte Carlo calculation. For every value of R , we generate a value for μ according to the probability $\pi_\mu(R)$. We then have μ initial clusters, each having c.m. energy E_i , whose value is specified by a randomly chosen variable x_i according to $E_i = x_i\sqrt{s}$, satisfying $0 < x_i < 1$ and $\sum_{i=1}^\mu x_i = 1$. These clusters undergo successive binary decays in a manner to be described below, and the distribution B_n^μ can then be calculated at the end of the branching process. Note that this procedure automatically gives rise to diffractive-like processes at large R , since $\mu = 1$ is the only important contribution to Eq. 3 in this case, and with $x_1 = 1$ the one cut-Pomeron gives rise to a two-cluster fragmentation process. At smaller R higher μ values also become important, so the sharing of $\sqrt{s}$ among the μ initial clusters forces the decay particles to populate the central region, resulting in a central plateau for the single-particle inclusive distribution. Momentum conservation is imposed in each cut-Pomeron, when the clusters undergo cascading.

The second issue of major importance is the nature of cluster decay. In a soft hadronic process no high virtuality is associated with any constituents. The valence

quarks that carry nearly half the incident momentum do not interact strongly with soft partons because of the large separation in rapidity; they hadronize into the leading particles in the fragmentation region. A description of the hadronization in the central region that does not give any preeminent role to any individual partons should be to consider the interacting partons collectively in terms of s-channel clusters that evolve in a self-similar way. This point of view has received some support from the recent interest in intermittency, which is essentially the phenomenology of self-similarity. As mentioned in the previous paragraph, the leading cluster and the centrally produced ones are all included in our approach, since for every R , μ is summed over all values, the $\mu = 1$ term contributing principally to the fragmentation region.

Since we want to describe the cluster decay by a self-similar process, we recall the statistical boost-trap model of Hagedorn[13], who has for many years emphasized the picture of fireball within fireballs in a self-consistent description. Indeed, it was Hagedorn who first pointed out that the mass spectrum of hadrons is an exponentially growing one: e^{m/T_0} , where T_0 is the Hagedorn temperature[14]. The distribution of cluster masses at temperature T is

$$\rho(m) \sim m^\alpha e^{(\frac{1}{T_0} - \frac{1}{T})m}, \quad (5)$$

where $T < T_0$ and we shall consider T to be a constant in the hadronic interaction. We adopt the general ideas of Hagedorn as hints on our cluster decays and use

$$\rho(m) \sim m^\alpha e^{-\beta m}, \quad (6)$$

as the universal mass distribution of clusters produced at all stages of the successive decays. In Eq. 6, α and β are parameters in our model to be adjusted to fit the

data.

Specifically, our procedure is the following. For each set of values of μ and E_i , $i = 1, \dots, \mu$ generated in our Monte Carlo simulation, we let the initial cluster in the i th cut Pomeron decay into two clusters whose masses m_1 and m_2 are determined randomly according to Eq. 6. Energy-momentum conservation then fixes their longitudinal momenta, their transverse momenta being neglected in this investigation. We then go to the rest frames of the daughter clusters and repeat the decay procedure, always using Eq. 6 to determine the subsequent cluster masses. The decay sequence is terminated when the mass reaches a value below $2m_\pi$. In this way we determine B_n^μ in Eq. 3. Because we have not considered the charge and flavors of the produced hadrons, we take all the final particles to be pions and regard the number of charged pions as $2/3$ of the total produced particles.

We have found that by choosing $\alpha = 0.1$ and $\beta = 0.01 \text{ GeV}^{-1}$ we obtain approximately Koba-Nielsen-Olesen(KNO) scaling for pp collision in the ISR region. The resultant multiplicity distribution agrees well with the data[15], as shown in Fig.1. The normalized moments $C_q = \langle n^q \rangle / \langle n \rangle^q$ are shown in Fig.2 as functions of $\sqrt{s}$. The average multiplicity $\langle n \rangle$, shown in Fig.3, is also in good agreement with data[15]. At higher energies minijet production must also be taken into consideration, so P_n is expected to get broader[2].

We have not fine-tuned the parameters to yield the best fit because our present calculation ignores the transverse momenta of clusters. We are encouraged by our results, since they indicate that the GBM, when implemented by cascade cluster decays in one dimension, can well describe the global data on P_n and $\langle n \rangle$. This is an improvement on our previous results, since we could not calculate $\langle n \rangle$ before. More importantly, we now have a realistic scheme of branching, which will facilitate further development into a comprehensive code. For that we need to improve on our consideration of the transverse degrees of freedom, the production of resonances, the

inclusion of charges and flavors, etc. After those improvements are made, we shall then be able to calculate more reliably the rapidity distribution and the associated quantities related to intermittency. The fact that this model treats the geometrical features of hadronic collisions properly and contains a self-similar scheme of particle production makes us feel quite hopeful that this approach will render a successful description of the soft interaction of hadrons.

References

- [1] Most of the models are described in review articles in *Hadronic Multiparticle Production*, edited by P. Carruthers(World Scientific, Singapore, 1988).
- [2] W. R. Chen and R. C. Hwa, Phys. Rev. **D36**, 760(1987); R. C. Hwa, Phys. Rev. **D37**, 1830(1988); X. N. Wang and R. C. Hwa, Phys. Rev. **D39**, 187(1989); W. R. Chen and R. C. Hwa, **D38**, 3394(1988); W. R. Chen, R. C. Hwa and X. N. Wang, Phys. Rev. **D38**, 3394(1988); X. N. Wang, Report No. LBL-28790, submitted to Phys. Rev. D.
- [3] T. T. Chou and C. N. Yang, Phys. Rev. Lett. **20**, 1213(1968); Phys. Rev. **170**, 1591(1968); L. Durand and R. Lipes, Phys. Rev. Lett. **20**, 637(1968).
- [4] R. J. Glauber, in *Lectures in Theoretical Physics*, edited by W. E. Brittin and L. G. Danham(Interscience, New York, 1959), Vol.I; R. J. Glauber and G. Mathiae, Nucl. Phys. **B21**, 135(1970).
- [5] See R. C. Hwa in Ref. [1].
- [6] A. Bialas and R. Peschanski, Nucl. Phys. **B273**, 703(1986).
- [7] For reviews see experimental papers in *Multiparticle Dynamics*, edited by A. Giovannini and W. Kittel(World Scientific, Singapore, 1990); B. Buschbeck and P. Lipa, Mod. Phys. Lett. **A4**, 1871(1989).
- [8] W. Ochs and J. Wosiek, Phys. Lett. **B214**, 617(1988).
- [9] V. A. Abramovski, V. V. Gribov and O. V. Kancheli, Yad. Fiz. **18**, 595(1973)[Sov. J. Nucl. Phys., **18**, 308(1974)].
- [10] V. Anisovich, M. N. Kobrinsky, J.Nyiri and Yu. M. Shabelski, *Quark Model and High Energy Collisions*(World Scientific, Singapore, 1985); V. M. Braun and Yu. M. Shabelski, Int. J. Mod. Phys. **A3**, 2417(1988).

- [11] A. Capella and J. Tran Thanh Van, Z. Phys. **C10**, 249(1981); see also A. Capella *et al.* in Ref. [1].
- [12] B. Andersson, G. Gustafson and B. Nilsson-Almqvist, Nucl. Phys. **B281**, 289(1987).
- [13] R. Hagedorn, Suppl. Nuovo Cimento **3**, 147(1965); Rivista Nuovo Cimento **6**, 1 (1983).
- [14] R. Hagedorn, Astron. and Astrophys. **5**, 184(1970).
- [15] G. J. Alner *et al.*, Phys. Lett. **B160**, 193(1985); **B167**, 476(1986).

Figure Captions

Fig. 1 KNO plot of the calculated result(in histogram) as compared to the data, taken from Ref. [15].

Fig. 2 Normalized moments of multiplicity distributions at various energies. The data are taken from Ref. [15].

Fig. 3 Comparision of calculated average multiplicity(solid line) with the data, taken from Ref. [15].

Figure 1

Figure 2

Figure 3

LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
INFORMATION RESOURCES DEPARTMENT
BERKELEY, CALIFORNIA 94720