

UCLA

UCLA Previously Published Works

Title

Anticholinergic burden in older psychiatric inpatients

Permalink

<https://escholarship.org/uc/item/8sc2j462>

Journal

Australasian Psychiatry, 26(4)

ISSN

1039-8562

Authors

Woo, Benjamin Kp

Agustines, Davin

Publication Date

2018-08-01

DOI

10.1177/1039856218765869

Copyright Information

This work is made available under the terms of a Creative Commons Attribution-NonCommercial-NoDerivatives License, available at

<https://creativecommons.org/licenses/by-nc-nd/4.0/>

Peer reviewed

Australasian Psychiatry
2018, Vol 26(4) 438–439
© The Royal Australian and
New Zealand College of Psychiatrists 2018
Reprints and permissions:
sagepub.co.uk/journalsPermissions.nav
journals.sagepub.com/home/apy



Anticholinergic burden in older psychiatric inpatients

Dear Sir

Lee et al.¹ reported a study to explore the prescribing patterns in a geriatric psychiatric inpatient unit and found that anticholinergic burden of medication was rarely considered. Also, from admission to discharge, the proportion of patients on any anticholinergic medication increased. Furthermore, only 10% of patients were free of anticholinergic medication upon discharge. These findings highlighted some points with implications for clinical practice.

First, older patients are more vulnerable to anticholinergic burden. Quality of care has several dimensions, including safety, effectiveness, and efficiency.² As the authors pointed out, 'there were no evidence of anticholinergic burden review in the form of electronic documentation,' future studies should evaluate the utility of incorporating an anticholinergic medication scale as part of the electronic health record. Reviewing such red flags may improve the quality of care and prevent iatrogenic complications amongst older inpatients.

Second, Lee et al.¹ found that the proportion of patients on at least one anticholinergic medication increased from admission to discharge, and suggested that clinicians should be vigilant when prescribing anticholinergic medications to older psychiatric patients. Past studies additionally pointed out that multidisciplinary teams utilizing interdisciplinary geriatric psychiatric and medicine care teams may decrease overuse of inappropriate medications in the elderly, including

anticholinergic medications.^{3–5} As such, quality care requires a multidisciplinary approach for older patients.

Third, polypharmacy is recognized as a major contributor to anticholinergic burden, and is associated with functional decline.⁴ The current study demonstrated that a majority of older patients at discharge suffered from definitive anticholinergic burden.¹ Since older psychiatric patients are frequently treated with anticholinergic medications, clinicians should monitor the anticholinergic burden carefully as a potential risk factor for decline in daily functioning.

Anticholinergic burden may cause psychiatric exacerbations and decline in functioning. The present study provided insight into the frequency and nature of anticholinergic burden in hospitalized older mental health inpatients. Clinicians should respond to the question raised by the authors, 'do we care [about anticholinergic burden],' with a resounding yes. Since older psychiatric inpatients have increased sensitivity to anticholinergic burden, clinicians should further explore the use of technology and interdisciplinary teams to improve the quality of care provided to this vulnerable population.

ORCID iD

Benjamin K.P. Woo  <https://orcid.org/0000-0002-0127-4850>

References

1. Lee MS, Kisely S, Zolotarev B, et al. Anticholinergic burden in older inpatients on psychotropic medication: do we care? *Australas Psychiatry* 2017; 25 :566–570.
2. Woo BK and Sultzer DL. A key step for quality care in psychiatric emergency settings. *Gen Hosp Psychiatry* 2009; 31: 503–504.
3. Walsh KA, O'Riordan D, Kearney PM, et al. Improving the appropriateness of prescribing in older patients: a systematic review and metaanalysis of pharmacists' interventions in secondary care. *Age Ageing* 2016; 45: 201–209.
4. Chan VT, Woo BK, Sewell DD, et al. Reduction of sub-optimal prescribing and clinical outcome for dementia patients in a senior behavioral health inpatient unit. *Int Psychogeriatr* 2009; 21: 195–199.
5. Woo BK, Daly JW, Allen EC, et al. Unrecognized medical disorders in older psychiatric inpatients in a senior behavioral health unit in a university hospital. *J Geriatr Psychiatry Neurol* 2003; 16: 121–125.

Benjamin KP Woo, Davin Agustines
Los Angeles, CA, USA

DOI: 10.1177/1039856218765869

Enabling interagency collaboration in providing comprehensive mental health care: the rise of the care coordinator

Dear Sir

Individuals with severe and persistent mental illness (SPMI) have multiple needs such as support to access general medical services, housing, community and domiciliary services, and employment services.¹ Recovery can be achieved only when all or most of these needs are met. Case managers tend not to have the time to support the non-clinical needs of their clients.²

Multiple needs of individuals with SPMI can be addressed by interagency collaboration, which has shown to have better outcomes for clients and services.¹ However, barriers such as competition for funding, conflicts related to areas of responsibility and integration strategies make collaborative efforts quite challenging.¹ As a result, individuals