

# UCSF

## UC San Francisco Previously Published Works

### Title

An adapted European LeukemiaNet genetic risk stratification for acute myeloid leukemia patients undergoing allogeneic hematopoietic cell transplant. A CIBMTR analysis

### Permalink

<https://escholarship.org/uc/item/8sg9d0v3>

### Journal

Bone Marrow Transplantation, 56(12)

### ISSN

0268-3369

### Authors

Jimenez Jimenez, Antonio M  
De Lima, Marcos  
Komanduri, Krishna V  
[et al.](#)

### Publication Date

2021-12-01

### DOI

10.1038/s41409-021-01450-3

Peer reviewed



Published in final edited form as:

*Bone Marrow Transplant.* 2021 December ; 56(12): 3068–3077. doi:10.1038/s41409-021-01450-3.

## An Adapted European LeukemiaNet Genetic Risk Stratification for Acute Myeloid Leukemia Patients Undergoing Allogeneic Hematopoietic Cell Transplant. A CIBMTR Analysis

A full list of authors and affiliations appears at the end of the article.

### Abstract

Cytogenetic and molecular abnormalities are known to influence post-transplant outcomes in acute myeloid leukemia (AML) but data assessing the prognostic value of combined genetic models in the HCT setting are limited. We developed an adapted European LeukemiaNet (aELN) risk classification based on available genetic data reported to the Center for International Blood and Marrow Transplant Research, to predict post-transplant outcomes in 2289 adult AML patients transplanted in first remission, between 2013 and 2017. Patients were stratified according to aELN into three groups: favorable (Fav, N=181), intermediate (IM, N=1185) and adverse (Adv, N=923). Univariate analysis demonstrated significant differences in 2-year overall survival (OS) (Fav: 67.7%, IM: 64.9% and Adv: 53.9%;  $p<0.001$ ); disease-free survival (DFS) (Fav: 57.8%, IM: 55.5% and Adv: 45.3;  $p<0.001$ ) and relapse (Fav: 28%, IM: 27.5% and Adv: 37.5%;  $p<0.001$ ). Multivariate analysis (MVA) revealed no differences in outcomes between the Fav and IM groups, thus they were combined. On MVA, patients in the Adv risk group had the highest risk of relapse (HR 1.47  $p<0.001$ ) and inferior DFS (HR 1.35  $p<0.001$ ) and OS (HR 1.39  $p<0.001$ ), even using myeloablative conditioning or in those without pre-HCT measurable-residual disease. Novel approaches to mitigate relapse in this high-risk group are urgently needed.

### INTRODUCTION

Allogeneic hematopoietic cell transplantation (HCT) is an effective post-remission strategy for patients with acute myeloid leukemia (AML) in first complete remission (CR1). Due to considerable morbidity and mortality risk, careful consideration of patient (1) and disease-specific (2, 3) characteristics is required. Traditionally, biological stratification of AML risk has been based on cytogenetic findings at diagnosis (3, 4). HCT in CR1 is considered for eligible patients with intermediate and high-risk cytogenetics whereas chemotherapy is recommended for those with favorable cytogenetics (2, 5, 6). Approximately 45% of AML cases have normal karyotype and cannot be subdivided based on cytogenetics; these patients have been assigned intermediate risk (4). Subsequent identification of genomic mutations in AML, such as *FLT3-ITD*, *NPM1*, *CEBPA*, *MLL/KMT2A*, *IDH1-2*, *DNMT3A*, *TET2*, *TP53* and *BCOR* amongst others, has proven valuable for subdividing cytogenetically normal (CN) AML into subsets with different outcomes (7).

**CORRESPONDING AUTHOR:** Antonio M. Jimenez Jimenez, MD, Assistant Professor of Medicine, Division of Transplantation and Cell Therapy, University of Miami, Miller School of Medicine, 1475 NW 12. Av, Miami, FL 33136, Tel./Fax: 305-243-1000/305-243-1223, amjimenez@med.miami.edu.

In 2010, the European LeukemiaNet (ELN2010) proposed a standardized prognostic system incorporating both cytogenetic and select molecular abnormalities (*CEBPA*, *NPM1* and *FLT3-ITD*) to distinguish 4 distinct genetic risk groups (8). Multiple studies confirmed the prognostic accuracy of ELN2010, applying it to patient cohorts receiving consolidation chemotherapy (9, 10) or HCT (11, 12). In 2017, the ELN updated and simplified their classification (ELN2017) into 3-groups of favorable, intermediate, and adverse risk (13). The new classification incorporated the addition of *RUNX1*, *ASXL1* and *TP53* mutations to the adverse risk group, the inclusion of biallelic (but not monoallelic) *CEBPA* mutations to the favorable group and stratification of *FLT3-ITD* on the basis of ITD-to-wild-type allelic ratio. Several groups have since applied ELN2017 to predict outcomes after chemotherapy (14–17) as well as compared the outcomes between ELN2010 and ELN2017 (14). Single institution studies have also evaluated the utility of ELN2017 in predicting post- HCT outcomes (18, 19), but large, multicenter studies evaluating the impact of a combined genetic model exclusively in the transplant setting are lacking.

We analyzed data from the Center for International Blood and Marrow Transplant Research (CIBMTR) to evaluate the prognostic ability of an adapted ELN2017 model to predict allogeneic HCT outcomes.

## MATERIALS AND METHODS

### Data Sources

Data was obtained from the CIBMTR which includes a voluntary network of over 500 transplantation centers worldwide that contribute detailed HCT data to a statistical center at the Medical College of Wisconsin in Milwaukee and the National Marrow Donor Program (NMDP®) Coordinating Center in Minneapolis. Participating centers are required to report all transplants consecutively; patients are followed longitudinally and compliance is monitored by on-site audits. Observational studies conducted by the CIBMTR are performed in compliance with all applicable federal regulations pertaining to the protection of human research subjects and approved by the National Marrow Donor Program® (NMDP)/Be The Match® Institutional Review Board. Protected Health Information used in the performance of such research is collected and maintained in CIBMTR's capacity as a Public Health Authority under the Health Insurance Portability Accountability Privacy Rule.

### Patient Selection

An initial cohort of 4777 adult patients (< 18 years) with a diagnosis of AML receiving their first allogeneic HCT between 2013 and 2017 and reported to the CIBMTR were included. Patients receiving a graft from a human leukocyte antigen (HLA)-matched sibling, fully matched unrelated donor, umbilical cord blood (UCB) graft and haploidentical donors were eligible for study. Patients with active disease (i.e., > 5% bone marrow blasts prior to transplantation or those with evidence of extra medullary disease), remission status beyond CR1, a diagnosis of acute promyelocytic leukemia and recipients of syngeneic or mismatched unrelated donor grafts were excluded.

Cytogenetic and molecular abnormalities present at the time of diagnosis or prior to initiation of the conditioning regimen were reported to the CIBMTR. When required, additional review of reported genetic data was performed by three reviewers (AJ, DW and KC) to adjudicate any uncertainties in classification. Cases with incomplete genetic data (N=152) were excluded. Additionally, subjects with incomplete research consent forms and those from embargoed centers were also excluded. Ultimately, a cohort of 2289 transplant recipients from 163 centers was analyzed.

### Cytogenetic and Molecular Classification

Patients were assessed for the presence of specific chromosomal and molecular abnormalities at diagnosis and prior to HCT. Cytogenetic data reported to the CIBMTR conformed to the International System of Cytogenetic Nomenclature (20). The definition of complex karyotype (≥ 3 abnormalities) and monosomal karyotype were made according to previously published criteria. (21, 22).

Comprehensive molecular information (i.e., *FLT3*-ITD, *NPM1*, *CEBPA*, *ASXL1*, *RUNX1*, *TP53*, *DNMT3A*, *BCR-ABL1*, *IDH1* and *IDH2* mutational status obtained by next-generation sequencing [NGS]) or PCR was reported to the CIBMTR beginning in 2013. Information regarding *FLT3*-ITD allelic ratio and *CEBPA* allelic status is not regularly reported to the CIBMTR, and was therefore not included in our classification. On the basis of available registry molecular and conventional cytogenetic data, we defined an adapted ELN genetic risk stratification (aELN) (Table 1) whereby patients were stratified into three distinct groups: favorable (Fav, N=181), intermediate (IM, N=1185) and adverse (Adv, N=923).

The aELN classified patients with mutated *NPM1* without *FLT3*-ITD and those with mutated *CEBPA* (without clarity of bi/mono allelic status) as favorable. Patients with mutated *NPM1* with *FLT3*-ITD (irrespective of allelic ratio) and those with wild type *NPM1* without *FLT3*-ITD were classified as intermediate. Patients with *FLT3*-ITD (irrespective of allelic ratio) and wild-type *NPM1* were classified as adverse. Other chromosomal and molecular abnormalities were classified according to the 2017 ELN genetic risk criteria.

### Study Endpoints and Variables

Disease-free survival (DFS, defined as time to relapse or death from any cause, with surviving patients in CR censored at last follow-up) was the study's primary endpoint. Secondary endpoints were: overall survival (OS, defined as time to death from any cause with surviving patients censored at last follow-up), non-relapse mortality (NRM, defined as death without preceding disease relapse/progression, relapse being a competing event), relapse (any reported events of leukemia relapse with NRM as competing event), acute and chronic graft vs. host disease (GVHD, with death as competing risk (23)). Patients were censored at subsequent HCT or last follow-up alive.

The study main effect was the influence of the aELN genetic risk group. Additional variables considered in the multivariate analysis (MVA) were: age at transplant, gender, race, Karnofsky performance score (KPS), HCT comorbidity index (HCT-CI), clinical onset of AML (*de novo* vs. transformed vs. therapy-related), time to achieve first remission,

measurable residual disease (MRD) status at transplantation, conditioning intensity, graft source, donor type and in vivo T-cell depletion (anti-thymocyte globulin [ATG] or alemtuzumab). Given intrinsic biologic and treatment-related differences between younger (i.e., <60 years) and older (i.e., ≥60 years) patients and previous reports demonstrating age-related differences in the distribution and impact of specific genetic abnormalities (9, 24), post-transplant outcomes were also analyzed separately between differing age cohorts. We also identified the most commonly occurring abnormalities (i.e., reported in at least >100 patients) within the adverse aELN genetic group, and evaluated specific outcomes for these larger individual subsets.

### Statistical Analysis

Univariate analysis was performed using the Kaplan-Meier Method and compared using the log-rank test for OS and DFS, while acute/chronic GVHD, NRM, and relapse used the cumulative incidence method considering competing risks, with comparisons performed using Gray's competing hazard method (25). MVA was performed using the Cox proportional hazard model (26) for OS and DFS. The main effect of the aELN risk group was retained in all models. The assumption of proportional hazards for each factor in the Cox model was tested by adding time-dependent covariates as necessary. When the test indicated differential effects over time (non-proportional hazards), models were constructed breaking the post-transplant time course into two periods, using the maximized partial likelihood method to find the most appropriate breakpoint. A backward stepwise model selection approach was used to identify significant risk factors. Factors which were significant at a 5% level were kept in the final model. Potential interaction between the main effect and significant co-variables was also tested for all endpoints. Adjusted probabilities of DFS and OS and adjusted cumulative incidence functions of NRM and relapse were calculated using the MVA models, stratified on main effect and weighted by the pooled sample proportion value for each prognostic factor. These adjusted probabilities estimate likelihood of outcomes in populations with similar prognostic factors. All analyses were done using the statistical package SAS version 9.4 (Cary, NC).

## RESULTS

### Demographics

Baseline clinical features of all 2289 patients are presented in Table 2. The median age at HCT was 57.2 years (IQR 19.2–74.2 years). Importantly, 41% (936 patients) were ≥60 years, 76% (n=1743) had *de novo* AML and 49% (1115) received a myeloablative conditioning (MAC) regimen. More patients in the Adv group (53% vs. 41.8% for Fav/IM patients,  $p<0.001$ ) had high HCT-CI. The frequency of secondary AML or therapy-related disease differed amongst groups (Fav 18.8%, IM 22.2% and Adv 26.9%;  $p=0.02$ ). Pre-transplant measurable residual disease (MRD) was also significantly different between groups (Fav 21%, IM 11.5% and Adv 22.9%;  $p<0.001$ ), however morphologic complete remission with incomplete blood count recovery (CRi) status was comparable for all cohorts ( $p=0.29$ ).

Of 923 patients with Adv risk, 380 (41.1%) had  $-5/\text{del}5q$ , or monosomy 7; 158 (17.1%) had wild-type *NPM1* with mutated *FLT3-ITD*; 139 (15.06%) carried a  $t(v;11q23.3)$ ; *KMT2A* rearrangement and 137 patients (14.84%) had a complex karyotype (Table 3). Complex karyotype included all other Adv-risk patients (without  $\text{del}5/\text{del}7$ , wt*NPM1* with *FLT3-ITD* or  $t(v;11q23.3)$  *KMT2A* rearrangement) having multiple ( $\geq 3$ ) chromosomal abnormalities.

**Transplant outcomes by aELN classification**—Overall, 1237 of 2289 patients (54%) were alive at last follow up with a median survival of 37 months (IQR, 24–51 months) and 1031 (83% of the survivors; 45% of the whole group) were alive and disease-free. Median follow up for survivors was 35 months.

### Disease Relapse

Cumulative incidence of relapse at two-years was 28% (95% confidence interval [CI], 21.5–35.1%), 27.5% (24.9–30.1%) and 37.5% (34.3–40.7%) for Fav, IM and Adv cohorts, respectively ( $p < 0.001$ ) (Table 4, Figure 1C).

MVA confirmed that Adv risk patients had higher relapse rates (HR for Adv vs. Fav/IM: 1.47, 95% CI 1.28–1.70,  $p < 0.0001$ ) with no differences observed between the Fav and IM groups; thus they were combined for subsequent analysis of other endpoints. Variables of interest considered in the MVA are listed in Supplemental Table S1.

Important variables independently associated with higher rates of relapse included: pre-transplant MRD (29.2% for MRD-negative [MRD<sup>neg</sup>] vs. 42% for MRD-positive [MRD<sup>pos</sup>]; HR: 1.47, 95% CI 1.23–1.76,  $p < 0.0001$ ); AML onset (29.9% for *de novo* vs. 40.1% for secondary and therapy-related disease; HR: 1.43, 95% CI 1.19–1.72,  $p = 0.0001$ ) and conditioning intensity (26.2% for MAC vs. 36.9% for non-myeloablative/reduced intensity regimens [NMA/RIC]; HR: 1.58, 95% CI 1.37–1.83,  $p < 0.0001$ , Supplemental Table S2).

While patients receiving RIC/NMA conditioning experienced higher rates of relapse, the impact of aELN Adv risk on relapse was observed for patients receiving either MAC or RIC/NMA ( $p < 0.001$  for both).

### Overall and Disease-Free Survival

There were significant differences in 2-year OS (Fav: 67.7% [95% CI, 60.5–74.6%], IM: 64.9% [62.1–67.6%] and Adv: 53.9% [50.6–57.2%];  $p < 0.001$ ) and DFS (Fav: 57.8% [95% CI, 50.3–65.2%], IM: 55.5% [52.6–58.4%] and Adv: 45.3% [42–48.6%];  $p < 0.001$ ) among aELN groups. (Table 4, Figure 1A, 1B). As above, Fav/IM were combined for further analysis.

The presence of pre-transplant MRD was associated with inferior two-year OS (63.7% for MRD<sup>neg</sup> vs. 46.6% for MRD<sup>pos</sup>,  $p < 0.001$ ) and DFS (54.4% for MRD<sup>neg</sup> vs. 37.4% for MRD<sup>pos</sup>,  $p < 0.001$ ). MAC led to superior two-year DFS (57.2% vs. 46% for NMA/RIC regimens,  $p < 0.001$ ) and OS (66% vs. 55.6% for MAC and NMA/RIC regimens,  $p < 0.001$ ).

Adv aELN was shown on MVA to be higher risk for both OS (HR for Adv vs. Fav/IM: 1.39, 95% CI 1.22–1.57,  $p < 0.001$ ) and DFS (HR for Adv vs. Fav/IM: 1.35, 95% CI 1.20–

1.51,  $p < 0.001$ , Table 5). Other variables independently associated with both OS and DFS included: age at HCT, pre-transplant MRD, donor type, clinical onset of AML and WBC at diagnosis ( $p = 0.01$  for all, Supplemental Table S2).

### Non-relapse Mortality and Graft-versus-Host Disease

There were no significant differences in NRM among aELN groups (two-year NRM: Fav: 14.2% [95% CI, 9.3–19.8%], IM: 17.1% [95% CI, 14.9–19.2%] and Adv: 17.3% [95% CI, 14.9–19.8%];  $p = 0.46$ ). Similarly, aELN classification had no impact on the incidence of grade 2–4 acute GVHD (Fav: 30.9% [95% CI, 24.4–37.9%], IM: 35.4% [32.7–38.2%] and Adv: 35% [32–38.2%];  $p = 0.42$ ) or chronic GVHD (Fav: 40% [32.7–47.4%], IM: 43.6% [40.7–46.6%] and Adv: 40.2% [37–43.5%];  $p = 0.44$ ).

### Comparison of outcomes by aELN risk group and age

There were no significant differences in relapse rates among age groups (HR for Fav/IM  $\geq 60$  vs.  $< 60$ : 1.19 [0.98–1.44]  $p = 0.08$ ; HR for Adv  $\geq 60$  vs.  $< 60$ : 1.17 [0.95–1.44]  $p = 0.15$ ), but MVA confirmed higher NRM for older patients ( $\geq 60$  y/o) in both Fav/IM (HR 1.40 [1.10–1.78]  $p = 0.007$ ) and Adv risk cohorts (HR 1.59 [1.18–2.13]  $p = 0.002$ ). Clinical outcomes were separately analyzed for two cohorts of younger patients (i.e., 18–39 vs. 40–59 years). MVA demonstrated inferior OS (HR 1.44 [1.18–1.75]  $p = 0.0003$ ) and DFS (HR 1.23 [1.03–1.46]  $p = 0.02$ ) for the 40–59 group. No significant differences in NRM or relapse were identified between these two younger cohorts. (Supplemental Table S2)

### Outcomes by Genetic subsets within the aELN Adv Risk Cohort

Genetic subset comparisons within the Adv-risk group showed that patients harboring monosomy 5, del(5q) or monosomy 7 (abnormal 5/7 subgroup) had inferior 2-year OS (42.6%,  $p < 0.001$ ) and DFS (35.2%,  $p < 0.001$ ), as well as higher rates of relapse (45%,  $p = 0.002$ ) when compared to other patients within the Adv risk cohort (OS 60%, DFS 50.8%, relapse 33.5%).

Patients in the abnormal 5/7 subgroup were older (median age 59.8 [IQR 18.92–75.84] vs. 51.9 [IQR 18.9–73.7] for other Adv-risk patients,  $p < 0.001$ ) and less often received MAC (45% MAC vs. 56.4% MAC for other Adv-risk patients  $p < 0.001$ ). Patients in the abnormal 5/7 subgroup also had a higher prevalence of pre-transplant MRD (28.4% vs. 19%,  $p = 0.003$ ) when compared to other Adv risk patients ( $p < 0.001$ ). MVA confirmed that abnormal 5/7 patients had higher relapse rates and inferior DFS and OS compared to other Adv risk patients ( $p < 0.0001$  for all, Table 3B).

## DISCUSSION

This large registry analysis of patients with AML undergoing allogeneic HCT demonstrates robust prognostic stratification by aELN in regard to relapse, DFS and OS. Inferior survival in the Adv-risk cohort appears to be driven by higher relapse rates and ensuing inferior DFS. There were no significant interactions between the main effect of aELN risk group and any significant risk factors. This confirms that the aELN classification is applicable to all patients receiving HCT in CR1, both younger ( $< 60$  years) and older ( $\geq 60$  years) and for

patients with *de novo* and transformed/secondary AML. Additionally, the impact of adverse genetics is not overcome by greater conditioning intensity.

Previous reports examining the prognostic ability of ELN2017 as compared to ELN2010 support the new combined genetic prognostic model for treatment of newly diagnosed AML patients (14–17) and a recent single institution analysis also evaluated the prognostic impact of ELN2017 following HCT (18). Some of these studies are limited by patient heterogeneity, small sample sizes or different consolidation strategies. We assessed the impact of a combined genetic model exclusively in the transplant setting and restricted our analysis to patients receiving allogeneic HCT in CR1. While the effect of genetic abnormalities continues to be important beyond CR1, prior relapse, particularly early relapse, carries a heavier prognostic weight than most genetic abnormalities (27).

Fav and IM risk groups were combined in our study after pairwise multivariable analysis demonstrated no significant differences in any outcomes. This could, in part, be explained by a smaller number of Fav-risk AML patients proceeding to HCT in CR1, but perhaps most importantly by the presence of non-genetic, adverse clinical features in the Fav risk cohort (i.e., elevated WBC at diagnosis, failed induction, pre-transplant MRD, etc.) which could have resulted in early transplantation referral. Thus, due to inherent clinical differences between Fav patients referred to HCT and those with newly diagnosed disease, aELN distinguished only two (and not three) distinct prognostic groups in the HCT setting.

Despite inferior outcomes compared to the Fav/IM group, the overall survival for Adv patients was still 52% at 2 years, a modest difference from the Fav/IM group; but most importantly, significantly better than reported without allogeneic HCT. Herold and colleagues recently validated the ELN2017 risk stratification among 1116 adult AML patients enrolled on two multicenter phase III trials of the German AML Cooperative Group (16). Two-year survival for Adv patients was 22.1% (20.6% following chemotherapy or autoHCT consolidation for patients in CR1). Several cooperative and single-institution studies have also reported very poor outcomes for Adv risk patients receiving post-remission therapies other than allogeneic HCT (14, 15). While clear differences in outcomes were noted between Adv and Fav/IM groups, prognostic separation was clear when either were compared to patients receiving non-transplant consolidation. It is not unreasonable to suggest that allogeneic transplantation could have a partial ‘equalizing’ effect amongst different genetic subsets.

We identified the most commonly occurring genetic abnormalities (i.e., >100 cases) within the Adv aELN group. As reported, (18, 28) we confirmed that patients carrying monosomy 5, del(5q) or monosomy 7 (abnormal 5/7) had significantly higher rates of post-transplant relapse and inferior survival, even when transplanted in CR1. Monosomy 7 was the most common individual abnormality reported, with worse outcomes noted in the context of monosomal karyotype. Patients in the abnormal 5/7 subgroup were older and received less aggressive conditioning; however, the adverse impact of 5/7 was seen across all age groups and also in patients receiving MAC.



Unfortunately we could not assess in our analysis the clinical impact of post-transplant maintenance and preemptive strategies in our cohort or elucidate the indications for HCT in Fav/IM risk patients. Additional study limitations were intrinsic to a retrospective cohort analysis, including incomplete genetic data for some patients and lack of uniformity for MRD assessment methods among multiple institutions.

The impact of pre-HCT MRD has been previously reported by several groups (29–32). The presence of MRD correlates with increased post-transplant relapse, particularly in patients receiving RIC/NMA regimens. While we observed a higher proportion of pre-HCT MRD in the Adv aELN group, the presence of MRDpos was independently associated with higher post-HCT relapse across the aELN subsets. The patients in this study were transplanted prior to the publication of the ELN consensus guidelines for AML MRD assessment however, even without interlaboratory or detection technology standardization, residual disease pre-HCT was consistently associated with more relapse.

In conclusion, this adapted ELN classification had prognostic value for survival and relapse after allogeneic HCT. Novel peri-transplant pre-emptive and therapeutic strategies to reduce relapse remain necessary for all groups, but particularly those within the Adv cohort, those at highest-risk.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Authors

Antonio M. Jimenez Jimenez<sup>1</sup>, Marcos de Lima<sup>2</sup>, Krishna V. Komanduri<sup>1</sup>, Trent P Wang<sup>1</sup>, Mei-Jie Zhang<sup>3,4</sup>, Karen Chen<sup>3</sup>, Hisham Abdel-Azim<sup>5</sup>, Muhammad Bilal Abid<sup>6</sup>, Mahmoud Aljurf<sup>7</sup>, Hassan Alkhateeb<sup>8</sup>, Amer Assal<sup>9</sup>, Ulrike Bacher<sup>10</sup>, Frédéric Baron<sup>11</sup>, Minoo Battiwalla<sup>12</sup>, Amer Beitinjaneh<sup>1</sup>, Nelli Bejanyan<sup>13</sup>, Vijaya Raj Bhatt<sup>14</sup>, Michael Byrne<sup>15</sup>, Jean-Yves Cahn<sup>16</sup>, Mitchell Cairo<sup>17</sup>, Paul Castillo<sup>18</sup>, Edward Copelan<sup>19</sup>, Zachariah DeFilipp<sup>20</sup>, Miguel Angel Diaz Perez<sup>21</sup>, Mahmoud Elsayy<sup>22</sup>, Robert Peter Gale<sup>23</sup>, Biju George<sup>24</sup>, Michael R. Grunwald<sup>25</sup>, Gerhard C. Hildebrandt<sup>26</sup>, William J. Hogan<sup>27</sup>, Christopher G. Kanakry<sup>28</sup>, Ankit Kansagra<sup>29</sup>, Mohamed A. Kharfan-Dabaja<sup>30</sup>, Nandita Khera<sup>31</sup>, Maxwell M. Krem<sup>32</sup>, Aleksandr Lazaryan<sup>13</sup>, Joseph Maakaron<sup>33</sup>, Rodrigo Martino<sup>34</sup>, Joseph McGuirk<sup>35</sup>, Fotios V. Michelis<sup>36</sup>, Giuseppe Milone<sup>37</sup>, Asmita Mishra<sup>13</sup>, Hemant S. Murthy<sup>30</sup>, Alberto Mussetti<sup>38,39</sup>, Sunita Nathan<sup>40</sup>, Taiga Nishihori<sup>13</sup>, Richard F. Olsson<sup>41,42</sup>, Neil Palmisiano<sup>43</sup>, Sagar Patel<sup>44</sup>, Ayman Saad<sup>45</sup>, Sachiko Seo<sup>46</sup>, Akshay Sharma<sup>47</sup>, Melhem Solh<sup>48</sup>, Leo F. Verdonck<sup>49</sup>, Baldeep Wirk<sup>50</sup>, Jean A. Yared<sup>51</sup>, Mark Litzow<sup>52</sup>, Partow Kebriaei<sup>53</sup>, Christopher S. Hourigan<sup>54</sup>, Wael Saber<sup>3</sup>, Daniel Weisdorf<sup>55</sup>

## Affiliations

<sup>1</sup>Division of Transplantation and Cellular Therapy, University of Miami Miller School of Medicine, Miami, FL

<sup>2</sup>Department of Medicine, Seidman Cancer Center, University Hospitals Case Medical Center, Cleveland, OH

<sup>3</sup>(CIBMTR)<sup>®</sup> Center for International Blood and Marrow Transplant Research, Department of Medicine, Medical College of Wisconsin, Milwaukee, WI

<sup>4</sup>Division of Biostatistics, Institute for Health and Society, Medical College of Wisconsin, Milwaukee, WI

<sup>5</sup>Division of Hematology, Oncology and Blood & Marrow Transplantation, Children's Hospital Los Angeles, University of Southern California Keck School of Medicine, Los Angeles, CA

<sup>6</sup>Divisions of Hematology/Oncology & Infectious Diseases, Department of Medicine, Medical College of Wisconsin, Milwaukee, WI

<sup>7</sup>Department of Oncology, King Faisal Specialist Hospital Center & Research, Riyadh, Saudi Arabia

<sup>8</sup>Mayo Clinic Rochester, Rochester, MN

<sup>9</sup>Columbia University Irving Medical Center, Department of Medicine, Bone Marrow Transplant and Cell Therapy Program, New York, NY

<sup>10</sup>Department of Hematology, Inselspital, Bern University Hospital, University of Bern, Switzerland

<sup>11</sup>CHU and University of Liege, Liege, Belgium

<sup>12</sup>Sarah Cannon Blood Cancer Network, Nashville, TN

<sup>13</sup>Department of Blood & Marrow Transplant and Cellular Immunotherapy (BMT CI), Moffitt Cancer Center, Tampa, FL

<sup>14</sup>The Fred and Pamela Buffett Cancer Center, University of Nebraska Medical Center, Omaha, NE

<sup>15</sup>Vanderbilt University Medical Center, Nashville, TN

<sup>16</sup>Department of Hematology, CHU Grenoble Alpes, Université Grenoble Alpes, Grenoble, France

<sup>17</sup>Division of Pediatric Hematology, Oncology and Stem Cell Transplantation, Department of Pediatrics, New York Medical College, Valhalla, NY

<sup>18</sup>UF Health Shands Children's Hospital, Gainesville, FL

<sup>19</sup>Department of Hematologic Oncology and Blood Disorders, Levine Cancer Institute, Atrium Health, Charlotte, NC

<sup>20</sup>Hematopoietic Cell Transplant and Cellular Therapy Program, Massachusetts General Hospital, Boston, MA

<sup>21</sup>Department of Hematology/Oncology, Hospital Infantil Universitario Niño Jesús, Madrid, Spain

- <sup>22</sup>Division of Hematology, Department of Medicine, Dalhousie University, Halifax, Canada
- <sup>23</sup>Haematology Research Centre, Department of Immunology and Inflammation, Imperial College London, London, UK
- <sup>24</sup>Christian Medical College, Vellore, India
- <sup>25</sup>Department of Hematologic Oncology and Blood Disorders, Levine Cancer Institute, Atrium Health, Charlotte, NC
- <sup>26</sup>Markey Cancer Center, University of Kentucky, Lexington, KY
- <sup>27</sup>Division of Hematology/BMT, Mayo Clinic, Rochester, MN
- <sup>28</sup>Experimental Transplantation and Immunotherapy Branch, Center for Cancer Research, National Cancer Institute, National Institutes of Health, Bethesda, MD
- <sup>29</sup>UT Southwestern Medical Center, Blood and Marrow Transplant Program, Dallas, TX
- <sup>30</sup>Division of Hematology-Oncology, Blood and Marrow Transplantation Program, Mayo Clinic, Jacksonville, FL
- <sup>31</sup>Department of Hematology/Oncology, Mayo Clinic, Phoenix, AZ
- <sup>32</sup>Markey Cancer Center, University of Kentucky College of Medicine, Lexington, KY
- <sup>33</sup>Division of Hematology, Oncology and Transplantation, Department of Medicine, University of Minnesota, Minneapolis, MN
- <sup>34</sup>Division of Clinical Hematology, Hospital de la Santa Creu i Sant Pau, Barcelona, Spain
- <sup>35</sup>Division of Hematologic Malignancies and Cellular Therapeutics, The University of Kansas Cancer Center
- <sup>36</sup>Allogeneic Blood and Marrow Transplant Program, Princess Margaret Cancer Centre, Toronto, Canada
- <sup>37</sup>Azienda Ospedaliera Universitaria Policlinico-San Marco, Catania, Italy
- <sup>38</sup>Hematology Department, Institut Català d' Oncologia – Hospitalet, Barcelona, Spain
- <sup>39</sup>IDIBELL-Institut Català d' Oncologia, l'Hospitalet de Llobregat, El Prat de Llobregat, Spain
- <sup>40</sup>Section of Bone Marrow Transplant and Cell Therapy, Rush University Medical Center, Chicago, IL
- <sup>41</sup>Department of Laboratory Medicine, Karolinska Institutet, Stockholm, Sweden
- <sup>42</sup>Centre for Clinical Research Sormland, Uppsala University, Uppsala, Sweden
- <sup>43</sup>Sidney Kimmel Cancer Center, Thomas Jefferson University, Philadelphia, PA

<sup>44</sup>Blood and Marrow Transplant Program, Huntsman Cancer Institute, University of Utah, Salt Lake City, UT

<sup>45</sup>Division of Hematology, Ohio State University, Columbus, OH

<sup>46</sup>Department of Hematology and Oncology, Dokkyo Medical University, Tochigi, Japan

<sup>47</sup>Department of Bone Marrow Transplantation and Cellular Therapy, St. Jude Children's Research Hospital, Memphis, TN

<sup>48</sup>The Blood and Marrow Transplant Group of Georgia, Northside Hospital, Atlanta, GA

<sup>49</sup>Department of Hematology/Oncology, Isala Clinic, Zwolle, The Netherlands

<sup>50</sup>Bone Marrow Transplant Program, Penn State Cancer Institute, Hershey, Pennsylvania

<sup>51</sup>Blood & Marrow Transplantation Program, Division of Hematology/Oncology, Department of Medicine, Greenebaum Comprehensive Cancer Center, University of Maryland, Baltimore, MD

<sup>52</sup>Division of Hematology and Transplant Center, Mayo Clinic Rochester, Rochester, MN

<sup>53</sup>Department of Stem Cell Transplantation, Division of Cancer Medicine, The University of Texas MD Anderson Cancer Center

<sup>54</sup>Laboratory of Myeloid Malignancies, National Heart, Lung, and Blood Institute, National Institutes of Health, Bethesda, MD

<sup>55</sup>Division of Hematology, Oncology and Transplantation, Department of Medicine, University of Minnesota, Minneapolis, MN

## ACKNOWLEDGEMENTS

Data sharing:

CIBMTR supports accessibility of research in accord with the National Institutes of Health (NIH) Data Sharing Policy and the National Cancer Institute (NCI) Cancer Moonshot Public Access and Data Sharing Policy. The CIBMTR only releases de-identified datasets that comply with all relevant global regulations regarding privacy and confidentiality.

Additional contributing co-authors from the writing committee:

Kehinde Adekola, Ibrahim Ahmed, Talha Badar, Ashish Bajel, Qaiser Bashir, Chris Bredeson, Bruce Camitta, Jan Cerny, Stefan Ciurea, Corey S. Cutler, Shahinaz Gadalla, Sid Ganguly, Saar Gill, Nasheed Hossain, Luis Isola, Mark Juckett, Mary Laughlin, Hillard Lazarus, Jong Wook Lee, Richard Lin, Hongtao Liu, Vikram Matthews, Shahram Mori, Reinhold Munker, Ryotaro Nakamura, Miguel-Angel Perales, Marjolein van der Poel, Uday Popat, Jeffrey Pu, Marcie Riches, Olle Ringdén, David Rizzieri, Mitchell Sabloff, Lynn Savoie, Bart Scott, Anthony Stein, Masumi Ueda, Celalettin Ustun, Eric Wong.

The CIBMTR is supported primarily by Public Health Service U24CA076518 from the National Cancer Institute (NCI), the National Heart, Lung and Blood Institute (NHLBI) and the National Institute of Allergy and Infectious Diseases (NIAID); U24HL138660 from NHLBI and NCI; OT3HL147741, and U01HL128568 from the NHLBI; HHS250201700006C, and HHS250201700007C from the Health Resources and Services Administration (HRSA); and N00014-20-1-2705 and N00014-20-1-2832 from the Office of Naval Research; Additional

federal support is provided by P01CA111412, R01CA152108, R01CA215134, R01CA218285, R01CA231141, R01AI128775, R01HL126589, R01HL129472, R01HL130388, R01HL131731, U01AI069197, U01AI126612, UG1HL06924, and BARDA. Support is also provided by Be the Match Foundation, Boston Children's Hospital, Dana Farber, St. Baldrick's Foundation, Stanford University, the Medical College of Wisconsin the National Marrow Donor Program, and from the following commercial entities: Actinium Pharmaceuticals, Inc.; Adienne SA; Allovir, Inc.; Amgen, Inc.; Angiocrine Bioscience; Astellas Pharma US; bluebird bio, Inc.; Bristol Myers Squibb Co.; Celgene Corp.; CSL Behring; CytoSen Therapeutics, Inc.; Daiichi Sankyo Co., Ltd.; ExcellThera; Fate Therapeutics; Gamida-Cell, Ltd.; Genentech Inc; Incyte Corporation; Janssen/Johnson & Johnson; Jazz Pharmaceuticals, Inc.; Kiadis Pharma; Kite, a Gilead Company; Kyowa Kirin; Legend Biotech; Magenta Therapeutics; Merck Sharp & Dohme Corp.; Millennium, the Takeda Oncology Co.; Miltenyi Biotec, Inc.; Novartis Pharmaceuticals Corporation; Omeros Corporation; Oncoimmune, Inc.; Orca Biosystems, Inc.; Pfizer, Inc.; Pharmacyclics, LLC; Sanofi Genzyme; Stemcyte; Takeda Pharma; Vor Biopharma; Xenikos BV. The views expressed in this article do not reflect the official policy or position of the National Institute of Health, the Department of the Navy, the Department of Defense, Health Resources and Services Administration (HRSA) or any other agency of the U.S. Government.

#### COMPETING INTERESTS

Dr. Assal reports grants and personal fees from Incyte Corporation, personal fees from Boston Biomedical, personal fees from Alpha Insights, outside the submitted work.

Dr. Bejanyan reports personal fees from Magenta therapeutics, outside the submitted work.

Dr. Cahn reports Advisory Boards with Agios, AbbVie, Otsuka, Race Oncology.

Dr. Bhatt reports personal fees from Agios, grants and personal fees from Incyte, personal fees from Takeda, personal fees from Partner Therapeutics, personal fees from Omeros, grants and personal fees from Abbvie, grants from Jazz, grants from National Marrow Donor Program, other from Oncoceutics, personal fees from Partnership for health analytic research, LLC, grants and other from Pfizer, personal fees from CSL Behring, grants from Tolero Pharmaceuticals, personal fees from Rigel Pharmaceuticals, other from Novartis, personal fees from Genentech, outside the submitted work.

Dr. de Lima reports grants from Pfizer, grants from Celgene, personal fees from Kadmon, personal fees from Pfizer, personal fees from Incyte, personal fees from BMS, outside the submitted work.

Dr. Gale is a consultant to BeiGene Ltd., Kite Pharma Inc., Fusion Pharma LLC, LaJolla NanoMedical Inc., Mingsight Pharmaceuticals Inc. and CStone Pharmaceuticals; Medical Director, FFF Enterprises Inc.; Partner, AZCA Inc.; Board of Directors, RakFond Foundation for Cancer Research Support; Scientific Advisory Board, Antegene Biotech LLC and StemRad Ltd.

Dr. Grunwald reports personal fees from Abbvie, personal fees from Agios, personal fees from Amgen, personal fees from Cardinal Health, personal fees from BMS/Celgene, personal fees from Daiichi Sankyo, personal fees and other from Incyte, personal fees from Merck, personal fees from Pfizer, personal fees from Premier, personal fees from Karius, other from Forma Therapeutics, other from Genentech/Roche, other from Janssen, personal fees from Astellas, personal fees from Trovogene, personal fees from Stemline, personal fees from Gilead, outside the submitted work.

Dr. Hildebrandt reports other from Pfizer, other from Kite Pharma, other from Incyte, other from Jazz Pharmaceuticals, other from Alexion Pharmaceutical, other from Karyopharm, other from Pharmacyclics, other from Takada, other from Jazz Pharmaceuticals, other from Kite Pharma, other from Incyte, other from Pfizer, other from Falk Foundation, other from Astellas Pharma, outside the submitted work.

Dr. Hourigan reports other from Sellas, outside the submitted work.

Dr. Kansagra reports other from Takeda, other from Jansen, other from Pfizer, other from Celgene/BMS, other from Sanofi, other from Alnylam, other from GSK, other from Karyopharm, other from Oncocept, outside the submitted work.

Dr. Kharfan-Dabaja reports other from Daiichi Sankyo, other from Pharmacyclics, outside the submitted work.

Dr. McGuirk reports grants and personal fees from AlloVir HCP, grants and personal fees from Juno Therapeutics, Inc, grants and personal fees from Gilead-Kite Pharmaceuticals, grants and other from Magenta Therapeutics, other from EcoR1 Cap, outside the submitted work.

Dr. Mishra reports grants from Novartis, from null, outside the submitted work.

Dr. Mussetti reports grants from Gilead, personal fees from Novartis, Bristol-Myers Squibb, outside the submitted work.

Dr. Patel reports personal fees from Kite Pharma, outside the submitted work.

Dr. Olsson reports personal fees from AstraZeneca, outside the submitted work.

Dr. Saad reports personal fees from Magenta Therapeutics, personal fees from Incyte Pharmaceuticals, personal fees from CareDx, outside the submitted work.

Dr. Sharma reports clinical trial salary support from Vertex Pharmaceuticals, CRISPR Therapeutics, Novartis paid to his institution, and personal consultancy fees from Spotlight Therapeutics, outside the submitted work.

All other authors declare no potential competing interests.


## REFERENCES

1. Sorror ML, Maris MB, Storb R, Baron F, Sandmaier BM, Maloney DG, et al. Hematopoietic cell transplantation (HCT)-specific comorbidity index: a new tool for risk assessment before allogeneic HCT. *Blood*. 2005;106(8):2912–9. [PubMed: 15994282]
2. Koreth J, Schlenk R, Kopecky KJ, Honda S, Sierra J, Djulbegovic BJ, et al. Allogeneic stem cell transplantation for acute myeloid leukemia in first complete remission: systematic review and meta-analysis of prospective clinical trials. *JAMA*. 2009;301(22):2349–61. [PubMed: 19509382]
3. Slovak ML, Kopecky KJ, Cassileth PA, Harrington DH, Theil KS, Mohamed A, et al. Karyotypic analysis predicts outcome of preremission and postremission therapy in adult acute myeloid leukemia: a Southwest Oncology Group/Eastern Cooperative Oncology Group Study. *Blood*. 2000;96(13):4075–83. [PubMed: 11110676]
4. Grimwade D, Hills RK, Moorman AV, Walker H, Chatters S, Goldstone AH, et al. Refinement of cytogenetic classification in acute myeloid leukemia: determination of prognostic significance of rare recurring chromosomal abnormalities among 5876 younger adult patients treated in the United Kingdom Medical Research Council trials. *Blood*. 2010;116(3):354–65. [PubMed: 20385793]
5. Dholaria B, Savani BN, Hamilton BK, Oran B, Liu HD, Tallman MS, et al. Hematopoietic Cell Transplantation in the Treatment of Newly Diagnosed Adult Acute Myeloid Leukemia: An Evidence-Based Review from the American Society of Transplantation and Cellular Therapy. *Biol Blood Marrow Transplant*. 2020.
6. Oliansky DM, Appelbaum F, Cassileth PA, Keating A, Kerr J, Nieto Y, et al. The role of cytotoxic therapy with hematopoietic stem cell transplantation in the therapy of acute myelogenous leukemia in adults: an evidence-based review. *Biol Blood Marrow Transplant*. 2008;14(2):137–80. [PubMed: 18215777]
7. Grossmann V, Schnittger S, Kohlmann A, Eder C, Roller A, Dicker F, et al. A novel hierarchical prognostic model of AML solely based on molecular mutations. *Blood*. 2012;120(15):2963–72. [PubMed: 22915647]
8. Dohner H, Estey EH, Amadori S, Appelbaum FR, Buchner T, Burnett AK, et al. Diagnosis and management of acute myeloid leukemia in adults: recommendations from an international expert panel, on behalf of the European LeukemiaNet. *Blood*. 2010;115(3):453–74. [PubMed: 19880497]
9. Mrozek K, Marcucci G, Nicolet D, Maharry KS, Becker H, Whitman SP, et al. Prognostic significance of the European LeukemiaNet standardized system for reporting cytogenetic and molecular alterations in adults with acute myeloid leukemia. *J Clin Oncol*. 2012;30(36):4515–23. [PubMed: 22987078]
10. Rollig C, Bornhauser M, Thiede C, Taube F, Kramer M, Mohr B, et al. Long-term prognosis of acute myeloid leukemia according to the new genetic risk classification of the European LeukemiaNet recommendations: evaluation of the proposed reporting system. *J Clin Oncol*. 2011;29(20):2758–65. [PubMed: 21632498]
11. Medeiros BC, Tian L, Robenson S, Laport GG, Johnston LJ, Shizuru JA, et al. European LeukemiaNet classification intermediate risk-1 cohort is associated with poor outcomes in adults with acute myeloid leukemia undergoing allogeneic hematopoietic cell transplantation. *Blood Cancer J*. 2014;4:e216. [PubMed: 24879117]

12. Oran B, Jimenez AM, De Lima M, Popat UR, Bassett R, Andersson B, et al. Age and Modified European LeukemiaNet Classification to Predict Transplant Outcomes: An Integrated Approach for Acute Myelogenous Leukemia Patients Undergoing Allogeneic Stem Cell Transplantation. *Biol Blood Marrow Transplant.* 2015;21(8):1405–12. [PubMed: 25840338]
13. Dohner H, Estey E, Grimwade D, Amadori S, Appelbaum FR, Buchner T, et al. Diagnosis and management of AML in adults: 2017 ELN recommendations from an international expert panel. *Blood.* 2017;129(4):424–47. [PubMed: 27895058]
14. Boddu PC, Kadia TM, Garcia-Manero G, Cortes J, Alfayez M, Borthakur G, et al. Validation of the 2017 European LeukemiaNet classification for acute myeloid leukemia with NPM1 and FLT3-internal tandem duplication genotypes. *Cancer.* 2019;125(7):1091–100. [PubMed: 30521114]
15. Harada Y, Nagata Y, Kihara R, Ishikawa Y, Asou N, Ohtake S, et al. Prognostic analysis according to the 2017 ELN risk stratification by genetics in adult acute myeloid leukemia patients treated in the Japan Adult Leukemia Study Group (JALSG) AML201 study. *Leuk Res.* 2018;66:20–7. [PubMed: 29360622]
16. Herold T, Rothenberg-Thurley M, Grunwald VV, Janke H, Goerlich D, Sauerland MC, et al. Validation and refinement of the revised 2017 European LeukemiaNet genetic risk stratification of acute myeloid leukemia. *Leukemia.* 2020;34(12):3161–72. [PubMed: 32231256]
17. Straube J, Ling VY, Hill GR, Lane SW. The impact of age, NPM1(mut), and FLT3(ITD) allelic ratio in patients with acute myeloid leukemia. *Blood.* 2018;131(10):1148–53. [PubMed: 29183886]
18. Grimm J, Jentzsch M, Bill M, Goldmann K, Schulz J, Niederwieser D, et al. Prognostic impact of the ELN2017 risk classification in patients with AML receiving allogeneic transplantation. *Blood Adv.* 2020;4(16):3864–74. [PubMed: 32810221]
19. Hansen DK, Nieder ML, Thompson Z, Kim J, Pidala JA, Nishihori T, et al. ELN 2017 Risk Classification Predicts Survival of AML Patients Receiving Allogeneic Hematopoietic Stem Cell Transplantation. *Biology of Blood and Marrow Transplantation.* 2019;25(3):S112.
20. McGowan-Jordan J ISCN 2016: An International System for Human Cytogenomic Nomenclature (2016); Recommendations of the International Standing Human Committee on Human Cytogenomic Nomenclature Including New Sequence-based Cytogenomic: Karger; 2016.
21. Breems DA, Van Putten WL, De Greef GE, Van Zelderen-Bhola SL, Gerssen-Schoorl KB, Mellink CH, et al. Monosomal karyotype in acute myeloid leukemia: a better indicator of poor prognosis than a complex karyotype. *J Clin Oncol.* 2008;26(29):4791–7. [PubMed: 18695255]
22. Byrd JC, Mrozek K, Dodge RK, Carroll AJ, Edwards CG, Arthur DC, et al. Pretreatment cytogenetic abnormalities are predictive of induction success, cumulative incidence of relapse, and overall survival in adult patients with de novo acute myeloid leukemia: results from Cancer and Leukemia Group B (CALGB 8461). *Blood.* 2002;100(13):4325–36. [PubMed: 12393746]
23. Przepiorka D, Weisdorf D, Martin P, Klingemann HG, Beatty P, Hows J, et al. 1994 Consensus Conference on Acute GVHD Grading. *Bone Marrow Transplant.* 1995;15(6):825–8. [PubMed: 7581076]
24. Thiede C, Koch S, Creutzig E, Steudel C, Illmer T, Schaich M, et al. Prevalence and prognostic impact of NPM1 mutations in 1485 adult patients with acute myeloid leukemia (AML). *Blood.* 2006;107(10):4011–20. [PubMed: 16455956]
25. Fine JP, Gray RJ. A Proportional Hazards Model for the Subdistribution of a Competing Risk. *Journal of the American Statistical Association.* 1999;94(446):496–509.
26. Cox DR. Regression Models and Life-Tables. *Journal of the Royal Statistical Society Series B (Methodological).* 1972;34(2):187–220.
27. Breems DA, Van Putten WL, Huijgens PC, Ossenkoppele GJ, Verhoef GE, Verdonck LF, et al. Prognostic index for adult patients with acute myeloid leukemia in first relapse. *J Clin Oncol.* 2005;23(9):1969–78. [PubMed: 15632409]
28. Pasquini MC, Zhang MJ, Medeiros BC, Armand P, Hu ZH, Nishihori T, et al. Hematopoietic Cell Transplantation Outcomes in Monosomal Karyotype Myeloid Malignancies. *Biol Blood Marrow Transplant.* 2016;22(2):248–57. [PubMed: 26327629]

29. Baron F, Labopin M, Ruggeri A, Sierra J, Robinson S, Labussiere-Wallet H, et al. Impact of detectable measurable residual disease on umbilical cord blood transplantation. *Am J Hematol.* 2020;95(9):1057–65. [PubMed: 32449964]
30. Buckley SA, Wood BL, Othus M, Hourigan CS, Ustun C, Linden MA, et al. Minimal residual disease prior to allogeneic hematopoietic cell transplantation in acute myeloid leukemia: a meta-analysis. *Haematologica.* 2017;102(5):865–73. [PubMed: 28126965]
31. Gilleece MH, Labopin M, Yakoub-Agha I, Volin L, Socie G, Ljungman P, et al. Measurable residual disease, conditioning regimen intensity, and age predict outcome of allogeneic hematopoietic cell transplantation for acute myeloid leukemia in first remission: A registry analysis of 2292 patients by the Acute Leukemia Working Party European Society of Blood and Marrow Transplantation. *Am J Hematol.* 2018;93(9):1142–52. [PubMed: 29981272]
32. Hourigan CS, Dillon LW, Gui G, Logan BR, Fei M, Ghannam J, et al. Impact of Conditioning Intensity of Allogeneic Transplantation for Acute Myeloid Leukemia With Genomic Evidence of Residual Disease. *J Clin Oncol.* 2020;38(12):1273–83. [PubMed: 31860405]





**Figure 1.** OS (A), DFS (B) and cumulative incidence of disease relapse (C) for AML transplant recipients within favorable/intermediate vs. adverse aELN cohorts  
 Abbreviations: DFS, disease-free survival; aELN, adapted European LeukemiaNet; OS, overall survival.

**Table 1.**

Adapted\* ELN (aELN) risk stratification

Favorable	t(8;21)(q22;q22.1); <i>RUNX1-RUNX1T1</i>
	inv(16)(p13.1q22) or t(16;16)(p13.1;q22); <i>CBFB-MYH11</i>
	Mutated <i>NPM1</i> without <i>FLT3</i> -ITD
	Mutated <i>CEBPA</i>
Intermediate	Mutated <i>NPM1</i> and <i>FLT3</i> -ITD
	Wild-type <i>NPM1</i> without <i>FLT3</i> -ITD
	t(9;11)(p21.3;q23.3); <i>MLLT3-KMT2A</i>
	Cytogenetic abnormalities not classified as favorable or adverse
Adverse	t(6;9)(p23;q34.1); <i>DEK-NUP214</i>
	t(v;11q23.3); <i>KMT2A</i> rearranged
	t(9;22)(q34.1;q11.2); <i>BCR-ABL1</i>
	inv(3)(q21.3q26.2) or t(3;3)(q21.3;q26.2); <i>GATA2,MECOM</i>
	-5 or del(5q); -7; -17/abn(17p); mutated <i>TP53</i>
	Complex karyotype ( ≥ 3 abnormalities)
	Monosomal karyotype
	Wild-type <i>NPM1</i> and <i>FLT3</i> -ITD
	Mutated <i>RUNX1</i> without good risk karyotype
	Mutated <i>ASXL1</i> without good risk karyotype

<sup>±</sup> *FLT3*-ITD allelic ratio and *CEBPA* mono/bi-allelic status not available

**Table 2.**

## Patient, disease and transplant characteristics

Characteristic	Favorable	Intermediate	Adverse	P Value
No. of patients	181	1185	923	
No. of centers	66	144	125	
<b>Patient related</b>				
Age at HCT - no. (%)				0.01 <sup>a</sup>
Median (range) years	57.3 (18.4–73.5)	58.1 (19.6–74)	56.2 (19–74.4)	
18–29	12 (6.6)	97 (8.2)	104 (11.3)	
30–59	88 (48.6)	96 (48.9)	473 (51.3)	
60–69	63 (34.8)	417 (35.2)	287 (31.1)	
≥70	18 (9.9)	92 (7.8)	59 (6.4)	
Gender - no. (%)				0.16 <sup>a</sup>
Male	95 (52.5)	615 (51.9)	517 (56)	
Female	86 (47.5)	570 (48.1)	406 (44)	
Race - no. (%)				< 0.001 <sup>a</sup>
Caucasian	138 (76.2)	942 (79.5)	718 (77.8)	
African-American	6 (3.3)	70 (5.9)	101 (10.9)	
Asian/Pacific Islander	27 (14.9)	110 (9.3)	70 (7.6)	
Other/missing	10 (5.6)	63 (5.3)	34 (3.7)	
Performance Score - no. (%)				0.09 <sup>a</sup>
<90	64 (35.4)	434 (36.6)	386 (41.8)	
≥90	115 (63.5)	739 (62.4)	524 (56.8)	
Missing	2 (1.1)	12 (1)	13 (1.4)	
HCT-CI - no. (%)				< 0.001 <sup>a</sup>
0	39 (21.5)	245 (20.7)	171 (18.5)	
1–2	62 (34.3)	409 (34.5)	249 (27)	
3+	74 (40.9)	497 (41.9)	489 (53)	
Missing	6 (3.3)	34 (2.9)	14 (1.5)	
<b>Disease related</b>				
White blood count at diagnosis, x10 <sup>9</sup> /L				< 0.001 <sup>a</sup>
Median (range)	13.9 (0–270.5)	7.2 (0–329.7)	5 (0.4–290.9)	
≤ 10	73 (40.3)	597 (50.4)	528 (57.2)	
10 – 100	86 (47.5)	396 (33.4)	275 (29.8)	
> 100	14 (7.7)	114 (9.6)	57 (6.2)	
Missing	8 (4.4)	78 (6.6)	63 (6.8)	
Clinical onset of AML - no. (%)				0.02 <sup>a</sup>
<i>De novo</i>	147 (81.2)	922 (77.8)	674 (73)	
Transformed from MDS/MPN	23 (12.7)	188 (15.9)	161 (17.4)	

Characteristic	Favorable	Intermediate	Adverse	P Value
Therapy related	11 (6.1)	75 (6.3)	88 (9.5)	
CRi status pre-HCT - no. (%)				0.29 <sup>a</sup>
Yes	49 (27.1)	328 (27.7)	269 (29.1)	
No	132 (72.9)	830 (70)	636 (68.9)	
Missing	0	27 (2.3)	18 (2)	
MRD pre-HCT <sup>+</sup> - no. (%)				< 0.001 <sup>a</sup>
No	128 (70.7)	917 (77.4)	666 (72.2)	
Yes	38 (21)	136 (11.5)	211 (22.9)	
Missing	15 (8.3)	132 (11.1)	46 (5)	
<b>Transplant related</b>				
Conditioning intensity - no. (%)				0.12 <sup>a</sup>
MAC	91 (50.3)	547 (46.2)	477 (51.7)	
RIC/NMA	90 (49.7)	637 (53.8)	446 (48.3)	
Missing	0	1 (0.1)	0	
Graft type - no. (%)				0.62 <sup>a</sup>
Bone marrow	34 (18.8)	173 (14.6)	135 (14.6)	
Peripheral blood	119 (65.7)	827 (69.8)	635 (68.8)	
Cord blood	28 (15.5)	185 (15.6)	153 (16.6)	
Donor type - no. (%)				0.68 <sup>a</sup>
HLA-identical sibling	49 (27.1)	325 (27.4)	223 (24.2)	
Other related	14 (7.7)	99 (8.4)	69 (7.5)	
Haploidentical	23 (12.7)	124 (10.5)	117 (12.7)	
HLA-Matched unrelated (8/8)	67 (37)	452 (38.1)	361 (39.1)	
Cord blood	28 (15.5)	185 (15.6)	153 (16.6)	
GVHD prophylaxis - no. (%)				0.24 <sup>a</sup>
Ex vivo T-cell depletion/CD34 selection	10 (5.5)	49 (4.1)	55 (6)	
PTCy	24 (13.3)	179 (15.1)	158 (17.1)	
TAC based	111 (61.3)	713 (60.2)	554 (60)	
CSA based	33 (18.2)	230 (19.4)	149 (16.1)	
Other	3 (1.7)	14 (1.2)	7 (0.8)	
In vivo T-cell depletion - no. (%)				0.46 <sup>a</sup>
No	137 (75.7)	944 (79.7)	733 (79.4)	
ATG/Alemtuzumab	44 (24.3)	241 (20.3)	190 (20.6)	
Year of HCT - no. (%)				0.15 <sup>a</sup>
2013	34 (18.8)	220 (18.6)	139 (15.1)	
2014	36 (19.9)	275 (23.2)	237 (25.7)	
2015	36 (19.9)	278 (23.5)	208 (22.5)	
2016	47 (26)	248 (20.9)	188 (20.4)	
2017	28 (15.5)	164 (13.8)	151 (16.4)	

Characteristic	Favorable	Intermediate	Adverse	P Value
Follow-up of survivors - median (range) months	24.6 (0.4–62.8)	35.3 (0.2–69.4)	35.1 (0.4–63.3)	

Hypothesis testing:

<sup>a</sup> Pearson chi-square test

<sup>+</sup> MRD was assessed by different methods including, but not limited to: next generation sequencing (NGS), polymerase chain reaction (PCR) testing, chromosomal / genomic microarray analysis, fluorescence in situ hybridization (FISH), karyotyping, flow cytometry.

Abbreviations: AML, acute myeloid leukemia; ATG, anti-thymocyte globulin; CMV, cytomegalovirus; CRi, complete remission with incomplete blood count recovery; CSA, cyclosporine; HCT, hematopoietic cell transplant; HiDAC, high-dose cytarabine; HMA, hypomethylating agent; IDAC, intermediate-dose cytarabine; MAC, myeloablative conditioning; MDS, myelodysplastic syndrome; aELN, adapted European LeukemiaNet; MMF, mycophenolate mofetil; MPN, myeloproliferative neoplasm; MRD, measurable residual disease; NMA, non-myeloablative; PTCy, post-transplant cyclophosphamide; RIC, reduced intensity conditioning; TAC, tacrolimus.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

**Table 3.**

**3A.** Adverse-risk patient characteristics by genetic subgroups **3B.** Multivariate Analysis of Outcomes.  
Comparison of abnormal 5/7 subgroup vs. Other Adverse aELN risk group\*

<b>3A.</b> Characteristic	Abnormal 5/7	* Others	P Value
No. of patients	380	543	
Age at HCT - no. (%)			< 0.001 <sup>a</sup>
Median (range)	59.81 (18.98–75.84)	51.92 (18.96–73.76)	
18–59	195 (51.3)	382 (70.3)	
>=60	185 (48.7)	161 (29.7)	
Conditioning intensity - no. (%)			< 0.001 <sup>a</sup>
MAC	171 (45)	306 (56.4)	
RIC/NMA	209 (55)	237 (43.6)	
MRD at time of HCT - no. (%)			0.003 <sup>a</sup>
No	255 (67.1)	411 (75.7)	
Yes	108 (28.4)	103 (19)	
Missing	17 (4.5)	29 (5.3)	
CRi status prior to conditioning - no. (%)			0.06 <sup>a</sup>
Yes	127 (33.4)	142 (26.2)	
No	246 (64.7)	390 (71.8)	
Missing	7 (1.8)	11 (2)	
<b>3B.</b>			
		<b>HR (95% CI)</b>	<b>p-value</b>
Overall survival		1.58 (1.32–1.90)	<0.0001
Disease-free survival		1.54 (1.30–1.83)	<0.0001
Relapse		1.59 (1.28–1.97)	<0.0001

\* Reference group. Includes wt*NPM1* with *FLT3*-ITD, t(v;11q23.3), *DEK-NUP214*, complex karyotype, *BCR-ABL1*, *GATA2*, *MECOM*, mutated *TP53*, non-del5/del7 monosomy, mutated *RUNX1/ASXL1* without good-risk karyotype

<sup>a</sup>Pearson chi-square test

Abbreviations: CRi, complete remission with incomplete blood count recovery; GVHD, graft-vs.-host disease; HCT, hematopoietic cell transplant; MAC, myeloablative conditioning; aELN, adapted European LeukemiaNet; MRD, measurable residual disease; NMA, non-myeloablative; RIC, reduced intensity conditioning.

**Table 4.**

Univariate outcomes by aELN risk group

Outcomes	Favorable (N = 181)		Intermediate (N = 1185)		Adverse (N = 923)		P Value
	N	Prob (95% CI)	N	Prob (95% CI)	N	Prob (95% CI)	
Overall survival *	181		1185		923		<0.001
		67.7 (60.5–74.6)%		64.9 (62.1–67.6)%		53.9 (50.6–57.2)%	
Disease free survival *	178		1170		910		<0.001
		57.8 (50.3–65.2)%		55.5 (52.6–58.4)%		45.3 (42–48.6)%	
Relapse *	178		1170		910		<0.001
		28 (21.5–35.1)%		27.5 (24.9–30.1)%		27.5 (24.9–30.1)%	
Non-relapse mortality *	178		1170		910		0.467
		14.2 (9.3–19.8)%		17.1 (14.9–19.3)%		17.3 (14.9–19.8)%	
Grade II-IV acute GVHD <sup>+</sup>	181		1171		913		0.423
		30.9 (24.4–37.9)%		30.9 (24.4–37.9)%		35 (32–38.2)%	
Chronic GVHD *	181		1177		918		0.442
		40 (32.7–47.4)%		43.6 (40.7–46.6)%		40.2 (37–43.5)%	

Outcomes at:

\* 2 years

<sup>+</sup> 100 days.

Abbreviations: GVHD, graft-vs.-host disease; aELN, adapted European LeukemiaNet.

**Table 5:**

Multivariate Analysis of outcomes. Comparison of Adverse vs. Favorable/Intermediate\* aELN risk groups

	<b>HR (95% CI)</b>	<b>p-value</b>
Overall survival	1.39 (1.22–1.57)	<0.001
Disease-free survival	1.35 (1.20–1.51)	<0.001
Relapse	1.47 (1.28–1.70)	<0.001
Non-relapse mortality	1.01 (0.81–1.19)	0.89
Grade II-IV acute GVHD	1.04 (0.91–1.19)	0.53
Chronic GVHD	0.95 (0.83–1.08)	0.42

\* Reference group

Abbreviations: GVHD, graft-vs.-host disease; aELN, adapted European LeukemiaNet.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript