

UCSF

UC San Francisco Previously Published Works

Title

Comparative Molecular Analysis of Gastrointestinal Adenocarcinomas

Permalink

<https://escholarship.org/uc/item/8st0r4rp>

Journal

Cancer Cell, 33(4)

ISSN

1535-6108

Authors

Liu, Yang
Sethi, Nilay S
Hinoue, Toshinori
[et al.](#)

Publication Date

2018-04-01

DOI

10.1016/j.ccell.2018.03.010

Peer reviewed



Published in final edited form as:

Cancer Cell. 2018 April 09; 33(4): 721–735.e8. doi:10.1016/j.ccell.2018.03.010.

Comparative Molecular Analysis of Gastrointestinal Adenocarcinomas

Yang Liu^{1,2,*}, Nilay S. Sethi^{1,2,*}, Toshinori Hinoue^{3,*}, Barbara G. Schneider^{4,*}, Andrew D. Cherniack^{1,2}, Francisco Sanchez-Vega⁵, Jose A. Seoane⁶, Farshad Farshidfar⁷, Reanne Bowlby⁸, Mirazul Islam^{1,2}, Jaegil Kim¹, Walid Chatila⁹, Rehan Akbani¹⁰, Rupa S. Kanchi¹⁰, Charles S. Rabkin¹¹, Joseph E. Willis¹², Kenneth K. Wang¹³, Shannon J. McCall¹⁴, Lopa Mishra¹⁵, Akinyemi I. Ojesina¹⁶, Susan Bullman¹⁷, Chandra Sekhar Pedomallu¹⁷, Alexander J. Lazar¹⁸, Ryo Sakai¹⁹, The Cancer Genome Atlas Research Network[#], Vésteinn Thorsson^{20,†}, Adam J. Bass^{1,2,21,†}, and Peter W. Laird^{3,†,§}

¹The Eli and Edythe L. Broad Institute of Massachusetts Institute of Technology and Harvard University, Cambridge, Massachusetts 02142, USA

²Department of Medical Oncology, Dana-Farber Cancer Institute, Boston, Massachusetts 02115, USA

³Center for Epigenetics, Van Andel Research Institute, Grand Rapids, Michigan 49503, USA

⁴Department of Medicine, Division of Gastroenterology, Vanderbilt University Medical Center, Nashville, Tennessee 37232, USA

⁵Computational Biology Center, Memorial Sloan Kettering Cancer Center, New York, New York 10065, USA

Correspondence: Peter W. Laird: Peter.Laird@vai.org Adam J. Bass: Adam_Bass@dfci.harvard.edu Vésteinn Thorsson:

Vesteinn.Thorsson@systemsbiology.org.

*These authors contributed equally

†These authors contributed equally

§Lead Contact.

#Network membership provided following the main text

AUTHOR CONTRIBUTIONS

Conceptualization: YL, NSS, TH, ADC, JAS, FF, RB, VT, AJB, PWL Data Analysis: YL, NSS, TH, ADC, FSV, JAS, FF, RB, MI, JK, WC, RA, RSK, VT Clinicopathologic Analysis: CSR, JEW, KKW, SJM, LM, AIO, SB, CSP, AJL Critical thinking: YL, NSS, TH, ADC, JAS, FF, RB, VT, AJB, PWL Writing - Original Draft: YL, NSS, TH, BGS, ADC, FF, VT, AJB, PWL Visualization: YL, NSS, TH, ADC, RB, MI, JK, WC, RA, RSK, RS, VT Writing - Review & Editing: YL, NSS, TH, BGS, ADC, VT, AJB, PWL

DECLARATION OF INTERESTS

Andrew Cherniack, Ashton C. Berger, and Galen Gao receive research support from Bayer Pharmaceuticals AG. Kenneth Wang serves on Advisory Board for Boston Scientific, Microtech, Olympus. Peter W. Laird is on the Scientific Advisory Board for AnchorDx. Josh M. Stuart is the Founder of Five3 Genomics and shareholder of Nantomics. Christina Yau is a part-time employee/consultant at NantOmics. Gordon Mills serves on the External Scientific Review Board of Astrazeneca. Daniel J. Weisenberger is a consultant for Zymo Research Corporation. Charles M. Perou is an equity stock holder, consultant, and Board of Director Member, of BioClassifier LLC and GeneCentric Diagnostics and is listed as inventor on patent applications on the Breast PAM50 and Lung Cancer Subtyping assays. Matthew Meyerson has research support from Bayer, is equity holder in, consultant for and Scientific Advisory Board chair for Origimed, and is inventor on patent for EGFR mutation diagnosis in lung cancer, licensed to LabCorp. Kyle R. Covington is an employee of Castle Biosciences Inc. Joel Tepper is a consultant at EMD Serono. Gordon Mills serves on the External Scientific Review Board of Astrazeneca. Anil Sood is on the SAB for Kiyatec and is a shareholder in BioPath. Beth Y. Karlan is on the Advisory Board for Invitae. Han Liang is a shareholder and scientific advisor of Precision Scientific Ltd. and Eagle Nebula Inc.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

⁶Department of Medicine, and Department of Genetics, Stanford University School of Medicine, Stanford, California 94305, USA

⁷Department of Oncology, Cumming School of Medicine, University of Calgary, Canada

⁸Canada's Michael Smith Genome Sciences Centre, BC Cancer Agency, Vancouver, BC V5Z 4S6, Canada

⁹Department of Epidemiology and Biostatistics, Memorial Sloan Kettering Cancer Center, New York, New York 10065, USA

¹⁰Department of Bioinformatics and Computational Biology, The University of Texas MD Anderson Cancer Center, Houston, Texas 77030, USA

¹¹National Cancer Institute, Bethesda, Maryland 20892, USA

¹²Department of Pathology, Case Western Reserve University, Cleveland, Ohio 44106, USA

¹³Division of Gastroenterology and Hepatology, Mayo Clinic, Rochester, Minnesota 55905, USA

¹⁴Department of Pathology, Duke University, Durham, North Carolina 27710, USA

¹⁵Center for Translational Research, Department of Surgery, George Washington University Cancer Center, Washington DC 20052, USA

¹⁶University of Alabama at Birmingham, Birmingham, AL 35294, USA, and HudsonAlpha Institute for Biotechnology, Huntsville, Alabama, USA

¹⁷Department of Medical Oncology, Dana-Farber Cancer Institute, Boston, Massachusetts 02215, USA

¹⁸Department of Pathology, The University of Texas MD Anderson Cancer Center, Houston, Texas 77030, USA

¹⁹PharmiWeb Solutions, Bracknell, RG12 1QB, United Kingdom

²⁰Institute for Systems Biology, Seattle, Washington 98109, USA


²¹Center for Cancer Genome Discovery, Dana-Farber Cancer Institute, Boston, Massachusetts 02115, USA

SUMMARY

We analyzed 921 adenocarcinomas of esophagus, stomach, colon and rectum to examine shared and distinguishing molecular characteristics of gastrointestinal tract adenocarcinomas (GIAC). Hypermutated (HM) tumors were distinct regardless of cancer type and comprised those enriched for insertions/deletions, representing microsatellite instability cases with epigenetic silencing of *MLH1* in the context of CpG Island Methylator Phenotype (CIMP), plus tumors with elevated single nucleotide variants (HM-SNV) associated with mutations in *POLE*. Tumors with chromosomal instability (CIN) were diverse, with gastroesophageal adenocarcinomas harboring fragmented genomes associated with genomic doubling and distinct mutational signatures. We identified a group of tumors in the colon and rectum lacking hypermutation and aneuploidy termed Genome Stable (GS) and enriched in DNA hypermethylation and mutations in *KRAS*, *SOX9* and *PCBP1*.

Graphical Abstract

Liu et al. analyze 921 gastrointestinal (GI) tract adenocarcinomas and find that hypermutated tumors are enriched for insertions/deletions, upper GI tumors with chromosomal instability harbor fragmented genomes, and a group of genome stable colorectal tumors are enriched in mutations in *SOX9* and *PCBP1*.



Keywords

Cancer; Tumor; Genomic; Esophagus; Stomach; Colon; Rectum; Colorectal; Methylation; Epigenetic

INTRODUCTION

Traditional classifications of tumors have utilized tissue of origin and histologic types. These categories have been refined with comprehensive molecular characterizations across large numbers of tumors. Adenocarcinomas of the gastrointestinal tract share similar endodermal developmental origins and exposure to common insults that promote tumor formation. We sought to evaluate molecular characteristics that distinguish gastrointestinal tract adenocarcinomas (GIAC) from other cancers and to investigate the molecular features of GIAC across anatomic boundaries to provide insight into the pathogenesis of these deadly malignancies.

Approximately 1.4 million people die each year worldwide from adenocarcinomas of the esophagus, stomach, colon or rectum (Arnold et al., 2015; Torre et al., 2016). Non-surgical treatment approaches have made only modest progress over the past half-century, inspiring efforts to better understand the biological basis of these cancers as a foundation for improving prevention, screening and therapy. Prior studies that separately evaluated GIAC of the upper (gastroesophageal) and lower (colorectal) GI tract found subgroups such as

chromosomal instability (CIN), microsatellite instability (MSI) and tumors with hypermethylation phenotypes. However, systematic efforts to characterize how shared molecular processes present differently across the GI tract have not been undertaken.

RESULTS

The Cancer Genome Atlas Network obtained fresh frozen tissues from 921 primary GIAC (79 esophageal, 383 gastric, 341 colon, and 118 rectal cancers) without prior chemotherapy or radiotherapy. All patients provided informed consent, and collections were approved by local Institutional Review Boards. Adjacent non-malignant tissues were obtained from 76 patients. We characterized samples by single-nucleotide polymorphism (SNP) array profiling for somatic copy-number alterations (SCNA), whole-exome sequencing, array-based DNA methylation profiling, messenger RNA (mRNA) sequencing, microRNA sequencing, and for a subset of samples, reverse-phase protein array (RPPA) profiling. Key characteristics of tumor samples are summarized in Table S1.

Shared Features of GIAC

We investigated whether GIAC share characteristic molecular features compared to other adenocarcinomas (Table S2). Joint analysis of GIAC together with adenocarcinomas from breast (n=1001), endometrium (506), cervix (24), bile ducts (33), lung (240), pancreas (183), prostate (381) and ovaries (503) revealed that GIAC clustered together by DNA hypermethylation profiles (Figure S1A), mRNA (Figure S1B), and Reverse Phase Protein Array data (RPPA) (Figure S1C). These results are consistent with integrated clustering analysis across multiple platforms of 10,000 TCGA tumors, which identified GIAC as a distinct group (Hoadley et al., 2018).

Genes mutated significantly more frequently in GIAC compared to non-GI adenocarcinomas (non-GIAC) included *FBXW7*, *SMAD2*, *SOX9*, and *PCBP1* (Figure 1A and Table S3). A GIAC-focused analysis revealed that *ATM*, *PZP*, *CACNA1C*, and *FBN3* were significantly mutated genes not previously reported in TCGA studies of single cancer types (Figure S1D and Table S3). We evaluated SCNA data to identify amplifications and deletions more common in GIAC than in non-GIAC (Figures 1B and S1E and Table S4). Arm-level gain of chromosome 13q was GIAC-specific (Figure S1F), noteworthy as this region containing tumor suppressor *RBI* is often deleted in non-GIAC. *CDX2* (13q12.2) and *KLF5* (13q22.1) encoding two transcription factors in this amplified region may contribute to GIAC pathogenesis. Other genes preferentially amplified in GIAC included *CDK6* (7q21.2), *GATA6* (18q11.2), *GATA4* (8p23.1), *EGFR* (7p11.2), *CD44* (11p13), *BCL2L1* (20q11.21), *FGFR1* (8p11.22), and *IGF2* (11p15.5). *APC* and *SOX9* deletions were observed preferentially in GIAC, as were frequent mutations in these genes.

GIAC displayed markedly higher frequencies of CpG island hypermethylation than did non-GIAC (Figure 1C, upper graphs). This finding is attributable in part to the high frequency of CpG Island Methylator Phenotype (CIMP) in GIAC, but was also evident in non-CIMP tumors. The average density of somatic mutations was also higher in GIAC. Clusters of tumors with high mutation densities were observed in gastric and colorectal GIAC as well as in breast and uterine non-GIAC (Figure 1C, middle graphs). Frequent SCNA were observed

in all GIAC, especially in esophageal adenocarcinomas (EAC), and ovarian and a subset of breast non-GIAC (Figure 1C, bottom graphs).

Gene expression analysis revealed 553 genes that were differentially expressed in GIAC compared to non-GIAC, after exclusion of genes that differed among corresponding normal tissues (Figure S1G and Table S5). Supervised multivariate orthogonal partial least squares-discriminant analysis ranked 51 of these 553 genes to have significantly higher expression in GIAC. Notably, these genes include several that have roles in gastrointestinal stem cell biology (e.g. *OLFM4*, *CD44*, and *KLF4*) and genes related to the EGFR signaling pathway (Figure S1G).

We next investigated whether genes encoding 139 transcription factors (TFs) that are important in GI development (Noah et al., 2011; Sherwood et al., 2009) displayed distinct gain- or loss-of-function events in GIAC compared to non-GIAC. Amplifications were considered gain-of-function (GOF) events, while deletions, epigenetic silencing, and nonsense or indel mutations were considered loss-of-function (LOF) events (Table S6). We found 33 transcription factor genes with GOF or LOF exceeding 5% in at least one GIAC tumor type (Figure 1D). *CDX2* encodes a homeobox transcription factor expressed early in endoderm development with evidence as either a lineage-survival oncogene (Salari et al., 2012) or a tumor-suppressor gene (Bonhomme et al., 2003) in colorectal cancers (CRC), depending on context, and is also a marker of intestinal metaplasia in Barrett's esophagus (Moons et al., 2004). Interestingly, we observed *CDX2* amplification in esophageal, colon, and rectal adenocarcinomas, but LOF in gastric cancers. Although amplifications in the genomic loci containing the stem-cell transcription factor *KLF5* gene were found in all GIAC, these amplifications were associated with increased stemness only in EAC based on a gene-expression signature (Malta et al., 2018) (Figure S1H).

Molecular Subtypes within GIAC

Other studies have relied on gene expression, oncogenic pathway, or histopathological criteria for subtype delineation among GIAC (Budinska et al., 2013; Cristescu et al., 2015; Dienstmann et al., 2017; Guinney et al., 2015; Roepman et al., 2014; Tan et al., 2011). We found that unsupervised clustering of GIAC using mRNA, miRNA, and RPPA data was strongly influenced by tissue type, thus complicating defining molecular groups spanning anatomic boundaries. By contrast, evaluation of mutations, copy-number alterations, and DNA methylation patterns yielded tumor subtypes spanning tissue boundaries (Figure S1A). Our subgroups are consistent with those identified by recent genomic research across GIAC (Cancer Genome Atlas Research Network, 2012, 2014, 2017; Cristescu et al., 2015; Secrier et al., 2016; Wang et al., 2014), and rely on molecular features generally evaluable by the clinical community.

A subgroup of tumors was characterized by a high Epstein-Barr Virus (EBV) burden, as previously determined via mRNA and miRNA analysis (Cancer Genome Atlas Research Network, 2014) (Figure 2A). EBV⁺ tumors, found only in the stomach (n = 30), display the most extensive hypermethylation of any tumor type in TCGA (see Figure S4.6 in (Cancer Genome Atlas Research Network, 2014). Hypermutated tumors (n = 157), defined by mutation density > 10 per megabase (Mb) (Figure S2A) were further substratified based on

the implied mechanism of replication error. MSI, arising from defective DNA mismatch repair, often yields insertion-deletion (indel) mutations in addition to single nucleotide variants (SNV) (Sia et al., 1997), whereas hotspot mutations in polymerase E (*POLE*) are associated with SNV-predominant profiles (Cancer Genome Atlas Research Network, 2012; Palles et al., 2013; Zhou et al., 2009) (Figure 2B). Hypermuted samples with an indel density of > 1 per Mb and an indel/SNV ratio > 1/150 consisted of almost all tumors with clinically-defined MSI (MSI, n = 138; 54% gastroesophageal or GE; 46% colorectal or CR) (Figure S2B). All other hypermutated samples were categorized as Hypermuted-SNV (HM-SNV, n = 19 (n = 11 with *POLE* mutations), (47% GE; 53% CR) (Figure 2B and S2B). The remaining two groups were distinguished by presence or absence of extensive SCNA (Figure S2C). Chromosomal instability (CIN) tumors (n = 625, 48% GE; 52% CR) exhibited marked aneuploidy, defined by a clonal deletion score (CDS, See STAR Methods) > 0.0249, which is largely determined by chromosome- and arm-level losses. By contrast, Genome stable (GS) n = 109, 47% GE; 53% CR) samples lacked such aneuploidy (Figure 2B and S2B).

We evaluated the relationship between our molecular subtypes and consensus molecular subtypes (CMS), which have been established for CRC based primarily on gene expression (Guinney et al., 2015). We applied the CMS classification system to the lower GI tumors in our study and found a significant association between the two groupings ($p < 2.2 \times 10^{-16}$), but with noteworthy differences (Figure S2D). The CMS1 - MSI Immune grouping did not discriminate MSI tumors from the HM-SNV tumors (Figure S2B). A substantial fraction of GS CRC were represented in the CMS3 - Metabolic subtype ($p = 1.6 \times 10^{-6}$), but the CMS subtype system appeared to be largely unable to distinguish CIN and GS (Figure S2D).

Our molecular groupings also correlated with key immune features of GIAC (Figure 2C and S2E). As previously reported, EBV⁺ tumors possessed the highest gene expression scores for CD8 T-cells, M1-macrophages, and IFN γ signatures (Figure 2C and S2E) (Derks et al., 2016; Koh et al., 2017). MSI tumors showed the next greatest IFN γ signature, consistent with reported immunogenicity of MSI tumors (Guinney et al., 2015). Moreover, MSI tumors displayed diverse immune signatures depending on their tissue of origin (Figures S2F and S2G); for example, checkpoint protein CD276 was significantly enriched in MSI CRC whereas *ENTPDI* was preferentially expressed in MSI gastroesophageal adenocarcinomas (GEA) (Figure S2G). HM-SNV also demonstrated heterogeneity in immune signature expression when comparing the upper and lower GI tract (Figure S2F). Of translational importance, an attenuation in HLA/antigen presentation (Figure S2F) and significant elevation in NK-cell gene expression was found in HM-SNV CRC (Figure S2H), suggesting that NK cells are found in a subset of tumors and are capable of anti-tumor responses (Wagner et al., 2017). The cytotoxic activity of NK cells is finely regulated by the integration of activating and inhibitor cues (Ljunggren and Malmberg, 2007), and cells lacking MHC expression often are subjected to NK cell cytotoxicity due to the absence of inhibitory cues mediated by killer-cell immunoglobulin-like receptor (KIR). These data suggest agents to enhance NK activity may be a therapeutic option for HM-SNV tumors.

Unsupervised clustering of DNA methylation data across GIAC using cancer-associated methylated sites (excluding CpGs with tissue-specific methylation) revealed extensive CpG

island methylation in the EBV⁺ gastric cancers, distinguishing these tumors (Figure 2D), as previously noted by us and others (Cancer Genome Atlas Research Network, 2014; Matsusaka et al., 2011; Wang et al., 2014). The remaining tumors could be characterized as those lacking a CIMP phenotype (Non-CIMP) and those displaying a low or high frequency of DNA hypermethylation (CIMP-L and CIMP-H, respectively).

Hypermutated tumors were located primarily in the central part of the GI tract, the distal stomach and proximal colon, whereas CIN tumors were more prevalent in the anatomic extremes, the esophagus and distal colon/rectum (Figure 2E) (Budinska et al., 2013). Although CIMP-H occurred throughout the upper GI tract and proximal colon (Figure 2F), epigenetic silencing of *MLH1*, responsible for MSI, was observed primarily in the distal stomach and proximal colon (Figure 2G). Within the proximal stomach and esophagus, only 4/29 (14%) of CIMP-H tumors exhibited *MLH1* epigenetic silencing and MSI, while 23 of the 29 (79%) were MSS and displayed the CIN phenotype (Figure 2H). In the lower GI tract, CIMP-H and MSI were largely absent in the descending colon and rectum (Figure 2E, 2H).

Analysis of Hypermethylation and Hypermutation

MSI tumors exhibited distinct expression features independent of tissue of origin, implying common biological features of this class of tumors (Figure S3A, S3B). Most sporadic MSI cases in both colorectal and gastric cancer arise as a consequence of epigenetic silencing of *MLH1* by promoter DNA hypermethylation (Herman et al., 1998; Leung et al., 1999) in the context of CIMP-H (Weisenberger et al., 2006). MSI tumors with *MLH1* methylation were associated with *BRAF*^{V600E} mutation only in the colon, not the stomach (Figure 3A). *KRAS* mutations were found primarily in CIMP-L tumors of the lower GI tract, whereas *KRAS* amplification was observed in upper GI tumors (Figure 3A). *TFAP2E* promoter methylation, which is associated with non-response to chemotherapy in colorectal cancer (Ebert et al., 2012), was found in a substantial fraction of CIMP-H tumors and in almost all EBV⁺ gastric cancers (Figure 3A). CIMP-H tumors showed near-ubiquitous methylation of the tumor suppressor *CDKN2A* in gastric and colon MSI tumors (Figure 3A and 3B). However, 39% of the CIMP-H tumors lacked *MLH1* silencing and MSI and instead included other classes of GIAC, most commonly CIN tumors in the proximal stomach/esophagus or rectum/descending colon (Figure 2H, 3B).

Given the tight associations between CIMP-H and MSI and their heterogeneity across anatomic boundaries, we studied the collection of tumors containing either of these features in more detail (Figure 3B). A portion of MSI cases lacking both *MLH1* methylation and the CIMP phenotype contained somatic mutations in *MLH1* or *MSH2*, indicating an alternative route to loss of DNA mismatch repair (Figure 3B, right side). These tumors were preferentially associated with mutations in *KRAS* rather than *BRAF*. A small number of MSI tumors (n=8) could not be explained by genetic or epigenetic inactivation of a mismatch repair gene.

Broadly, the MSI group of CRC harbored lower WNT signatures than did other CRC (Figure S3C, S3D), a finding that may be attributable to a reduced reliance of CIMP-H tumors on WNT signaling. Among MSI CRC, those arising in the context of CIMP-H have a lower percentage of *APC* mutation (28%) than those arising in either CIMP-L (78%) or non-

CIMP (58%) (Fisher Exact $p = 0.0091$). This finding holds true for MSS CIMP-H tumors as well, and is discussed in the GS subtype section below. CIMP-H MSI CRC showed a reduced combined frequency of either *APC* or *CTNNB1* mutations and decreased WNT gene expression signatures compared to non-CIMP-H MSI cases, and were more similar to upper GI MSI tumors in their lower reliance on WNT activation (Figure 3C). Despite the reduced frequency of *APC* and *CTNNB1* mutations, MSI CIMP-H tumors displayed overall greater mutational densities and arose at an older age of onset than did non-CIMP-H MSI cases or upper GI MSI cases (Figure 3C).

We investigated the genes silenced by promoter hypermethylation in the molecular subgroups (Figure 3D, 3E, and Table S7). Pathway analysis of epigenetically silenced genes among all subgroups revealed enrichment for genes encoding DNA binding proteins and transcription factors, consistent with previous findings of enrichment for stem-cell Polycomb Target Genes (Widschwendter et al., 2007). We identified 135 genes silenced in at least 25% of the upper or lower GI MSI tumors and compared their relative frequency of silencing and frequency of several key gene mutations (Figure 3E). *HUNK*, a negative regulator of intestinal cell proliferation (Reed et al., 2015), was found to be frequently silenced in MSI tumors. Another frequently silenced gene, *ELOVL5*, lies within the locus with germline variants most significantly linked to survival of CRC patients (Phipps et al., 2016).

Molecular Features of the CIN Subtype

The landscape of SCNAs revealed a more finely fragmented genome in GEA compared to CRC, despite an overall similar pattern of affected regions of the genome (Figure 4A). Evaluation of SCNA distribution, categorized by both focality and intensity, revealed higher prevalence of focal copy-number events within the CIN GEA population (Figure 4B and S4A). The difference between upper and lower GI was greater for focal amplifications than for deletions (Figure 4B), primarily evident in high-amplitude focal amplifications (Figure S4A). We developed a score that captures the quantity and intensity of focal high-level amplicons (see STAR Methods). CIN tumors with a higher score were designated CIN-Focal (CIN-F) whereas those with a lower score, and therefore low-amplitude, broader amplicons, were called CIN-Broad (CIN-B) (Figure 4C). The distribution of these two classes of CIN differed between upper and lower GIAC, with CIN GEA displaying 74% CIN-F and 26% CIN-B, and CRC showing reversed proportions consisting of 22% CIN-F and 78% CIN-B (Figure 4C). Despite this difference between upper and lower GI tumors, the ratios of CIN-B/CIN-F did not vary anatomically within upper GI tumors or within the lower GI tract tumors (Figure S4B). Notably, in addition to the higher prevalence of CIN-F in upper GIAC, such CIN-F GEA displayed a higher intensity in the focal-amplification score compared to their CIN-F CRC counterparts (Figure S4C). CIN-F GEA was associated with advanced tumor stage, underscoring its potential clinical significance (Figure 4D).

Although CIMP frequency displayed an anatomic gradient within upper GI (Figure S4D), we found no correlation of CIMP class with arm-level or focal SCNA in CIN (Figure S4E). CIN-F GEA demonstrated significantly more whole-genome duplication (WGD) than did CIN-B GEA, 68% vs. 42% (Figure 4E and S4F), with evidence of two or more genome doublings (WGD2) in 18% of CIN-F, compared to 7% of CIN-B in upper GI CIN tumors.

WGD2 was associated with poor survival in GEA, independent of age and stage (Figure S4G). However, the strong association of genomic doubling and CIN-F was not observed in CRC, despite similar total rates of genome duplication (Figure S4H; 59% in lower GI and 61% in upper GI).

CIN-F GEA sustained significantly more frequent focal amplification of genes encoding receptor tyrosine kinases (RTKs), *KRAS* and cell cycle mediators (Figure 4F). In contrast, CIN-B GEA more commonly sustained activating mutations of oncogenes (e.g. *KRAS* and *ERBB2*) than did GEA-CIN-F tumors (Figure 4F). *ERBB2* amplifications significantly co-occurred with *CCNE1* amplifications ($p = 0.039$) and trended towards co-occurrence with gains in chr. 20q/*SRC* ($p = 0.0692$). Intriguingly, activating mutations in *ERBB2* co-occurred with *ERBB2* amplifications ($p = 0.0087$). CIN-B GEA harbored more frequent somatic inactivation of tumor suppressors related to cell cycle regulation (e.g. *CDKN2A*), WNT-pathway activation (e.g. *APC*), and TGF β regulation (e.g. *SMAD2* and *SMAD4*) than CIN GEA-F. By contrast, CIN-F GEA showed a higher frequency of *TP53* mutations (Figure 4F; 76% vs. 54%) and higher rates of oncogene amplifications (Figure 4G).

Among lower GI CIN tumors, the differences in somatic mutations and copy-number alterations found in CIN-F and CIN-B tumors were modest (Figure S4I), although CIN-F did associate with poorer survival in CRC (Cox regression $p = 0.0053$, adjusted for stage, age, and molecular subtype). We identified amplifications including *CDX2*, *ERBB2*, and *CCND2* enriched in these tumors. Consistent with the different patterns of CIN between upper and lower GI cancers, we found that *ERBB2*⁺ CRC not only harbor lower CIN-F scores (Figure S4J) but also fewer co-occurring genomic alterations than *ERBB2*⁺ GEA (Figure S4K). These findings are consistent with efficacy in CRC of *ERBB2* therapy without chemotherapy (Sartore-Bianchi et al., 2016), compared to *ERBB2*⁺ GEA, which often carry co-occurring amplified oncogenes implicated in *de novo* resistance (Janjigian et al., 2018; Kim et al., 2014; Sartore-Bianchi et al., 2016).

CIN-B and CIN-F CRC displayed comparable rates of *APC* and *KRAS* mutations (Figure S4I; *APC*: 79% vs 87%; *KRAS*: 35 vs 44%). However, *PIK3CA* mutations and TGF β pathway alterations were more common in CIN-B CRC than in CIN-F CRC (Figure S4I). Both groups of CIN CRC had somatic patterns more closely resembling the CIN-B GEA group, in which oncogenes were activated more commonly by mutation than by amplification. These data suggest that the preponderance of early *APC* loss and selection for mutational activation of oncogenes like *KRAS* may precede a form of aneuploidy and transformation distinct from the catastrophic aneuploidy and resulting oncogene amplification occurring in GEA (Figure 4G).

Among CIN CRC, we observed more frequent CIMP, primarily CIMP-L, in proximal, right-sided CIN tumors and less frequent CIMP in distal, left-sided ones (Figure S5A). Arm-level SCNAs were significantly less frequent in CIMP⁺ CIN CRC (Wilcoxon $p = 2.7 \times 10^{-9}$), despite the lack of an overall difference in focal alterations (Figure S5B). Among chromosome arms, gain of 20q was most enriched in non-CIMP CIN CRC, with a mean copy-number gain of 1.8 (ploidy-adjusted), compared to 1.1 in CIMP⁺ CIN CRC (Figure S5C). By contrast, except for *TP53*, which was more frequently mutated in non-CIMP CIN

tumors, the frequency of somatic mutations was significantly higher in CIMP⁺ CIN CRC (Figure S5D), notably affecting the TGF β pathway and key oncogenes including *KRAS*/*NRAS*/*BRAF* and *PIK3CA*. Dichotomizing CIN CRC tumors by CIMP status thus showed parallels to the division of upper GI CIN tumors by CIN-F/CIN-B status. CIMP⁺ CIN CRC, like CIN-B GEA, harbored more oncogene mutations (Figure 5A). Taken together, these data suggest that CIMP status may play an important role in shaping evolution of CIN tumors in the lower GI tract, and to a lesser extent in the upper GI tract.

Molecular Features of the GS Subtype

Although CRC are classically divided between hypermutated/MSI and CIN (Bijlsma et al., 2017), we detected a population of CRC lacking both aneuploidy/CIN and hypermutation, a group we classified as GS (Figure 5B). Unlike with MSI, these GS CRC shared few underlying biologic features with GS in upper GIAC. As we reported earlier (Cancer Genome Atlas Research Network, 2014), upper GI GS tumors are enriched for the diffuse-type gastric cancer (65.7%) and commonly harbor mutations in *CDH1* and *RHOA* (Figure S5E). Thus, upper GI GS, like EBV⁺ tumors, represent an essentially unique entity confined to stomach.

GS CRC shared features of other CRC; like the CIN CRC, GS CRC shared a predilection for loss of *APC* (GS 81% vs CIN 85%, Figure S5F). GS CRC were more common in ascending and transverse colon (Figure 2E) and when compared to the CIN CRC, showed significant enrichment for the CIMP-L phenotype (79% vs 40%, $p = 1.2 \times 10^{-9}$, Figure 5B) and for the CMS3 metabolic consensus molecular subtype ($p = 1.6 \times 10^{-6}$, Figure S2D) (Guinney et al., 2015). Despite having fewer SCNAs, a subset of GS CRC showed amplifications of *IGF2* (q value < 0.05) (Figure S5G). MAPK pathway mutations were more common in these tumors, with *KRAS*, *NRAS* or *BRAF* mutated in 69%, 10%, and 9% of tumors, respectively, and with *PIK3CA* mutations present in 43%, compared to 22% of CIN CRC (Figure 5A). Consistent with the relative lack of aneuploidy, *TP53* mutations were less common (16%) in GS compared to CIN tumors (80%) (Figure S5E). However, we observed enrichment for somatic mutations in *SOX9*, which encodes a transcription factor, and in *PCBP1*, which encodes an RNA-binding protein that regulates splicing, mRNA stability, and translation (Leffers et al., 1995) (Figure 5A, 5C, S5H). *SOX9*, mutated in 29% of GS CRC, encodes a WNT-regulated transcription factor that controls cell fate and crypt homeostasis in intestinal development (McConnell et al., 2011; Nandan et al., 2014). GS CRC with mutations in *SOX9* also had more frequent somatic mutations in the TGF β pathway genes, including *PCBP1* (Figure S5I). Our mutation analysis within GS revealed highly clustered missense mutations in the KH domain of *PCBP1* in 13% of GS CRC, raising the potential for a GOF event (Figure 5C). Interestingly, overexpression of wild-type *PCBP1* was associated with oxaliplatin resistance in CRC (Guo et al., 2010).

Overall, GS CRC had more frequent mutations in the TGF β pathway, RAS/RAF genes, and *PIK3CA* than did CIN CRC (Figure S5F). Comparison of GS CRC to CIN CRC revealed a progressive gradation of features between non-CIMP CIN CRC, CIMP-H or CIMP-L CIN CRC and GS CRC (Figure 5A). These data suggest a pathway to cancer in the colorectum in which APC-mutant cells, typically containing the CIMP-L phenotype, are able to undergo

transformation by sustaining additional pathogenic mutations without the need for p53 loss or aneuploidy (Figure 5D).

We had noted earlier that CIMP-H MSI tumors appeared to rely less on WNT signaling. CIMP-H MSS tumors also display reduced rates of *APC* mutation (47%) compared to CIMP-L (87%) or non-CIMP (86%) (Fisher Exact $p = 0.00066$). These findings suggest an alternative CRC pathway that is not initiated by mutation of *APC*, but rather by an epigenetic aberration causing CIMP-H. If *MLH1* is silenced in the context of CIMP-H, then the tumor would become MSI, whereas if *MLH1* is not affected, the tumor would develop along the CIN pathway to give rise to CIMP-H MSS CIN tumors (Figure 5D). Non-hypermuted CRC from the right-sided (ascending/transverse) colon revealed significantly higher rates of *KRAS*, *PIK3CA* and *SOX9* mutation than those from the left-sided (descending) colon/rectum (Figure 5E).

Mutational Signatures in GIAC

MSI and POLE signatures dominated the total mutational signature scores among GIAC as a consequence of the high mutational burden in MSI and POLE-deficient tumors in the MSI and lower-GI HM-SNV groups, respectively (Figures S6A and S6B). Signature discovery following removal of hypermutated cases revealed a BRCA signature (COSMIC signature 3), two APOBEC signatures, a signature resembling COSMIC signature 17 with common AA>AC transversions, and a signature dominated by C>T transitions at CpG dinucleotides (COSMIC signature 1) (Figures 6A, 6B, 6C, 6D, 6E and 6F) (Alexandrov et al., 2013; Bignell et al., 2010). The APOBEC signatures contributed minimally to the mutational profile across GIAC (Figures 6B, 6C, S6C and S6D), but the other three signatures had substantial activity in non-hypermuted GIAC with the AA>AC signature limited to upper GIAC (Figures 6B, 6C, 6E, and S6D). A recent study, identified the existence of the BRCA signature in gastric cancers that lacked mutations in *BRCA1* and *BRCA2* (Alexandrov et al., 2015). We confirmed the presence of BRCA signature activity in GIAC, with significant enrichment of somatic and germline mutations in several homologous recombination genes such as *BRCA1*, *BRCA2* and *PALB2* (Figure S6E). BRCA signature activity was also significantly enriched in tumors with epigenetic silencing of *BRCA1* or *RAD51C*, including within EBV⁺ GCs (Figure S6F). We observed a significant association between BRCA signature activity and upper GI cancers, particularly the CIN subtype (Figure 6D). The BRCA signature was associated only with focal SCNA events (Figure S6G), which are likely initiated by double-strand breaks. The AA>AC signature was also enriched in upper-GI CIN (Figures 6E, S6C and S6D), most notably in the tubular esophagus (Figure S6D). Moreover, this mutational signature was enriched in CIN-F and *TP53*-mutated upper GI CIN tumors (Figure 6E). The AA>AC signature lacks a known etiology, but its association with GEA and its correlation with higher CIN-F scores raises the possibility that this signature reflects a process that contributes to greater focal aneuploidy observed in GEA compared to CRC and differences in oncogene profiles between upper and lower GIAC (Figure 7).

The CpG>TpG pattern, often termed the “aging signature”, was the most common signature among all tumors, but it was especially frequent in right-sided CRC (Figure S6C). This signature is thought to arise from spontaneous hydrolytic deamination of 5-methylcytosine,

and is consolidated as a persistent mutation if it occurs during DNA replication. Hence, this signature tracks the cumulative number of cell divisions and aging. Although we observed an association with CIMP status (Figure 6F), we do not believe that this is explained by a simple quantitative difference in DNA methylation. The CIMP hypermethylation is measured primarily at promoter CpG islands, which are unmethylated in normal cells and thus do not sustain many CpG>TpG mutations prior to acquisition of methylation and clonal expansion, whereas the mutation signatures were obtained by exome sequencing of gene bodies, which are generally highly methylated. The association between CIMP status and CpG>TpG signature may reflect the fact that CIMP tumors require more cell divisions to progress and thus acquire more CpG>TpG mutations over time.

DISCUSSION

GIAC originate from columnar epithelium with a shared endodermal origin and display a spectrum of common molecular features such as aneuploidy and microsatellite instability that span anatomic boundaries. GIAC are enriched for activation of the WNT signaling pathway, particularly in the lower GI tract, consistent with the importance of WNT in GI development (Schepers and Clevers, 2012). We found that CIMP-H CRC appeared less dependent on canonical WNT signaling mutations and pathways. GIAC also displayed a predisposition for disruptions in TGF β and SMAD signaling components. TGF β signaling helps to maintain intestinal stem cell equilibrium, promoting growth during development, but controlling self-renewal in adult epithelium (Mishra et al., 2005).

The vast majority of sporadic MSI tumors arise as a consequence of promoter methylation of *MLH1* in the context of CIMP-H. Methylation profiles of CIMP-H tumors are quite consistent throughout the GI tract. However, *MLH1* silencing within CIMP-H is much more anatomically restricted, primarily observed in the distal stomach and proximal (ascending and transverse) colon, but notably uncommon in proximal upper GI cancers. The epithelia of the distal stomach and proximal colon appear more susceptible to oncogenic effects of *MLH1* silencing. High rates of epithelial cell turnover with accompanying DNA replication may more effectively consolidate replication-associated errors in these sections of the GI tract. This hypothesis is consistent with the tumor spectrum observed with germline mutations in mismatch repair genes, leading to increased risk of cancers arising in highly proliferative tissues (Lynch et al., 2015). In this scenario, stochastic promoter methylation of *MLH1* from CIMP-H would provide less selective advantage when arising in the less proliferative sections of the GI tract.

CIMP-H GIAC possessed other differences in molecular features between various anatomic locations. *BRAF*^{V600E} mutations occurred almost exclusively in CIMP-H tumors of the ascending colon and were absent from otherwise similar CIMP-H GEAs. In addition, some colorectal CIMP-H tumors with similar DNA methylation profiles lacked *BRAF*^{V600E} mutations, a finding inconsistent with the proposed role for *BRAF*^{V600E} mutation as a cause of CIMP-H (Fang et al., 2016). Alternatively, CIMP may provide a permissive environment for *BRAF*^{V600E} mutation, perhaps by silencing pathways involved in oncogene-induced senescence and apoptosis (Hinoue et al., 2009). Despite the large overlap of CIMP-H and MSI in GIAC, our data revealed that this co-occurrence occurs predominantly in the distal

stomach and ascending colon. The etiology for CIMP-H tumors commonly progressing via a CIN pathway in proximal GE and distal CRC is not established.

CIN is a common feature of GIAC and other tumors (Cancer Genome Atlas Research Network, 2012, 2014, 2017; Hoadley et al., 2014). Despite the deleterious effect on cellular and organismal fitness (Sheltzer et al., 2011; Sheltzer et al., 2017; Torres et al., 2007; Williams et al., 2008), CIN with its resultant aneuploidy remains the predominant molecular subtype among GIAC, found most frequently in the proximal upper and distal lower GI tract (Dulak et al., 2012). Unlike tumors with MSI, CIN tumors had more discrepant molecular features between upper and lower GI cancers. Most striking was the preponderance of focal, high amplitude SCNAs, especially amplifications, in GEA. Within CIN GEA, we found that tumors with high CIN-F scores had a strong association with prior genome doubling, a process associated with CIN (Ganem et al., 2007). Amplifications in CIN-F GEA commonly targeted mitogen pathway components, cell cycle regulators, and lineage survival transcription factors, whereas CIN-B and GS tumors more frequently carried activating mutations in these pathways.

A notable finding was the predilection in CIN-B GEA for alterations in tumor suppressors such as *CDKN2A*, *APC* and *SMAD4*. These findings suggest that the marked aneuploidy found within the CIN-F GEA is less apt to occur in precursors with pathogenic alterations other than *TP53*. One explanation is that precursors with already altered oncogenes/tumor suppressors have less requirement for more ‘catastrophic’ aneuploidy to simultaneously abrogate multiple such checkpoints. By contrast, such marked instability could facilitate transformation in precursors with p53 loss without as many other preexisting pathogenic alterations. Indeed, although p53 loss alone is not sufficient to promote aneuploidy (Bunz et al., 2002), several lines of evidence support its necessity, most likely by circumventing p53-dependent cell cycle arrest in response to damage by reactive oxygen species (ROS) (Guo et al., 2010), to mutations in ataxia telangiectasia (*ATM*) (Li et al., 2010), or to spindle assembly checkpoint (SAC) activation (Thompson and Compton, 2010). Given these data, the lesser rates of CIN-F in lower GI CIN tumors (compared to CIN tumors of the upper GI tract) may be a consequence of *APC* loss as an early event in colorectal neoplasia, thus leading to *TP53* mutation rarely occurring in the absence of a prior *APC* loss. Instead, we noted that CIMP status likely has a stronger influence on the features of CIN CRC, with CIMP being associated with mutations in *KRAS* and in tumor suppressor pathways such as TGF β . Aneuploid CIMP tumors in the lower GI tract showed lower rates of SCNA, but a greater number of oncogenic mutations compared to non-CIMP. Both upper and lower GI CIN tumors were also associated with the BRCA mutational signature. However, the propensity for greater CIN-F in upper GIAC correlated with the AA>AC mutational signature, a signature of unknown etiology, previously reported in upper GI tumors (Dulak et al., 2013).

Our exploration of the role of CIMP in shaping the features of CIN in CRC became linked with our finding of a GS CRC subtype falling outside the classic CIN/MSI CRC dichotomy. This GS subtype may partially overlap with the previously identified Microsatellite and Chromosome Stable (MACS) CRC (Chan et al., 2001), while showing important differences. The MACS phenotype is an independent predictor of poor outcome (Banerjee et

al., 2009), in contrast, GS CRC are enriched for earlier stage tumors. MACS tumors have an elevated proportion of early onset cases (Chan et al., 2001), whereas GS CRC have a higher mean age at diagnosis than CIN cases. Like MACS, HM-SNV cases are microsatellite and genome stable, and also arise in younger patients, so it is possible that some early-onset MACS tumors may have represented unrecognized HM-SNV tumors. The GS CRC overlap with a subgroup identified by gene expression clustering as CMS3 (Guinney et al., 2015) and commonly displaying CIMP-L. Many features enriched in CIMP CIN CRC compared to non-CIMP CIN CRC were even more prevalent in GS CRC. Moreover, we found these tumors to have recurrent mutations in *SOX9* and *PCBP1*. While the presence of frameshift mutations of *SOX9* implies LOF, truncating mutations in *SOX9* are overexpressed in primary tumor specimens (Javier et al., 2016), making their functional significance unclear. GS CRC with mutations in *SOX9* also had more frequent somatic mutations in TGF β pathway genes, including *PCBP1*, which impacts TGF β signaling by regulating Smad3-associated alternative splicing (Tripathi et al., 2016). Given the strikingly low frequency of *TP53* mutations in GS CRC, the presence of *SOX9* and *PCBP1* mutations may cooperate with *APC* and *KRAS* mutation to facilitate transformation, despite lack of hypermutation and low levels of aneuploidy.

Our findings also bear some relevance to the evolving field of immunotherapy, which already has established efficacy in MSI tumors. The HM-SNV tumors, which display a large SNV burden in the setting of *POLE* mutations, did not harbor the equivalent CD8 or IFN γ signatures as did the MSI tumors, perhaps suggesting that indel mutations may better generate neoantigens than do SNVs. The strong signatures in EBV⁺ tumors suggest a potential for immune checkpoint inhibition in this subset. The reason for consistently higher IFN γ signatures in upper GI compared to lower GI adenocarcinomas when stratified by molecular subtype is less obvious and may simply indicate that GEA are more immunogenic than CRC, results consistent with the presence of clinical responses to PD1 inhibitor monotherapy in MSS GEA, but not in CRC (Jin and Yoon, 2016; Muro et al., 2016).

In summary, these results highlight how processes such as DNA hypermethylation and CIN can manifest themselves in different ways across related tissues. In some instances, as with DNA hypermethylation in upper-GI vs. lower-GI MSI tumors, such differences can be subtle. However, as the exploration of CIN indicates, other processes can lead to substantially different molecular outcomes across these regions. Provision of greater detail in the various manifestations of molecular defects may reveal new opportunities for targeted therapies for these cancers. Furthermore, these data highlight how consideration of molecular subtypes as well as organ of origin will be essential in the study and treatment of cancer.

STAR METHODS

CONTACT FOR RESOURCE SHARING

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Peter W. Laird (Peter.Laird@vai.org). Sequence data hosted at the GDC is under controlled access. Details for gaining access can be found at (<https://gdc.cancer.gov/access-data/data-access-processes-and-tools>).

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Human Subjects and Tumor Data Selection—Molecular data were obtained as part of the Cancer Genome Atlas Project, from patients untreated by chemo- or radiation therapy and who provided informed consent; tissue collection was approved by the local Institutional Review Boards (IRBs) as noted below. GIAC cases (n=921) were selected as follows. Of the 559 Upper GI cases (171 ESCA and 388 STAD) in (Cancer Genome Atlas Research Network, 2017), 90 were excluded as ESCC and two as undifferentiated (TCGA-2H-A9GQ, TCGA-VR-A8Q7). Of the remaining 467 Upper GI adenocarcinomas, 462 (79 ESCA, 383 STAD) cases have molecular data available from the five TCGA core platforms (RNASeq, miRNASeq, DNA Methylation, SNP6, and mutation calls). We used germline DNA from blood or nonmalignant gastrointestinal tissue as a reference for detecting somatic alterations. For lower GI, all available TCGA COAD and READ cases were considered, but cases bearing the BCR annotation “Redacted” were excluded, as were cases with Notification: ‘Unacceptable Prior Treatment’ or ‘Item does not meet study protocol’. Review of COAD and READ pathology reports led to the exclusion of three additional COAD cases from this study (TCGA-AA-A022: Pathology report indicates poorly-differentiated carcinoma of the neuroendocrine type; TCGA-AA-A02R: Pathology report shows poorly-differentiated carcinoma with positivity for both S-100 and chromogranin, and focal synaptophysin; and TCGA-AZ-6607: Pathology report indicates this is likely to be a pancreaticobiliary primary tumor metastasizing to colon. The remaining 459 lower GI cases (341 COAD and 118 READ) with molecular data available for the five platforms were retained.

A group of 2,871 non-GIAC cases was constructed from TCGA tumor types BRCA, CESC, CHOL, LUAD, OV, PAAD, PRAD and UCEC, comprising all cases meeting the established criteria of the PanCancer Atlas Consortium (exclusion of Redacted, ‘Unacceptable Prior Treatment’ or ‘Item does not meet study protocol’ and cases with no molecular data). For BRCA, CHOL, PRAD, and OV, and UCEC cases annotated as problematic by Expert Pathology Review (marked as `AWG_excluded_because_of_pathology` in the PanCancerAtlas Merged Annotation File) were excluded. For CESC, LUAD, and PAAD, further exclusions were made based on case review, as follows: CESC, retain only adenocarcinomas; LUAD, exclude samples without histology; PAAD, exclude samples with cellularity < 20%.

Demographic data for patients are as follows: GIAC (60.3% male, median age 68 years, range 29–90 years); Non-GIAC (21.3% male; median age 61 years, range 25 to 90 years).

TCGA Project Management collected necessary human subjects documentation to ensure the project complies with 45-CFR-46 (the “Common Rule”). The program has obtained documentation from every contributing clinical site to verify that IRB approval has been obtained to participate in TCGA. Such documented approval may include one or more of the following:

- An IRB-approved protocol with Informed Consent specific to TCGA or a substantially similar program. In the latter case, if the protocol was not TCGA-specific, the clinical site PI provided a further finding from the IRB that the already-approved protocol is sufficient to participate in TCGA.

- A TCGA-specific IRB waiver has been granted.
- A TCGA-specific letter that the IRB considers one of the exemptions in 45-CFR-46 applicable. The two most common exemptions cited were that the research falls under 46.102(f)(2) or 46.101(b)(4). Both exempt requirements for informed consent, because the received data and material do not contain directly identifiable private information.
- A TCGA-specific letter that the IRB does not consider the use of these data and materials to be human subjects research. This was most common for collections in which the donors were deceased.

METHOD DETAILS

Sample Processing—RNA and DNA were extracted from tumor and adjacent normal tissue specimens using a modification of the DNA/RNA AllPrep kit (Qiagen). The flow-through from the Qiagen DNA column was processed using a mirVana miRNA Isolation Kit (Ambion). This latter step generated RNA preparations that included RNA <200 nt suitable for miRNA analysis. DNA was extracted from blood using the QiaAmp blood midi kit (Qiagen). Each specimen was quantified by measuring Abs260 with a UV spectrophotometer or by PicoGreen assay. DNA specimens were resolved by 1% agarose gel electrophoresis to confirm high molecular weight fragments. A custom Sequenom SNP panel or the AmpFISTR Identifier (Applied Biosystems) was utilized to verify tumor DNA and germline DNA were derived from the same patient. Five hundred nanograms of each tumor and normal DNA were sent to Qiagen for REPLI-g whole genome amplification using a 100 µg reaction scale. Only specimens yielding a minimum of 6.9 µg of tumor DNA, 5.15 µg RNA, and 4.9 µg of germline DNA were included in this study. RNA was analyzed via the RNA6000 Nano assay (Agilent) for determination of an RNA Integrity Number (RIN), and only the cases with RIN >7.0 were included in this study.

Pathology Review—All samples were systematically evaluated by gastroenterological pathologists to confirm the histopathologic diagnosis and any variant histology according to the most recent World Health Organization (WHO) classification (International Agency for Research on Cancer, 2010). All tumor samples were assessed for tumor content (percent tumor nuclei), Tumor samples were evaluated for the presence and extent of inflammatory infiltrate as well as the type of the infiltrating cells in the tumor microenvironment (lymphocytes, neutrophils, eosinophils, histiocytes, plasma cells). Any non-concordant diagnoses among the pathologists were re-reviewed and resolution achieved after discussion.

DNA Sequencing data—Exome capture was performed using Agilent SureSelect Human All Exon 50 Mb according to the manufacturers' instructions. Briefly, 0.5–3 micrograms of DNA from each sample were used to prepare the sequencing library through shearing of the DNA followed by ligation of sequencing adaptors. All whole exome (WES) and whole genome (WGS) sequencing was performed on the Illumina HiSeq platform. Paired-end sequencing (2 × 101 bp for WGS and 2 × 76 bp for WE) was carried out using HiSeq sequencing instruments; the resulting data was analyzed with the current Illumina pipeline. Basic alignment and sequence QC was done with the Picard and Firehose pipelines at the

Broad Institute. Sequencing data were processed using two consecutive pipelines: **(1) Sequencing data processing pipeline (“Picard pipeline”)**. Picard (<http://picard.sourceforge.net/>) uses the reads and qualities produced by the Illumina software for all lanes and libraries generated for a single sample (either tumor or normal) and produces a single BAM file (<http://samtools.sourceforge.net/SAM1.pdf>) representing the sample. The final BAM file stores all reads and calibrated qualities along with their alignments to the genome. **(2) Cancer genome analysis pipeline (“Firehose pipeline”)**. Firehose (<http://www.broadinstitute.org/cancer/cga/Firehose>) takes the BAM files for the tumor and patient-matched normal samples and performs analyses including quality control, local realignment, mutation calling, small insertion and deletion identification, rearrangement detection, coverage calculations and others as described briefly below. The pipeline represents a set of tools for analyzing massively parallel sequencing data for both tumor DNA samples and their patient-matched normal DNA samples. Firehose uses GenePattern (Reich et al., 2006) as its execution engine for pipelines and modules based on input files specified by Firehose. The pipeline contains the following steps:

Quality control. This step confirms identity of individual tumor and normal to avoid mix-ups between tumor and normal data for the same individual.

Local realignment of reads. This step realigns reads at sites that potentially harbor small insertions or deletions in either the tumor or the matched normal, to decrease the number of false positive single nucleotide variations caused by misaligned reads.

Identification of somatic single nucleotide variations (SSNVs) – This step detects candidate SSNVs using a statistical analysis of the bases and qualities in the tumor and normal BAMs, using Mutect (Cibulskis et al., 2013).

Identification of somatic small insertions and deletions – In this step, putative somatic events were first identified within the tumor BAM file and then filtered out using the corresponding normal data, using Indelocator (Ratan et al., 2015).

Mutation Data—A series of quality-control filters according to the MC3 MAF were applied to mutations: (1) A filter for artificial CC>CA mutations caused by sample oxidation (8-oxoguanine) was applied to remove potential CC>CA artifacts (Costello et al., 2013); (2) Variants that were frequently observed in the Exome Aggregation Consortium (<http://exac.broadinstitute.org>) were excluded; (3) mutations with evidence of strand bias were excluded; (4) mutations with “ndp” labels were excluded; (5) duplicated mutations due to redundant tumor or normal samples were excluded. Somatic mutation calling was focused on coding mutations spanning missense and nonsense mutations, in-frame and frame-shift indels, and mutations that occurred on splice site, start codon, or stop codon.

The MutSig2CV (Cancer Genome Atlas Research Network, 2011) was applied to the quality-controlled mutation data to evaluate significance of mutated genes and estimate mutation densities of samples. MutSig2CV combines evidence from background mutation rate, clustering of mutations on hotspots and conservation of mutated sites to calculate the false discovery rates (q values). Genes of q value < 0.1 were declared significant.

Microsatellite Instability—DNA samples were evaluated for Microsatellite Instability using the MSI-Mono-Dinucleotide assay, which examines four mononucleotide repeat loci (polyadenine tracts BAT25, BAT26, BAT40 and transforming growth factor receptor type II) and three dinucleotide repeat loci (CA repeats in D2S123, D5S346 and D17S250).

Somatic Copy Number Alterations—DNA from each tumor or germline sample was hybridized to Affymetrix SNP 6.0 arrays using protocols from the Genome Analysis Platform of the Broad Institute as previously described (McCarroll et al., 2008). From raw CEL files, Birdseed was used to infer a preliminary copy-number at each probe locus (Korn et al., 2008). For each tumor, genome-wide copy-number estimates were refined using tangent normalization, in which tumor signal intensities are divided by signal intensities from the linear combination of all normal samples that are most similar to the tumor. This linear combination of normal samples tends to match the noise profile of the tumor better than any set of individual normal samples, thereby reducing the contribution of noise to the final copy-number profile. Individual copy-number estimates then underwent segmentation using Circular Binary Segmentation (Olshen et al., 2004). Segmented copy-number profiles for tumor and matched control DNAs were analyzed using Ziggurat Deconstruction, an algorithm that parsimoniously assigns a length and amplitude to the set of inferred copy-number changes underlying each segmented copy number profile, and the analysis of broad copy-number alterations was then conducted as previously described (Mermel et al., 2011). Significant focal copy-number alterations were identified from segmented data using GISTIC 2.0 (Mermel et al., 2011). Allelic copy number, regions of homozygous deletions, whole genome doubling and purity and ploidy estimates were calculated using the ABSOLUTE algorithm (Carter et al., 2012).

Copy ratios of the genomic segments were adjusted by purity and ploidy using the In Silico Admixture Removal (ISAR) method (Carter et al., 2012). The tumor purity and ploidy were estimated with ABSOLUTE (Absolute quantification of somatic DNA alterations in human cancer (Carter et al., 2012)). GISTIC 2.0 (Mermel et al., 2011) was used to identify significant genomic regions, and q values that were smaller than 0.1 were considered significant. The gene under selective pressure in each significant amplification/deletion peak was manually curated with consideration of the common fragile sites (CFS). The gene-level copy numbers were obtained from GISTIC, and the gene was considered as amplified or deleted if the gene-level copy number change (ploidy-adjusted) was larger than 2 or smaller than -1.3, respectively. Whole-genome doubling (WGD) calls, absolute allelic copy numbers, and clonal statuses of the SCNAs were all obtained from ABSOLUTE.

Aneuploidy Scores—The aneuploidy scores were calculated to quantify various kinds of aneuploidy in terms of length and magnitude of the copy-number events including segment gains and losses. The aneuploidy scores in this study were obtained as follows: (1) the original copy ratios of the genomic segments were adjusted by purity and ploidy using the ISAR method as noted above; (2) GISTIC 2.0 was used to deconstruct the ISAR-adjusted copy-number profile into SCNA events (discrete copy-number alterations), and each SCNA event could be categorized based on its length and magnitude (with details below); (3) for each category of SCNA events, e.g., focal amplifications, the corresponding aneuploidy

score was calculated as $\log_{10}(1 + n)$, where n is the number of events in that category. Similar approaches to the aneuploidy scores in principle were applied in a recent study (Davoli et al., 2017) as well as in our previous study (Dulak et al., 2012). The categories of SCNA events were defined as (1) Arm-level events: the relative SCNA length (as a proportion to the arm length) $I_{arm} > 0.5$, and the absolute value of amplitude $|m| > 0.3$, and the threshold of 0.3 was applied to remove low copy ratio changes that were likely noise; (2) Focal events: $I_{arm} < 0.5$, $|m| > 0.3$; (3) Focal amplifications: $I_{arm} < 0.5$, $m > 0.3$; (4) Focal deletions: $I_{arm} < 0.5$, $m < -0.3$; (5) High-level focal amplifications: $I_{arm} < 0.5$, $m > 1$; (6) Deep-level focal deletions: $I_{arm} < 0.5$, $m < -1$. This method serves as a quantification of different types of genomic aneuploidy, and it is different from the gene-level amplification and deletion mentioned above, where conservative thresholds (2 and -1.3) for the gene-level copy number (not SCNA events) were applied to define functional alterations of the genes.

CIN-Focal Score—We developed a CIN-Focal (CIN-F) score to capture the most focal high-level amplicons (MFAs), which are likely to be functional gains of specific genomic regions that were subject to positive selection during cancer evolution. Based on the deconstructed copy-number events from GISTIC 2.0, we defined those MFAs as $l < 3$ Mb and $m > 2$, where l is the length of the amplicon in mega-bases, and m is the event amplitude as mentioned above. Given each of those amplicons, the CIN-F score of a tumor was first calculated as the weighted sum of the magnitude m_a of each amplicon a (weighted by its length l_a), and then log-transformation was applied:

$$S_{\text{CIN-F}} = \log_{10} \left(1 + \sum_a l_a \cdot m_a \right)$$

Because m_a is the ploidy-adjusted amplitude of copy-number gain, $l_a \cdot m_a$ is theoretically proportional to the relative amount of DNA (compared to the total cancer DNA) of the amplicon a , so that the CIN-F score corresponds to the amount of additional DNA within the MFAs. An alternative metric to CIN-F score is simply the total number of MFAs in a genome regardless of the lengths and amplitudes of the MFAs. The CIN-F score showed a binomial distribution in the upper GI cancers. We used kernel density estimation of Gaussian kernels (R statistical software, the “density” function) to set the threshold for dichotomization at the local minimum of estimated density of the CIN-F score, and this analysis yielded a threshold of $S_{\text{CIN-F}} = 0.438$. The CIN tumors was then dichotomized into CIN-F and CIN-B as shown in Figure 4C.

Clonal Deletion Score (CDS)—To identify tumors with chromosomal instability, we developed a score, termed the Clonal Deletion Score, or CDS, which quantifies the number of clonally deleted genomic regions in each tumor’s genome. The CDS of each tumor was calculated using absolute allelic copy numbers of genomic segments of the tumor. For each genomic segment, the absolute allelic copy numbers are denoted as q_1 and q_2 for the two alleles with lower and higher copy number, respectively. If (1) the segment is a deletion, i.e., $q_1 < q_2$, and $q_1 + q_2 < \bar{p}$, where \bar{p} is the average tumor ploidy; and (2) the deletion is clonal, i.e., q_1 is a clonal copy number according to ABSOLUTE; then the clonal deletion effect (CDE) of the segment is calculated as:

$$\text{CDE} = 2 \left(1 - \frac{q_1 + q_2}{\tau} \right)$$

If a segment does not satisfy the above criteria, the CDE of that segment is zero. The copy number of the higher allele q_2 was incorporated so as to diminish the CDE when there was a gain of the higher allele, e.g., copy-neutral loss of heterozygosity (LOH). Given the CDE of each segment s , the CDS of a tumor is the average of CDE weighted by the lengths of the segments:

$$\text{CDS} = \sum_s w_s \cdot \text{CDE}_s, w_s = \frac{l_s}{\sum_s l_s}$$

where l_s is the length of a segment. The CDSs from the GI adenocarcinomas showed a clear bimodal separation. The kernel density estimation approach as mentioned above was used to set the threshold for dichotomization of CDS. A threshold of CDS = 0.0249 was then applied for the binary CIN/GS classification (Figure 2B and S2B), which corresponds to distinct copy-number profiles as shown in Figure S2C.

Mutational Signatures—Mutational signatures were identified from SNVs using a Bayesian version of the non-negative matrix factorization method as described previously (Kim et al., 2016). The mutations were deconvoluted into distinct mutational signatures based on the number of mutations partitioned by 6 base substitutions (C>A, C>G, C>T, T>A, T>C, and T>G) and 16 possible combinations of neighboring bases that resulted in 96 possible types of mutations. A 96-by-M matrix of mutation counts (M is the number of samples) was constructed as the input data for signature discovery. Cosine similarity was used to evaluate the resemblance of the identified signatures with the COSMIC signatures (<http://cancer.sanger.ac.uk/cosmic/signatures>). For each sample, the estimated number of mutations from a signature was used as the intensity of that signature. A two-stage strategy of mutational signature discovery was performed in this study to achieve more accuracy in the identification of signatures. In the first stage, all samples were used to identify the signatures. In the second stage, the analysis was performed only for the non-hypermutated cases with the MSI and POLE signatures removed from the mutation counts to facilitate identification of signatures in the non-HM population.

Stemness Index—We used one-class logistic regression (Sokolov et al., 2016) to derive a stemness index based on a gene expression signature derived from embryonic and differentiated cells from the PCBC dataset (Daily et al., 2017; Salomonis et al., 2016) and applied this to GIAC samples using Spearman correlations between the model's weight vector and the GIAC sample's expression profile (Malta et al., 2018).

Differential Gene Expression Analysis between GIAC and non-GI AC—To identify genes differentially abundant in GIAC versus non-GI AC, excluding genes that are differentially expressed between normal GI tissue compared to normal non-GI tissue, we needed to use external gene expression data normal tissues. We selected 4 gastrointestinal

(esophagus, stomach, colon-transverse, and colon-sigmoid) and 5 non-gastrointestinal (breast, lung, ovary, prostate, and uterine) normal tissue types through GTEx repository of normal tissues (Consortium et al., 2017) (<https://www.gtportal.org/home/datasets>, GTEx Version 7), and utilized their RNA-sequencing expression dataset. Normalized expression values for both TCGA tumor and GTEx normal tissue cases were calculated by robust scaling (on values between 2.5 and 97.5 percentile) and winsorizing of each gene's expression (mean \pm 3 standard deviations) in the respective case population of tumoral or normal cases. Gastrointestinal and non-gastrointestinal normal tissues were selected based on the matching with composition of available GI and non-GI adenocarcinomas in TCGA PanCancer project. Orthogonal partial least squares-discriminant analysis (OPLS-DA) was used to discover a subgroup of genes (n=671) that were not differentially expressed in GI and non-GI normal tissues, but were members of our list of differentially expressed genes between GI adenocarcinomas (GIAC) and non-GI adenocarcinomas (Non-GI AC). Significance was determined by absolute loading in the OPLS discriminant analysis of higher than 0.05. The genes for which expression was highly associated with the stromal class of GI tumors identified by the method described in Isella et al (Isella et al., 2015) (n=118) were excluded from further analysis (absolute loading higher than 0.05). By utilizing an OPLS-DA model comparing GIAC and non-GI AC cases, the remaining 553 genes were ranked by their loadings toward overexpression in GIAC. Results were depicted in two heatmaps illustrating the normalized expression values for the selected genes in both GIAC and non-GI AC tissues (first heatmap), and normal GI and non-GI tissues (second heatmap).

Selection of transcription factors for gain- and loss-of-function studies—We used multiple sources to select 139 transcription factors (TFs) that are important in GI development. We first identified 40 TFs in the Gene Ontology (GO) database based on the intersection of two GO terms, *RNA polymerase II transcription factor activity, sequence-specific DNA binding* (GO: 0000981) and *digestive tract development* (GO: 0048565), in *Homo sapiens*. Further, we collected 24 TFs from the review by Noah et al. on human intestinal development and differentiation (Noah et al., 2011). Additionally, 93 genes were identified in the study in which Sherwood and colleagues used microarray and dynamic immunofluorescence technologies to profile gene expression during mouse endodermal organ formation (Sherwood et al., 2009). Finally, we also included nine other TFs that were significantly mutated in GIAC. In all, we examined 139 genes (taking the union of the four gene lists and removing genes with missing platform data).

DNA methylation data—Illumina Infinium DNA methylation arrays [including both HumanMethylation27 (HM27) and HumanMethylation450 (HM450)] were used to assay 921 GIAC and 76 adjacent non-malignant tissues. Level 3 data from two generations of Illumina Infinium DNA methylation arrays were combined and further normalized between platforms using a probe-by-probe proportional rescaling method as outlined below to yield a final common set of 22,601 probes with comparative methylation levels between platforms. During data generation, a single technical replicate of the same cell line control sample from either of two different DNA extractions (TCGA-07-0227/TCGA-AV-A03D) was included on each plate as a control, and measured 44/198 times and 12/169 times on HM27 and HM450,

respectively. These repeated measurements were therefore used for rescaling of the HM27 data to be comparable to HM450. For each probe within each platform, we computed the median β value across all technical replicates of each of the two TCGA IDs. We then combined the two extractions by taking the mean of the two medians obtained for each of the two replicate TCGA IDs, and obtained a single summarized DNA methylation read-out (β value) for the corresponding probe i for each platform, noted as $\overline{\text{Beta}}_{\text{hm27},i}$ and $\overline{\text{Beta}}_{\text{hm450},i}$, respectively. We then applied a constrained (within the range of 0 to 1 for β values) linear rescaling of the HM27 data for each probe and for each patient's sample using $\overline{\text{Beta}}_{\text{hm27},i}$ and $\overline{\text{Beta}}_{\text{hm450},i}$. When the HM27 β value of a patient's sample j for probe i was smaller than the mean of median replicate samples on the HM27 for that probe, we linearly rescaled the HM27 β value $\text{Beta}_{\text{hm27},i,j}$ in the $(0, \text{Beta}_{\text{hm27},i,j})$ space; and when $\text{Beta}_{\text{hm27},i,j}$ was greater, we linearly rescaled the HM27 beta value $\text{Beta}_{\text{hm27},i,j}$ in the $(\text{Beta}_{\text{hm27},i,j}, 1)$ space; This translates into the following mathematical computation:

$$\begin{aligned} \text{Beta}_{\text{hm450},i,j} &= \text{Beta}_{\text{hm27},i,j} * (\overline{\text{Beta}}_{\text{hm450},i} / \overline{\text{Beta}}_{\text{hm27},i}), \text{ if } \text{Beta}_{\text{hm27},i,j} < \overline{\text{Beta}}_{\text{hm27},i}; \text{ and} \\ \text{Beta}_{\text{hm450},i,j} &= 1 - (1 - \text{Beta}_{\text{hm27},i,j}) * ((1 - \overline{\text{Beta}}_{\text{hm450},i}) / (1 - \overline{\text{Beta}}_{\text{hm27},i})), \text{ if} \\ \text{Beta}_{\text{hm27},i,j} &> \overline{\text{Beta}}_{\text{hm27},i} \end{aligned}$$

After the between-platform normalization, we further excluded 779 probes that still showed a consistent platform difference (mean β value difference greater than or equal to 0.1) in six or more tumor types.

Unsupervised clustering analysis of DNA methylation data—Unsupervised clustering analyses of DNA methylation data were performed based on promoter CpG sites that did not exhibit tissue-specific DNA methylation in normal tissues and blood cells (mean β value < 0.2 for each tissue type), but acquired methylated in tumors.

GIAC and non-GI AC (Figure S1A): We analyzed DNA methylation profiles of 3,759 adenocarcinomas including 921 GI adenocarcinomas and 2,828 non-GI adenocarcinomas representing 12 disease types (four GIAC and eight non-GI AC). We also included data from 333 histologically normal tumor-adjacent tissue specimens corresponding each disease type (BRCA $n=101$, PRAD $n=39$, OV $n=12$, CEAD $n=1$, UCEC $n=43$, EAC/GAC $n=33$, COAD $n=37$, READ $n=6$, CHOL $n=9$, PAAD $n=10$, LUAD $n=42$). We first used the data from the normal tissues and leukocytes to select CpG sites that lacked tissue-specific DNA methylation (mean β value < 0.2 in any tissue type and β value > 0.3 in no more than five samples across the entire set). We then performed clustering analysis of the adenocarcinomas using 2,783 CpG sites that were hypermethylated (β value > 0.3) in more than 10% within any of the 12 disease types. To minimize the influence of tumor purity on a clustering result, we dichotomized the data using a β value of > 0.3 to define positive DNA methylation and < 0.3 to specify lack of methylation. We applied hierarchical clustering with Ward's method to cluster the distance matrix computed with the Jaccard index. Heatmap was generated based on the original β values for 1,000 loci (a subset of 2,783 loci) with the highest standard deviation in DNA methylation measurements among all adenocarcinomas.

GIAC (Figure 2D): We analyzed DNA methylation profiles of 921 GIAC and 77 (33 gastric and 44 colorectal) histologically normal tumor-adjacent tissue specimens. The precise locations within the GI organs from which the normal-adjacent tissue specimens were excised are not available. Unsupervised clustering of GIAC was performed based on 2,845 gene promoter loci unmethylated in normal tissues and leukocytes (mean β value < 0.2 in both normal gastric and colorectal tissues) but methylated (β value > 0.3) more than 5% in at least one of the GIAC tumor types. To minimize the influence of tumor purity, we dichotomized the data into 0's and 1's using a β value threshold of 0.3. The optimal number of clusters was assessed based on 80% probe and tumor resampling over 1,000 iterations of hierarchical clustering for $K = 2, 3, 4, \dots, 20$ using the binary distance metric for clustering and Ward's method for linkage as implemented in the R/Bioconductor ConsensusClusterPlus package. The heatmap was generated using the original β values. The probes were displayed based on the order of unsupervised hierarchical clustering of the β values using the Euclidean distance metric and Ward's linkage method.

The union of MSI and CIMP-H GIAC (Figure 3B): We used 158 tumors (93 GEA and 65 CRC) that were classified as either CIMP-H or MSI and 44 normals (12 stomach and 32 colorectal), which were assayed on the HM450 platform. Unsupervised and dichotomized clustering was performed using 35,436 sites lacking DNA methylation in normal tissues (mean β value < 0.2 in both normal gastric and colorectal tissues) and methylated (β value > 0.3) more than 10% in any of the tumor type. Heatmap was generated based on the top 10% of the most variably hypermethylated sites across 158 GIAC.

GIAC DNA hypermethylation subtypes—We chose seven GIAC DNA methylation clusters defined by the consensus clustering. For further integrative analyses, we focused on four prominent clusters showing a high frequency of cancer-associated DNA hypermethylation. We found that the gastroesophageal (GEA) and colorectal adenocarcinomas (CRC) largely clustered separately. Among GEA, EBV⁺ gastric cancers stood out from all the rest by their extensive DNA hypermethylation (cluster 4) and were designated as **EBV-CIMP** as previous study (Cancer Genome Atlas Research Network, 2014). Cluster 5 is significantly enriched for MSI tumors originated in both stomach and colon. It included well-known CIMP-High CRC associated with *BRAF*^{V600E} mutations and MSI-associated Gastric-CIMP described previously (Cancer Genome Atlas Research Network, 2014; Weisenberger et al., 2006). We classified these tumors as **GIAC CIMP-H**, as having a higher prevalence of DNA hypermethylation than all the other clusters with the exception of EBV-CIMP. Further, we named cluster 6 as **CRC CIMP-L** that exhibited features consistent with CIMP-Low subtype previously described (Cancer Genome Atlas Research Network, 2012). It had a significant association with *KRAS* mutations ($p < 2.2 \times 10^{-16}$ [vs. CRC in other groups], Fisher's exact test). Among GEA, cluster 1 was enriched for esophageal tumors ($p = 8.0 \times 10^{-8}$ [vs. GEAs in other groups]), and also had a mean DNA hypermethylation frequency slightly higher than that in CRC CIMP-L and other GEA clusters (cluster 2 and 3). We specified these tumors as **GEA-CIMP-L**. These tumors showed frequent epigenetic silencing of tumor suppressor genes including *CDKN2A* and *MGMT* ($p = 1.5 \times 10^{-10}$ and $p = 1.5 \times 10^{-11}$, respectively, [vs. GEA clusters 2 and 3]).

Identification of epigenetically silenced genes—We used 775 GIAC and 44 adjacent non-malignant tissues assayed on the HM450 platform. Probes located within potential promoter regions (1500 bp flanking regions upstream and downstream of Transcription Start Sites (TSSs) of all transcripts annotated by UCSC) were examined for evidence of epigenetic silencing. We removed the CpG sites that were methylated in normal tissues and blood cells (mean β value > 0.2 for each tissue type). In order to remove the effect of tissue specificity on gene expression, we z-score-transformed log₂ gene expression data first within each cancer type. The z-scores were derived using the mean and standard deviation calculated with the unmethylated tumors only, defined as those with a β value of (0, 0.2). Samples across all the cancer types were then pooled. For each probe/gene pair, we chose the probes that exhibited epigenetic silencing with the following criteria: 1) at least 8 samples ($>1\%$ of all tumors) were observed with a β value of 0.3 or above (defined as the methylated group); 2) mean z-score of the methylated group was lower than -1.65 ; 3) FDR-corrected p value according to one-side *t*-test on z-scores was lower than 0.001 between the unmethylated and methylated groups. Probes surviving these steps were retained to call epigenetic silencing events based on DNA methylation profiles for each sample. If there were multiple probes associated with the same gene, a sample identified as epigenetically silenced at more than half the probes for the corresponding gene was also labeled as epigenetically silenced at the gene level.

CDKN2A epigenetic silencing calls were made using the exon-level RNA-seq data. *CDKN2A* DNA methylation status was assessed in each sample, based on the probe (cg13601799) located in the p16INK4 promoter CpG island. p16INK4 expression was determined by the log₂(RPKM+1) level of its first exon (chr9: 21974403–21975132). The epigenetic silencing calls for each sample were made by evaluating a scatter plot showing an inverse association between DNA methylation and expression. For *RAD51C*, there was no common probe between HM27 and HM450 that was located in the promoter region. However, probe cg14837411 from HM27 and probe cg27221688 from HM450 were only 100bp apart, and both correlated with gene expression. Therefore, we combined them in determining the silencing status of this gene. Samples with a β value of 0.2 or above for either probe were designated as cases with epigenetic silencing.

DNA hypermethylation frequency in GIAC and Non-GI AC—We identified a set of 13,809 CpG sites that were unmethylated in normal tissues and blood cells (mean β value < 0.2 for each tissue type). For each CpG locus, tumors with a β value of 0.3 or greater were designated as methylated, and tumors with a β value of lower than 0.3 were designated as unmethylated. We then calculated the percentage of loci that were methylated among the loci investigated in each tumor.

Methods for integrative pathway analysis—We evaluated somatic mutations and copy-number changes relevant to well-studied signaling pathways curated in previous TCGA publications. Oncogenic relevance was assessed using OncoKB, a knowledge base for the oncogenic effects of cancer genes, that is manually curated by researchers and physicians at Memorial Sloan Kettering (Chakravarty et al., 2017). Specifically, a mutation was counted and included in the diagrams if either (1) it had been reported as a recurrent

alteration in COSMIC (Forbes et al., 2011) or (2) it had been labeled as oncogenic or likely oncogenic in OncoKB. Amplifications and deep deletions were based on GISTIC calls and reflect a change of more than half of the baseline gene copies. The actual list of oncogenic and likely oncogenic alterations is regularly updated based on the literature; the most recent version can be retrieved online from the OncoKB public website (www.oncokb.org) or visualized when viewing the data in the cBioPortal (www.cbioportal.org). For known oncogenes, only genetic alterations inferred to be activating were considered; for tumor suppressor genes, only alterations inferred to be inactivating were considered.

QUANTIFICATION AND STATISTICAL ANALYSIS

We used Fisher's exact test for independence between two categorical variables throughout the analyses. Wilcoxon rank-sum test was performed for any independence test between a continuous variable and a binary categorical variable. For any test between two continuous variables or any association test that needed to be adjusted by covariates, a (multiple) linear model was fitted to evaluate the significance of coefficients, and analysis of variance was used to calculate the proportion of variance explained by each variable. Non-negative variables that were heavily right-skewed, which included the aneuploidy scores, CIN-F score, number of MFAs, and the intensities of mutational signatures, were log-transformed (with a pseudo-count of 1 added) for appropriate fitting of multiple linear models. For the association test between aneuploidy scores and BRCA signature, the arm-level score and focal score were simultaneously included as explanatory variables in the multiple linear model. The association test between BRCA signature and PARP1 expression (log-transformed) was adjusted by the copy number of PARP1. The intensity of the CpG>TpG signature was modeled by multiple linear regression with explanatory variables of upper/lower GI, molecular subtype, age, and CIMP status as an ordinal variable. A logistic regression model was fitted when the response variable was binary. The test between the CIN-F score and clinical stage was performed using an ordered logit model as the clinical stage was considered an ordinal variable, and the p values were calculated using normal approximation. The association test between number of MFAs and the CRC stromal subtype was performed using negative-binomial regression that models the sparse number of MFAs, so as to increase statistical power. Cox regression was used for survival analysis to evaluate the significance of the variables. All statistical analyses in this study were performed using the R statistical software (<https://www.r-project.org>).

DATA AND SOFTWARE AVAILABILITY

The raw data, processed data and clinical data can be found at the legacy archive of the GDC (<https://portal.gdc.cancer.gov/legacy-archive/search/f>) and the PancanAtlas publication page (<https://gdc.cancer.gov/about-data/publications/pancanatlas>). The mutation data can be found here: (<https://gdc.cancer.gov/about-data/publications/mc3-2017>). TCGA data can also be explored through the Broad Institute FireBrowse portal (<http://gdac.broadinstitute.org>) and the Memorial Sloan Kettering Cancer Center cBioPortal (<http://www.cbioportal.org>). Details for software availability are in the Key Resource Table.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank all patients who contributed to this study. This work was supported by the Intramural Research Program and the following grants from the United States National Institutes of Health: U24 CA143799, U24 CA143835, U24 CA143840, U24 CA143843, U24 CA143845, U24 CA143848, U24 CA143858, U24 CA143866, U24 CA143867, U24 CA143882, U24 CA143883, U24 CA144025, U54 HG003067, U54 HG003079, U54 HG003273 and P30 CA16672.

References

- Alexandrov LB, Nik-Zainal S, Siu HC, Leung SY, Stratton MR. A mutational signature in gastric cancer suggests therapeutic strategies. *Nat Commun.* 2015; 6:8683. [PubMed: 26511885]
- Alexandrov LB, Nik-Zainal S, Wedge DC, Aparicio SA, Behjati S, Biankin AV, Bignell GR, Bolli N, Borg A, Borresen-Dale AL, et al. Signatures of mutational processes in human cancer. *Nature.* 2013; 500:415–421. [PubMed: 23945592]
- Arnold M, Soerjomataram I, Ferlay J, Forman D. Global incidence of oesophageal cancer by histological subtype in 2012. *Gut.* 2015; 64:381–387. [PubMed: 25320104]
- Banerjea A, Hands RE, Powar MP, Bustin SA, Dorudi S. Microsatellite and chromosomal stable colorectal cancers demonstrate poor immunogenicity and early disease recurrence. *Colorectal Dis.* 2009; 11:601–608. [PubMed: 18637931]
- Bignell GR, Greenman CD, Davies H, Butler AP, Edkins S, Andrews JM, Buck G, Chen L, Beare D, Latimer C, et al. Signatures of mutation and selection in the cancer genome. *Nature.* 2010; 463:893–898. [PubMed: 20164919]
- Bijlsma MF, Sadanandam A, Tan P, Vermeulen L. Molecular subtypes in cancers of the gastrointestinal tract. *Nat Rev Gastroenterol Hepatol.* 2017; 14:333–342. [PubMed: 28400627]
- Bonhomme C, Duluc I, Martin E, Chawengsaksophak K, Chenard MP, Keding M, Beck F, Freund JN, Domon-Dell C. The Cdx2 homeobox gene has a tumour suppressor function in the distal colon in addition to a homeotic role during gut development. *Gut.* 2003; 52:1465–1471. [PubMed: 12970140]
- Budinska E, Popovici V, Tejpar S, D'Ario G, Lapique N, Sikora KO, Di Narzo AF, Yan P, Hodgson JG, Weinrich S, et al. Gene expression patterns unveil a new level of molecular heterogeneity in colorectal cancer. *J Pathol.* 2013; 231:63–76. [PubMed: 23836465]
- Bunz F, Fauth C, Speicher MR, Dutriaux A, Sedivy JM, Kinzler KW, Vogelstein B, Lengauer C. Targeted inactivation of p53 in human cells does not result in aneuploidy. *Cancer Res.* 2002; 62:1129–1133. [PubMed: 11861393]
- Cancer Genome Atlas Research Network. Integrated genomic analyses of ovarian carcinoma. *Nature.* 2011; 474:609–615. [PubMed: 21720365]
- Cancer Genome Atlas Research Network. Comprehensive molecular characterization of human colon and rectal cancer. *Nature.* 2012; 487:330–337. [PubMed: 22810696]
- Cancer Genome Atlas Research Network. Comprehensive molecular characterization of gastric adenocarcinoma. *Nature.* 2014; 513:202–209. [PubMed: 25079317]
- Cancer Genome Atlas Research Network. Integrated genomic characterization of oesophageal carcinoma. *Nature.* 2017; 541:169–175. [PubMed: 28052061]
- Carter SL, Cibulskis K, Helman E, McKenna A, Shen H, Zack T, Laird PW, Onofrio RC, Winckler W, Weir BA, et al. Absolute quantification of somatic DNA alterations in human cancer. *Nat Biotechnol.* 2012; 30:413–421. [PubMed: 22544022]
- Chakravarty D, Gao J, Phillips S, Kundra R, Zhang H, Wang J, Rudolph JE, Yaeger R, Soumerai T, Nissan MH, et al. OncoKB: A Precision Oncology Knowledge Base. *JCO Precision Oncology.* 2017:1–16.

- Chan TL, Curtis LC, Leung SY, Farrington SM, Ho JW, Chan AS, Lam PW, Tse CW, Dunlop MG, Wyllie AH, et al. Early-onset colorectal cancer with stable microsatellite DNA and near-diploid chromosomes. *Oncogene*. 2001; 20:4871–4876. [PubMed: 11521198]
- Chu J, Sadeghi S, Raymond A, Jackman SD, Nip KM, Mar R, Mohamadi H, Butterfield YS, Robertson AG, Birol I. BioBloom tools: fast, accurate and memoryefficient host species sequence screening using bloom filters. *Bioinformatics*. 2014; 30:3402–3404. [PubMed: 25143290]
- Cibulskis K, Lawrence MS, Carter SL, Sivachenko A, Jaffe D, Sougnez C, Gabriel S, Meyerson M, Lander ES, Getz G. Sensitive detection of somatic point mutations in impure and heterogeneous cancer samples. *Nat Biotechnol*. 2013; 31:213–219. [PubMed: 23396013]
- Cibulskis K, McKenna A, Fennell T, Banks E, DePristo M, Getz G. ContEst: estimating cross-contamination of human samples in next-generation sequencing data. *Bioinformatics*. 2011; 27:2601–2602. [PubMed: 21803805]
- Consortium, G.T., Laboratory, D.A., Coordinating Center -Analysis Working G, Statistical Methods groups-Analysis Working G, Enhancing G.g, Fund NIHC, Nih/Nci, Nih/Nhgri, Nih/Nimh, Nih/Nida, et al. Genetic effects on gene expression across human tissues. *Nature*. 2017; 550:204–213. [PubMed: 29022597]
- Costello M, Pugh TJ, Fennell TJ, Stewart C, Lichtenstein L, Meldrim JC, Fostel JL, Friedrich DC, Perrin D, Dionne D, et al. Discovery and characterization of artifactual mutations in deep coverage targeted capture sequencing data due to oxidative DNA damage during sample preparation. *Nucleic Acids Res*. 2013; 41:e67. [PubMed: 23303777]
- Cristescu R, Lee J, Nebozhyn M, Kim KM, Ting JC, Wong SS, Liu J, Yue YG, Wang J, Yu K, et al. Molecular analysis of gastric cancer identifies subtypes associated with distinct clinical outcomes. *Nat Med*. 2015; 21:449–456. [PubMed: 25894828]
- Daily K, Ho Sui SJ, Schriml LM, Dexheimer PJ, Salomonis N, Schroll R, Bush S, Keddache M, Mayhew C, Lotia S, et al. Molecular, phenotypic, and sample-associated data to describe pluripotent stem cell lines and derivatives. *Sci Data*. 2017; 4:170030. [PubMed: 28350385]
- Davoli T, Uno H, Wooten EC, Elledge SJ. Tumor aneuploidy correlates with markers of immune evasion and with reduced response to immunotherapy. *Science*. 2017; 355
- Derks S, Liao X, Chiaravalli AM, Xu X, Camargo MC, Solcia E, Sessa F, Fleitas T, Freeman GJ, Rodig SJ, et al. Abundant PD-L1 expression in Epstein-Barr Virusinfected gastric cancers. *Oncotarget*. 2016; 7:32925–32932. [PubMed: 27147580]
- Dienstmann R, Vermeulen L, Guinney J, Kopetz S, Tejpar S, Tabernero J. Consensus molecular subtypes and the evolution of precision medicine in colorectal cancer. *Nat Rev Cancer*. 2017; 17:79–92. [PubMed: 28050011]
- Dulak AM, Schumacher SE, van Lieshout J, Imamura Y, Fox C, Shim B, Ramos AH, Saksena G, Baca SC, Baselga J, et al. Gastrointestinal adenocarcinomas of the esophagus, stomach, and colon exhibit distinct patterns of genome instability and oncogenesis. *Cancer Res*. 2012; 72:4383–4393. [PubMed: 22751462]
- Dulak AM, Stojanov P, Peng S, Lawrence MS, Fox C, Stewart C, Bandla S, Imamura Y, Schumacher SE, Shefler E, et al. Exome and whole-genome sequencing of esophageal adenocarcinoma identifies recurrent driver events and mutational complexity. *Nat Genet*. 2013; 45:478–486. [PubMed: 23525077]
- Ebert MP, Tanzer M, Balluff B, Burgermeister E, Kretzschmar AK, Hughes DJ, Tetzner R, Lofton-Day C, Rosenberg R, Reinacher-Schick AC, et al. TFAP2E-DKK4 and chemoresistance in colorectal cancer. *N Engl J Med*. 2012; 366:44–53. [PubMed: 22216841]
- Fang M, Hutchinson L, Deng A, Green MR. Common BRAF(V600E)-directed pathway mediates widespread epigenetic silencing in colorectal cancer and melanoma. *Proc Natl Acad Sci U S A*. 2016; 113:1250–1255. [PubMed: 26787892]
- Forbes SA, Bindal N, Bamford S, Cole C, Kok CY, Beare D, Jia M, Shepherd R, Leung K, Menzies A, et al. COSMIC: mining complete cancer genomes in the Catalogue of Somatic Mutations in Cancer. *Nucleic Acids Res*. 2011; 39:D945–950. [PubMed: 20952405]
- Ganem NJ, Storchova Z, Pellman D. Tetraploidy, aneuploidy and cancer. *Curr Opin Genet Dev*. 2007; 17:157–162. [PubMed: 17324569]

- Glover TW, Wilson TE, Arlt MF. Fragile sites in cancer: more than meets the eye. *Nat Rev Cancer*. 2017; 17:489–501. [PubMed: 28740117]
- Guinney J, Dienstmann R, Wang X, de Reynies A, Schlicker A, Soneson C, Marisa L, Roepman P, Nyamundanda G, Angelino P, et al. The consensus molecular subtypes of colorectal cancer. *Nat Med*. 2015; 21:1350–1356. [PubMed: 26457759]
- Guo Z, Kozlov S, Lavin MF, Person MD, Paull TT. ATM activation by oxidative stress. *Science*. 2010; 330:517–521. [PubMed: 20966255]
- Herman JG, Umar A, Polyak K, Graff JR, Ahuja N, Issa JP, Markowitz S, Willson JK, Hamilton SR, Kinzler KW, et al. Incidence and functional consequences of hMLH1 promoter hypermethylation in colorectal carcinoma. *Proc Natl Acad Sci U S A*. 1998; 95:6870–6875. [PubMed: 9618505]
- Hinoue T, Weisenberger DJ, Pan F, Campan M, Kim M, Young J, Whitehall VL, Leggett BA, Laird PW. Analysis of the association between CIMP and BRAF in colorectal cancer by DNA methylation profiling. *PLoS One*. 2009; 4:e8357. [PubMed: 20027224]
- Hoadley KA, Yau C, Hinoue T, Wolf DM, et al. Cell-of-Origin Patterns Dominate the Molecular Classification of 10,000 Tumors from 33 Types of Cancer. *Cell*. 2018 Under Review.
- Hoadley KA, Yau C, Wolf DM, Cherniack AD, Tamborero D, Ng S, Leiserson MD, Niu B, McLellan MD, Uzunangelov V, et al. Multiplatform analysis of 12 cancer types reveals molecular classification within and across tissues of origin. *Cell*. 2014; 158:929–944. [PubMed: 25109877]
- International Agency for Research on Cancer. WHO Classification of Tumours of the Digestive System. 4. 2010.
- Isella C, Terrasi A, Bellomo SE, Petti C, Galatola G, Muratore A, Mellano A, Senetta R, Cassenti A, Sonetto C, et al. Stromal contribution to the colorectal cancer transcriptome. *Nat Genet*. 2015; 47:312–319. [PubMed: 25706627]
- Janjigian YY, Sanchez-Vega F, Jonsson P, Chatila WK, Hechtman JF, Ku GY, Riches JC, Tuvy Y, Kundra R, Bouvier N, et al. Genetic Predictors of Response to Systemic Therapy in Esophagogastric Cancer. *Cancer Discov*. 2018; 8:49–58. [PubMed: 29122777]
- Javier BM, Yaeger R, Wang L, Sanchez-Vega F, Zehir A, Middha S, Sadowska J, Vakiani E, Shia J, Klimstra D, et al. Recurrent, truncating SOX9 mutations are associated with SOX9 overexpression, KRAS mutation, and TP53 wild type status in colorectal carcinoma. *Oncotarget*. 2016; 7:50875–50882. [PubMed: 27248473]
- Jin Z, Yoon HH. The promise of PD-1 inhibitors in gastro-esophageal cancers: microsatellite instability vs. PD-L1. *J Gastrointest Oncol*. 2016; 7:771–788. [PubMed: 27747091]
- Kandoth C, McLellan MD, Vandin F, Ye K, Niu B, Lu C, Xie M, Zhang Q, McMichael JF, Wyczalkowski MA, et al. Mutational landscape and significance across 12 major cancer types. *Nature*. 2013; 502:333–339. [PubMed: 24132290]
- Kim J, Fox C, Peng S, Pusung M, Pectasides E, Matthee E, Hong YS, Do IG, Jang J, Thorner AR, et al. Preexisting oncogenic events impact trastuzumab sensitivity in ERBB2-amplified gastroesophageal adenocarcinoma. *J Clin Invest*. 2014; 124:5145–5158. [PubMed: 25401468]
- Kim J, Mouw KW, Polak P, Braunstein LZ, Kamburov A, Tiao G, Kwiatkowski DJ, Rosenberg JE, Van Allen EM, D'Andrea AD, et al. Somatic ERCC2 mutations are associated with a distinct genomic signature in urothelial tumors. *Nat Genet*. 2016; 48:600–606. [PubMed: 27111033]
- Koh J, Ock CY, Kim JW, Nam SK, Kwak Y, Yun S, Ahn SH, Park DJ, Kim HH, Kim WH, et al. Clinicopathologic implications of immune classification by PD-L1 expression and CD8-positive tumor-infiltrating lymphocytes in stage II and III gastric cancer patients. *Oncotarget*. 2017; 8:26356–26367. [PubMed: 28412752]
- Korn JM, Kuruvilla FG, McCarroll SA, Wysoker A, Nemesh J, Cawley S, Hubbell E, Veitch J, Collins PJ, Darvishi K, et al. Integrated genotype calling and association analysis of SNPs, common copy number polymorphisms and rare CNVs. *Nat Genet*. 2008; 40:1253–1260. [PubMed: 18776909]
- Kostic AD, Ojesina AI, Pedamallu CS, Jung J, Verhaak RG, Getz G, Meyerson M. PathSeq: software to identify or discover microbes by deep sequencing of human tissue. *Nat Biotechnol*. 2011; 29:393–396. [PubMed: 21552235]
- Lawrence MS, Stojanov P, Mermel CH, Robinson JT, Garraway LA, Golub TR, Meyerson M, Gabriel SB, Lander ES, Getz G. Discovery and saturation analysis of cancer genes across 21 tumour types. *Nature*. 2014; 505:495–501. [PubMed: 24390350]

- Le DT, Durham JN, Smith KN, Wang H, Bartlett BR, Aulakh LK, Lu S, Kemberling H, Wilt C, Luber BS, et al. Mismatch repair deficiency predicts response of solid tumors to PD-1 blockade. *Science*. 2017; 357:409–413. [PubMed: 28596308]
- Leffers H, Dejgaard K, Celis JE. Characterisation of two major cellular poly(rC)-binding human proteins, each containing three K-homologous (KH) domains. *Eur J Biochem*. 1995; 230:447–453. [PubMed: 7607214]
- Leung SY, Yuen ST, Chung LP, Chu KM, Chan AS, Ho JC. hMLH1 promoter methylation and lack of hMLH1 expression in sporadic gastric carcinomas with high-frequency microsatellite instability. *Cancer Res*. 1999; 59:159–164. [PubMed: 9892201]
- Li B, Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinformatics*. 2011; 12:323. [PubMed: 21816040]
- Li M, Fang X, Baker DJ, Guo L, Gao X, Wei Z, Han S, van Deursen JM, Zhang P. The ATM-p53 pathway suppresses aneuploidy-induced tumorigenesis. *Proc Natl Acad Sci U S A*. 2010; 107:14188–14193. [PubMed: 20663956]
- Ljunggren HG, Malmberg KJ. Prospects for the use of NK cells in immunotherapy of human cancer. *Nat Rev Immunol*. 2007; 7:329–339. [PubMed: 17438573]
- Lynch HT, Snyder CL, Shaw TG, Heinen CD, Hitchins MP. Milestones of Lynch syndrome: 1895–2015. *Nat Rev Cancer*. 2015; 15:181–194. [PubMed: 25673086]
- Malta, TM., Sokolov, A., Gentles, AJ., et al. *Cancer Cell Under Review*. 2018.
- Matusaka K, Kaneda A, Nagae G, Ushiku T, Kikuchi Y, Hino R, Uozaki H, Seto Y, Takada K, Aburatani H, et al. Classification of Epstein-Barr virus-positive gastric cancers by definition of DNA methylation epigenotypes. *Cancer Res*. 2011; 71:7187–7197. [PubMed: 21990320]
- McCarroll SA, Kuruvilla FG, Korn JM, Cawley S, Nemes J, Wysoker A, Shapero MH, de Bakker PI, Maller JB, Kirby A, et al. Integrated detection and population-genetic analysis of SNPs and copy number variation. *Nat Genet*. 2008; 40:1166–1174. [PubMed: 18776908]
- McConnell BB, Kim SS, Yu K, Ghaleb AM, Takeda N, Manabe I, Nusrat A, Nagai R, Yang VW. Kruppel-like factor 5 is important for maintenance of crypt architecture and barrier function in mouse intestine. *Gastroenterology*. 2011; 141:1302–1313. 1313 e1301–1306. [PubMed: 21763241]
- Mermel CH, Schumacher SE, Hill B, Meyerson ML, Beroukhi R, Getz G. GISTIC2.0 facilitates sensitive and confident localization of the targets of focal somatic copy-number alteration in human cancers. *Genome Biol*. 2011; 12:R41. [PubMed: 21527027]
- Mishra L, Shetty K, Tang Y, Stuart A, Byers SW. The role of TGF-beta and Wnt signaling in gastrointestinal stem cells and cancer. *Oncogene*. 2005; 24:5775–5789. [PubMed: 16123810]
- Moons LM, Bax DA, Kuipers EJ, Van Dekken H, Haringsma J, Van Vliet AH, Siersema PD, Kusters JG. The homeodomain protein CDX2 is an early marker of Barrett's oesophagus. *J Clin Pathol*. 2004; 57:1063–1068. [PubMed: 15452161]
- Muro K, Chung HC, Shankaran V, Geva R, Catenacci D, Gupta S, Eder JP, Golan T, Le DT, Burtness B, et al. Pembrolizumab for patients with PD-L1-positive advanced gastric cancer (KEYNOTE-012): a multicentre, open-label, phase 1b trial. *Lancet Oncol*. 2016; 17:717–726. [PubMed: 27157491]
- Nandan MO, Ghaleb AM, Liu Y, Bialkowska AB, McConnell BB, Shroyer KR, Robine S, Yang VW. Inducible intestine-specific deletion of Kruppel-like factor 5 is characterized by a regenerative response in adult mouse colon. *Dev Biol*. 2014; 387:191–202. [PubMed: 24440658]
- Noah TK, Donahue B, Shroyer NF. Intestinal development and differentiation. *Exp Cell Res*. 2011; 317:2702–2710. [PubMed: 21978911]
- Olshen AB, Venkatraman ES, Lucito R, Wigler M. Circular binary segmentation for the analysis of array-based DNA copy number data. *Biostatistics*. 2004; 5:557–572. [PubMed: 15475419]
- Palles C, Cazier JB, Howarth KM, Domingo E, Jones AM, Broderick P, Kemp Z, Spain SL, Guarino E, Salguero I, et al. Germline mutations affecting the proofreading domains of POLE and POLD1 predispose to colorectal adenomas and carcinomas. *Nat Genet*. 2013; 45:136–144. [PubMed: 23263490]

- Phipps AI, Passarelli MN, Chan AT, Harrison TA, Jeon J, Hutter CM, Berndt SI, Brenner H, Caan BJ, Campbell PT, et al. Common genetic variation and survival after colorectal cancer diagnosis: a genome-wide analysis. *Carcinogenesis*. 2016; 37:87–95. [PubMed: 26586795]
- Ramos AH, Lichtenstein L, Gupta M, Lawrence MS, Pugh TJ, Saksena G, Meyerson M, Getz G. Oncotator: cancer variant annotation tool. *Hum Mutat*. 2015; 36:E2423–2429. [PubMed: 25703262]
- Ratan A, Olson TL, Loughran TP Jr, Miller W. Identification of indels in next-generation sequencing data. *BMC Bioinformatics*. 2015; 16:42. [PubMed: 25879703]
- Reed KR, Korobko IV, Ninkina N, Korobko EV, Hopkins BR, Platt JL, Buchman V, Clarke AR. Hunk/Mak-v is a negative regulator of intestinal cell proliferation. *BMC Cancer*. 2015; 15:110. [PubMed: 25881306]
- Reich M, Liefeld T, Gould J, Lerner J, Tamayo P, Mesirov JP. GenePattern 2.0. *Nat Genet*. 2006; 38:500–501. [PubMed: 16642009]
- Roepman P, Schlicker A, Tabernero J, Majewski I, Tian S, Moreno V, Snel MH, Chresta CM, Rosenberg R, Nitsche U, et al. Colorectal cancer intrinsic subtypes predict chemotherapy benefit, deficient mismatch repair and epithelial-to-mesenchymal transition. *Int J Cancer*. 2014; 134:552–562. [PubMed: 23852808]
- Salari K, Spulak ME, Cuff J, Forster AD, Giacomini CP, Huang S, Ko ME, Lin AY, van de Rijn M, Pollack JR. CDX2 is an amplified lineage-survival oncogene in colorectal cancer. *Proc Natl Acad Sci U S A*. 2012; 109:E3196–3205. [PubMed: 23112155]
- Salomonis N, Dexheimer PJ, Omberg L, Schroll R, Bush S, Huo J, Schriml L, Ho Sui S, Keddache M, Mayhew C, et al. Integrated Genomic Analysis of Diverse Induced Pluripotent Stem Cells from the Progenitor Cell Biology Consortium. *Stem Cell Reports*. 2016; 7:110–125. [PubMed: 27293150]
- Sartore-Bianchi A, Trusolino L, Martino C, Bencardino K, Lonardi S, Bergamo F, Zagonel V, Leone F, Depetris I, Martinelli E, et al. Dual-targeted therapy with trastuzumab and lapatinib in treatment-refractory, KRAS codon 12/13 wild-type, HER2-positive metastatic colorectal cancer (HERACLES): a proof-of-concept, multicentre, openlabel, phase 2 trial. *Lancet Oncol*. 2016; 17:738–746. [PubMed: 27108243]
- Schepers A, Clevers H. Wnt signaling, stem cells, and cancer of the gastrointestinal tract. *Cold Spring Harb Perspect Biol*. 2012; 4:a007989. [PubMed: 22474007]
- Secrier M, Li X, de Silva N, Eldridge MD, Contino G, Bornschein J, MacRae S, Grehan N, O'Donovan M, Miremadi A, et al. Mutational signatures in esophageal adenocarcinoma define etiologically distinct subgroups with therapeutic relevance. *Nat Genet*. 2016; 48:1131–1141. [PubMed: 27595477]
- Sheltzer JM, Blank HM, Pfau SJ, Tange Y, George BM, Humpton TJ, Brito IL, Hiraoka Y, Niwa O, Amon A. Aneuploidy drives genomic instability in yeast. *Science*. 2011; 333:1026–1030. [PubMed: 21852501]
- Sheltzer JM, Ko JH, Replogle JM, Habibe Burgos NC, Chung ES, Meehl CM, Sayles NM, Passerini V, Storchova Z, Amon A. Single-chromosome Gains Commonly Function as Tumor Suppressors. *Cancer Cell*. 2017; 31:240–255. [PubMed: 28089890]
- Sherwood RI, Chen TY, Melton DA. Transcriptional dynamics of endodermal organ formation. *Dev Dyn*. 2009; 238:29–42. [PubMed: 19097184]
- Sia EA, Kokoska RJ, Dominska M, Greenwell P, Petes TD. Microsatellite instability in yeast: dependence on repeat unit size and DNA mismatch repair genes. *Mol Cell Biol*. 1997; 17:2851–2858. [PubMed: 9111357]
- Sokolov A, Paull EO, Stuart JM. One-Class Detection of Cell States in Tumor Subtypes. *Pac Symp Biocomput*. 2016; 21:405–416. [PubMed: 26776204]
- Tan IB, Ivanova T, Lim KH, Ong CW, Deng N, Lee J, Tan SH, Wu J, Lee MH, Ooi CH, et al. Intrinsic subtypes of gastric cancer, based on gene expression pattern, predict survival and respond differently to chemotherapy. *Gastroenterology*. 2011; 141:476–485. 485 e471–411. [PubMed: 21684283]
- Thompson SL, Compton DA. Proliferation of aneuploid human cells is limited by a p53-dependent mechanism. *J Cell Biol*. 2010; 188:369–381. [PubMed: 20123995]

- Torre LA, Siegel RL, Ward EM, Jemal A. Global Cancer Incidence and Mortality Rates and Trends-- An Update. *Cancer Epidemiol Biomarkers Prev.* 2016; 25:16–27. [PubMed: 26667886]
- Torres EM, Sokolsky T, Tucker CM, Chan LY, Boselli M, Dunham MJ, Amon A. Effects of aneuploidy on cellular physiology and cell division in haploid yeast. *Science.* 2007; 317:916–924. [PubMed: 17702937]
- Tripathi V, Sixt KM, Gao S, Xu X, Huang J, Weigert R, Zhou M, Zhang YE. Direct Regulation of Alternative Splicing by SMAD3 through PCBP1 Is Essential to the Tumor-Promoting Role of TGF-beta. *Mol Cell.* 2016; 64:1010.
- Wagner JA, Rosario M, Romee R, Berrien-Elliott MM, Schneider SE, Leong JW, Sullivan RP, Jewell BA, Becker-Hapak M, Schappe T, et al. CD56bright NK cells exhibit potent antitumor responses following IL-15 priming. *J Clin Invest.* 2017; 127:4042–4058. [PubMed: 28972539]
- Wang K, Singh D, Zeng Z, Coleman SJ, Huang Y, Savich GL, He X, Mieczkowski P, Grimm SA, Perou CM, et al. MapSplice: accurate mapping of RNA-seq reads for splice junction discovery. *Nucleic Acids Res.* 2010; 38:e178. [PubMed: 20802226]
- Wang K, Yuen ST, Xu J, Lee SP, Yan HH, Shi ST, Siu HC, Deng S, Chu KM, Law S, et al. Whole-genome sequencing and comprehensive molecular profiling identify new driver mutations in gastric cancer. *Nat Genet.* 2014; 46:573–582. [PubMed: 24816253]
- Weisenberger DJ, Siegmund KD, Campan M, Young J, Long TI, Faasse MA, Kang GH, Widschwendter M, Weener D, Buchanan D, et al. CpG island methylator phenotype underlies sporadic microsatellite instability and is tightly associated with BRAF mutation in colorectal cancer. *Nat Genet.* 2006; 38:787–793. [PubMed: 16804544]
- Widschwendter M, Fiegl H, Egle D, Mueller-Holzner E, Spizzo G, Marth C, Weisenberger DJ, Campan M, Young J, Jacobs I, et al. Epigenetic stem cell signature in cancer. *Nat Genet.* 2007; 39:157–158. [PubMed: 17200673]
- Williams BR, Prabhu VR, Hunter KE, Glazier CM, Whittaker CA, Housman DE, Amon A. Aneuploidy affects proliferation and spontaneous immortalization in mammalian cells. *Science.* 2008; 322:703–709. [PubMed: 18974345]
- Xiao Y, Freeman GJ. The microsatellite unstable subset of colorectal cancer is a particularly good candidate for checkpoint blockade immunotherapy. *Cancer Discov.* 2015; 5:16–18. [PubMed: 25583798]
- Zhou Q, Talvinen K, Sundstrom J, Elzagheid A, Pospiech H, Syvaoja JE, Collan Y. Mutations/ polymorphisms in the 55 kDa subunit of DNA polymerase epsilon in human colorectal cancer. *Cancer Genomics Proteomics.* 2009; 6:297–304. [PubMed: 20065316]

CONSORTIUM AUTHOR LIST

The members of The Cancer Genome Atlas Research Network for this project are:

NCI/NHGRI Project Team

Samantha J. Caesar-Johnson, John A. Demchok, Ina Felau, Martin L. Ferguson, Carolyn M. Hutter, Melpomeni Kasapi, Heidi J. Sofia, Roy Tarnuzzer, Zhining Wang, Liming Yang, Jean C. Zenklusen, Jiashan (Julia) Zhang

TCGA DCC

Sudha Chudamani, Jia Liu, Laxmi Lolla, Rashi Naresh, Todd Pihl, Qiang Sun, Yunhu Wan, Ye Wu

Genome Data Analysis Centers (GDACs)

The Broad Institute

Juok Cho, Timothy DeFreitas, Scott Frazer, Nils Gehlenborg, Gad Getz, David I. Heiman, Jaegil Kim, Michael S. Lawrence, Pei Lin, Sam Meier, Michael S. Noble, Gordon Saksena, Doug Voet, Hailei Zhang

Institute for Systems Biology

Brady Bernard, Nyasha Chambwe, Varsha Dhankani, Theo Knijnenburg, Roger Kramer, Kalle Leinonen, Yuexin Liu, Michael Miller, Sheila Reynolds, Ilya Shmulevich, Vesteinn Thorsson, Wei Zhang

MD Anderson Cancer Center

Rehan Akbani, Bradley M. Broom, Apurva M. Hegde, Zhenlin Ju, Rupa S. Kanchi, Anil Korkut, Jun Li, Han Liang, Shiyun Ling, Wenbin Liu, Yiling Lu, Gordon B. Mills, Kwok-Shing Ng, Arvind Rao, Michael Ryan, Jing Wang, John N. Weinstein, Jiexin Zhang

Memorial Sloan Kettering Cancer Center

Adam Abeshouse, Joshua Armenia, Debyani Chakravarty, Walid K. Chatila, Ino de Bruijn, Jianjiong Gao, Benjamin E. Gross, Zachary J. Heins, Ritika Kundra, Konnor La, Marc Ladanyi, Augustin Luna, Moriah G. Nissan, Angelica Ochoa, Sarah M. Phillips, Ed Reznik, Francisco Sanchez-Vega, Chris Sander, Nikolaus Schultz, Robert Sheridan, S. Onur Sumer, Yichao Sun, Barry S. Taylor, Jioajiao Wang, Hongxin Zhang

Oregon Health and Science University

Pavana Anur, Myron Peto, Paul Spellman

University of California Santa Cruz

Christopher Benz, Joshua M. Stuart, Christopher K. Wong, Christina Yau

University of North Carolina at Chapel Hill

D. Neil Hayes, Joel S. Parker, Matthew D. Wilkerson

Genome Characterization Centers (GCC)

BC Cancer Agency

Adrian Ally, Miruna Balasundaram, Reanne Bowlby, Denise Brooks, Rebecca Carlsen, Eric Chuah, Noreen Dhalla, Robert Holt, Steven J.M. Jones, Katayoon Kasaian, Darlene Lee, Yussanne Ma, Marco A. Marra, Michael Mayo, Richard A. Moore, Andrew J. Mungall, Karen Mungall, A. Gordon Robertson, Sara Sadeghi, Jacqueline E. Schein, Payal Sipahimalani, Angela Tam, Nina Thiessen, Kane Tse, Tina Wong

The Broad Institute

Ashton C. Berger, Rameen Beroukhim, Andrew D. Cherniack, Carrie Cibulskis, Stacey B. Gabriel, Galen F. Gao, Gavin Ha, Matthew Meyerson, Gordon Saksena, Steven E. Schumacher, Juliann Shih

Harvard

Melanie H. Kucherlapati, Raju S. Kucherlapati

Johns Hopkins

Stephen Baylin, Leslie Cope, Ludmila Danilova

University of Southern California

Moiz S. Bootwalla, Phillip H. Lai, Dennis T. Maglinte, David J. Van Den Berg, Daniel J. Weisenberger

University of North Carolina at Chapel Hill

J. Todd Auman, Saianand Balu, Tom Bodenheimer, Cheng Fan, D. Neil Hayes, Katherine A. Hoadley, Alan P. Hoyle, Stuart R. Jefferys, Corbin D. Jones, Shaowu Meng, Piotr A. Mieczkowski, Lisle E. Mose, Joel S. Parker, Amy H. Perou, Charles M. Perou, Jeffrey Roach, Yan Shi, Janae V. Simons, Tara Skelly, Matthew G. Soloway, Donghui Tan, Umadevi Veluvolu, Matthew D. Wilkerson

Van Andel Research Institute

Toshinori Hinoue, Peter W. Laird, Hui Shen

Genome Sequencing Centers (GSC)**Baylor College of Medicine**

Michelle Bellair, Kyle Chang, Kyle Covington, Chad J. Creighton, Huyen Dinh, HarshaVardhan Doddapaneni, Lawrence A. Donehower, Jennifer Drummond, Richard A. Gibbs, Robert Glenn, Walker Hale, Yi Han, Jianhong Hu, Viktoriya Korchina, Sandra Lee, Lora Lewis, Wei Li, Xiuping Liu, Margaret Morgan, Donna Morton, Donna Muzny, Jireh Santibanez, Margi Sheth, Eve Shinbrot, Linghua Wang, Min Wang, David A. Wheeler, Liu Xi, Fengmei Zhao

The Broad Institute

Carrie Cibulskis, Stacy B. Gabriel, Julian Hess

Washington University at St. Louis

Elizabeth L. Appelbaum, Matthew Bailey, Matthew G. Cordes, Li Ding, Catrina C. Fronick, Lucinda A. Fulton, Robert S. Fulton, Cyriac Kandoth, Elaine R. Mardis, Michael D. McLellan, Christopher A. Miller, Heather K. Schmidt, Richard K. Wilson

Biospecimen Core Resource

The International Genomics Consortium

Daniel Crain, Erin Curley, Johanna Gardner, Kevin Lau, David Mallery, Scott Morris, Joseph Paulauskis, Robert Penny, Candace Shelton, Troy Shelton, Mark Sherman, Eric Thompson, Peggy Yena

Nationwide Children's Organization

Jay Bowen, Julie M. Gastier-Foster, Mark Gerken, Kristen M. Leraas, Tara M. Lichtenberg, Nilsa C. Ramirez, Lisa Wise, Erik Zmuda

Tissue Source Sites

Australian Prostate Cancer Research Center

Niall Corcoran, Tony Costello, Christopher Hovens

Barretos Cancer Hospital

Andre L. Carvalho, Ana C. de Carvalho, José H. Fregnani, Adhemar Longatto-Filho, Rui M. Reis, Cristovam Scapulatempo-Neto, Henrique C. S. Silveira, Daniel O. Vidal

Barrow Neurological Institute

Andrew Burnette, Jennifer Eschbacher, Beth Hermes, Ardene Noss, Rosy Singh

Baylor College of Medicine

Matthew L. Anderson, Patricia D. Castro, Michael Ittmann

BC Cancer Agency

David Huntsman

BioreclamationIVT

Bernard Kohl, Xuan Le, Richard Thorp

Boston Medical Center

Chris Andry, Elizabeth R. Duffy

Botkin Hospital

Vladimir Lyadov, Oxana Paklina, Galiya Setdikova, Alexey Shabunin, Mikhail Tavobilov

Brain Tumor Center at the University of Cincinnati Gardner Neuroscience Institute

Christopher McPherson, Ronald Warnick

Brigham and Women's Hospital

Ross Berkowitz, Daniel Cramer, Colleen Feltmate, Neil Horowitz, Adam Kibel, Michael Muto, Chandrajit P. Raut

Capital Biosciences, Inc

Andrei Malykh

Case Comprehensive Cancer CenterJill S. Barnholtz-Sloan, Wendi Barrett, Karen Devine, Jordonna Fulop, Quinn T. Ostrom,
Kristen Shimmel, Yingli Wolinsky**Case Western Reserve School of Medicine**

Andrew E. Sloan

Catholic University of the Sacred Heart

Agostino De Rose, Felice Giuliante

Cedars-Sinai Medical Center

Marc Goodman, Beth Y. Karlan

Central Arkansas Veterans Healthcare System

Curt H. Hagedorn

Centura Health

John Eckman, Jodi Harr, Jerome Myers, Kelinda Tucker, Leigh Anne Zach

Chan Soon-Shiong Institute of Molecular Medicine at WindberBrenda Deyarmin, Hai Hu, Leonid Kvecher, Caroline Larson, Richard J. Mural, Stella
Somari**Charles University**

Ales Vicha, Tomas Zelinka

Christiana Care Health System

Joseph Bennett, Mary Iacocca, Brenda Rabeno, Patricia Swanson

CHU of Montreal

Mathieu Latour

CHU of Quebec

Louis Lacombe, Bernard Têtu

CHU of Quebec, Laval University Research Center of Chus

Alain Bergeron

Cleveland Clinic Foundation

Mary McGraw, Susan M. Staugaitis

Columbia University

John Chabot, Hanina Hibshoosh, Antonia Sepulveda, Tao Su, Timothy Wang

Cureline, Inc

Olga Potapova, Olga Voronina

Curie Institute

Laurence Desjardins, Odette Mariani, Sergio Roman-Roman, Xavier Sastre, Marc-Henri Stern

Dana-Farber Cancer Institute

Feixiong Cheng, Sabina Signoretti

Dignity Health Mercy Gilbert Medical Center

Jennifer Eschbacher

Duke University Medical Center

Andrew Berchuck, Darell Bigner, Eric Lipp, Jeffrey Marks, Shannon McCall, Roger McLendon, Angeles Secord, Alexis Sharp

Emory University

Madhusmita Behera, Daniel J. Brat, Amy Chen, Keith Delman, Seth Force, Fadlo Khuri, Kelly Magliocca, Shishir Maithel, Jeffrey J. Olson, Taofeek Owonikoko, Alan Pickens, Suresh Ramalingam, Dong M. Shin, Gabriel Sica, Erwin G. Van Meir, Hongzheng Zhang

Erasmus Medical Center

Wil Eijckenboom, Ad Gillis, Esther Korpershoek, Leendert Looijenga, Wolter Oosterhuis, Hans Stoop, Kim E. van Kessel, Ellen C. Zwarthoff

Foundation of the Carlo Besta Neurological Institute, IRCCS

Chiara Calatozzolo, Lucia Cuppini, Stefania Cuzzubbo, Francesco DiMeco, Gaetano Finocchiaro, Luca Mattei, Alessandro Perin, Bianca Pollo

Fred Hutchinson Cancer Research Center

Chu Chen, John Houck, Pawadee Lohavanichbutr

Friedrich-Alexander-University

Arndt Hartmann, Christine Stoehr, Robert Stoehr, Helge Taubert, Sven Wach, Bernd Wullich

Greater Poland Cancer Center

Witold Kycler, Dawid Murawa, Maciej Wiznerowicz

Greenville Health System Institute for Translational Oncology Research

Ki Chung, W. Jeffrey Edenfield, Julie Martin

Gustave Roussy institute

Eric Baudin

Harvard University

Glenn Bublely, Raphael Bueno, Assunta De Rienzo, William G. Richards

Henry Ford Health System

Ana deCarvalho, Steven Kalkanis, Tom Mikkelsen, Tom Mikkelsen, Houtan Noushmehr,
Lisa Scarpace

Hospices Civils de Lyon

Nicolas Girard

Hospital Clinic

Marta Aymerich, Elias Campo, Eva Giné, Armando López Guillermo

Hue Central Hospital

Nguyen Van Bang, Phan Thi Hanh, Bui Duc Phu

Human Tissue Resource Network

Yufang Tang

Huntsman Cancer Institute

Howard Colman, Kimberley Evason

Icahn School of Medicine at Mount Sinai

Peter R. Dottino, John A. Martignetti

Imperial College London

Hani Gabra

Indivumed GmbH

Hartmut Juhl

Institute of Human Virology Nigeria

Teniola Akeredolu

Institute of Urgent Medicine

Serghei Stepa

The International Genomics Consortium

Daniel Crain, Erin Curley, Johanna Gardner, David Mallery, Scott Morris, Joseph Paulauskis, Robert Penny, Candace Shelton, Troy Shelton, Eric Thompson

John Wayne Cancer Institute

Dave Hoon

Keimyung University

Keunsoo Ahn, Koo Jeong Kang

Ludwigh Maximilians University Munich

Felix Beuschlein

Maine Medical Center

Anne Breggia

Massachusetts General Hospital

Michael Birrer

Mayo Clinic

Debra Bell, Mitesh Borad, Alan H. Bryce, Erik Castle, Vishal Chandan, John Cheville, John A. Copland, Michael Farnell, Thomas Flotte, Nasra Giama, Thai Ho, Michael Kendrick, Jean-Pierre Kocher, Karla Kopp, Catherine Moser, David Nagorney, Daniel O'Brien, Brian Patrick O'Neill, Tushar Patel, Gloria Petersen, Florencia Que, Michael Rivera, Lewis Roberts, Robert Smallridge, Thomas Smyrk, Melissa Stanton, R. Houston Thompson, Michael Torbenson, Ju Dong Yang, Lizhi Zhang

McGill University Health Center

Fadi Brimo

MD Anderson Cancer Center

Jaffer A. Ajani, Ana Maria Angulo Gonzalez, Carmen Behrens, Jolanta Bondaruk, Russell Broaddus, Bradley Broom, Bogdan Czerniak, Bitu Esmaeli, Junya Fujimoto, Jeffrey Gershenwald, Charles Guo, Alexander J. Lazar, Christopher Logothetis, Funda Meric-Bernstam, Funda Meric-Bernstam, Cesar Moran, Lois Ramondetta, David Rice, Anil Sood, Pheroze Tamboli, Timothy Thompson, Patricia Troncoso, Anne Tsao, Ignacio Wistuba

Melanoma Institute Australia

Candace Carter, Lauren Haydu, Peter Hersey, Valerie Jakrot, Hojabr Kakavand, Richard Kefford, Kenneth Lee, Georgina Long, Graham Mann, Michael Quinn, Robyn Saw, Richard Scolyer, Kerwin Shannon, Andrew Spillane, Jonathan Stretch, Maria Synott, John Thompson, James Wilmott

Memorial Sloan Kettering Cancer Center

Hikmat Al-Ahmadie, Timothy A. Chan, Ronald Ghossein, Anuradha Gopalan, Victor Reuter, Samuel Singer, Bhuvanesh Singh

Ministry of Health of Vietnam

Nguyen Viet Tien

Molecular Response

Thomas Broudy, Cyrus Mirsaidi, Praveen Nair

Nancy N. and J.C. Lewis Cancer & Research Pavilion at St. Joseph's/Candler

Paul Drwiega, Judy Miller, Jennifer Smith, Howard Zaren

National Cancer Center Korea

Joong-Won Park

National Cancer Hospital of Vietnam

Nguyen Phi Hung

National Cancer Institute

Electron Kebebew, W. Marston Linehan, Adam R. Metwalli, Karel Pacak, Peter A. Pinto, Mark Schiffman, Laura S. Schmidt, Cathy D. Vocke, Nicolas Wentzensen, Robert Worrell, Hannah Yang

National Cancer Center of Georgia

Armaz Mariamidze

Norfolk & Norwich University Hospital

Marc Moncrieff

NYU Langone Medical Center

Chandra Goparaju, Jonathan Melamed, Harvey Pass

Ohio State University

Mohamed H. Abdel-Rahman, Dina Aziz, Sue Bell, Colleen M. Cebulla, Amy Davis, Rebecca Duell, J. Bradley Elder, Joe Hilty, Bahavna Kumar, James Lang, Norman L. Lehman, Randy Mandt, Phuong Nguyen, Robert Pilarski, Karan Rai, Lynn Schoenfield, Kelly Senecal, Paul Wakely

Oncology Institute

Natalia Botnariuc, Irina Caraman, Mircea Cernat, Inga Chemencedji, Adrian Clipca, Serghei Doruc, Ghenadie Gorincioi, Sergiu Mura, Maria Pirtac, Irina Stancul, Diana Tcaciuc

Ontario Tumour Bank

Monique Albert, Iakovina Alexopoulou, Angel Arnaout, John Bartlett, Jay Engel, Sebastien Gilbert, Jeremy Parfitt, Harman Sekhon

The Oregon Clinic

Paul Hansen

Oregon Health & Science University

George Thomas

Papworth Hospital NHS Foundation Trust

Doris M. Rassl, Robert C. Rintoul

Providence Health and Services

Carlo Bifulco, Raina Tamakawa, Walter Urba

QIMR Berghofer Medical Research Institute

Nicholas Hayward

Radboud Medical University Center

Henri Timmers

Regina Elena National Cancer Institute

Anna Antenucci, Francesco Facciolo, Gianluca Grazi, Mirella Marino, Roberta Merola

Reinier de Graaf Hospital

Ronald de Krijger

René Descartes University

Anne-Paule Gimenez-Roqueplo

Research Center of Chus Sherbrooke, Québec

Alain Piché

Research Institute of the McGill University Health Centre

Simone Chevalier, Ginette McKercher

The Research Institute at Nationwide Children's Hospital

Nilsa Ramirez

Rockefeller University

Kivanc Birsoy

Rose Ella Burkhardt Brain Tumor and Neuro-Oncology Center

Gene Barnett, Cathy Brewer, Carol Farver, Theresa Naska, Nathan A. Pennell, Daniel Raymond, Cathy Schilero, Kathy Smolenski, Felicia Williams

Roswell Park Cancer Institute

Carl Morrison

Rush University

Jeffrey A. Borgia, Michael J. Liptay, Mark Pool, Christopher W. Seder

Saarland University

Kerstin Junker

Sage Bionetworks

Larsson Omberg

Saint-Petersburg City Clinical Oncology Hospital

Mikhail Dinkin, George Manikhas

Sapienza University of Rome

Domenico Alvaro, Maria Consiglia Bragazzi, Vincenzo Cardinale, Guido Carpino, Eugenio Gaudio

Spectrum Health

David Chesla, Sandra Cottingham

St. Petersburg Academic University RAS

Michael Dubina, Fedor Moiseenko

Stanford University

Renumathy Dhanasekaran

Technical University of Munich

Karl-Friedrich Becker, Klaus-Peter Janssen, Julia Slotta-Huspenina

Tufts Medical Center

Ronald Lechan, James Powers, Arthur Tischler

University of Alabama at Birmingham Medical Center

William E. Grizzle, Katherine C. Sexton

UC Cancer Institute

Alison Kastl

UCSF-Helen Diller Family Comprehensive Cancer Center

Joel Henderson, Sima Porten

University Health Network

Sylvia L. Asa

University Hospital of Giessen and Marburg

Jens Waldmann

University Hospital in Wurzburg, Germany

Martin Fassnacht

University Hospital Essen

Dirk Schadendorf

University Hospitals Case Medical Center

Marta Couce

University Medical Center Hamburg-Eppendorf

Markus Graefen, Hartwig Huland, Guido Sauter, Thorsten Schlomm, Ronald Simon, Pierre Tennstedt

University of Abuja Teaching Hospital

Oluwole Olabode

University of Arizona

Mark Nelson

University of Calgary

Oliver Bathe

University of California

Peter R. Carroll, June M. Chan, Philip Disaia, Pat Glenn, Robin K. Kelley, Charles N. Landen, Joanna Phillips, Michael Prados, Jeffrey Simko, Karen Smith-McCune, Scott VandenBerg

University of Chicago Medicine

Kevin Roggin

University of Cincinnati

Ashley Fehrenbach, Ady Kendler

University of Cincinnati Cancer Institute

Suzanne Sifri, Ruth Steele

University of Colorado Cancer Center

Antonio Jimeno

University of Dundee

Francis Carey, Ian Forgie

University of Florence

Massimo Mannelli

University of Hawaii Cancer Center

Michael Carney, Brenda Hernandez

University of Heidelberg

Benito Campos, Christel Herold-Mende, Christin Jungk, Andreas Unterberg, Andreas von Deimling

University of Iowa Hospital & Clinics

Aaron Bossler, Joseph Galbraith, Laura Jacobus, Michael Knudson, Tina Knutson, Deqin Ma, Mohammed Milhem, Rita Sigmund

University of Kansas Cancer Center

Eryn M. Godwin, Sara Kendall, Cassandra Shipman

University of Kansas Medical Center

Andrew K. Godwin, Rashna Madan, Howard G. Rosenthal

University of Maryland School of Medicine

Clement Adebamowo, Sally N. Adebamowo

University of Melbourne

Alex Boussioutas

University of Michigan

David Beer, Carol Bradford, Thomas Carey, Thomas Giordano, Andrea Haddad, Jeffrey Moyer, Lisa Peterson, Mark Prince, Laura Rozek, Gregory Wolf

University of Montreal

Anne-Marie Mes-Masson, Fred Saad

University of New Mexico

Therese Bocklage

University of Oklahoma

Lisa Landrum, Robert Mannel, Kathleen Moore, Katherine Moxley, Russel Postier, Joan Walker, Rosemary Zuna

University of Pennsylvania

Michael Feldman, Federico Valdivieso

University of Pittsburgh

Rajiv Dhir, James Luketich

University of Puerto Rico

Edna M. Mora Pinero, Mario Quintero-Aguilo

University of São Paulo

Carlos Gilberto Carlotti Junior, Jose Sebastião Dos Santos, Rafael Kemp, Ajith Sankarankuty, Daniela Tirapelli, Natália D. Aredes

University of Sheffield Western Bank

James Catto

University of Washington

Kathy Agnew, Elizabeth Swisher

University of Western Australia

Jenette Creaney, Bruce Robinson

University of Wisconsin School of Medicine and Public Health

Carl Simon Shelley

UQ Thoracic Research Centre

Rayleen Bowman, Kwun M. Fong, Ian Yang

Valley Health System

Robert Korst

Vanderbilt University Medical Center

W. Kimryn Rathmell

Walter Reed National Medical Center

J. Leigh Fantacone-Campbell, Jeffrey A. Hooke, Albert J. Kovatich, Craig D. Shriver

Washington University

John DiPersio, Bettina Drake, Ramaswamy Govindan, Sharon Heath, Timothy Ley, Brian Van Tine, Peter Westervelt

Weill Cornell Medical College

Mark A. Rubin

Yonsei University College of Medicine

Jung Il Lee

Pan-Gastrointestinal Cancer Analysis Working Group Participants**Baylor College of Medicine**

Yumeng Wang, David A. Wheeler

BC Cancer Agency

Reanne Bowlby, Andrew J. Mungall

The Broad Institute

Adam J. Bass, Rameen Beroukhim, Susan Bullman, Andrew D. Cherniack, Mirazul Islam, Jaegil Kim, Yang Liu, Sam Meier, Chandra Sekhar Pedamallu, Steven E. Schumacher, Nilay S. Sethi, Juliann Shih

Case Western Reserve School of Medicine

Joseph E. Willis

Columbia University

Evan O. Pauli

Dana-Farber Cancer Institute

Adam J. Bass, Andrew D. Cherniack, Yang Liu, Nilay S. Sethi

Duke University Medical Center

Katherine S. Garman, Shannon J. McCall

George Washington University

Lopa Mishra, *Sobia Zaida*

Greater Poland Cancer Center

Maciej Wiznerowicz

Harvard University

Melanie H. Kucherlapati, Raju S. Kucherlapati, Hiroyuki Yoshida, Artem Sokolov

Henry Ford Health System

Tathiane M. Malta, Houtan Noushmehr

Institute for Systems Biology

William J. R. Longabaugh, Michael Miller, Ilya Shmulevich, Vésteinn Thorsson

Massachusetts General Hospital

Hiroyuki Yoshida

Mayo Clinic

Kenneth K. Wang

MD Anderson Cancer Center

Rehan Akbani, Apurva M. Hegde, Rupa S. Kanchi, Alexander J. Lazar, Shiyun Ling, Yuexin Liu, André Schultz, John N. Weinstein

Memorial Sloan Kettering Cancer Center

Francisco Sanchez-Vega

National Cancer Institute

Charles S. Rabkin

Stanford University

Christina Curtis, Andrew J. Gentles, Olivier Gevaert, Haruka Itakura, Jose A. Seoane

University of Alabama at Birmingham Medical Center

Akinyemi Ojesina

University of Calgary

Farshad Farshidfar

University of Kansas Medical Center

Andrew K. Godwin

University of Lausanne, Lausanne, Switzerland

Giovanni Cirello, Marco Mina

University of North Carolina at Chapel Hill

Margaret L. Gulley, Joel E. Tepper

University of Pennsylvania

Anil K. Rustgi

University of Southern California

Daniel J. Weisenberger

University of Texas Southwestern Medical Center

Havish S. Kantheti

Van Andel Research Institute

Toshinori Hinoue, Peter W. Laird, Hui Shen

Vanderbilt University Medical Center

Barbara G. Schneider

Institution Addresses

Australian Prostate Cancer Research Center, Epworth Hospital, VIC, Australia

Baylor College of Medicine, Department of Pathology and Immunology, One Baylor Plaza, Houston, TX 77030, USA

Baylor College of Medicine, Department of Obstetrics and Gynecology, One Baylor Plaza, Houston, Texas 77030, USA

Baylor College of Medicine, One Baylor Plaza, Houston, TX 77030, USA

Barretos Cancer Hospital, Av: Antenor Duarte Villela, 1331, Barretos, São Paulo, Brazil

Barrow Neurological Institute, St. Joseph's Hospital and Medical Center, Phoenix, AZ 85013, USABC Cancer Agency, 675 W 10th Ave, Vancouver, BC V5Z 1L3, Canada

Beth Israel Deaconess Medical Center, Harvard University Medical School, Boston, MA 02215, USA

BioreclamationIVT, 99 Talbot Blvd Chestertown, MD 21620, USA

Boston Medical Center, Boston, MA 02118, USA

Botkin Hospital, 2-y Botkinskiy pr-d, 5, Moskva, Russia, 125284

Brain Tumor Center at the University of Cincinnati Gardner Neuroscience Institute, and Department of Neurosurgery, University of Cincinnati College of Medicine, and Mayfield Clinic, 260 Stetson Street, Suite 2200, Cincinnati, OH, 45219, USA

Brain Tumor Center at the University of Cincinnati Neuroscience Institute, and Department of Neurosurgery, University of Cincinnati College of Medicine, and Mayfield Clinic, 234 Goodman Street, Cincinnati, OH, 45219, USA

Brain Tumor and Neuro-oncology Center, Department of Neurosurgery, University Hospitals Case Medical Center, Case Western Reserve School of Medicine, 11100 Euclid

Ave, Cleveland, OH, 44106, USA
Brigham and Women's Hospital, 75 Francis St, Boston
MA 02115, USA

Brigham and Women's Hospital, Division of Surgical Oncology, Department of Surgery, 75
Francis Street, Boston, MA 02115, USA

Brigham and Women's Hospital, Harvard Medical School, Department of Surgery, Boston,
MA 02115, USA

Capital Biosciences, Inc., 900 Clopper Rd, Suite 120, Gaithersburg, MD 20878, USA

Case Comprehensive Cancer Center, 11100 Euclid Ave - Wearn 152, Cleveland, OH
44106-5065, USA

Catholic University of the Sacred Heart, Hepatobiliary Surgery Unit, A. Gemelli Hospital,
Largo Agostino Gemelli 8, 00168 Rome, Italy

Cedars-Sinai Medical Center, 8700 Beverly Boulevard, Suite 290 West MOT, Los Angeles,
CA 90048, USA

Central Arkansas Veterans Healthcare System, Little Rock, AR 72205, USA

CHU of Quebec, Laval University Research Center of Chus 2705, boul. Laurier Bureau
TR72

QUÉBEC, Quebec G1V 4G2

Centura Health, 9100 E Mineral Cir, Centennial, CO 80112, USA

Chan Soon-Shiong Institute of Molecular Medicine at Windber, Windber, PA 15963, USA

Charles University, Czech Republic

CHU of Quebec, Hôtel-Dieu de Quebec-University Laval, 11 cote du palais, Quebec City,
G1R 2J6

CHUM, Montreal, Qc, Canada.

Cleveland Clinic Taussig Cancer Institute, 9500 Euclid Avenue, Cleveland, OH 44195, USA

Clinic of Urology and Pediatric Urology, Saarland University, Homburg, Germany.

Columbia University, Department of Surgery, New York, NY 10032, USA

Columbia University, Department of Pathology and Cell Biology, New York, NY 10032,
USA

Columbia University Medical Center, Molecular Pathology Shared Resource of Herbert
Irving Comprehensive Cancer Center of Columbia University, New York, NY 10032, USA

Comprehensive Cancer Center Tissue Procurement Shared Resource, Cooperative Human Tissue Network Midwestern Division, Dept. of Pathology, Human Tissue Resource Network, The Ohio State University, 410 West 10th Ave, Doan Hall, Room E413A, Columbus, OH 43210, USA

Cureline, Inc., 290 Utah Ave, Ste 300, South San Francisco, CA 94080, USA

Dana-Farber Cancer Institute, 450 Brookline Ave, Boston, MA 02215, USA

Dardinger Neuro-Oncology Center, Department of Neurosurgery, James Comprehensive Cancer Center and The

Dignity Health Mercy Gilbert Medical Center, 3555 S Val Vista Dr, Gilbert, AZ 85297, USA

Duke University Medical Center, 177 MSRB, Box 3156, Durham, NC 27710, USA

Duke University Medical Center, Gynecologic Oncology, Box 3079, Durham, NC 27710, USA

Duke University School of Medicine, Department of Pathology, Durham, NC 27710, USA

Emory University, Departments of Neurosurgery and Hematology and Medical Oncology, School of Medicine and Winship Cancer Institute, 1365C Clifton Road. N.E., Atlanta, GA 30322, USA

Erasmus Medical Center, Wytemaweg 80, 3015 CN, Rotterdam, The Netherlands

Erasmus University Medical Center Rotterdam, Cancer Institute, Wytemaweg 80, 3015CN, Rotterdam, the Netherlands

The Foundation of the Carlo Besta Neurological Institute, IRCCS via Celoria 11, 20133, Milan, Italy

Fred Hutchinson Cancer Research Center, 1100 Fairview Ave N, Seattle, WA 98109, USA

Fred Hutchinson Cancer Research Center, Program in Epidemiology, Seattle, WA 98109, USA

Friedrich-Alexander-University Erlangen-Nuremberg, Division Molecular Urology, Department of Urology and Pediatric Urology, University Hospital Erlangen, 91054 Erlangen, Germany

Greater Poland Cancer Center, Garbary 15, 61-866 Poznań, Poland

Greenville Health System Institute for Translational Oncology Research, 900 West Faris Road, Greenville, SC 29605, USA

Harvard University, Cambridge, MA 02138, USA

Havener Eye Institute, The Ohio State University Wexner Medical Center, 915 Olentangy River Rd, Columbus, OH 43212, USA

Henry Ford Hospital, 2799 West Grand Blvd, Detroit, MI 48202, USA

Hermelin Brain Tumor Center, Henry Ford Health System, 2799 W Grand Blvd, Detroit, MI 48202, USA

Hospices Civils de Lyon, CARDIOBIOTEC, Lyon F-69677, France

Hospital Clinic, Villarroel 180, Barcelona, Spain, 08036

Hue Central Hospital, Hue, Vietnam

Human Tissue Resource Network, Dept. of Pathology, College of Medicine, 1615 Polaris Innovation Ctr, 2001 Polaris, Columbus, OH 43240, USA

Huntsman Cancer Institute, Univ. of Utah, 2000 Circle of Hope, Salt Lake City, UT 84112, USA

Icahn School of Medicine at Mount Sinai, Department of Genetics & Genomic Sciences, 1 Gustave L. Levy Place, New York, NY 10029, USA

Icahn School of Medicine at Mount Sinai, Department of Obstetrics/Gynecology and Reproductive Sciences, 1 Gustave L. Levy Place, New York, NY 10029, USA

Indivumed GmbH, 20251 Hamburg, Germany

René Descartes University, Hospital Européen Georges Pompidou, 20 rue Leblanc, 75015, Paris, France

Curie Institute, 26 rue Ulm, 75005 Paris, France

Gustave Roussy Institute of Oncology, 39 Rue Camille Desmoulins 94805, Villejuif, France

Imperial College London, Department of Surgery and Cancer, Du Cane Road London W12 0NN, UK

Institute of Human Virology Nigeria, Abuja, Nigeria

Institute of Pathology, Technical University of Munich, Trogerstr. 18, 83675 Munich, Germany

Institute of Pathology, University Hospital Erlangen, Friedrich-Alexander-University Erlangen-Nuremberg, 91054 Erlangen, Germany

Institute of Urgent Medicine, Republic of Moldova

Regina Elena National Cancer Institute Irccs - Ifo, Via Elio Chianesi 53, 00144, Rome, Italy

John Wayne Cancer Institute, 2200 Santa Monica Blvd, Santa Monica, CA 90404, USA

Keimyung University, Daegu, South Korea

Klinikum rechts der Isar, Technical University of Munich, Dept. of Surgery, Ismaninger Str. 22, 81675 Munich, Germany

Knight Comprehensive Cancer Institute, Oregon Health & Science University, Portland, OR 97239, USA

Ludwig Maximilians University Munich, Ziemssenstrasse 1, D-80336, Munich, Germany

Maine Medical Center, 22 Bramhall St., Portland, ME 04102, USA

Martini-Clinic, Prostate Cancer Center, University Medical Center Hamburg-Eppendorf, Martinistr. 52, D-20246 Hamburg, Germany

Massachusetts General Hospital, 55 Fruit Street, Boston, MA 02114, USA

Mayo Clinic, 5777 E Mayo Blvd, Phoenix, AZ 85054, USA

Mayo Clinic, 4500 San Pablo Road, Jacksonville, FL 32224, USA

Mayo Clinic, 200 First St. SW, Rochester, MN 55905, USA

Mayo Clinic Arizona, Department of Urology, 5779 E. Mayo Blvd, Phoenix AZ 85054, USA

Mayo Clinic Arizona, Department of Hematology and Medical Oncology, 5779 E. Mayo Blvd, Phoenix AZ 85054, USA

McGill University Health Center. 1001 Decarie Blvd, Montreal, QC, Canada H4A 3J1

MD Anderson Cancer Center, Department of Pathology, 1515 Holcombe Blvd. Unit 0085, Houston, TX 77030, USA

MD Anderson Cancer Center, Life Science Plaza Building, 2130 W. Holcombe Blvd, Unit 2951, Office: LSP9.4029, Houston, TX 77030, USA

M.D. Anderson Cancer Center, Orbital Oncology & Ophthalmic Plastic Surgery, Department of Plastic Surgery, 1515 Holcombe Blvd, Unit 1488 Houston, TX 77030, USA

MD Anderson Cancer Center, Departments of Pathology & Translational Molecular Pathology, 1515 Holcombe Blvd--Unit 85, Houston, TX 77030, USA

Melanoma Institute Australia, North Sydney, NSW, Australia 2060

Memorial Sloan Kettering Cancer Center, Department of Pathology, 1275 York Avenue, New York, NY 10065, USA

Memorial Sloan Kettering Cancer Center, Center for Molecular Oncology, 1275 York Avenue, New York, NY 10065, USA

Ministry of Health of Vietnam, Hanoi, Vietnam

Molecular Response, 11011 Torreyana Road, San Diego, CA 92121, USA

Nancy N. and J.C. Lewis Cancer & Research Pavilion at St. Joseph's/Candler, 225 Candler Drive, Savannah, GA 31405, USA

National Cancer Hospital of Vietnam

National Cancer Institute, 31 Center Dr, Bethesda, MD 20892, USA

National Cancer Institute, Division of Cancer Epidemiology and Genetics, 9609 Medical Center Dr., Bethesda MD 20892, USA

National Cancer Institute, Urologic Oncology Branch, Center for Cancer Research, Building 10, Room 1-5940, Bethesda, MD 20892-1107, USA

National Cancer Center Korea, Center for Liver Cancer, 323 Ilsan-ro, Ilsan dong-gu, Goyang, Gyeonggi 10408, South Korea

Norfolk & Norwich University Hospital, Norwich, UK. NR4 7UY

NYU Langone Medical Center, Cardiothoracic Surgery, 530 First Avenue, 9V, New York, NY 10016, USA

The Ohio State University Comprehensive Cancer Center, 320 W 10th Avenue, Columbus, OH 43210, USA

The Ohio State University Wexner Medical Center, 2012 Kenny Rd, Columbus, OH 43221, USA

The Ohio State University Medical Center, Department of Neurological Surgery, 320 W 10th Ave, Columbus, OH, 43210, USA

The Ohio State University Wexner Medical Center, Department of Pathology, Doan Hall N337B and N308, 410 West 10th Ave., Columbus, OH 43210-1267, USA

Oncology Institute, Republic of Moldova

Ontario Tumor Bank - Hamilton site, St. Joseph's Healthcare Hamilton, Hamilton, Ontario L8N 3Z5, Canada

Ontario Tumor Bank - Kingston site, Kingston General Hospital, Kingston, Ontario K7L 5H6, Canada

Ontario Tumor Bank – Ottawa site, The Ottawa Hospital, Ottawa, Ontario K1H 8L6, Canada.

Ontario Tumor Bank, London Health Sciences Centre, London, Ontario N6A 5A5, Canada

Ontario Tumor Bank, Ontario Institute for Cancer Research, Toronto, Ontario M5G 0A3, Canada

Papworth Hospital NHS Foundation Trust, Cambridge CB23 3RE, UK

QIMR Berghofer Medical Research Institute, Herston, QLD, Australia

Radboud Medical University Center, Geert Grooteplein-Zuid 10, Nijmegen, the Netherlands

Regina Elena National Cancer Institute, 00144 Rome, Italy

Reinier de Graaf Hospital, Reinier de Graafweg 5, 2625AD, Delft, the Netherlands

Research Institute of the McGill University Health Centre, McGill University, Montréal, Québec, Canada

Research Center Of Chus Sherbrooke, Québec aile 9, porte 6, 3001 12e Avenue Nord, Sherbrooke, QC J1H 5N4, Canada

Ribeirão Preto Medical School - FMRP, Department of Surgery and Anatomy, University of São Paulo, Brazil, 14049-900

Robert J. Tomsich Pathology & Laboratory Medicine Institute, Lerner Research Inst, Dept. of Pathology, Cleveland Clinic Foundation, Cleveland, OH 44195, USA

Rockefeller University, 1230 York Ave, New York, NY 10065, USA

Rose Ella Burkhardt Brain Tumor and Neuro-Oncology Center ND4-52A, Cleveland Clinic Foundation, 9500 Euclid Ave, Cleveland, OH 44195, USA

Rose Ella Burkhardt Brain Tumor and Neuro-Oncology Center, 9500 Euclid Avenue - CA51, Cleveland, OH 44195, USA

Rose Ella Burkhardt Brain Tumor and Neuro-Oncology Center, Department of Neurosurgery, Neurological and

Roswell Park Cancer Institute. Elm & Carlton Streets, Buffalo, NY 14263, USA

Rush University Medical Center, Department of Cardiovascular and Thoracic Surgery. Suite 774 Professional Office Building, 1735 W. Harrison St., Chicago, IL 60612, USA

Rush University Medical Center, Department of Pathology, Department of Cell and Molecular Medicine. 570 Jelke Southcenter, 1750 W. Harrison St., Chicago, IL 60612, USA

Sage Bionetworks, Seattle, WA 98109, USA

Saint-Petersburg City Clinical Oncology Hospital, 56 Veteranov prospect, Saint-Petersburg, 198255, Russia

Sapienza University of Rome, Piazzale Aldo Moro 5, 00185 Rome, Italy

Sir Peter MacCallum, Department of Oncology, University of Melbourne, Parkville, 3050, Victoria, Australia

St. Joseph's/Candler Hospital, Department of Pathology, 5353 Reynolds St., Savannah, GA 31405, USA

St. Petersburg Academic University RAS, 8/3 Khlopin Str., St. Petersburg, 194021, Russia

Spectrum Health, Department of Pathology, 35 Michigan NE, Grand Rapids, MI 49503, USA

Stanford University, Palo Alto, CA, USA, USA

Stephenson Cancer Center, University of Oklahoma, Oklahoma City, OK 73104, USA

Tayside Tissue Bank, University of Dundee, Scotland UK DD1 9SY

The International Genomics Consortium, 445 N. 5th Street, Phoenix, AZ 85004, USA

The Oregon Clinic, 1111 NE 99th Ave, Portland, OR 97220, USA

The Prince Charles Hospital, UQ Thoracic Research Centre, Australia 4032

The Research Institute at Nationwide Children's Hospital, 700 Children's Drive, Columbus, OH 43205, USA

Tufts Medical Center, 800 Washington St., Boston, MA 02111, USA

UCSF-Helen Diller Family Comprehensive Cancer Center, 550 16th St., Mission Hall WS 6532 Box 3211, San Francisco, CA 94143, USA

University Hospital of Giessen and Marburg, Badingerstrasse 3, 35044, Marburg, Germany

University Hospital in Würzburg, Germany, Oberdürrbacher Strasse 6, 97080, Würzburg, Germany

University Health Network, 200 Elizabeth Street, Toronto ON M5G 2C4 Canada

University Hospital Essen, University Duisburg-Essen, German Cancer Consortium, Hufelandstr. 55; 45239 Essen, Germany

University Medical Center Hamburg-Eppendorf, Martinistr. 52, D-20246 Hamburg, Germany

University of Abuja Teaching Hospital, Gwagalada, FCT, Nigeria

University of Alabama at Birmingham Medical Center, 401 Beacon Pkwy W, Birmingham, AL 35209, USA

University of Arizona, Tucson AZ 85721, USA

University of Calgary, Departments of Surgery and Oncology, 1331 - 29th St NW, Calgary, AB, T2N 4N2, Canada

University of California San Francisco, 2340 Sutter St Rm S 229, San Francisco, CA 94143, USA

University of California, Irvine, 333 City Boulevard West, Suite 1400, Orange, CA 92868, USA

University of Chicago Medicine, 5841 S. Maryland Ave., Room G-216, MC 5094|Chicago, IL 60637, USA

University of Cincinnati Cancer Institute, Brain Tumor Clinical Trials, 200 Albert Sabin Way, Suite 1012, Cincinnati, OH 45267, USA

University of Cincinnati Cancer Institute, 200 Albert Sabin Way, Suite 1012, Cincinnati, OH 45267-0502, USA

University of Cincinnati, UC Health University Hospital, Dept. of Pathology & Laboratory Medicine, 234 Goodman Street, Cincinnati, OH 45219-0533, USA

University of Cincinnati Cancer Institute, Holmes Bldg., 200 Albert Sabin Way, Ste 1002, Cincinnati, OH 45267-0502, USA

University of Colorado Cancer Center, Aurora, CO 80111, USA

University of Dundee, Scotland UK DD1 9SY

University of Florence, Viale Pieraccini 6, 50139 Firenze, Italy

University of Hawaii Cancer Center, 701 Ilalo Street, Honolulu, HI 96813, USA

University of Heidelberg, Dept. Neuropathology, INF 224, 69120 Heidelberg, Germany

University of Heidelberg, Dept. Neurosurgery, INF 400, 69120 Heidelberg, Germany

University of Heidelberg, Division of Neurosurgical Research, Dept. Neurosurgery, INF 400, 69120 Heidelberg, Germany

University Hospitals Cleveland Medical Center, Division of Neuropathology, Department of Pathology, Providence Health and Services, Cleveland, OH 44106, USA

University of Iowa Hospital & Clinics, 200 Hawkins Drive, Clinical Trials-Data Management, 11510 PFP, Iowa City, IA 52242, USA

University of Iowa Hospital & Clinics, 200 Hawkins Drive, Hematology/Oncology, C32 GH, Iowa City, IA 52242, USA

University of Iowa Hospital & Clinics, 200 Hawkins Drive, ICTS-Informatics, 272 MRF, Iowa City, IA 52242, USA

University of Iowa Hospital & Clinics, 200 Hawkins Drive, Medicine Administration, 380 MRC, Iowa City, IA 52242, USA

University of Iowa Hospital & Clinics, 200 Hawkins Drive, Molecular Pathology, B606 GH, Iowa City, IA 52242, USA

University of Iowa Hospital & Clinics, 200 Hawkins Drive, Pathology, SW247 GH, Iowa City, IA 52242, USA

University of Kansas Cancer Center, 3901 Rainbow Blvd, Kansas City, KS 66160, USA

University of Kansas Medical Center, Kansas City, KS 66160, USA

University of Kansas Medical Center, Department of Pathology and Laboratory Medicine, Kansas City, KS 66206, USA

University of Kansas Medical Center Department of Orthopedic Surgery, 3901 Rainbow Boulevard, Kansas City, KS 66160, USA

University of Lausanne, Department of Computational Biology, Lausanne, Switzerland

University of Maryland School of Medicine, Department of Epidemiology and Public Health, Baltimore MD 21201, USA

University of Michigan, 500 S State St, Ann Arbor, MI 48109, USA

University of Michigan, Department of Surgery, Ann Arbor, MI 48109, USA

University of Montreal, 2900 Edouard Mont petit Blvd, Montreal, QC H3T 1J4, Canada

University of New Mexico, Albuquerque, NM 87131, USA

University of Pennsylvania, Philadelphia, PA 19104, USA

University of Pittsburgh, Department of Cardiothoracic Surgery, 200 Lothrop St, Suite C-800, Pittsburgh, PA 15213, USA

University of Pittsburgh, Department of Pathology, Pittsburgh, PA 15213, USA

University of Puerto Rico Comprehensive Cancer Center Biobank, Celso Barbosa St. Medical Center Area, San Juan, PR 00936

University of Sheffield Western Bank, Sheffield S10 2TN, UK

University of Washington, Seattle, WA 98105, USA

University of Western Australia School of Medicine, National Center for Asbestos Related Research, Nedlands, WA, Australia 6009

University of Wisconsin School of Medicine and Public Health, Department of Medicine, 1685 Highland Avenue, Madison, WI 53705, USA

Valley Health System, 1 Valley Health Plaza, Paramus, NJ 07652, USA

Vanderbilt University Medical Center, 1211 Medical Center Dr, Nashville, TN 37232, USA

Walter Reed National Military Medical Center, Clinical Breast Care Project, Murtha Cancer Center, Uniformed Services University, Bethesda, MD 20889, USA

Walter Reed National Military Medical Center, Murtha Cancer Center, Uniformed Services University, Bethesda, MD 20889, USA

Washington University School of Medicine, 600 S. Taylor Ave, St. Louis, MO 63110, USA

Washington University in St. Louis, Department of Medicine, 660 S. Euclid Ave., CB 8066, St. Louis, MO 63110, USA

Weill Cornell Medical College, New York, NY 10065, USA

Yonsei University College of Medicine, Department of Medicine, Seoul, Republic of Korea

SIGNIFICANCE

Adenocarcinomas of the gastrointestinal tract share not only a poor prognosis but also conserved molecular features. Hypermuted tumors display diverse immune features depending on tissue origin and molecular subtype, with implications for targeted immunotherapeutics. Upper GI tumors with chromosomal instability display a fine genome fragmentation enriched for high amplitude, focal somatic copy number alterations associated with whole genome doubling, specific mutational signatures, and advanced stage. We identified a genome stable molecular subtype among colorectal cancers with an elevated frequency of recurrent mutations in SOX9 and PCBP1.

HIGHLIGHTS

- GI adenocarcinomas comprised five molecular subtypes: EBV, MSI, HM-SNV, CIN, and GS
- Hypermutated tumors had diverse immune features varying by tissue and subtype
- CIN tumors displayed more fragmented copy number alterations in the upper GI tract
- Genome-stable CRC subtype was enriched for recurrent mutations in *SOX9* and *PCBP1*




Figure 1. Genomic Features of Gastrointestinal Adenocarcinomas

(A) Significantly mutated genes in gastrointestinal adenocarcinomas (GIAC) indicated by green circles, significantly mutated genes identified in other adenocarcinomas (non-GIAC) indicated by red circles, and genes identified as significantly mutated in all adenocarcinomas indicated by white circles. (B) Genes identified as significantly amplified (left) or deleted (right) in GIAC compared to in non-GIAC. (C) DNA hypermethylation frequency (top), mutation density (middle), and arm-level and focal copy-number events (bottom) in GIAC and non-GI AC. (D) Percent GOF or LOF events in developmental transcription factors by cancer type. See also Figure S1 and Tables S1–S6.




Figure 2. Molecular Subtypes of Gastrointestinal Adenocarcinomas

(A) Flowchart of molecular subtypes: Epstein-Barr virus (EBV)-positive (red); hypermutated-single-nucleotide variant predominant (HM-SNV) (gold); microsatellite instability (MSI) (blue); chromosomal instability (CIN) (purple); and genomically stable (GS) (green). (B) 3D plot of GIAC by SNV density, indel density, and clonal deletion score (CDS). Tumors annotated as Upper GI (crosses) and lower GI (circles) and color-coded by subtypes. (C) IFN γ pathway score (top) and CD8⁺ T-cell score (adjusted for total leukocytes; bottom) by subtypes stratified by upper vs lower GI. Horizontal bars indicate median values, boxes represent interquartile range, and whiskers indicate values within 1.5 times interquartile range. (D) Unsupervised analysis of DNA methylation across GIAC. (E, F) Distribution of subtypes (E) and CIMP subgroups (F) across anatomic regions. (G, H) Distribution of *MLH1/CDKN2A* silencing (G) and subtypes (H) in CIMP-H tumors by anatomic region. See also Figure S2.




Figure 3. Analysis of MSI and CIMP Tumors

(A) Methylation subtypes with four CIMPs: EBV-CIMP (red), CIMP-high (blue), GEA-CIMP low (yellow), CRC-CIMP low (green) with alterations of indicated genes. (B) Methylation profiles of union of CIMP-high and MSI tumors with *MLH1* silencing, *KRAS*, *BRAF*, *MLH1* and *MSH2* mutations. (C) Features of MSI tumors stratified by upper vs. lower GI and by CIMP-high status. Horizontal bars indicate median values, boxes represent interquartile range, and whiskers indicate values within 1.5 times interquartile range. (D) Unique and overlapping epigenetically silenced genes (>25%) in upper GI (top left), upper GI tumors excluding EBV⁺ (top right), lower GI (bottom, left), and MSI (bottom, right). (E) Frequency of silencing (black) and mutation (blue) of select genes in upper GI MSI (vertical axis) vs. lower GI MSI tumors (horizontal axis). See also Figure S3 and Table S7.




Figure 4. Molecular Features of the CIN Subtype in Upper GI

(A) Copy-number heatmap of non-hypermuted GIAC with amplification (red) and deletion (blue) with upper GI CIN tumors (top), CIN CRC (middle) and GS (bottom). (B) Plots of arm-level and focal copy-number events in CIN tumors by upper and lower GI tract. Horizontal bars indicate median values, boxes represent interquartile range, and whiskers indicate values within 1.5 times interquartile range. (C) Distribution of CIN-F (CIN-Focal) score by upper and lower GI CIN tumors. CIN-B denotes CIN-Broad. (D) Distribution of CIN-F score by clinical stage in Upper GI. Horizontal bars indicate median values, boxes represent interquartile range, and whiskers indicate values within 1.5 times interquartile range. (E) Whole genome doubling (WGD) in CIN-F and CIN-B tumors in the upper GI tract; WGD1 indicates one WGD, and WGD2 indicates >WGD (F) Frequency of distinct classes of somatic alterations in RAS and receptor tyrosine kinases (RTK; *KRAS*, *PIK3CA*, *BRAF*, *ERBB3*, *ERBB2*, *NRAS*, *EGFR*, *FGFR1*, *FGFR2*), cell cycle (CC; *FBXW7*, *CCNE1*, *CDK6*, *CDKN2A*, *CDKN1B*, *CCND1*, *CCND2*) and tumor suppressor genes (TSG) including WNT(*APC*, *RNF43*, *SOX9*, *TCF7L2*, *CTNNB1*), TGF β : *TGFBR2*, *ACVR2A*, *ACVR1B*, *SMAD4*, *SMAD2*, *SMAD*) and *TP53* in upper GI CIN-F and CIN-B tumors (G) Schematic model of CIN-F and CIN-B pathogenesis in upper GI. See also Figure S4.




Figure 5. Molecular Features of CIN and GS Colorectal Cancer

(A) Frequency of somatic alterations in indicated genes or pathways in non-CIMP CIN, CIMP-H/L CIN, and GS lower GI tumors. (B) SCNAs (top), mutation density (middle), and CIMP classes (bottom) across subtypes in lower GI tract. Horizontal bars indicate median values, boxes represent interquartile range, and whiskers indicate values within 1.5 times interquartile range. (C) Distribution of somatic mutations in *SOX9* and *PCBP1* in lower GI GS. (D) Schematic model of pathogenesis of molecular subtypes in lower GI. (E) Frequency of mutations in indicated genes in lower GI CIN/GS stratified anatomically. See also Figure S5.




Figure 6. Gastrointestinal Adenocarcinoma Mutational Signatures

(A) Mutation signatures in non-hypermethylated GIAC displayed by substitution class and sequence immediately 3' and 5' to the mutated base. (B) Key molecular features of GIAC by anatomical distribution. (C) Intensities of mutational signatures in CIN and GS subtypes by upper and lower GI. (D) BRCA signature in CIN and GS tumors in the upper and lower GI tract. (E) AA>AC signature stratified by CIN-F and CIN-B (top) and *TP53* mutation (bottom) in upper GI CIN tumors. (F) CpG>TpG signature in CIN and GS tumors in upper and lower GI stratified by CIMP status. For all boxplots, horizontal bars indicate median values, boxes represent interquartile range, and whiskers indicate values within 1.5 times interquartile range. See also Figure S6.




Figure 7. Integrated Molecular Comparison of Somatic Alterations Across GIAC Molecular Subtypes

(A–C) Alterations in select genes and pathways including RTK/RAS/PI-3K (A), TP53, cell cycle (B), and WNT/TGFβ (C). Deep deletions representing loss of more than half of the gene copies for the given ploidy of the tumor, blue; amplifications, red; missense mutations in the COSMIC repository, green; nonsense or frameshift mutations, black. Percentage of somatic alteration is indicated by numbers to the left of each gene box and divided by upper (U) and lower GI (L).

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
RPPA antibodies	RPPA Core Facility, MD Anderson Cancer Center	https://www.mdanderson.org/research/research-resources/core-facilities/functional-proteomics-rppa-core.html
Biological Samples		
Tumor and normal tissue and blood samples	TCGA Network	https://portal.gdc.cancer.gov/legacy-archive/
Critical Commercial Assays		
DNA/RNA AllPrep kit	Qiagen	Cat# 80204
mirVana miRNA Isolation kit	Ambion	Cat# AM1560
QiaAmp blood midi kit	Qiagen	Cat# 51185
AmpFISTR Identifier kit	Applied Biosystems	Cat# A30737
RNA6000 nano Assay	Agilent	Cat# 5067-1511
SureSelect Human All Exon 50 Mb	Agilent	Cat# G3370f
Genome-Wide Human SNP Array 6.0	Affymetrix	Cat# 901150
Illumina Barcoded Paired-End Library Preparation kit	Illumina	http://www.hgsc.bcm.edu/sites/-default/files/documents/-Illumina_Barcoded_Paired-End_Capture_Library_Preparation.pdf
TruSeq PE Cluster Generation kit	Illumina	PE-401-3001
Phusion PCR Supermix HiFi (2X)	New England Biolabs	Cat# M0531L
HumanMethylation450	Infinium	Cat# WG-314-1002
HumanMethylation450	Infinium	Cat# WG-311-2201
miRNA TruSeq kit	Illumina	Cat# RS-122-2001
Deposited Data		
Raw genomic and clinical data	NCI Genomic Data Commons	https://portal.gdc.cancer.gov/legacy-archive/
MC3 mutation annotation file	NCI Genomic Data Commons	https://gdc.cancer.gov/about-data/publications/mc3-2017
Processed data files	NCI Genomic Data Commons	https://gdc.cancer.gov/about-data/publications/pancanatlas
Software and Algorithms		
Broad Institute QC on BAM files - ContEst	(Cibulskis et al., 2011)	PMID: 21803805
Broad Institute Mutation Calling - MuTect	(Cibulskis et al., 2013)	PMID: 23396013
Broad Institute small indel Calling - Indelocator		https://www.broadinstitute.org/cancer/cga/indelocator
Broad Institute Mutation/Indel Annotation - Oncotator	(Ramos et al., 2015)	PMID: 25703262
Mutation Significance Analysis - MutSigCV	(Lawrence et al., 2014)	PMID: 24390350
RNA, DNaseq classifier - BioBloomTools(v1.2.4.b)	(Chu et al., 2014)	PMID: 25143290
Broad Institute - PathSeq	(Kostic et al., 2011)	PMID: 21552235
RNA read assembly - MapSplice 0.7.4	(Wang et al., 2010)	PMID: 20802226
Gene expression quantification - RSEM	(Li and Dewey, 2011)	PMID: 21816040
Copy number estimation	NA	http://archive.broadinstitute.org/cancer/cga/copynumber_pipeline
Significant focal copy number change - GISTIC 2.0	(Mermel et al., 2011)	http://software.broadinstitute.org/software/cprg/?q=node/31
Purity, ploidy, genome doubling - ABSOLUTE	(Carter et al., 2012)	http://archive.broadinstitute.org/cancer/cga/absolute
Cluster analysis - ConsensusClusterPlus	(Wilkerson and Hayes, 2010)	http://bioconductor.org/packages/release/bioc/html/ConsensusClusterPlus.html
Mbatch (EB++)	NA	http://bioinformatics.mdanderson.org/main/TCGABatchEffects:Overview