

UC San Diego

UC San Diego Previously Published Works

Title

Nonoblique effects in the Zbb^- vertex from extended technicolor dynamics

Permalink

<https://escholarship.org/uc/item/8sw33186>

Journal

Physical Review Letters, 69(4)

ISSN

0031-9007

Authors

Chivukula, R Sekhar
Selipsky, Stephen B
Simmons, Elizabeth H

Publication Date

1992-07-27

DOI

10.1103/physrevlett.69.575

Peer reviewed

Non-Oblique Effects in the $Zb\bar{b}$ Vertex from ETC Dynamics

R. Sekhar Chivukula^{a*}, Stephen B. Selipsky^{a*} and Elizabeth H. Simmons^{b*}

^a Boston University
Department of Physics
590 Commonwealth Avenue
Boston, MA 02215

^b Lyman Laboratory of Physics
Harvard University
Cambridge, MA 02138

ABSTRACT

Extended technicolor theories generate potentially large corrections to the $Zb\bar{b}$ vertex which can be observed in current experiments at LEP.

There exist no compelling or even consistent theories to explain the origin of the diverse masses and mixings of the quarks and leptons. In this regard, the origin of the large top mass is particularly puzzling. In technicolor models [1], this large top mass is presumably the result of extended technicolor (ETC) [2] dynamics at relatively low energy scales¹. Since the magnitude of the KM matrix element $|V_{tb}|$ is very nearly one, $SU(2)_W$ gauge invariance insures that the ETC dynamics which generates the top mass also couples to the left-handed component of the bottom. In this note, we point out that this dynamics produces potentially large “non-oblique” [5] effects at the $Zb\bar{b}$ vertex. In particular, if $m_t \gtrsim 100$ GeV and no effect is visible with data currently being obtained at LEP, theories in which the ETC and weak interactions commute (*i.e.* in which the ETC gauge bosons are $SU(2)_W$ singlets) can be ruled out, with the same confidence as models with excessive flavor changing neutral currents.

If the top mass is generated by the exchange of an $SU(2)_W$ neutral ETC gauge boson, then this boson carries technicolor and couples with strength g_{ETC} to the current

$$\xi \bar{\psi}_L^i \gamma^\mu T_L^{iw} + \frac{1}{\xi} \bar{t}_R \gamma^\mu U_R^w, \quad (1)$$

where $\psi_L = (t, b)_L$ is the left-handed tb doublet, $T_L = (U, D)_L$ is a left-handed technifermion weak doublet, and U_R is a corresponding right-handed technifermion weak singlet. The indices i and w are for $SU(2)_W$ and technicolor, respectively. The constant ξ is an ETC gauge-group dependent Clebsch and is expected to be of order one. At energies lower than the mass (M_{ETC}) of the ETC gauge boson, the effects of its exchange may be approximated by local four-fermion operators. In particular, the top mass arises from an operator coupling the left- and right- handed pieces of the current (1)

$$-\frac{g_{ETC}^2}{M_{ETC}^2} (\bar{\psi}_L^i \gamma^\mu T_L^{iw}) (\bar{U}_R^w \gamma_\mu t_R) + \text{h.c.} . \quad (2)$$

This may be Fierzed into a product of technicolor singlet densities

$$2 \frac{g_{ETC}^2}{M_{ETC}^2} (\bar{\psi}_L^i t_R) (\bar{U}_R T_L^i) + \text{h.c.} . \quad (3)$$

In what follows we will assume (for simplicity) that there is only one doublet of technifermions, that the strong technicolor interactions respect an $SU(2)_L \times SU(2)_R$ chiral

¹ This is true so long as there are no additional light scalar particles coupling to ordinary and technifermions [3] [4].

symmetry, and therefore that the technicolor F constant (analogous to f_π in QCD) is $v \approx 250$ GeV. Using the rules of naive dimensional analysis [6] we find the top quark mass is

$$m_t = \frac{g_{ETC}^2}{M_{ETC}^2} \langle \bar{U}U \rangle \approx \frac{g_{ETC}^2}{M_{ETC}^2} (4\pi v^3) . \quad (4)$$

Equivalently, the scale of the ETC dynamics responsible for generating the top mass is

$$M_{ETC} \approx 1.4 \text{ TeV} \cdot g_{ETC} \left(\frac{100 \text{ GeV}}{m_t} \right)^{\frac{1}{2}} . \quad (5)$$

In the absence of fine tuning [3] and as long as $g_{ETC}^2 v^2 / M_{ETC}^2 < 1$ (or, equivalently, when $m_t / (4\pi v)$ is small), the ETC interactions may be treated as a small perturbation on the technicolor dynamics and our estimates are self consistent.

These dimensional estimates are typically modified in “walking technicolor” models [7] where there is an enhancement of operators of the form (3) due to a large anomalous dimension of the technifermion mass operator. The enhancement is important for the ETC interactions responsible for light fermion masses (for which M_{ETC} must be quite high), but will not be numerically significant in the case of the top quark because the ETC scale (5) associated with the top quark is so low. Hence, the results in “walking” theories are expected to be similar to those presented below.

Consider the four-fermion operator² arising from the left-handed part of the current (1)

$$-\xi^2 \frac{g_{ETC}^2}{M_{ETC}^2} (\bar{\psi}_L^i \gamma^\mu T_L^{iw}) (\bar{T}_L^{jw} \gamma_\mu \psi_L^j) . \quad (6)$$

This may be Fierzed into the form of a product of technicolor singlet currents and includes

$$-\frac{\xi^2}{2} \frac{g_{ETC}^2}{M_{ETC}^2} (\bar{\psi}_L \gamma^\mu \tau^a \psi_L) (\bar{T}_L \gamma_\mu \tau^a T_L) , \quad (7)$$

where g_{ETC} and M_{ETC} are as in eqn. (3) and the τ^a are weak isospin Pauli matrices. We will show that this operator can generate sizeable deviations in the predictions for the $Zb\bar{b}$ coupling. There are also operators involving products of weak singlet left-handed currents, but these operators will not affect the $Zb\bar{b}$ coupling.

² Ref. [8] lists possible four-fermion operators arising from ETC exchange, with emphasis on potentially dangerous ETC contributions to $\delta\rho$.

Our analysis of the effects of operator (7) proceeds along the lines of ref. [9]. Adopting an effective chiral Lagrangian description appropriate below the technicolor chiral symmetry breaking scale, we may replace the technifermion current by a sigma-model current [10]:

$$(\bar{T}_L \gamma_\mu \tau^a T_L) = \frac{v^2}{2} \text{Tr} (\Sigma^\dagger \tau^a i D_\mu \Sigma) , \quad (8)$$

where $\Sigma = \exp(2i\tilde{\pi}/v)$ transforms as $\Sigma \rightarrow L\Sigma R^\dagger$ under $SU(2)_L \times SU(2)_R$, and the covariant derivative is

$$\partial_\mu \Sigma + i \frac{e}{s_\theta \sqrt{2}} (W_\mu^+ \tau^+ + W_\mu^- \tau^-) \Sigma + i \frac{e}{s_\theta c_\theta} Z_\mu \left(\frac{\tau_3}{2} \Sigma - s_\theta^2 [Q, \Sigma] \right) + ie A_\mu [Q, \Sigma] . \quad (9)$$

In unitary gauge $\Sigma = 1$ and operator (7) becomes

$$\frac{\xi^2}{2} \frac{g_{ETC}^2 v^2}{M_{ETC}^2} \bar{\psi}_L \left(\frac{e}{s_\theta c_\theta} \not{Z} \frac{\tau_3}{2} + \frac{e}{s_\theta \sqrt{2}} (W^+ \tau^+ + W^- \tau^-) \right) \psi_L . \quad (10)$$

This yields a correction

$$\delta g_L = -\frac{\xi^2}{2} \frac{g_{ETC}^2 v^2}{M_{ETC}^2} \frac{e}{s_\theta c_\theta} (I_3) = \frac{\xi^2}{4} \frac{m_t}{4\pi v} \cdot \frac{e}{s_\theta c_\theta} \quad (11)$$

to the tree-level $Zb\bar{b}$ couplings $g_L = \frac{e}{s_\theta c_\theta} (I_3 - Qs_\theta^2) = \frac{e}{s_\theta c_\theta} (-\frac{1}{2} + \frac{1}{3}s_\theta^2)$ and $g_R = \frac{e}{s_\theta c_\theta} (\frac{1}{3}s_\theta^2)$.

The $Zb\bar{b}$ width consequently shifts by an amount

$$\frac{\delta \Gamma}{\Gamma_{b\bar{b}}} \approx \frac{2g_L \delta g_L}{g_L^2 + g_R^2} \approx -3.7\% \cdot \xi^2 \left(\frac{m_t}{100 \text{ GeV}} \right) . \quad (12)$$

Note that the shift is *linear* in m_t ; it is also a non-oblique correction applying only to the $Zb\bar{b}$ width. As the standard model prediction for $\Gamma_{b\bar{b}}$ is 378 MeV, we see that (12) leads to a reduction of

$$14 \text{ MeV} \cdot \xi^2 \left(\frac{m_t}{100 \text{ GeV}} \right) \quad (13)$$

in both $\Gamma_{b\bar{b}}$ and Γ_{had} (the hadronic width). By way of comparison, the leading m_t -dependent term of the $Zb\bar{b}$ vertex correction in the one Higgs standard model is [11]

$$\delta g_L = -2I_3 \left(\frac{m_t}{4\pi v} \right)^2 \cdot \frac{e}{s_\theta c_\theta} \quad (14)$$

giving an m_t -dependance in the $Zb\bar{b}$ width of

$$\frac{\delta \Gamma}{\Gamma_{b\bar{b}}} \approx -0.5\% \left(\frac{m_t}{100 \text{ GeV}} \right)^2 \quad (15)$$

and a correspondingly smaller effect in the hadronic width. For $m_t \gtrsim 100$ GeV, (15) is quite a good estimate of the full one-loop standard model result. For a 100 GeV top, (15) corrects $\Gamma_{b\bar{b}}$ and Γ_{had} by approximately 2 MeV.

Experiments at LEP currently measure $\Gamma_{b\bar{b}}$ to an accuracy of 5% [12] and Γ_{had} to 12 MeV [13], so a shift on the order of (12) cannot currently be excluded. The measurement of $\Gamma_{b\bar{b}}$ should eventually reach 2% [12], at which point it will be possible to distinguish between the results (12) and (15). The similarly large Wtb vertex contribution in eqn. (10) is much more difficult to observe without detailed studies of the top quark.

So far, we have assumed that the ETC and weak interactions commute. In theories with weak-charged gauge bosons, we can make no definite predictions. As before, the ETC boson responsible for generating the top mass can contribute to the $Zb\bar{b}$ vertex. For example, the operator (3) can arise from the exchange of a weak-doublet ETC gauge boson which couples T_L to t_L^c (the field which is charge-conjugate to t_R) and ψ_L to U_L^c . Such a gauge boson will give rise to the $SU(2)_{L+R}$ triplet operator $(\bar{U}_R \gamma^\mu U_R)(\bar{\psi}_L \gamma_\mu \psi_L)$. In addition, there may be technicolor-neutral weak-triplet ETC bosons contributing directly to an operator of the form (7). In both of these cases, we would generically expect effects on the $Zb\bar{b}$ coupling to be of the same order of magnitude as those already described, but the size and sign of the total shift (12) will be model-dependent.

It is interesting to note that a correction to the $Zb\bar{b}$ vertex linear in m_t can also occur in models [14] where fermion masses arise from mixing of ordinary fermions and technibaryons. In this case, the b_L and b_R are partly technibaryons and, as in QCD, the axial technibaryon coupling is renormalized. Then the left- and right-handed couplings receive a correction of the form

$$\delta g_L = (g_L - g_R) \left(\frac{g_A - 1}{2} \right) \sin^2 \alpha , \quad (16)$$

and

$$\delta g_R = (g_R - g_L) \left(\frac{g_A - 1}{2} \right) \sin^2 \beta , \quad (17)$$

where g_A is the axial current renormalization, while α and β are the mixing angles relating the left- and right-handed components of the mass eigenstate b field to the corresponding gauge eigenstate b and technibaryon fields. In this model, the mass of the top is

$$m_t \approx m_{TB} \sin \alpha \sin \gamma , \quad (18)$$

where m_{TB} is the mass of a technibaryon and γ is the mixing angle for the right-handed top. If $\sin \gamma$ and $\sin \alpha$ are of the same order of magnitude, $\sin \alpha \approx \sqrt{m_t/m_{TB}}$ and

$$\frac{\delta\Gamma}{\Gamma} \approx (g_A - 1) \frac{m_t}{m_{TB}} . \quad (19)$$

In a QCD-like theory with $g_A \approx 1.25$, and for $m_t = 100$ GeV and $m_{TB} = 1$ TeV, this results in an effect of order $+2.5\%$.

Acknowledgements

We thank Andrew Cohen, Howard Georgi, David Kaplan, Joe Kroll, Kenneth Lane, Jenny Thomas, and Bing Zhou for useful conversations and comments. R.S.C. acknowledges the support of an NSF Presidential Young Investigator Award and of an Alfred P. Sloan Foundation Fellowship. S.B.S. and E.H.S. acknowledge the support of Texas National Research Laboratory Commission SSC Fellowship Awards. This work was supported in part under NSF contracts PHY-9057173 and PHY-8714654 and under DOE contract DE-AC02-89ER40509 and by funds from the Texas National Research Laboratory Commission under grants RGFY91B6 and RGFY9106.

References

- [1] S. Weinberg, Phys. Rev. D13 (1976) 974 and Phys. Rev. D19 (1979) 1277;
L. Susskind, Phys. Rev. D20 (1979) 2619.
- [2] S. Dimopoulos and L. Susskind, Nucl. Phys. B155 (1979) 237;
E. Eichten and K. Lane, Phys. Lett. B90 (1980) 125.
- [3] T. Appelquist, T. Takeuchi, M. Einhorn and L.C.R. Wijewardhana, Phys. Lett. B220 (1989) 223;
T. Takeuchi, Phys. Rev. D40 (1989) 2697;
V.A. Miransky and K. Yamawaki, Mod. Phys. Lett. A4 (1989) 129;
R. S. Chivukula, A. G. Cohen, and K. Lane, Nucl. Phys. B343 (1990) 554.
- [4] E. H. Simmons, Nucl. Phys. B312 (1989) 253.
- [5] B.W. Lynn, M.E. Peskin and R.G. Stuart, SLAC-PUB-3725 (1985); in *Physics at LEP*, Yellow Book CERN 86-02, Vol. I p.90.
- [6] A. Manohar and H. Georgi, Nucl. Phys. B234 (1984) 189.
- [7] B. Holdom, Phys. Lett. B105 (1985) 301;
T. Appelquist, D. Karabali and L.C.R. Wijewardhana, Phys. Rev. D35 (1987) 2605;
M. Bando, T. Morozumi, H. So and K. Yamawaki, Phys. Rev. Lett. 59 (1987) 389;
V.A. Miransky, Nuovo Cimento 90A (1985) 149.
- [8] T. Appelquist, M.J. Bowick, E. Cohler and A.I. Hauser, Phys. Rev. D31 (1985) 1676.
- [9] L. Randall and R.S. Chivukula, Nucl. Phys. B326 (1989) 1.
- [10] H. Georgi, *Weak Interactions and Modern Particle Theory*, (Benjamin-Cummings, Menlo Park, 1984), p.77.
- [11] A.A. Akhundov, D.Yu. Bardin and T. Riemann, Nucl. Phys. B276 (1986) 1 ;
W. Beenakker and W. Hollik, Z. Phys. C40 (1988) 141;
J. Bernabeu, A. Pich and A. Santamaria, Phys. Lett. B200 (1988) 569;
B.W. Lynn and R.G. Stuart, Phys. Lett. B252 (1990) 676.
- [12] J. Kroll, XXVIIth Rencontres de Moriond “Electroweak Interactions and Unified Theories”, March 1992, to be published
- [13] The LEP Collaborations, Phys. Lett. B276 (1992) 247.
- [14] D.B. Kaplan, Nucl. Phys. B365 (1991) 259.