

Lawrence Berkeley National Laboratory

Lawrence Berkeley National Laboratory

Title

First search for extraterrestrial neutrino-induced cascades with IceCube

Permalink

<https://escholarship.org/uc/item/8sz0d7fv>

Author

Kiryluk, Joanna

Publication Date

2009-08-05

Peer reviewed

**First search for extraterrestrial neutrino-induced cascades
with
IceCube**

**Joanna Kiryluk
The IceCube Collaboration**

This work was supported by the Director, Office of Science, Office of Basic Energy Sciences, of the U.S. Department of Energy under Contract No. DE-AC02-05CH11231.

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor The Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or The Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or The Regents of the University of California.

First search for extraterrestrial neutrino-induced cascades with IceCube

Joanna Kiryluk* for the IceCube Collaboration†

* Lawrence Berkeley National Laboratory and University of California Berkeley, Berkeley, CA 94720, USA

† see special section of these proceedings

Abstract. We report on the first search for extraterrestrial neutrino-induced cascades in IceCube. The analyzed data were collected in the year 2007 when 22 detector strings were installed and operated. We will discuss the analysis methods used to reconstruct cascades and to suppress backgrounds. Simulated neutrino signal events with a E^{-2} energy spectrum, which pass the background rejection criteria, are reconstructed with a resolution $\Delta(\log E) \sim 0.27$ in the energy range from ~ 20 TeV to a few PeV. We present the range of the diffuse flux of extra-terrestrial neutrinos in the cascade channel in IceCube within which we expect to be able to put a limit.

Keywords: extraterrestrial, neutrino, IceCube

I. INTRODUCTION

IceCube is a 1 km^3 Cherenkov detector under construction at the South Pole. Its primary goals are to detect high energy extra-terrestrial neutrinos of all flavors in a wide energy range, from ~ 100 GeV to ~ 100 EeV, search for their sources, for example active galactic nuclei and gamma ray bursts, and to measure their diffuse flux. When complete, the IceCube detector will be composed of 4800 Digital Optical Modules (DOMs) on 80 strings spaced 125 m apart. In addition there will be 6, more densely populated, Deep Core strings inside the IceCube detector volume. The array covers an area of one km^2 at depths from 1.45 to 2.45 km below the surface [1].

High energy neutrinos are detected by observing the Cherenkov radiation from secondary particles produced in neutrino interactions inside or near the detector. Muon neutrinos in charged current (CC) interactions are identified by the final state muon track [2]. Electron and tau neutrinos in CC interactions, as well as all flavor neutrinos initiating neutral current (NC) interactions are identified by observing electromagnetic or hadronic showers (cascades). A 10 TeV cascade triggers IceCube optical modules out to a radius of about 130 m. Cascades can be reconstructed with good energy resolution, but limited pointing resolution. The good energy resolution and low background from atmospheric neutrinos make cascades attractive for diffuse extraterrestrial neutrino searches [3].

We present expected sensitivities for the diffuse flux of extra-terrestrial neutrinos in the cascade channel in IceCube. This work uses data collected in 2007 with

the 22 strings that were deployed in IceCube at that time. The total livetime amounts to 270 days. Ten per cent of the data were used as a "burn" sample to develop background rejection criteria. The results, after unblinding, will be based on the remaining 90% of the data, about 240 days.

II. DATA AND ANALYSIS

Backgrounds from atmospheric muons, produced in interactions of cosmic rays with nuclei in the Earth's atmosphere form a considerable complication in all neutrino searches in IceCube. A filtering chain developed using Monte Carlo simulations of muon background and neutrino signal was used to reject these backgrounds online and offline.

The atmospheric muon background was simulated with CORSIKA [4]. In addition to the single muon events, which form the dominant background, an appropriate number of overlaying events was passed through the IceCube trigger and detector simulator to obtain a sample of coincident muons. The coincident muon events make a few per cent contribution to the total trigger rate. The signal, electron neutrino events, was simulated using an adapted version of the Monte Carlo generator ANIS [5] for energies from 40 GeV to 1 EeV and with a E^{-2} energy spectrum.

All estimates for the number of signal events later in the text assume an E^{-2} spectrum and flux strength of:

$$\Phi_{model} = 1.0 \times 10^{-6} (E/\text{GeV})^{-2} / (\text{GeV s sr cm}^2). \quad (1)$$

A. Online filtering

The main physics trigger is a "simple multiplicity trigger" (SMT), requiring photon signals in at least 8 DOMs, with the additional requirement of accompanying hits in any of the ± 2 neighboring DOMs, each above a threshold of 1/6 single photoelectron signal and within a $5 \mu\text{s}$ coincidence window. Averaging over seasonal changes of the trigger rate for IC22 was 550 Hz. The mean SMT rate is generally well reproduced by Monte Carlo simulation, which gives 565 Hz. Assuming the flux given in Eq. 1, approximately 2.7×10^3 electron neutrino events and $\sim 1 \times 10^{10}$ background event are expected to trigger the detector in 240 days.

The backgrounds are suppressed online with first-guess reconstruction algorithms [6]. A first guess track fit assumes that all hits can be projected onto a line, and that a particle producing those hits travels with velocity v_{line} . In addition a simple cut on sphericity




Fig. 1. The reconstructed center-of-gravity (COG) x after online filtering. Data is shown as continuous lines, background Monte Carlo is shown as dashed lines. Monte Carlo data is normalized to the experimental number of events.

of the events ($\text{EvalRatio}_{\text{Tot}}$) is used to select events with hit topology consistent with cascades. Cut values used in online filter are given in Table I. In the case of cascades, the online filter reduced the SMT trigger rate to ~ 20 Hz, or 3.5 % of the total trigger rate. Monte Carlo studies show that the filter retains about 70% of the simulated signal and rejects 97.5% of the simulated background that trigger the detector. The Monte Carlo simulation thus underestimates the overall rate observed in the data. Otherwise main characteristics are well reproduced, Fig. 1 which shows the reconstructed center-of-gravity (COG) x position. The COG is calculated for each event as the signal amplitude weighted mean of all hit DOM positions.

B. Offline filtering

The data, after online filtering and transfer from the South Pole, were passed through more sophisticated algorithms to reconstruct both muon tracks and cascades. This reconstruction uses the maximum-likelihood reconstruction algorithms described in [2], [6].

Several cuts were applied sequentially, and the intermediate data sets are identified as different levels. Level-1 is the trigger level and events passing the online filtering correspond to Level-2. The rates at different levels are summarized in Table I.

At Level-3 events were selected with (i) a reconstructed track zenith angle greater than 73° and (ii) a difference $\text{Llh}(\text{track}) - \text{Llh}(\text{cascade}) > -16.2$ in the likelihood parameters of the track and cascade reconstructions to select cascade-like events. This selection was optimized for the combined efficiency ($\sim 80\%$) in both atmospheric[7] and extra-terrestrial neutrino searches and keeps the data at this level common to both analyses. At Level-4 we require that all cascades originate inside the detector. In IceCube many muon tracks that radiate energetic bremsstrahlung or produce

hits in DOMs close to the detector edges can mimic uncontained cascades. To remove this background of partial bright muon events we require that the four earliest hits in the event are inside the fiducial volume of the detector. The boundaries of the fiducial volume in x - y are shown in Fig.2 as continuous lines. In the z direction only an upper boundary was used. It was set at the position of the 8th DOM from the top. Approximately 1% of the background events (data and Monte Carlo) and $\sim 13\%$ of the Monte Carlo signal events after online filtering pass Level-3 selections and satisfy the fiducial volume requirement.

At Level-5 we require that the number of hit DOMs (N_{Ch}) is greater than 20, that the reconstructed track zenith angle exceeds 69° , and that the event duration, defined as a time difference between the last and first hit DOM, is less than $5 \mu\text{s}$. The later cut removes long events, which are mostly coincident double or triple muon events typically with a high multiplicity of hit




Fig. 2. a) The y versus x positions of the strings in the IC22 detector configuration. b) The reconstructed center-of-gravity (COG) y versus x position from IC22 data after online filtering. The continuous lines show the boundaries of the fiducial volume, which is used in the analyses to restrict the position of the first hits in the event.




Fig. 3. The fill-ratio versus the distance D (defined in the text) for signal Monte Carlo (left) muon background Monte Carlo (middle) and the data (right) for events with $\text{COG-}z > -100\text{ m}$ (top) and $\text{COG-}z < -100\text{ m}$ (bottom). The dashed lines show the background cut at level 7 used in the analysis.

DOMs.

At Level-6 we require that the reconstructed cascade vertex positions $x(y)$ and $\text{COG-}x(y)$ agree to within 60 meters, and that the reduced track and cascade reconstruction likelihood ratio is less than 0.95. For each event we apply the two track reconstruction algorithm and require that the reconstructed tracks coincide to within $1\mu\text{s}$. This selection mostly removes background events with coincident muon tracks which are well separated in time.

At Level-7 stringent selections are made on the DOM multiplicity and the fill-ratio. The fill-ratio quantifies the fraction of hit DOMs within a sphere around the reconstructed cascade vertex position with a radius $2 \times D$, where D is the average displacement of the reconstructed cascade vertex with respect to the positions of the hit DOMs in the event. The fill-ratio versus the distance D for signal Monte Carlo, muon background Monte Carlo, and the data for events with $\text{COG-}z > -100\text{ m}$ and $\text{COG-}z < -100\text{ m}$ is shown in Fig.3. The presently used version of background Monte Carlo is in good agreement with the data for the top part of the detector, but not for the bottom part of the detector. In the bottom part of the detector, the clear ice (less absorption than at the top of the detector) makes some muons look like cascade (spherical shape and high DOM multiplicity). After applying the cuts on the fill-ratio and the distance D , as shown by the dashed lines in Fig.3, 135 events from the data burn sample and 11 background Monte Carlo events remained. Almost all of them originate in the bottom part of the detector, as shown in Fig. 3. Remaining 11 Monte Carlo background

events correspond to an expected ~ 90 events for the 240 days of the IC22 run.

We placed a final Level-8 cut on the reconstructed energy, $\log E_{\text{reco}} > 4.2$, which rejects all remaining background events in the data burn sample and in the background Monte Carlo.

III. RESULTS

The expected number of signal events (N_{Signal}) from a diffuse flux with a strength of $10^{-6}(\text{E}/\text{GeV})^{-2}/(\text{GeV} \cdot \text{s} \cdot \text{sr} \cdot \text{cm}^2)$ is 52 ν_e events for 240 days of livetime. Signal simulations show that events that pass all background rejection criteria are in the energy range from $\sim 20\text{ TeV}$ to a few PeV (with a mean energy of $\sim 160\text{ TeV}$). The energy resolution is $\Delta(\log E) \sim 0.27$, the x and y position resolution is ~ 10 meters. The z position resolution is worse, 25 m, because of a small fraction of events that originated below the detector where no fiducial volume cut was applied.

A burn sample of $\sim 10\%$ of the total IC22 data set and the background Monte Carlo sample were used in developing background rejection criteria. The selections are such that all events in the burn sample and all background Monte Carlo events are rejected, whereas a significant fraction of the signal Monte Carlo events are retained.

The model rejection factor (MRF) defined as: $\text{MRF} = \langle \mu_{90} \rangle / N_{\text{Signal}}$, will be used to determine the flux limit:

$$\Phi_{\text{limit}} = \text{MRF} \times f(E), \quad (2)$$

where $f(E)$ is given by Eq.1.

TABLE I
EVENT RATES AT DIFFERENT SELECTION LEVELS FOR EXPERIMENTAL DATA (BURN SAMPLE), ATMOSPHERIC MUONS BACKGROUND MONTE CARLO AND ν_e SIGNAL MONTE CARLO ASSUMING THE FLUX $\Phi_{model} = 1.0 \times 10^{-6} (\text{E}/\text{GeV})^{-2} / (\text{GeV} \cdot \text{s} \cdot \text{sr} \cdot \text{cm}^2)$

Level	Selection Criteria	Exp Data	Tot Bg MC	Signal MC ν_e
1	Trigger	580 Hz	565 Hz	$2.7 \times 10^3 \times (240 \text{ days})^{-1}$
2	$v_{\text{line}} < 0.25$ and $\text{EvalRatio}_{\text{TOI}} > 0.109$	20 Hz	14 Hz	$1.8 \times 10^3 \times (240 \text{ days})^{-1}$
3	$\text{Zenith} > 73^\circ$ and $\text{Llh}(\text{track}) - \text{Llh}(\text{cascade}) > -16.2$	4 Hz	2.8 Hz	$1.3 \times 10^3 \times (240 \text{ days})^{-1}$
4	Fiducial Volume (Fig.2)	0.3 Hz	0.15 Hz	$240 \times (240 \text{ days})^{-1}$
5	$NCh > 20$ and $\text{Zenith}_{32\text{iter}} > 69^\circ$ and $\text{EvtLength} < 5\mu\text{s}$	0.02 Hz	0.01 Hz	$165 \times (240 \text{ days})^{-1}$
6	$ \text{RecoX} - \text{COGX} < 60\text{m}$ and $ \text{RecoY} - \text{COGY} < 60\text{m}$ and $\text{ReducedLlh}(\text{track}) / \text{ReducedLlh}(\text{cascade}) > 0.95$ and $\text{RecoTrack1}(\text{Time}) - \text{RecoTrack2}(\text{Time}) < -1\mu\text{s}$	0.011 Hz	0.004 Hz	$161 \times (240 \text{ days})^{-1}$
7	Fill-Ratio (Fig.3)	6.8×10^{-5} Hz	4.3×10^{-6} Hz	$68 \times (240 \text{ days})^{-1}$
8	$NCh > 60$ and $\log E_{\text{reco}} > 4.2$	0	0	$52 \times (240 \text{ days})^{-1}$

The analysis is limited by the currently available background Monte Carlo sample. It is not possible to subtract the simulated residual background contribution with sufficient precision. Thus the sensitivities for the diffuse flux of extraterrestrial neutrino signal, defined as the average upper limit at 90% CL and absence of signal [9], cannot be determined. To give an order of magnitude for the limit, a conservative estimate making no assumptions on background would be $4 \times 10^{-8} (5 \times 10^{-7}) (\text{E}/\text{GeV})^{-2} / (\text{GeV} \cdot \text{s} \cdot \text{sr} \cdot \text{cm}^2)$ for a hypothetical number of observed events after unblinding of 0 (20).

Enclosing, we expect the flux limit from this analysis to be of the same order as the limit on the diffuse flux $\Phi_{\text{limit}} = 1.3 \times 10^{-7} (\text{E}/\text{GeV})^{-2} / (\text{GeV} \cdot \text{s} \cdot \text{sr} \cdot \text{cm}^2)$ [10] in the cascade channel as obtained from 5 years of AMANDA data. Additional background Monte Carlo events are being generated and systematic uncertainties are currently being studied.

This work was supported in part by the U.S. National Science Foundation under Grant 0653266 and the Department of Energy under contract DE-AC-02-05CH11231.

REFERENCES

- [1] IceCube, A. Achterberg *et al.*, *Astropart. Phys.* **26**, 155 (2006); R. Abbasi *et al.* *Nucl. Instrum. Meth.* **A601** (2009) 294.
- [2] AMANDA, J. Ahrens *et al.*, *Nucl. Instrum. Meth.* **A524** (2004) 169.
- [3] M. Kowalski, *JCAP* 05 (2005) 010.
- [4] D. Heck *et al.*, *Tech. Rep. FZKA*, 6019 (1998).
- [5] A. Gazizov and M. Kowalski, *Computer Physics Communications* Vol. 172 **3** (2005) 203.
- [6] AMANDA, J. Ahrens *et al.*, *Phys. Rev.* **D67** (2003) 012003.
- [7] M. D'Agostino (for the IceCube Collaboration), *A search for atmospheric neutrino-induced cascades with IceCube*, these proceedings.
- [8] IceCube, J. Ahrens *et al.*, *IceCube Preliminary Design Document* (2001).
- [9] G. Feldman and R. Cousins, *Phys. Rev.* **D57** (1998) 3873.
- [10] O. Tarasova, M. Kowalski and M. Walter (for the IceCube Collaboration), proceedings of the 30th International Cosmic Ray Conference (ICRC 2007), Merida, Yucatan, Mexico, 3-11 Jul 2007; arXiv:0711.0353 [astro-ph] pages 83-86.