

Lawrence Berkeley National Laboratory

Recent Work

Title

NEUTRAL PARTICLE PRODUCTION IN n+p AND pp COLLISIONS AT 100 GeV/c

Permalink

<https://escholarship.org/uc/item/8t8980kw>

Authors

Alston-Garnjost, Margaret

Erwin, J

Klems, J.H.

et al.

Publication Date

1974-08-01

Submitted to Physical Review Letters

LBL-3311
Preprint

RECEIVED
LAWRENCE
BERKELEY LABORATORY

JUL 25 1975

LIBRARY AND
DOCUMENTS SECTION

NEUTRAL PARTICLE PRODUCTION IN π^+p AND
pp COLLISIONS AT 100 GeV/c


Margaret Alston-Garnjost, J. Erwin, J. H. Klems,
W. Ko, R. L. Lander, D. E. Pellett, and P. M. Yager

August 1974

Prepared for the U.S. Energy Research and
Development Administration under Contract W-7405-ENG-48

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. [REDACTED]*



LBL-3311

LBL-3311
C.P.

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

-111-

Neutral Particle Production in π^+p and pp
Collisions at 100 GeV/c

Margaret Alston-Garnjost

Lawrence Berkeley Laboratory
University of California
Berkeley, California 94720*

and

J. Erwin, J. H. Klems, W. Ko, R. L. Lander
D. E. Pellett, P. M. Yager

Department of Physics
University of California at Davis
Davis, California 95616†

April 1975

ABSTRACT

We have measured the inclusive cross sections for γ , K_s^0 , Λ and $\bar{\Lambda}$ production in π^+p and pp interactions at 100 GeV/c and compared various inclusive distributions of the produced γ and K_s^0 .

We have used a sample of 100,000 pictures of the FNAL 30-inch hydrogen bubble chamber to compare the production of neutral particles (γ , K_s^0 , Λ , $\bar{\Lambda}$) in π^+p and pp interactions at 100 GeV/c. The chamber was exposed to an unseparated tagged beam of positive particles.¹ This allows us to compare π^+ and proton induced reactions with minimum biases.

The film was scanned twice, yielding 2418 events which had at least one Vee visible within a restricted fiducial volume in the chamber. The overall scanning efficiency for the two scans was 96%. The events on about half the film were measured on an LBL spiral reader, the others on an LBL Franckenstein. The measurements were processed using the programs POOH and TVGP, and the position of the beam track in the chamber was matched with data from the beam tagging system to identify the beam track. The incident particle was unambiguously identified as a π^+ or proton in 78% of the events. The program SQUAW was then used to fit the Vees associated with unambiguously identified beam tracks; in each fit the neutral was constrained to come from the measured production vertex. Events measured on the spiral reader were re-examined on a scan table and those considered to be possibly measureable were remeasured on a Franckenstein. The overall efficiency of the two measurements for Vees produced by an ambiguously identified π^+ or proton was 89%; for one Franckenstein measurement it was 82%. The scanning and measuring efficiencies were assumed to be the same for all events.

The identity of each neutral track producing a Vee was assigned, where possible, by using the ionization information from the special scan. Ambiguous events were selected on the basis of the confidence level of the kinematic fit and the transverse momentum of the decay tracks with respect to the line of flight of the neutral. A fiducial volume smaller than that used in scanning was imposed. Also, γ conversion pairs were rejected if they were within 4 cm of the production vertex and strange particle decays within 2 cm. The number of events remaining

after all acceptance criteria were met are given in Table I. The contamination of the γ 's is negligible and for K_s^0 , Λ and $\bar{\Lambda}$ is estimated to be less than 2, 5 and 10% respectively.

The events were weighted for escape from the chamber, decay close to the production vertex, and neutral decay modes, the average weights were 58, 3.0, 2.7, and 4.2 for γ , K_s^0 , Λ and $\bar{\Lambda}$ respectively. To investigate our detection efficiency, we looked at the inclusive distributions in the rapidity variable y plotted in Fig. 1 (the Λ and $\bar{\Lambda}$ distribution are not shown because of space limitations). We note that the γ and K_s^0 distributions for pp interactions are approximately symmetric about $y = 0$, indicating that our detection efficiency for these processes is similar in both the forward and backward hemisphere. This forward-backward similarity is lacking in observed Λ distributions because many of the Λ 's are produced peripherally and those in the forward hemisphere are very fast and leave the chamber before they decay. For $\bar{\Lambda}$'s it appears that our detection efficiency is adequate in the forward hemisphere. To calculate the total inclusive cross-sections for γ , K_s^0 and $\bar{\Lambda}$ production and for $\pi^+p \rightarrow \Lambda$ we used all events. For the reaction $pp \rightarrow \Lambda$ we used only the backward going Λ 's and the symmetry of the reaction. The neutral inclusive cross-sections are presented in Table I, where we normalize to the measured total cross sections.² The errors include uncertainties in the scanning, measuring and beam tagging efficiencies. The neutral inclusive pp cross-sections agree well with those of Chapman et al.³ at 102 GeV/c. The π^+p cross-sections for K_s^0 , Λ and $\bar{\Lambda}$ production when compared with those of Sturtebeck et al.⁴ at 18.5 GeV/c all show an increase with energy, a trend established at lower energies.⁵ For K_s^0 production the rise is rapid from 1.1 ± 0.07 mb to 4.1 ± 0.4 mb. On the other hand, the Λ cross-section has risen slowly from 0.73 ± 0.05 mb to 1.1 ± 0.2 mb; however, it must be remembered that our inclusive cross-section is really a lower limit because of our poor detection efficiency in the forward hemisphere. Increasing cross-sections with incident momentum are observed in pp and

$\pi^- p$ interactions,^{6,7,8} however, the rise in K_s^0 production is much less pronounced than the rise we see in the $\pi^+ p$ reaction. Comparison of our $\pi^+ p$ and pp cross-sections is interesting because it shows that the production of γ 's and $\bar{\Lambda}$'s is roughly in the ratio of the total cross-sections, whereas K_s^0 's are produced with nearly equal probability in the two reactions, and Λ 's are produced relatively prolifically in pp collisions.

To study in detail the dependence of the neutral particle production upon its longitudinal momentum, we show in Fig. 1 distributions in rapidity (y). Note, since $(1/\sigma_T) (d\sigma/dy) = (1/N_T) (dN/dy)$, all distributions have been normalized by the total events in our data sample reducing biases and making comparison easier. The errors are statistical only. The γ distributions for $\pi^+ p$ and pp reactions show similar shapes in the backward hemisphere. In the forward hemisphere there is a suggestion that the shapes are different and the ratio is greater than 1.0. The K_s^0 distributions show that the ratio (π^+/p) is greater than one both backwards and forwards.

Next we look at the dependence on the neutral particle transverse momentum squared, p_T^2 . We see in Fig. 2 the expected exponential fall with increasing p_T^2 and the slope decreasing with increasing neutral particle mass. We note also that a single exponential term will not describe either γ distributions because the lowest bin contains too many events. The average p_T^2 is given in Table I. The shapes of the $\pi^+ p$ and pp induced reactions are similar.

In Fig. 3 we show the average number of π^0 and K_s^0 produced per inelastic interaction as a function of the number of charged prongs produced (n_{ch}). Here we assume that all γ 's come from π^0 's and that $\sigma_{\pi^0} = \sigma_\gamma/2$. We see that the average number of π^0 's is similar for $\pi^+ p$ and pp interactions and rises more or less linearly, as observed in pp collisions at several energies.^{8,9} The average number of K_s^0 shows only weak dependence on the number of charged prongs. For pp interactions the distribution is fairly flat for $n_{ch} > 6$, an effect already seen in pp collisions at 102 and 205 GeV/c⁴ and in $\pi^- p$ at 205 GeV/c.⁶ The $\pi^+ p$ induced

distribution shows some excess at low multiplicity.

We have compared the inclusive cross sections for neutral particle production in π^+p and pp reactions at 100 GeV/c. We find that when we normalize by the total cross sections the γ production is similar in the two reactions with a slight excess of events in the forward hemisphere for the π^+p interaction, and a depletion backwards. The K_s^0 cross section is rising much more rapidly with energy than it does in pp and π^-p induced reactions.

We thank the staffs of the 30-inch Bubble Chamber and Neutrino Laboratory at the Fermi National Accelerator Laboratory for their assistance. The aid of the Proportional Hybrid System Consortium is gratefully acknowledged. We also thank the scanning and measuring staffs the University of California, Davis, and the Lawrence Berkeley Laboratory.

Footnotes and References

* Work done under the auspices of the U.S. Atomic Energy Commission.

† Work supported in part by U.S. Atomic Energy Commission Contract AT(04-3)-34 PA 191.

1. J. Erwin, J.H. Klems, W. Ko, R.L. Lander, D.E. Pellett, P.M. Yager, and M. Alston-Garnjost, Phys. Rev. Lett. 32, 254 (1974). We made two separate runs of 58,000 and 46,000 pictures. In the first run we tagged the ($\pi^+ + \mu^+$) and K^+ separately giving a beam composition of 51% p, 44% π^+ , 4% μ^+ , and 1% K^+ . In the second run no distinction was made between π^+ and K^+ resulting in a beam composition 50% p, 46% π^+ and K^+ , and 4% μ .
2. A.S. Carroll et al., Phys. Rev. Lett. 33, 928 (1974) and Phys. Rev. Lett. 33, 932 (1974).
3. J.W. Chapman, J. Cooper, N. Green, B.P. Roe, A.A. Seidl, J.C. Vander Velde, C.M. Bromberg, D. Cohen, T. Ferbel, P. Slattery. Physics Lett. 47B, 465 (1973).
4. P.H. Strutebeck, N.M. Cason, J.M. Bishop, N.N. Biswas, V.P. Kenney, W.D. Shepard. Phys. Rev. D, 9, 608 (1974).
5. D.J. Crennel, H.A. Gordon, M.L. Ioffredo and K.W. Lai, Phys. Rev. Lett. 28, 643 (1972).
6. pp reaction: H. Boggild et al., Nucl. Phys. B 57, 77 (1973), G. Charlton et al., Phys. Rev. Lett. 29, 1759 (1972), G. Charlton et al., Phys. Rev. Lett 30, 574 (1973), and F.T. Dao et al., Phys. Rev. Lett. 30, 1151 (1973).
7. π^-p reaction: D. Bogart, R. Hanft, F.R. Huson, D. Ljung, C. Pascaud, S. Pruss, W.M. Smart, H.H. Bingham, D.M. Chew, B.Y. Daugeras, W.B. Fretter, G. Goldhaber, W.R. Graves, A.D. Johnson, J.A. Kadyk, L. Stutte, G.H. Trilling, F.C. Winkelmann, and G.P. Yost. Paper submitted to XVII International Conference on High Energy Physics, London 1974. (Unpublished)

Footnotes and References (cont.)

8. For review articles on pp and π^-p production of neutral particles at high energy see D. Ljung, AIP Conference Proceedings No. 14, American Institute of Physics, 464 (1973) and J. Whitmore, Physics Reports 10, 273 (1974).
9. W.T. Ko, Proceedings of the XVII International Conference on High Energy Physics, London, I-76 (1974).

Table 1 Production of Neutral Particles

	<u>Reaction</u>	<u>γ</u>	<u>K_s^0</u>	<u>Λ</u>	<u>$\bar{\Lambda}$</u>
Number of Vees	$\pi^+ p$	350	190	57(6f) [*]	4
	pp	492	223	135(13f) [*]	11
Inclusive Cross Section (mb)	$\pi^+ p$	136±11	4.1±0.4	1.1±0.2	0.15±0.08
	pp	174±13	3.8±0.3	3.5±0.4	0.24±0.07
Ratio $\sigma(\pi^+ p)/\sigma(pp)$ ⁺		0.78±0.09	1.08±0.14	≥0.32	~0.6
$\langle P_T \rangle$	$\pi^+ p$	0.054±0.005	0.30±0.03	0.23±0.06	0.42±0.10
	pp	0.059±0.005	0.23±0.02	0.32±0.04	0.25±0.06

* f = number detected in forward hemisphere

+ $\sigma_T(\pi^+ p)/\sigma_T(pp) = 0.61$


Figure Captions

FIG. 1 - Inclusive distributions in center of mass rapidity.

$$y_{c.m.} = 0.5 \ln (E + p_L)/(E - p_L).$$

FIG. 2 - Inclusive distributions in transverse momentum, p_T^2 .

FIG. 3 - Average number of neutrals produced.




XBL 754-989

Fig. 1


$\gamma + \text{ANYTHING}$

$K_S^0 + \text{ANYTHING}$



XBL 754-990

Fig. 2



XBL 751-113

Fig. 3

LEGAL NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Energy Research and Development Administration, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

TECHNICAL INFORMATION DIVISION
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720