

UC Davis

UC Davis Previously Published Works

Title

Saccharomyces cerevisiae Gene Expression during Fermentation of Pinot Noir Wines at an Industrially Relevant Scale

Permalink

<https://escholarship.org/uc/item/8tn4k3c1>

Journal

Applied and Environmental Microbiology, 87(11)

ISSN

0099-2240

Authors

Reiter, Taylor
Montpetit, Rachel
Byer, Shelby
et al.

Publication Date

2021-05-11

DOI

10.1128/aem.00036-21

Peer reviewed



Saccharomyces cerevisiae Gene Expression during Fermentation of Pinot Noir Wines at an Industrially Relevant Scale

 Taylor Reiter,^{a,b,c} Rachel Montpetit,^b Shelby Byer,^b Isadora Frias,^b Esmeralda Leon,^d Robert Viano,^d Michael McLoughlin,^d Thomas Halligan,^d Desmon Hernandez,^d Ron Runnebaum,^{b,d}  Ben Montpetit^{a,b}

^aFood Science Graduate Group, University of California, Davis, Davis, California, USA

^bDepartment of Viticulture and Enology, University of California, Davis, Davis, California, USA

^cDepartment of Population Health and Reproduction, University of California, Davis, Davis, California, USA

^dDepartment of Chemical Engineering, University of California, Davis, Davis, California, USA

ABSTRACT *Saccharomyces cerevisiae* metabolism produces ethanol and other compounds during the fermentation of grape must into wine. Thousands of genes change expression over the course of a wine fermentation, allowing *S. cerevisiae* to adapt to and dominate the fermentation environment. Investigations into these gene expression patterns previously revealed genes that underlie cellular adaptation to the grape must and wine environments, involving metabolic specialization and ethanol tolerance. However, the majority of studies detailing gene expression patterns have occurred in controlled environments that may not recapitulate the biological and chemical complexity of fermentations performed at production scale. Here, an analysis of the *S. cerevisiae* RC212 gene expression program is presented, drawing from 40 pilot-scale fermentations (150 liters) using Pinot noir grapes from 10 California vineyards across two vintages. A core gene expression program was observed across all fermentations irrespective of vintage, similar to that of laboratory fermentations, in addition to novel gene expression patterns likely related to the presence of non-*Saccharomyces* microorganisms and oxygen availability during fermentation. These gene expression patterns, both common and diverse, provide insight into *Saccharomyces cerevisiae* biology critical to fermentation outcomes under industry-relevant conditions.

IMPORTANCE This study characterized *Saccharomyces cerevisiae* RC212 gene expression during Pinot noir fermentation at pilot scale (150 liters) using industry-relevant conditions. The reported gene expression patterns of RC212 are generally similar to those observed under laboratory fermentation conditions but also contain gene expression signatures related to yeast-environment interactions found in a production setting (e.g., the presence of non-*Saccharomyces* microorganisms). Key genes and pathways highlighted by this work remain undercharacterized, indicating the need for further research to understand the roles of these genes and their impact on industrial wine fermentation outcomes.

KEYWORDS *Saccharomyces cerevisiae*, fermentation, gene expression

Saccharomyces cerevisiae is most often the dominant fermentative organism during vinification. As a domesticated species, it has evolved specialized metabolic strategies to assimilate sugars in grape must and to transform them into ethanol, thereby outcompeting other microorganisms during fermentation (1). During this process, *S. cerevisiae* encounters a dynamic stress landscape. In early fermentation, sources of stress include high sugar concentrations (osmotic stress), low pH (acid stress), decreasing oxygen (hypoxia), the presence of other organisms that compete for nutrients or produce inhibitory compounds, and sulfur dioxide additions that are used to inhibit spoilage organisms. As fermentation progresses, nutrients become limiting (starvation), the temperature may rise

Citation Reiter T, Montpetit R, Byer S, Frias I, Leon E, Viano R, McLoughlin M, Halligan T, Hernandez D, Runnebaum R, Montpetit B. 2021. *Saccharomyces cerevisiae* gene expression during fermentation of Pinot noir wines at an industrially relevant scale. Appl Environ Microbiol 87:e00036-21. <https://doi.org/10.1128/AEM.00036-21>.

Editor Edward G. Dudley, The Pennsylvania State University

Copyright © 2021 Reiter et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Ben Montpetit, benmontpetit@ucdavis.edu.

Received 8 January 2021

Accepted 15 March 2021

Accepted manuscript posted online 19 March 2021

Published 11 May 2021

or be kept low (heat/cold stress), and ethanol concentrations rise (ethanol stress). However, through a coordinated gene expression response, *S. cerevisiae* adapts to these stresses and most often continues fermentation until the must contains no residual sugar.

High-throughput gene expression profiling (e.g., microarray and high-throughput RNA sequencing [RNA-seq]) has offered a window into the metabolic strategies used by *S. cerevisiae* during fermentation to adapt to and to dominate fermentation environments. Previous research reported expression changes in >2,000 genes during fermentation (2–4). In early fermentation, this is marked by expression of gene products that support biosynthetic processes and acquisition of abundant nutrient resources (2, 3). As fermentation progresses, nitrogen limitation, phosphate limitation, and/or ethanol accumulation can trigger a transition to a nonproliferative state (i.e., stationary phase), which involves remodeling the gene expression program to support cellular adaptation to the changing environment with continued metabolism (2, 3). Toward the end of fermentation, relief of nitrogen catabolite repression (2) and increased expression of nitrogen recycling genes (2, 5) is observed, which can be accompanied by further remodeling of the translational machinery and increased oxidative metabolism (5, 6). As ethanol concentrations rise through the end of fermentation, a gradual transcriptome response to ethanol stress is also observed (3). This response overlaps, but appears distinct from, the environmental stress response (ESR) seen in laboratory yeast (2, 3, 7), which may be related to the multitude of simultaneous stresses experienced by the yeast at the end of wine fermentation. Indeed, the majority of genes with stress response elements in their promoters are expressed at the end of fermentation (8).

Through the associated metabolic processes that consume and produce a multitude of compounds, *S. cerevisiae* gene expression in response to environmental factors is related to overall fermentation kinetics and wine sensory outcomes. For example, fermentations can become sluggish or stuck when *S. cerevisiae* inadequately adapts to stresses encountered in the wine fermentation environment (9). In addition, altered gene expression likely underlies different wine sensory characteristics in fermentations conducted with different industrial yeast strains (10, 11). To impact wine quality, genetic strategies have been applied in attempts to alter the expression of flavor-associated genes (12), which have achieved variable levels of success. Consequently, further study of the *S. cerevisiae* gene expression program across fermentation is required to understand the yeast-environment relationship and how these interactions may be controlled to alter fermentation outcomes.

Given the importance of the yeast-environment interaction in determining gene expression, a major consideration with respect to collecting such data is the fermentation conditions used. To date, the majority of gene expression surveys have profiled fermentations that deviate in one or more ways from the industrial conditions in which most fermentations take place. For example, hundreds to thousands of liters of grape must are fermented to wine at industrial scales, while milliliter to liter volumes are commonly used in laboratory studies of gene expression (2–5, 13–16). Industry-scale fermentations also have different kinetics, compared with laboratory-scale fermentations (4, 15, 17), and are less aromatic due to differences in hydrodynamics (15, 18). Similarly, dissolved oxygen levels differ at laboratory scale, compared with industry scale (4), which can impact fermentation outcomes (19, 20). Possibly reflecting these different environments, at the end of fermentation the expression of key genes involved in amino acid transport and other core metabolic processes have been shown to differ between laboratory and industrial fermentations (4). Consequently, we propose that the physical and chemical differences in laboratory-scale versus industry-scale wine fermentations are important factors to consider when analyzing gene expression patterns across fermentation.

Another major consideration when conducting gene expression studies is that most studies investigate the fermentative capability of *S. cerevisiae* in monoculture using sterile synthetic media or filter-sterilized grape must (2–5, 12, 21). These controlled studies are important and allow connections between the media, gene expression, and wine outcomes to be made (12) but do not recapitulate the complexity of a natural grape must that varies in

parameters such as nitrogen composition, pH, and phenolic and elemental profiles (22–25). In addition, these experiments lack the diverse grape must microbiome that is a contributing component of wine fermentations (26–40). These are all parameters that shape the fermentation environment and the metabolic response of *S. cerevisiae*.

Interspecies interactions are a critical component of the fermentation environment that informs the biology and behavior of *S. cerevisiae* during fermentation. It has been shown that non-*Saccharomyces* yeast impact the metabolism of *S. cerevisiae* through direct and indirect interactions (41–43), leading to faster resource acquisition by *S. cerevisiae* in early fermentation and altered metabolism of vitamins and minerals (42–45). While research is still needed to describe the impact of a diverse microbial consortia on *S. cerevisiae* during fermentation (46, 47), it remains that industrial fermentations are not sterile and involve diverse microorganisms (30, 36, 37, 39, 40). Even in fermentations treated with sulfur dioxide (SO₂) to control microbial spoilage organisms, native fungi and bacteria are metabolically active during fermentation (40, 48, 49). This makes profiling *S. cerevisiae* gene expression among diverse microbial consortia important, as it will lead to a better understanding of the principles that govern *S. cerevisiae* gene expression and metabolism during fermentation.

Here, to begin to address the impact of an industrial wine fermentation environment on *S. cerevisiae* gene expression, the inherent variability found in industrial fermentations was incorporated to determine the *S. cerevisiae* RC212 gene expression program across chemically and biologically diverse Pinot noir grape musts. Specifically, time-series RNA-seq was used to capture the gene expression profiles of RC212 during 40 inoculated primary fermentations at pilot scale (150 liters) using California Pinot noir grapes from 10 vineyards across two vintages. Using differential expression across the continuous variable Brix, the core gene expression program used by *S. cerevisiae* during these fermentations was observed.

RESULTS AND DISCUSSION

Conditions and rates of fermentation. Pinot noir grapes were harvested from the same 10 vineyards in California during the 2017 and 2019 vintages for wine production at the University of California, Davis, Teaching and Research Winery (Fig. 1A). To standardize fermentations, grapes from the same Pinot noir clone and rootstock were harvested at the same ripeness (~24 Brix, using total soluble solids as a proxy for sugar concentration). Duplicate fermentations that used the grape material from each vineyard were sampled for a total of 40 fermentations (20 from each vintage) at industry-relevant scales using the same wine-making protocol. Each fermentation was inoculated with the commercial wine strain *S. cerevisiae* RC212 and sampled to collect cells for gene expression analysis (3'-biased transcriptome sequencing [3'-Tag-seq]) at 16 h (exponential phase/early fermentation), 64 h (stationary phase/mid-fermentation), and 112 h (decline phase/end of fermentation) postinoculation (Fig. 1B). While sampling times were standardized across fermentations, the rates of fermentation varied, resulting in samples being collected across a range of Brix values (Fig. 1C). Differences in fermentation rates likely reflect diversity in the starting material and different fermentation outcomes, which were also demonstrated in sensory studies performed on wines produced from these vineyard sites in previous vintages (50).

Consistent whole-transcriptome remodeling occurs during fermentation, independent of vintage. The 3'-Tag-seq data from all 10 sites were combined and used to assess differential expression along the continuous variable Brix (Fig. 1C). This approach allowed incorporation and comparison of fermentations with disparate Brix values at mid-fermentation (Fig. 1B). These results define a core vineyard-independent gene expression program of RC212 during California Pinot noir fermentations. Under this model, log₂ fold change values represent the changes in gene expression for each 1-unit decrease of Brix. Therefore, a positive log₂ fold change corresponds to a gene that increased in expression as fermentation progressed, while a negative log₂ fold change value corresponds to a gene that decreased in expression as fermentation progressed (see examples in Fig. 2A). After assessment of differential expression, the




FIG 1 California vineyard locations and fermentation patterns. (A) Map displaying the six AVAs in which the 10 study vineyards are located. (B) Fermentation curves reflecting the change in Brix over fermentation. Brix is a measure of total soluble solids that is used as a proxy for sugar concentration in grapes, grape must, and wine. (C) Brix at the time of sampling for each RNA-seq sample, relative to inoculation. While samples were taken at the same absolute time, fermentations proceeded at different rates, leading to different Brix values in each fermentation.

differentially expressed genes were intersected across vintages to determine consistent changes that were vintage independent. From this analysis, 971 genes decreased expression as Brix decreased, while 1,026 genes increased expression as fermentation progressed (Fig. 2B; also see Data Set S1 in the supplemental material). Each vintage also showed unique differential gene expression patterns, which may occur due to vintage-specific differences in fermentation. However, these data were generated at different times and newly developed methods were applied (unique molecular identifier [UMI] barcoding) (see Materials and Methods) for sequencing of the 2019 samples; therefore, it is suspected that the higher number of differentially expressed genes in the 2017 vintage may reflect differences in the quality of the sequencing data. Nonetheless, the large fraction of shared differentially expressed genes suggests that a core gene expression program is followed independent of vintage.

Of the genes that are differentially expressed in fermentation and shared across vintage, many are known to function in wine fermentation and are central to yeast growth, metabolism, and cell survival (Fig. 2B and C and Table 1). A strong signature of growth early in fermentation that included cellular investment in ribosome biogenesis, metabolism of lipids, purines, and amino acids, and cell division machinery was observed (Fig. 2C; also see Fig. S1 and S2). These processes, coupled with enrichment




FIG 2 Transcriptome remodeling in fermentation is consistent across fermentations and vintages. (A) Graphs (y axis, normalized log gene expression counts; x axis, decrease in Brix over time) showing gene expression patterns in the 2017 and 2019 vintages, involving genes (Continued on next page)

of associated pathways involved in RNA transcription and transport, reflect energy use for cell growth and proliferation associated with log-phase growth occurring in early fermentation. Further in fermentation, changes in ribosomal machinery gene expression occurred, as reported in previous studies (51) (Fig. 2C), reflecting a transition to a nonproliferative metabolic state. Late in fermentation, this was accompanied by changes in gene expression linked to nutrient limitation, altered metabolism, and entry into meiosis (Fig. 2C; also see Fig. S3 and S4), which included gene expression patterns consistent with hallmark isoform switches in hexose transporters and glycolytic enzymes that occur as concentrations of glucose or fructose change (52) (Fig. 2B and C). For example, *HXT1* encodes a low-affinity glucose transporter that was more strongly expressed at the beginning of fermentation, when glucose is abundant. *HXT4* has a high affinity for glucose and is expressed when glucose concentrations are low (53), which was also observed in our data, as *HXT4* expression increased in late fermentation. Importantly, the pathways that were identified as enriched in early and late fermentation align with expectations based on previous research and the known biology of *S. cerevisiae* during fermentation (2–5, 13–16). This highlights the core processes that previous research efforts have defined and provides confidence that the analysis methods employed in these pilot-scale fermentations capture these biologically important transitions.

Beyond these previously defined core gene expression patterns, gene expression signatures indicative of less understood processes within these fermentations were also observed, which may be linked to the industry-like environment in which these studies were performed. These observations are discussed below.

Nutrient limitation in early fermentation. While gene expression data support logarithmic growth at 16 h postinoculation (Table 1; also see Fig. S1 and S2), at this early time point there is also evidence for the expression of genes that are typically upregulated in response to nutrient limitation. *PHO5* and *PHO89* encode phosphate transporters that are induced during phosphate starvation (54), both of which are expressed in early fermentation, along with *PHO90*. Phosphate limitation can cause stuck fermentations, because phosphate is critical for cellular function as a component of ATP, nucleotides, sugars, lipids, and macromolecules such as proteins (51, 55). Given that all of these Pinot noir fermentations went to completion and that the majority of glucose was converted to ethanol after the 16-h time point, induction of genes encoding phosphate transporters early in fermentation is not likely associated with phosphate starvation. Instead, it may be a response to the presence of non-*Saccharomyces* yeast, as cocultivation of *S. cerevisiae* with *Torulaspota delbrueckii* led to the induction of a gene encoding a high-affinity phosphate transporter (*PHO84*) after only 3 h of fermentation (42). Enological coculture of *S. cerevisiae* with organisms such as *Hanseniaspora guilliermondii* and *Brettanomyces* spp. has also been linked to induction of genes involved in vitamin biosynthesis in fermentation (56, 57), which could be indicative of increased nutrient competition and depletion of some nutrients early in fermentation. Induction of genes that encode enzymes involved in biosynthesis of B vitamins in early fermentation, including *BIO2* (biotin biosynthesis), *RIB3* and *RIB4* (riboflavin biosynthesis), *PAN6* (pantothenate synthesis), *SPE3* and *SPE4* (pantothenic acid synthesis), and *MIS1* and *FOL1* (folate biosynthesis) was observed. In addition, *THI21* was induced, which is involved in thiamine biosynthesis. As with phosphate, this may be related to the presence of metabolically active non-*Saccharomyces* microorganisms that were detected in all of these fermentations (58). We expect that continued work using industry-like fermentations across grape varieties and yeast strains, as well as controlled

FIG 2 Legend (Continued)

more highly expressed early (*HXT1*) and late (*HXT4*) in fermentation or constitutively (*ADH1*) across fermentation. (B) Upset plot showing the intersection of genes that are more highly expressed at the beginning and end of fermentation in each vintage, using a \log_2 fold change cutoff value of 1. The majority of genes are consistently expressed across fermentations and vintages. (C) Proteomaps depicting Gene Ontology pathways (left) and genes (right) that are more highly expressed in early (top) and late (bottom) fermentation. The sizes of individual genes reflect the associated \log_2 fold change values. Note that, for presentation purposes, not all genes that are significantly expressed are depicted. See Data Set S1 in the supplemental material for a complete list of genes.

TABLE 1 Genes differentially expressed throughout fermentation shared across vintages

Cellular process and gene(s)	Expressed at highest levels	Gene product function
Constitutively expressed in fermentation		
Glycolysis and fermentation		
<i>HXT3</i>	Late	Hexose transporter induced by both high and low glucose concentrations.
<i>PFK1, PFK2</i>	<i>PFK2</i> , early	Phosphofructokinases that catalyze the first irreversible reaction specific to glycolysis, producing fructose-1,6-bisphosphate from fructose-6-phosphate.
<i>ADH1 to ADH5</i>	<i>ADH2 to ADH4</i> , early; <i>ADH5</i> , late	Alcohol dehydrogenase isoenzymes. The dominant fermentative alcohol is <i>ADH1</i> (138), responsible for reoxidation of NADH to NAD ⁺ , which is a required cofactor in the metabolism of glyceraldehyde-3-phosphate in glycolysis. <i>ADH2</i> is a nondominant isoenzyme of alcohol dehydrogenase. It is typically repressed by glucose and converts ethanol to acetaldehyde (138), and it is overexpressed in some wine strains (139).
Expressed at higher levels in early fermentation in both 2017 and 2019 vintages		
Glycolysis and fermentation		
<i>HXT1</i>		Low-affinity hexose transporter.
<i>HXX2</i>		Hexokinase that phosphorylates glucose in the first irreversible step leading to glycolysis.
<i>PFK1, PFK2</i>		Phosphofructokinases that catalyze the first irreversible reaction specific to glycolysis, producing fructose-1,6-bisphosphate from fructose-6-phosphate.
Acetate metabolism		
<i>ALD4 to ALD6</i>		Aldehyde dehydrogenase isoenzymes that produce acetate as a byproduct when acetaldehyde is metabolized. <i>ALD6</i> encodes the main isoenzyme responsible for acetate production in wine (98). Aldehyde dehydrogenase isoenzymes <i>ALD4</i> and <i>ALD5</i> are expressed when ethanol is the carbon source and are not typically associated with wine fermentation. Acetate contributes the majority of volatile acidity associated with negative organoleptic properties in wine (140).
<i>PDR12</i>		Plasma membrane ABC transporter that is required for development of resistance to weak organic acids, including acetate (141).
Cell cycle		
<i>BAT1</i>		Expressed in logarithmic phase (142).
<i>CWP1</i>		Expressed in the S/G ₂ phase of the cell cycle (142).
Nitrogen metabolism		
<i>GNP1</i>		High-affinity glutamine permease that also transports leucine, serine, threonine, cysteine, methionine, and asparagine.
<i>MUP1</i>		High-affinity methionine permease that is also involved in cysteine transport.
<i>CAR1, CAR2</i>		Involved in arginine catabolism. Arginine is the most abundant amino acid in grape must after proline (68) and is used in protein synthesis during fermentation (143).
<i>YPQ1</i>		Vacuolar transporter for arginine and lysine. Unused arginine is stored in the vacuole for later use (143).
Ehrlich pathway		
<i>BAT1, ARO8</i>		Catalyzes transamination of amino acids, the product of which cannot be redirected to central carbon metabolism and so is excreted as fusel acid or fusel alcohol (144). Overexpression of <i>BAT1</i> increases the concentrations of isoamyl alcohol, its acetate ester, and isobutanol in wine (144).
<i>PDC1</i>		Catalyzes α -keto decarboxylation.
Glycerol biosynthesis		
<i>GPP1</i>		Cleaves phosphate from glycerol-3-phosphate.
Expressed at higher levels in late fermentation in both 2017 and 2019 vintages		
Nitrogen limitation		
<i>GAT1, DAL80</i>		Transcriptional activator (<i>GAT1</i>) and repressor (<i>DAL80</i>) of genes under nitrogen catabolite repression. Expression is inversely correlated, and the detection of both genes as induced in late fermentation likely indicates tight transcriptional regulation of nitrogen metabolism.
<i>DAL2 to DAL5, DAL80, DAL82</i>		Catalyze allantoin degradation, and expression is under nitrogen catabolite repression.
<i>MEP2</i>		Ammonia permease, and expression is under nitrogen catabolite repression.

(Continued on next page)

TABLE 1 (Continued)

Cellular process and gene(s)	Expressed at highest levels	Gene product function
<i>GAP1</i>		Amino acid permease.
<i>PTR2</i>		Peptide permease.
<i>AVT3, AVT4</i>		Vacuolar amino acid exporters that mobilize internal nitrogen stores for cell maintenance during stationary phase. Expression is under nitrogen catabolite repression.
Ubiquitin-mediated selective protein degradation		
<i>RPN4</i>		Transcription factor that induces expression of proteasome genes.
<i>TMC1</i>		Effector of proteotoxic stress that is induced by nitrogen limitation, weak acids, and misfolded proteins and is a target of <i>RPN4</i> .
<i>UBC8, VID24</i>		Negative regulators of fructose-1,6-bisphosphate through ubiquitination (<i>UBC8</i>) and vacuolar targeting (<i>VID24</i>).
<i>UBC1, UBC5, UBC7, UBC13</i>		Ubiquitin-conjugating enzymes.
<i>UBI4, CUZ1</i>		Involved in the ubiquitin-proteasome pathway.
Autophagy		
<i>ATG2, ATG4, ATG7 to ATG12, ATG14, ATG32, ATG40</i>		Proteins involved in autophagy. Autophagy is a key response to nutritional limitation that allows cells to maintain homeostasis (145). Nitrogen starvation leads to the largest autophagic response in yeast.
Ehrlich pathway		
<i>GRE2</i>		Final step of the pathway in which fusel aldehydes are oxidized or reduced to fusel acids or alcohols (144).
Carbon limitation		
<i>SNF3</i>		Plasma membrane low-glucose sensor involved in regulating glucose transport.
<i>SKS1</i>		Serine/threonine kinase involved in the adaptation to low glucose levels via <i>SNF3</i> -independent signaling.
<i>PGM2</i>		Phosphoglucomutase that catalyzes a key step in hexose metabolism and is induced in response to glucose limitation and ethanol stress (146).
<i>HXK1</i>		Hexokinase that phosphorylates glucose or fructose in the first irreversible step leading to glycolysis, under glucose-induced repression.
<i>HXT4, HXT6</i>		Hexose transporters required at the end of alcoholic fermentation.
Trehalose and glycogen		
<i>TSL1, NTH1, ATH1</i>		Involved in trehalose synthesis (<i>TSL1</i>) and degradation (<i>NTH1</i> and <i>ATH1</i>). Trehalose acts as a storage carbohydrate for cell maintenance under nongrowth conditions (147, 148), bolsters membrane integrity by displacing ethanol (148), and protects proteins from denaturation (149). Trehalose recycling is an important component of stress response (150).
<i>GLG1, GSY1, GSY2, IGD1, GPH1, SGA1, GAC1, GIP1, YPI1, PIG2, GLC8</i>		Involved in glycogen accumulation (<i>GLG1, GSY1, GSY2</i> , and <i>IGD1</i>), degradation (<i>GPH1</i> and <i>SGA1</i>), and metabolism (<i>GAC1, GIP1, YPI1, PIG2</i> , and <i>GLC8</i>). Glycogen accumulates during nutrient abundance and is metabolized during stationary phase and nutrient deprivation (147). Glycogen recycling is an important component of stress response (150).
Cell wall integrity		
<i>PIR3, SED1, SLT2</i>		Target genes of the cell wall response to ethanol. <i>PIR3</i> is required for cell wall stability and is induced in part by <i>SLT2</i> . <i>SED1</i> is a stress-induced cell wall structural protein (<i>SED1</i>) (151).
<i>PKH2, YPS1, PST1, KRE1</i>		General cell wall integrity response.

fermentations using reconstituted microbial consortiums, will be critical for understanding the relevance of these gene expression signatures to wine fermentation outcomes. If understood, such interactions could potentially be addressed through timely nutrient additions to a fermentation to achieve desired outcomes.

Evidence of varied gene expression patterns linked to oxygen exposure during fermentation. A wine fermentation is generally regarded as an anaerobic process, given that the carbon dioxide (CO₂) produced as a by-product of ethanol fermentation protects must from dissolved oxygen (59). However, within anaerobiosis, there is an important distinction between hypoxic (low-oxygen) and anoxic (no-oxygen) conditions. In a fermentation, it is expected that molecular oxygen (O₂) is introduced into the grape must by handling processes, including pump-overs, that may introduce small amounts of dissolved oxygen into industrial-scale tanks (60). Stratification within a fermentation may also expose

local cell populations to different oxygen environments, leading to yeast cell populations undergoing different anaerobic processes. In this study, gene expression patterns were consistent with different populations of cells experiencing varied levels of oxygen exposure during fermentation. For example, the yeast cell wall undergoes remodeling in response to oxygen availability, which is accomplished in part by regulated expression of cell wall mannoproteins encoded by *CWP1*, *CWP2*, *DAN1*, and *TIR1* to *TIR4* (61). Specifically, expression of *DAN1* and *TIR1* to *TIR4* occurs reciprocally with expression of *CWP1* and *CWP2*, with the *CWP* genes being expressed under aerobic conditions and *DAN1* and *TIR1* to *TIR4* under anaerobic conditions (61). *DAN1* expression is known to be repressed under aerobic conditions by four independent regulatory mechanisms (62). Interestingly, expression of both *CWP1* and *DAN1* and *TIR1* to *TIR4* was observed in early fermentation samples. Similarly, in early fermentations, both *HYP2* and *ANB1* were expressed. These paralogous genes encode translation elongation factor eIF5A and are part of a family of paired genes for which oxygen induces the aerobic isoform and represses the hypoxic isoform (63). *HYP2* is expressed during aerobic growth, while *ANB1* is expressed during hypoxic growth and is tightly regulated by the presence of oxygen (64). Together, these gene expression patterns indicate varied gene expression programs within yeasts that may be explained by differing levels of oxygen exposure.

Among late expressed genes, oxygen-regulated paired isoforms, including *COX5A* and *COX5B*, which encode a subunit of cytochrome *c* oxidase, were also expressed. Modulated expression of these two isoforms allows *S. cerevisiae* to produce holoenzymes with different catalytic properties in response to oxygen (65). *COX5A* expression declines between 1 and 5 $\mu\text{mol/liter O}_2$ and is undetectable below 0.25 $\mu\text{mol/liter O}_2$, while *COX5B* is undetectable until 0.25 $\mu\text{mol/liter O}_2$ (63). Simultaneous induction of both transcripts at the end of fermentation is again consistent with cells experiencing varied levels of dissolved oxygen in fermentation (60). In contrast, of the oxygen-regulated isoform pair *CYC1* and *CYC7* (63), only expression of the hypoxic isoform *CYC7* was detected at the end of fermentation. The breakpoint between expression of isoforms occurs at a higher concentration of 0.5 $\mu\text{mol/liter O}_2$ for *CYC1* and *CYC7*, compared with *COX5A* and *COX5B* (63), which may indicate that dissolved oxygen levels did not exceed 0.5 $\mu\text{mol/liter}$ and thus were not permissive for expression of *CYC1*.

In late fermentation, induction of pathways such as glycerol degradation and proline metabolism, which require oxygen, was also observed. Glycerol is a compatible solute involved in combating osmotic stress and redox balance and is primarily produced in early fermentation (66). *GCY1*, which encodes a glycerol dehydrogenase used under microaerobic conditions (67), was induced, as was *RSF2*, a transcriptional regulator of genes that encode proteins required for glycerol-based growth. Proline metabolism genes *PUT1*, *PUT2*, and *PUT4* were also expressed at the end of fermentation. Although proline is an abundant amino acid in grape must, it is a nonpreferred nitrogen source of yeast and requires oxygen to be metabolized (68). It was further observed that *PUT1* and *PUT2* were induced in a sealed laboratory wine fermentation but proline was not metabolized, given the absence of oxygen (2). Expression of *PUT1*, *PUT2*, and *PUT4* is regulated by nitrogen catabolite repression (69) and the presence of proline in the absence of other nitrogen sources (70) but is not regulated by the presence of oxygen. Intracellular proline accumulation also protects *S. cerevisiae* from reactive oxygen species (ROS) associated with ethanol-rich environments (71). While it possible that glycerol and proline were metabolized in late fermentation with oxygen ingress, other processes, such as nutrient limitation and oxidative stress, may also explain the induction of these genes.

Taken together, the gene expression data presented herein raise various questions about a distributed gradient of oxygen (hypoxia and anoxia) in the fermentation environment that may induce varied gene expression across the cell population. This could lead to yeast subpopulations with varied metabolic outputs or different levels of ethanol tolerance, due to the role of oxygen in these processes (72, 73). In the future, single-cell sequencing technologies combined with continuously monitored dissolved oxygen assays

may help resolve these questions. From a production perspective, in industrial fermentations, even those that employ pump-over systems and thus maintain mixing and better homogeneity, there is a gradient of dissolved oxygen in the fermentation tank, with higher oxygen concentrations toward the top of the vessel (60). This suggests that heterogeneous gene expression profiles in response to oxygen would likely exist in these environments too. This is also an important fact to consider, because oxygen additions during fermentation are known to influence both fermentation and sensory outcomes. For example, in late fermentation, a single oxygen pulse increases the rate of fermentation mediated by ergosterol biosynthesis (72). Similarly, oxygen additions at different stages of fermentation differentially impact the formation of wine aroma compounds such as volatile thiols and esters; however, this appears to occur in a strain-dependent manner (73). This knowledge, combined with the impact of oxygen addition on fermentation outcomes, raises the idea that timely addition of oxygen may be a way to control fermentations rates and formation of wine aromas, which would be a tool easily accessible to winemakers.

Mitochondria and fermentation. In late fermentation, there was striking enrichment of pathways involved in mitochondrial biogenesis and function, as well as oxidative phosphorylation, among differentially expressed genes (Fig. 2C; also see Fig. S3 and S4). Substantial metabolic investment in mitochondrial systems suggests a critical role for mitochondria late in fermentation. What that role is remains unclear, however, as limited research has been conducted on the mitochondria during enological fermentation (74, 75). While some studies that profiled the transcriptomes of primary fermentations either found no evidence for or made no comment regarding enrichment for oxidative metabolism at the end of fermentation, many studies found induction of mitochondrial genes, particularly those encoding proteins involved in oxidative phosphorylation. These studies included fermentations conducted under nitrogen limitation (6), lipid limitation (76), and standard laboratory conditions (3). Interestingly, under lipid limitation, oxidative phosphorylation was induced in the exponential phase of growth, as opposed to the end of fermentation (76). Given the role of membrane lipid composition in combating ethanol-induced membrane permeability (77) and the accumulation of ROS during ethanol exposure (78), induction of the respiratory chain may mitigate ROS, which are abundant at the end of fermentation. Nonetheless, the recurrence of these gene expression patterns in our studies and previous laboratory experiments suggests that cells are investing in mitochondrial systems during fermentation.

One potential reason for late induction of mitochondrial systems is that glucose limitation relieves the Crabtree effect. This may lead to induction of oxidative phosphorylation genes that change metabolism in a nutrient-limited environment to one that generates the largest amount of ATP per unit of glucose (79). In this way, an investment in mitochondrial infrastructure during late fermentation may be a starvation adaptation in which *S. cerevisiae* uses oxidative phosphorylation to harness the largest fraction of energy possible from the remaining carbohydrate sources. However, this strategy is predicated on the availability of molecular oxygen, which is required for the induction and function of the respiratory apparatus (80, 81). A second reason for mitochondrial gene expression may be related to the fact that meiosis- and sporulation-related genes were enriched at the end of fermentation (Fig. 2C; also see Fig. S3 and S4). Induction of meiosis likely occurs to produce spores resistant to the challenges of nutrient limitation and stress (82). Interestingly, mitochondrial biomass is a predictor of meiosis (83), and components of the respiratory chain are required for initiation of sporulation (84), providing another potential process that may underlie mitochondrial investment in late fermentation. Related to this fact, a propensity for yeast to undergo meiosis at this stage of vinification underlies fast adaptive genomic evolution of *S. cerevisiae* (85), suggesting that this may be an important acquired trait that allows yeast to successfully survive the wine environment.

Mitochondria also fulfill other critical roles in fermentation that are unrelated to respiration. For example, mitochondria play a role in sterol uptake and transport under strictly anaerobic conditions (86), and mitochondria quench ROS, especially during ethanol stress (87). While we did not observe induction of specific genes related to sterol biology

and we found induction of different genes related to ROS, compared to those identified previously (see below), these processes may also be linked to increased mitochondrial gene expression. Regardless of the role played by mitochondria in late fermentation, the striking and consistent induction of these genes in fermentations signals that more research is needed to understand the role of mitochondria in fermentation.

Thioredoxin and glutathione system activity throughout fermentation. The reducing environment of the cytosol in *S. cerevisiae* is key to various cellular functions, including deoxyribonucleoside triphosphate synthesis and the elimination of toxic compounds, including oxidants generated through cellular metabolism (88, 89). Key to maintaining redox balance are the thioredoxin (TRX) and glutathione (GSH) thiol reductase systems. For example, proper redox homeostasis is required to maintain the redox status of cysteine residues, which are essential for the function of numerous enzymes, protein receptors, and transcription factors. Similarly, redox homeostasis within cells aids to balance pools of reduced and oxidized pyridine nucleotide cofactors (NAD⁺/H and NADP⁺/H) that are essential to numerous metabolic reactions. ROS can alter this redox balance, causing oxidative stress and direct or indirect ROS-mediated damage of nucleic acids, proteins, and lipids. While typically associated with respiratory metabolism, ROS can be generated throughout fermentation, particularly by superoxide anions and peroxides (78, 90, 91). ROS may also be created by acetaldehyde, an intermediate in ethanol production (92).

In early fermentation, genes involved in the TRX system, such as *TRX1* and *TRR1*, were induced. Expressed targets of *TRX1* included *RNR1* to *RNR4* (93), genes encoding ribonucleotide diphosphate reductases required for DNA synthesis and cell cycle progression, as well as *MET16*, which encodes an enzyme required for sulfate assimilation (94). We further observed genes encoding Trx1 target peroxidases (*TSA1*) and peroxiredoxins (*AHP1*) that are constitutively expressed throughout fermentation along with superoxide dismutases (*SOD1* and *SOD2*). An additional source of ROS is peroxisomes, which may generate hydrogen peroxide in early fermentation via β -oxidation of fatty acids. *CTA1*, which encodes a peroxisomal catalase, and *ANT1*, which encodes a peroxisomal transporter involved in β -oxidation of fatty acids, were expressed in early fermentation. A major factor used to maintain redox balance is NADPH, which provides reducing potential for the TRX system. It has been shown that metabolic intermediates in glycolysis can be rerouted to the pentose phosphate pathway to generate NADPH in response to oxidative stress (95–97). In this study, the pentose phosphate pathway was enriched among genes expressed in early fermentation (Fig. 2C; also see Fig. S1 and S2), which includes *GND1*, encoding an enzyme that catalyzes NADPH regeneration and is required for the oxidative stress response. Other expressed genes that encode enzymes acting downstream of *GND1* in the pentose phosphate pathway included *RPE1*, *TLK1*, *TLK2*, and *TAL1*.

Central to the GSH thiol reductase system is GSH, an abundant tripeptide that is conserved throughout eukaryotic and prokaryotic cells, with a critical role in redox control, but its physiological role is both diverse and debated (97). Genes encoding enzymes involved in the degradation (*DUG1* and *DUG2*), import (*OPT1*), and biosynthesis (*GSH1* and *GSH2* in the 2017 vintage) of GSH were expressed in early fermentation. Additional generation of NADPH in early fermentation may be supported by the transformation of isocitrate to α -ketoglutarate via *IDP1* in the mitochondria and export via *YMH2*, as both genes were also expressed. Genes encoding aldehyde dehydrogenases (*ALD5* and *ALD6*) are similarly expressed in early fermentation, and both may regenerate NADPH through the transformation of acetaldehyde to acetate. *ALD6* is the dominant isoenzyme responsible for acetate production in wine (98).

Genes involved in GSH-mediated ROS mitigation were also induced in late fermentation. For example, a gene encoding cytosolic glutaredoxin (*GRX1*) was expressed in late fermentation. Unlike glutaredoxins in other species (e.g., mammals), yeast glutaredoxins do not function as deglutathionylase enzymes (99). Instead, induction of *GRX1* increases resistance to hydroperoxides by catalytically reducing hydroperoxides

through GSH conjugation and using the reducing power of NADPH (100). In addition, the cytosolic peroxidase GPX1 was expressed. GPX1 uses both GSH and TRX, in combination with NADPH, for reducing power (101). *GPX1* is known to be expressed with glucose and nitrogen starvation (102), which coincides with peak peroxide formation in yeast during wine fermentation (90). While the gene expression data in this study support a role for cytoplasmic GSH during late fermentation, genes encoding mitochondrial peroxidin (*PRX1*) and TRX (*TRX3*) were also expressed. Prx1 buffers the mitochondria from oxidative stress and is reductively protected by GSH, TRX reductase (Trr2), and Trx3 (103). Taken together, these results suggest that cytoplasmic and mitochondrial systems may be integral to combating increased oxidative stress at the end of fermentation.

GSH is also important for maintenance of cellular functions via other systems. For example, methylglyoxal, a reduced derivative of pyruvic acid, is a by-product of glycolysis that may account for up to 0.3% of glycolytic carbon flux in *S. cerevisiae* (104). GLO2, an enzyme that catalyzes methylglyoxal degradation in a GSH-dependent manner, was expressed in late fermentation, as were GSH-independent systems involved in the degradation of methylglyoxal (GRE2 and GRE3). Genes that encode proteins involved in GSH homeostasis were also expressed at the end of fermentation, including *GEX1*, which encodes a proton/GSH antiporter (105, 106). *GEX1* is known to be induced during oxidative stress (105) and modulates formation of the aromatic thiol 3-mercaptohexan-1-ol from its glutathionylated precursor in wines such as Sauvignon blanc (106). Conversely, induction of a gene that encodes an enzyme that cleaves GSH (*GCG1*) was observed and may be involved in apoptotic signaling via ROS accumulation (107).

Together, these gene expression patterns highlight how intertwined redox homeostasis is with almost all core metabolic processes in *S. cerevisiae*, as most pathways require oxidation or reduction by a pyridine nucleotide cofactor during at least one reaction. For example, NAD⁺/H and NADP⁺/H participate in 740 and 887 biochemical reactions through interactions with 433 and 462 enzymes, respectively (108). It is also well documented that experimental perturbation of both NAD⁺/H and NADP⁺/H leads to changes in aroma compounds in wine and other fermented beverages (109–112). The observations presented here, conserved across many Pinot noir fermentations, involving genes engaged in redox balance and mitigation of oxidative stress via thiol reductase systems offer further evidence for the importance of these systems. These findings provide motivation for future studies of these systems in the context of wine production, which would include control measures to aid cellular control of redox and to mitigate oxidative cellular stress.

Stress-associated gene expression during fermentation. During fermentation, *S. cerevisiae* has to adapt to a continually changing stress landscape. Macronutrients and micronutrients become limiting as ethanol concentrations increase and, as discussed above, production of acetaldehyde and other metabolic processes generate oxidative stress. To accommodate this dynamic environment, *S. cerevisiae* wine strains express genes that overlap, but are distinct from, the stress response of laboratory strains (7, 113). In accordance with previous studies (2, 3), a partial overlap was observed between genes expressed in fermentation and those involved in the ESR in laboratory strains. Specifically, 16 ESR genes were expressed at the beginning of fermentation and 78 ESR genes were expressed at the end of fermentation. This matches observations in synthetic must, in which stress genes were induced upon entry into stationary phase (2). Stress-related genes expressed at the beginning of fermentation were enriched for Gene Ontology pathways involving carbohydrate (mannose, fructose, glucose, and hexose) transmembrane transport and NADP regeneration (Fig. 3A), while stress-related genes expressed at the end of fermentation were enriched for oxidation-reduction processes, generation of precursor metabolites and energy, energy reserve metabolic processes, and glycogen metabolic processes (Fig. 3B).

A recent study investigated the fermentation of Riesling grape must at laboratory scale without the addition of oxygen (3). Using microarray analysis at five time points




FIG 3 Pathways enriched among genes differentially expressed across fermentation that are shared with the ESR. (A) Of 16 genes that overlap the ESR and are expressed in early fermentation, pathways related to carbohydrate metabolism were enriched. (B) Of 78 genes that overlap the ESR and are expressed later in fermentation, pathways related to oxidation-reduction and carbohydrate metabolism were enriched. GeneRatio refers to the fraction of genes in an enriched gene set that were present in the tested set.

in fermentation, the authors defined a fermentation stress response (FSR) as those genes that are induced at any point in fermentation and do not return to baseline (3). The FSR is differentiated from the ESR and the common stress response because adaptation over time through gene expression returning to prestress transcription levels is not observed (3, 7, 113). Of the 223 genes induced in the FSR, 84 were observed to be expressed in mid-fermentation or late fermentation. Of these 84 genes, 43 overlap genes expressed in other stress responses, as defined previously (3), including 16 with the ESR and 14 with the common stress response. Of the 41 genes that overlap the FSR, many were related to the challenging nutrient environment in wine, including glucose limitation (*NRG1*, *SKS1*, *HXT6*, and *VID24*), nitrogen limitation (*MEP2*, *GAP1*, *PTR2*, *AVT4*, and *VBA2*), vitamin limitation (*MCH5* and *VHR1*), and stress caused by heat, salt, protein misfolding, and cell wall defects (*GAC1*, *RPI1*, *JID1*, and *PSR2*). This suggests that multiple stress pathways are simultaneously activated by the challenging environment that *S. cerevisiae* encounters in wine fermentation, which likely defines the




FIG 4 Normalized log gene expression counts for genes involved in mannitol transport and degradation. *HXT13* (A and C) and *MAN2* (B and D) expression in the 2017 (A and B) and 2019 (C and D) vintages is graphed. *MAN2* and *HXT13* were the most-expressed genes at the end of fermentation in 2019 and fell behind only *HXT4* in the 2017 vintage. Gray lines indicate a linear model fit to normalized counts.

described FSR. Many genes identified in the FSR and expressed in this study during fermentation remain uncharacterized (*YPR152C*, *YBR085C-A*, *YDL024C*, *YDR042C*, *YMR244W*, and *YLL056C*), offering gene targets for future investigations related to adaptation to the fermentation and wine environment.

Polyol metabolism in late fermentation. Polyols, also called sugar alcohols, have recently been shown to be produced by non-*Saccharomyces* yeasts and by fructophilic lactic acid bacteria such as *Lactobacillus kunkeei* during fermentation (114, 115). Combined with other spoilage organism-associated metabolites, these compounds can have a significant impact on wine quality (116). Mannitol is one such polyol and a nonpreferred sugar that can be metabolized by *S. cerevisiae* (117–119). In *S. cerevisiae*, transporters encoded by *HXT13* and *HXT15* to *HXT17* were found to facilitate mannitol and sorbitol transport (118). In this study, the mannitol transporter gene *HXT13* was induced in both vintages, along with the mannitol dehydrogenase gene *MAN2*, which together indicate that mannitol may be present and metabolized by *S. cerevisiae* at the end of fermentation (Fig. 4). In line with this, although eukaryotic transcriptional profiling via 3'-Tag-seq was performed (see Materials and Methods), *L. kunkeei* transcripts were detected in some fermentations in both the 2017 and 2019 vintages (58), which is one potential source of mannitol production. These data raise the possibility of mannitol consumption by *S. cerevisiae*, demonstrating metabolic flexibility for carbon sources late in fermentation.

Notably, *L. kunkeei* can influence *S. cerevisiae* metabolism beyond the expression of genes for nonpreferred carbon sources. Via production of acetic acid and possibly

other compounds, *L. kunzei* has been shown to induce the [GAR⁺] prion phenotype in *S. cerevisiae*, thereby shifting carbon metabolism away from hexoses (120, 121). Given that the presence of *L. kunzei* RNA was detected in the 2017 vintage, the presence of the [GAR⁺] phenotype was tested for in the 2019 vintage via cell culture (121). The [GAR⁺] prion was not detected in any fermentation tested in the 2019 vintage. While the absence of the [GAR⁺] phenotype in the 2019 vintage does not preclude its presence in the 2017 vintage, consistent gene expression for mannitol transport and degradation in both vintages suggests that *S. cerevisiae* may be metabolizing mannitol in these Pinot noir fermentations due to the presence of non-*Saccharomyces* organisms, including *L. kunzei*.

Vintage-specific differences. From this data analysis, there were 778 genes and 385 genes differentially expressed in the 2017 and 2019 vintages, respectively. The majority of these genes were members of pathways enriched among all fermentations (Fig. 2C; also see Fig. S1 to S4). Using Gene Ontology enrichment analysis, no molecular function, cellular compartment, or biological process was enriched in either vintage that was not enriched in both vintages. This suggests that these differences may be largely due to sequencing depth or variations in the gene expression within these pathways and not differences in the overall biology of *S. cerevisiae*. Still, signatures indicative of vintage-specific effects, some of which may impact the sensory attributes of wine, were observed. For example, glycerol is an important fermentation by-product that can contribute to the mouth feel of wine (122). *S. cerevisiae* uses glycerol biosynthesis to generate NAD⁺, a required cofactor for glycolysis, when NAD⁺ levels are not sufficiently replenished through fermentation (123). During glycerol biosynthesis, enzymes encoded by *GPD1* and *GPD2* convert dihydroxyacetone phosphate into glycerol-3-phosphate (124). Both *GPD1* and *GPD2* were expressed in early fermentation in the 2017 vintage but not in the 2019 vintage. A second example involves genes encoding the fluoride transporters Fex1 and Fex2, which were expressed in late fermentation across all fermentations in the 2019 vintage. Fluoride is a toxic anion that *S. cerevisiae* exports via two plasma membrane transporters to avoid cell damage (125), which in excess can cause slow or stuck fermentation (126). Although fluoride is ubiquitous in terrestrial and aquatic environments (125), application of the insecticide Cryolite, which contains fluoride, has caused problematic fermentations in California vineyards (126). Currently, the reasons for these vintage-specific gene expression patterns are not known.

Finally, it was observed that genes of currently unknown function were differentially expressed in the two vintages assayed. Using a log₂ fold change cutoff value of 2, 14 genes in the 2017 vintage and 7 genes in the 2019 vintage were of unknown function. Across both vintages, more genes of unknown function were expressed in late fermentation than in early fermentation (10 in 2017 and 5 in 2019). Knowledge of the specific pathways expressed in late fermentation due to the stressful, nutrient-limited conditions offers clues to the potential functions of these genes that could be explored in future work.

Conclusion. In this study, a gene expression analysis from 40 pilot-scale fermentations of California Pinot noir wine using grapes from 10 vineyard sites and two vintages is presented. The fermentations were diverse, with different kinetics, initial chemical conditions, and microbial communities (58). Yet among this diversity, a core gene expression program by *S. cerevisiae* that is largely consistent with that observed at laboratory scale was detected (2–4). Given that there were many genes consistently expressed across these Pinot noir fermentations from diverse vineyards, members of this core fermentation gene program represent strong candidates for future study to impact wine outcomes, e.g., through manipulating the redox balance (109–112). Excitingly, this includes a large number of genes with unknown function that, through investigation, may provide new insights into the biology of *S. cerevisiae*.

The largest deviations from benchtop fermentations are likely attributed to activities of non-*Saccharomyces* organisms, but more research is needed to understand these complex ecological interactions and their impact on fermentation. The gene expression signatures around oxygen presence and metabolic availability also warrant

further research, particularly into the role of the mitochondria in late fermentation (3, 6, 76). While few vintage-specific differences were detected between fermentations, we expect that there are vineyard-site specific deviations from the consistent patterns of gene expression described here. Given the variability in fermentation kinetics with respect to time of sampling, new methods will likely be needed to resynchronize stages of fermentation to enable cross-vineyard comparisons (4). Future work is also needed to extend these observations to other grape varieties and *S. cerevisiae* wine strains, which will define both the shared and unique facets of the core gene expression program in *S. cerevisiae* linked to these variables. With such information, the impact of an industrial wine fermentation environment on *S. cerevisiae* gene expression can be addressed and approaches that can be used to manage commercial fermentation outcomes can be defined.

MATERIALS AND METHODS

Grape preparation and fermentation. The wine-making protocol used in this study was described previously (25, 50). The grapes used in this study originated from 10 vineyards in six American Viticulture Areas (AVAs) in California. All grapes were Pinot noir clone 667 rootstock 101-14. Grapes were harvested at approximately 24 Brix, and the fruit was transported to the University of California, Davis, Teaching and Research Winery for fermentation. Separate fermentations were performed for grapes from each site, with two fermentations per site, totaling 20 fermentations per vintage (40 fermentations in total). After harvest, the fruit was separated into one-half-ton macrobins on harvest day, and Inodose SO₂ (potassium metabisulfite and potassium bicarbonate) was added to achieve SO₂ levels of 40 ppm. The bins were stored in a 14°C cold room until destemming and division of the fruit into temperature-jacket-controlled tanks. N₂ sparging of the tank headspace was performed prior to fermentation, and the tanks were sealed with a rubber gasket. Grapes were cold soaked at 7°C for 3 days, with SO₂ additions made on day 2 of the cold soak to maintain a level of 40 ppm total SO₂. On the morning of day 3, musts were warmed for inoculation to 21°C with programmed pump-overs used to hold the tank at a constant temperature. Once the musts reached that temperature (~2 to 3 h), the musts were inoculated. For inoculation, *S. cerevisiae* RC212 (Lallemand) was reconstituted with Superstart Rouge (Laffort) at 20 g/hl and the must was inoculated with 25 g/hl yeast. Superstart Rouge is a yeast prepate for active dry yeast rehydration. Fermentation progress was determined by measuring Brix with a density meter (Anton Paar 35 DMA). At approximately 24 h after inoculation, the nitrogen content in the fermentations was adjusted by adding diammonium phosphate (DAP), according to the formula (target yeast-assimilable nitrogen [YAN] – 35 mg/liter – initial YAN)/2, and Nutristart (Laffort) using 25 g/hl. Nitrogen was adjusted only if the target YAN level was below 250 mg/liter based on measures of ammonia and free α -amino nitrogen content (Gallery automated photometric analyzer; Thermo Fisher Scientific). Approximately 48 h after fermentation, fermentation temperatures were permitted to increase to 27°C and DAP was added as described previously. Fermentations ran to completion when Brix was <0. Fermentations were sampled for Brix measurements and RNA isolation at 16, 64, and 112 h relative to inoculation. To ensure uniform sampling, a pump-over was performed 10 min prior to sampling of each tank. For RNA samples, 12 ml of juice was obtained and centrifuged at 4,000 rpm for 5 min. The supernatant was discarded, and the pellet was frozen in liquid nitrogen. Samples were stored at –80°C until RNA extraction.

RNA extraction and sequencing. Frozen yeast pellets were thawed on ice, resuspended in 5 ml Nanopure water, and centrifuged at 2,000 × *g* for 5 min, and the supernatant was aspirated. RNA was extracted using the Quick RNA fungal/bacterial miniprep kit, including DNase I column treatment (catalog number R2014; Zymo Research). Samples were eluted in 30 μ l of molecular grade water and assessed for concentration and quality via a NanoDrop spectrophotometer and RNA gel electrophoresis. Sample concentrations were adjusted to 200 ng/ μ l, and 20 μ l was sent for sequencing. 3'-Tag-seq (Lexogen QuantSeq) was used in both the 2017 and 2019 vintages, with the addition of UMI barcodes in 2019. The University of California, Davis DNA Technologies Core performed all library preparation and sequencing.

Differential expression analysis. Samples were processed according to manufacturer's recommendations (Lexogen). First, the first 12 bp was hard trimmed from each read and Illumina TruSeq adapters and poly(A) tails were removed. Next, STAR was used to align the reads against the *S. cerevisiae* S288C genome (reference genome R64 [GenBank accession number [GCF_000146045.2](https://www.ncbi.nlm.nih.gov/nuccore/GCF_000146045.2)]) with parameters –outFilterType BySJout –outFilterMultimapNmax 20 –alignSJoverhangMin 8 –alignSJDBoverhangMin 1 –outFilterMismatchNmax 999 –outFilterMismatchNoverLmax 0.6 –alignIntronMin 20 –alignIntronMax 1000000 –alignMatesGapMax 1000000 –outSAMattributes NH HI NM MD –outSAMtype BAM SortedByCoordinate (127). For the 2019 vintage, UMI-tools was used to deduplicate alignments (128). Reads mapping to each open reading frame were quantified using htseq-count (129). Counts were imported into R and filtered to mRNA transcripts. To prepare for differential expression, the edgeR function calcNormFactors was used with default parameters (130). The limma package was used for differential expression by building a model using the decrease in Brix from initial Brix values; the data were prepared for linear modeling with the voom function, and a linear model was built for each gene with the lmFit function (131). Any gene with an adjusted *P* value of <0.05 was considered significant. To combat batch effects from different library preparation techniques used for the 2017 and 2019 vintages, differential

expression was performed separately on counts from each vintage. The union of expressed and repressed genes was taken between vintages to generate the final set of differentially expressed genes. Expressed and repressed genes were visualized using proteomaps (132), and the intersection of differentially expressed genes between vintages was visualized using the R package ComplexUpset (<https://github.com/krassowski/complex-upset>). Gene set enrichment analysis was performed for genes that were expressed and repressed in both vintages against the Gene Ontology (ont = ALL) and Kyoto Encyclopedia of Genes and Genomes (organism = sce) databases using the R package clusterProfiler (133).

Detection of *Lactobacillus kunkeei* in RNA-seq reads. 3'-Tag-seq sequences the tail end of transcripts that contain poly(A) tails. The majority of transcripts with poly(A) tails are eukaryotic in origin but, given that bacteria perform polyadenylation as a degradation signal (134), a very small subset of transcripts may originate from bacteria. *Lactobacillus kunkeei* was identified in RNA-seq reads using sourmash gather (135, 136). Using all *L. kunkeei* genomes available in GenBank (6 August 2019), sourmash signatures were generated for each using a k-mer size of 31 and a scaled value of 100. Sourmash index was then used to generate a database of *L. kunkeei* genomes, and this database was queried using signatures for each RNA-seq sample. To validate findings from sourmash gather, BWA-MEM was used with default parameters to map a subset of samples against the best-matching *L. kunkeei* genome (137).

Culturing *Saccharomyces cerevisiae* for [GAR⁺] prion detection. To ascertain whether the [GAR⁺] prion state was detectable in wine fermentations, yeast were cultured for the prion as performed previously (121). Yeast peptone-based medium containing the designated carbon source was used, such as YPD (1% yeast extract, 2% peptone, 2% agar, 2% glucose), YPG (1% yeast extract, 2% peptone, 2% agar, 2% glycerol), or GGM [1% yeast extract, 2% peptone, 2% agar, 2% glycerol, 0.05% D-(+)-glucosamine hydrochloride]. Yeast from fermentations were inoculated into each well of a 96-well plate containing 200 μ l liquid YPD plus 34 g/ml chloramphenicol, and then yeast were grown at 30°C for 48 h. Yeast were pinned to YPG or GGM plates and grown at 30°C for 4 days.

AVA map construction. The AVA map featured in Fig. 1 was constructed from the University of California, Davis, library AVA project (<https://github.com/UCDavisLibrary/ava>).

Data availability. RNA-seq data are available in the Sequence Read Archive (SRA) under accession number PRJNA680606. All analysis code is available at github.com/montpetitlab/Reiter_et_al_2020_GEacrossBrix.

SUPPLEMENTAL MATERIAL

Supplemental material is available online only.

SUPPLEMENTAL FILE 1, PDF file, 0.1 MB.

SUPPLEMENTAL FILE 2, XLSX file, 1.1 MB.

ACKNOWLEDGMENTS

We thank all past and current members of the Runnebaum and Montpetit laboratories for their support of this work, as well as the students and staff of the University of California, Davis, Teaching and Research Winery.

T.R. was supported by the Harry Baccigaluppi Fellowship, Horace O. Lanza Scholarship, Louis R. Gomberg Fellowship, Margrit Mondavi Fellowship, Haskell F. Norman Wine and Food Fellowship, Chaîne des Rôtisseurs Scholarship, and Carpenter Memorial Fellowship. We recognize major support from Jackson Family Wines, in addition to support from Lallemand Inc.

REFERENCES

1. Querol A, Fernández-Espinar MT, Li del Olmo M, Barrio E. 2003. Adaptive evolution of wine yeast. *Int J Food Microbiol* 86:3–10. [https://doi.org/10.1016/S0168-1605\(03\)00244-7](https://doi.org/10.1016/S0168-1605(03)00244-7).
2. Rossignol T, Dulau L, Julien A, Blondin B. 2003. Genome-wide monitoring of wine yeast gene expression during alcoholic fermentation. *Yeast* 20:1369–1385. <https://doi.org/10.1002/yea.1046>.
3. Marks VD, Ho Sui SJ, Erasmus D, Van Der Merwe GK, Brumm J, Wasserman WW, Bryan J, Van Vuuren HJ. 2008. Dynamics of the yeast transcriptome during wine fermentation reveals a novel fermentation stress response. *FEMS Yeast Res* 8:35–52. <https://doi.org/10.1111/j.1567-1364.2007.00338.x>.
4. Rossouw D, Jolly N, Jacobson D, Bauer FF. 2012. The effect of scale on gene expression: commercial versus laboratory wine fermentations. *Appl Microbiol Biotechnol* 93:1207–1219. <https://doi.org/10.1007/s00253-011-3564-0>.
5. Backhus LE, DeRisi J, Brown PO, Bisson LF. 2001. Functional genomic analysis of a commercial wine strain of *Saccharomyces cerevisiae* under differing nitrogen conditions. *FEMS Yeast Res* 1:111–125. <https://doi.org/10.1111/j.1567-1364.2001.tb00022.x>.
6. Mendes-Ferreira A, Del Olmo M, García-Martínez J, Jiménez-Martí E, Mendes-Faia A, Pérez-Ortín JE, Leao C. 2007. Transcriptional response of *Saccharomyces cerevisiae* to different nitrogen concentrations during alcoholic fermentation. *Appl Environ Microbiol* 73:3049–3060. <https://doi.org/10.1128/AEM.02754-06>.
7. Gasch AP, Spellman PT, Kao CM, Carmel-Harel O, Eisen MB, Storz G, Botstein D, Brown PO. 2000. Genomic expression programs in the response of yeast cells to environmental changes. *Mol Biol Cell* 11:4241–4257. <https://doi.org/10.1091/mbc.11.12.4241>.
8. Puig S, Pérez-Ortín JE. 2000. Stress response and expression patterns in wine fermentations of yeast genes induced at the diauxic shift. *Yeast* 16:139–148. [https://doi.org/10.1002/\(SICI\)1097-0061\(20000130\)16:2<139:AID-YEA512>3.0.CO;2-J](https://doi.org/10.1002/(SICI)1097-0061(20000130)16:2<139:AID-YEA512>3.0.CO;2-J).
9. Bisson LF. 1999. Stuck and sluggish fermentations. *Am J Enol Viticult* 50:107–119.
10. Gutiérrez A, Beltran G, Warringer J, Guillamón JM. 2013. Genetic basis of variations in nitrogen source utilization in four wine commercial yeast strains. *PLoS One* 8:e67166. <https://doi.org/10.1371/journal.pone.0067166>.
11. Patel S, Shibamoto T. 2002. Effect of different strains of *Saccharomyces cerevisiae* on production of volatiles in Napa Gamay wine and Petite Sirah wine. *J Agric Food Chem* 50:5649–5653. <https://doi.org/10.1021/jf020337f>.

12. Rossouw D, Naes T, Bauer FF. 2008. Linking gene regulation and the exo-metabolome: a comparative transcriptomics approach to identify genes that impact on the production of volatile aroma compounds in yeast. *BMC Genomics* 9:530. <https://doi.org/10.1186/1471-2164-9-530>.
13. Riou C, Nicaud J-M, Barre P, Gaillardin C. 1997. Stationary-phase gene expression in *Saccharomyces cerevisiae* during wine fermentation. *Yeast* 13:903–915. [https://doi.org/10.1002/\(SICI\)1097-0061\(199708\)13:10<903::AID-YEA145>3.0.CO;2-1](https://doi.org/10.1002/(SICI)1097-0061(199708)13:10<903::AID-YEA145>3.0.CO;2-1).
14. Rossouw D, Bauer FF. 2009. Comparing the transcriptomes of wine yeast strains: toward understanding the interaction between environment and transcriptome during fermentation. *Appl Microbiol Biotechnol* 84:937–954. <https://doi.org/10.1007/s00253-009-2204-4>.
15. Casalta E, Aguera E, Picou C, Rodríguez-Bencomo J-J, Salmon J-M, Sablayrolles J-M. 2010. A comparison of laboratory and pilot-scale fermentations in winemaking conditions. *Appl Microbiol Biotechnol* 87:1665–1673. <https://doi.org/10.1007/s00253-010-2652-x>.
16. Walker ME, Nguyen TD, Liccioli T, Schmid F, Kalatzis N, Sundstrom JF, Gardner JM, Jiranek V. 2014. Genome-wide identification of the fermentome; genes required for successful and timely completion of wine-like fermentation by *Saccharomyces cerevisiae*. *BMC Genomics* 15:52. <https://doi.org/10.1186/1471-2164-15-52>.
17. Cadière A, Aguera E, Caillé S, Ortiz-Julien A, Dequin S. 2012. Pilot-scale evaluation the enological traits of a novel, aromatic wine yeast strain obtained by adaptive evolution. *Food Microbiol* 32:332–337. <https://doi.org/10.1016/j.fm.2012.07.010>.
18. Vila I. 1998. Les levures aromatiques en vinification: évaluation de ce caractère par l'analyse sensorielle et l'analyse chimique: déterminisme biochimique des facteurs responsables. PhD thesis. Montpellier 2 University, Montpellier, France.
19. Du Toit W, Marais J, Pretorius I, Du Toit M. 2006. Oxygen in must and wine: a review. *S Afr J Enol Viticult* 27:76–94. <https://doi.org/10.21548/27-1-1610>.
20. Orellana M, Aceituno FF, Slater AW, Almonacid LI, Melo F, Agosin E. 2014. Metabolic and transcriptomic response of the wine yeast *Saccharomyces cerevisiae* strain EC1118 after an oxygen impulse under carbon-sufficient, nitrogen-limited fermentative conditions. *FEMS Yeast Res* 14:412–424. <https://doi.org/10.1111/1567-1364.12135>.
21. Varela C, Cárdenas J, Melo F, Agosin E. 2005. Quantitative analysis of wine yeast gene expression profiles under winemaking conditions. *Yeast* 22:369–383. <https://doi.org/10.1002/yea.1217>.
22. Roullier-Gall C, Boutegrabet L, Gougeon RD, Schmitt-Kopplin P. 2014. A grape and wine chemodiversity comparison of different appellations in Burgundy: vintage vs terroir effects. *Food Chem* 152:100–107. <https://doi.org/10.1016/j.foodchem.2013.11.056>.
23. Vilanova M, Rodríguez I, Canosa P, Otero I, Gamero E, Moreno D, Talaverano I, Valdés E. 2015. Variability in chemical composition of *Vitis vinifera* cv Mencia from different geographic areas and vintages in Ribeira Sacra (NW Spain). *Food Chem* 169:187–196. <https://doi.org/10.1016/j.foodchem.2014.08.015>.
24. Ramos MC, Martínez de Toda F. 2019. Variability of Tempranillo grape composition in the Rioja DOCa (Spain) related to soil and climatic characteristics. *J Sci Food Agric* 99:1153–1165. <https://doi.org/10.1002/jsfa.9283>.
25. Grainger C, Yeh A, Byer S, Hjelmeland A, Lima MM, Runnebaum RC. 2021. Vineyard site impact on the elemental composition of Pinot noir wines. *Food Chem* 334:127386. <https://doi.org/10.1016/j.foodchem.2020.127386>.
26. Bokulich NA, Thorngate JH, Richardson PM, Mills DA. 2014. Microbial biogeography of wine grapes is conditioned by cultivar, vintage, and climate. *Proc Natl Acad Sci U S A* 111:E139–E148. <https://doi.org/10.1073/pnas.1317377110>.
27. Bokulich NA, Collins TS, Masarweh C, Allen G, Heymann H, Ebeler SE, Mills DA. 2016. Associations among wine grape microbiome, metabolome, and fermentation behavior suggest microbial contribution to regional wine characteristics. *mBio* 7:e00631-16. <https://doi.org/10.1128/mBio.00631-16>.
28. David V, Terrat S, Herzine K, Claisse O, Rousseaux S, Tourdot-Maréchal R, Masneuf-Pomarede I, Ranjard L, Alexandre H. 2014. High-throughput sequencing of amplicons for monitoring yeast biodiversity in must and during alcoholic fermentation. *J Ind Microbiol Biotechnol* 41:811–821. <https://doi.org/10.1007/s10295-014-1427-2>.
29. Pinto C, Pinho D, Cardoso R, Custódio V, Fernandes J, Sousa S, Pinheiro M, Egas C, Gomes AC. 2015. Wine fermentation microbiome: a landscape from different Portuguese wine appellations. *Front Microbiol* 6:905. <https://doi.org/10.3389/fmicb.2015.00905>.
30. Wang C, García-Fernández D, Mas A, Esteve-Zarzoso B. 2015. Fungal diversity in grape must and wine fermentation assessed by massive sequencing, quantitative PCR and DGGE. *Front Microbiol* 6:1156. <https://doi.org/10.3389/fmicb.2015.01156>.
31. Garofalo C, Russo P, Beneduce L, Massa S, Spano G, Capozzi V. 2016. Non-*Saccharomyces* biodiversity in wine and the “microbial terroir”: a survey on Nero di Troia wine from the Apulian region, Italy. *Ann Microbiol* 66:143–150. <https://doi.org/10.1007/s13213-015-1090-5>.
32. Mezzasalma V, Sandionigi A, Bruni I, Bruno A, Lovicu G, Casiraghi M, Labra M. 2017. Grape microbiome as a reliable and persistent signature of field origin and environmental conditions in Cannonau wine production. *PLoS One* 12:e0184615. <https://doi.org/10.1371/journal.pone.0184615>.
33. Mezzasalma V, Sandionigi A, Guzzetti L, Galimberti A, Grando MS, Tardaguila J, Labra M. 2018. Geographical and cultivar features differentiate grape microbiota in northern Italy and Spain vineyards. *Front Microbiol* 9:946. <https://doi.org/10.3389/fmicb.2018.00946>.
34. Singh P, Santoni S, This P, Péros J-P. 2018. Genotype-environment interaction shapes the microbial assemblage in grapevine’s phyllosphere and carposphere: an NGS approach. *Microorganisms* 6:96. <https://doi.org/10.3390/microorganisms6040096>.
35. Liu D, Chen Q, Zhang P, Chen D, Howell KS. 2020. The fungal microbiome is an important component of vineyard ecosystems and correlates with regional distinctiveness of wine. *mSphere* 5:e00534-20. <https://doi.org/10.1128/mSphere.00534-20>.
36. Jolly N, Augustyn O, Pretorius I. 2003. The occurrence of non-*Saccharomyces cerevisiae* yeast species over three vintages in four vineyards and grape musts from four production regions of the Western Cape, South Africa. *S Afr J Enol Viticult* 24:35–42. <https://doi.org/10.21548/24-2-2640>.
37. Ghosh S, Bagheri B, Morgan HH, Divol B, Setati ME. 2015. Assessment of wine microbial diversity using ARISA and cultivation-based methods. *Ann Microbiol* 65:1833–1840. <https://doi.org/10.1007/s13213-014-1021-x>.
38. Wang C, Esteve-Zarzoso B, Cocolin L, Mas A, Rantsiou K. 2015. Viable and culturable populations of *Saccharomyces cerevisiae*, *Hanseniaspora uvarum* and *Starmerella bacillaris* (synonym *Candida zemplinina*) during Barbera must fermentation. *Food Res Int* 78:195–200. <https://doi.org/10.1016/j.foodres.2015.10.014>.
39. Bagheri B, Bauer F, Setati M. 2016. The diversity and dynamics of indigenous yeast communities in grape must from vineyards employing different agronomic practices and their influence on wine fermentation. *S Afr J Enol Viticult* 36:243–251. <https://doi.org/10.21548/36-2-957>.
40. Bagheri B, Bauer FF, Cardinali G, Setati ME. 2020. Ecological interactions are a primary driver of population dynamics in wine yeast microbiota during fermentation. *Sci Rep* 10: 4911. <https://doi.org/10.1038/s41598-020-61690-z>.
41. Brou P, Taillandier P, Beaufort S, Brandam C. 2018. Mixed culture fermentation using *Torulaspora delbrueckii* and *Saccharomyces cerevisiae* with direct and indirect contact: impact of anaerobic growth factors. *Eur Food Res Technol* 244:1699–1710. <https://doi.org/10.1007/s00217-018-3095-3>.
42. Curiel JA, Morales P, Gonzalez R, Tronchoni J. 2017. Different non-*Saccharomyces* yeast species stimulate nutrient consumption in *S. cerevisiae* mixed cultures. *Front Microbiol* 8:2121. <https://doi.org/10.3389/fmicb.2017.02121>.
43. Alonso-del-Real J, Pérez-Torrado R, Querol A, Barrio E. 2019. Dominance of wine *Saccharomyces cerevisiae* strains over *S. kudriavzevii* in industrial fermentation competitions is related to an acceleration of nutrient uptake and utilization. *Environ Microbiol* 21:1627–1644. <https://doi.org/10.1111/1462-2920.14536>.
44. Tronchoni J, Curiel JA, Morales P, Torres-Pérez R, Gonzalez R. 2017. Early transcriptional response to biotic stress in mixed starter fermentations involving *Saccharomyces cerevisiae* and *Torulaspora delbrueckii*. *Int J Food Microbiol* 241:60–68. <https://doi.org/10.1016/j.jifoodmicro.2016.10.017>.
45. Shekawat K, Patterson H, Bauer FF, Setati ME. 2019. RNA-seq based transcriptional analysis of *Saccharomyces cerevisiae* and *Lachancea thermotolerans* in mixed-culture fermentations under anaerobic conditions. *BMC Genomics* 20:145. <https://doi.org/10.1186/s12864-019-5511-x>.
46. Conacher CG, Rossouw D, Bauer F. 2019. Peer pressure: evolutionary responses to biotic pressures in wine yeasts. *FEMS Yeast Res* 19:foz072. <https://doi.org/10.1093/femsyr/foz072>.
47. Bordet F, Joran A, Klein G, Roullier-Gall C, Alexandre H. 2020. Yeast-yeast interactions: mechanisms, methodologies and impact on composition. *Microorganisms* 8:600. <https://doi.org/10.3390/microorganisms8040600>.
48. Egli C, Edinger W, Mitrakul C, Henick-Kling T. 1998. Dynamics of indigenous and inoculated yeast populations and their effect on the sensory character of Riesling and Chardonnay wines. *J Appl Microbiol* 85:779–789. <https://doi.org/10.1046/j.1365-2672.1998.00521.x>.

49. Bartowsky EJ. 2009. Bacterial spoilage of wine and approaches to minimize it. *Lett Appl Microbiol* 48:149–156. <https://doi.org/10.1111/j.1472-765X.2008.02505.x>.
50. Cantu A, Lafontaine S, Frias I, Sokolowsky M, Yeh A, Lestringant P, Hjelmeland A, Byer S, Heymann H, Runnebaum RC. 2021. Investigating the impact of regionality on the sensorial and chemical aging characteristics of Pinot noir grown throughout the US West coast. *Food Chem* 337:127720. <https://doi.org/10.1016/j.foodchem.2020.127720>.
51. Bisson LF. 2019. Gene expression in yeasts during wine fermentation, p 165–209. In Romano P, Ciani M, Fleet GH (ed), *Yeasts in the production of wine*. Springer, New York, NY.
52. Rodicio R, Heinisch JJ. 2017. Carbohydrate metabolism in wine yeasts, p 189–213. In König H, Uden G, Fröhlich J (ed), *Biology of microorganisms on grapes, in must and in wine*, 2nd ed. Springer International Publishing, Cham, Switzerland.
53. Ozcan S, Johnston M. 1996. Two different repressors collaborate to restrict expression of the yeast glucose transporter genes *HXT2* and *HXT4* to low levels of glucose. *Mol Cell Biol* 16:5536–5545. <https://doi.org/10.1128/mcb.16.10.5536>.
54. Ogawa N, DeRisi J, Brown PO. 2000. New components of a system for phosphate accumulation and polyphosphate metabolism in *Saccharomyces cerevisiae* revealed by genomic expression analysis. *Mol Biol Cell* 11:4309–4321. <https://doi.org/10.1091/mbc.11.12.4309>.
55. Ljungdahl PO, Daignan-Fornier B. 2012. Regulation of amino acid, nucleotide, and phosphate metabolism in *Saccharomyces cerevisiae*. *Genetics* 190:885–929. <https://doi.org/10.1534/genetics.111.133306>.
56. Barbosa C, Mendes-Faia A, Lage P, Mira NP, Mendes-Ferreira A. 2015. Genomic expression program of *Saccharomyces cerevisiae* along a mixed-culture wine fermentation with *Hanseniaspora guilliermondii*. *Microb Cell Fact* 14:124. <https://doi.org/10.1186/s12934-015-0318-1>.
57. Kosel J, Cadež N, Schuller D, Carreto L, Franco-Duarte R, Raspor P. 2017. The influence of *Dekkera bruxellensis* on the transcriptome of *Saccharomyces cerevisiae* and on the aromatic profile of synthetic wine must. *FEMS Yeast Res* 17:fox018. <https://doi.org/10.1093/femsyr/fox018>.
58. Reiter T, Montpetit R, Byer S, Frias I, Leon E, Viano R, McLoughlin M, Halligan T, Hernandez D, Figueroa-Balderas R, Cantu D, Steenwerth K, Runnebaum R, Montpetit B. 2021. Transcriptomics provides a genetic signature of vineyard site and offers insight into vintage-independent inoculated fermentation outcomes. *mSystems* 6:e00033-21. <https://doi.org/10.1128/mSystems.00033-21>.
59. Devatine A, Chiciuc I, Mietton-Peuchot M. 2011. The protective role of dissolved carbon dioxide against wine oxidation: a simple and rational approach. *OENO One* 45:189–197. <https://doi.org/10.20870/oeno-one.2011.45.3.1494>.
60. Moenne MI, Saa P, Laurie VF, Pérez-Correa JR, Agosin E. 2014. Oxygen incorporation and dissolution during industrial-scale red wine fermentations. *Food Bioprocess Technol* 7:2627–2636. <https://doi.org/10.1007/s11947-014-1257-2>.
61. Abramova N, Sertil O, Mehta S, Lowry CV. 2001. Reciprocal regulation of anaerobic and aerobic cell wall mannoprotein gene expression in *Saccharomyces cerevisiae*. *J Bacteriol* 183:2881–2887. <https://doi.org/10.1128/JB.183.9.2881-2887.2001>.
62. Sertil O, Kapoor R, Cohen BD, Abramova N, Lowry CV. 2003. Synergistic repression of anaerobic genes by Mot3 and Rox1 in *Saccharomyces cerevisiae*. *Nucleic Acids Res* 31:5831–5837. <https://doi.org/10.1093/nar/gkg792>.
63. Kwast KE, Burke PV, Poyton RO. 1998. Oxygen sensing and the transcriptional regulation of oxygen-responsive genes in yeast. *J Exp Biol* 201:1177–1195.
64. Wöhl T, Klier H, Ammer H, Lottspeich F, Magdolen V. 1993. The *HYP2* gene of *Saccharomyces cerevisiae* is essential for aerobic growth: characterization of different isoforms of the hypusine-containing protein Hyp2p and analysis of gene disruption mutants. *Mol Gen Genet* 241:305–311. <https://doi.org/10.1007/BF00284682>.
65. Burke PV, Poyton RO. 1998. Structure/function of oxygen-regulated isoforms in cytochrome c oxidase. *J Exp Biol* 201:1163–1175.
66. Remize F, Cambon B, Barnavon L, Dequin S. 2003. Glycerol formation during wine fermentation is mainly linked to Gpd1p and is only partially controlled by the HOG pathway. *Yeast* 20:1243–1253. <https://doi.org/10.1002/yea.1041>.
67. Jung J-Y, Kim T-Y, Ng C-Y, Oh M-K. 2012. Characterization of GCY1 in *Saccharomyces cerevisiae* by metabolic profiling. *J Appl Microbiol* 113:1468–1478. <https://doi.org/10.1111/jam.12013>.
68. Kliewer WM. 1970. Free amino acids and other nitrogenous fractions in wine grapes. *J Food Sci* 35:17–21. <https://doi.org/10.1111/j.1365-2621.1970.tb12358.x>.
69. ter Schure EG, van Riel NAW, Verrips CT. 2000. The role of ammonia metabolism in nitrogen catabolite repression in *Saccharomyces cerevisiae*. *FEMS Microbiol Rev* 24:67–83. <https://doi.org/10.1111/j.1574-6976.2000.tb00533.x>.
70. Huang HL, Brandriss MC. 2000. The regulator of the yeast proline utilization pathway is differentially phosphorylated in response to the quality of the nitrogen source. *Mol Cell Biol* 20:892–899. <https://doi.org/10.1128/mcb.20.3.892-899.2000>.
71. Takagi H, Taguchi J, Kaino T. 2016. Proline accumulation protects *Saccharomyces cerevisiae* cells in stationary phase from ethanol stress by reducing reactive oxygen species levels. *Yeast* 33:355–363. <https://doi.org/10.1002/yea.3154>.
72. Rosenfeld E, Beauvoit B, Blondin B, Salmon J-M. 2003. Oxygen consumption by anaerobic *Saccharomyces cerevisiae* under enological conditions: effect on fermentation kinetics. *Appl Environ Microbiol* 69:113–121. <https://doi.org/10.1128/aem.69.1.113-121.2003>.
73. Tarko T, Duda-Chodak A, Sroka P, Siuta M. 2020. The impact of oxygen at various stages of vinification on the chemical composition and the antioxidant and sensory properties of white and red wines. *Int J Food Sci* 2020:7902974. <https://doi.org/10.1155/2020/7902974>.
74. O'Connor-Cox ESC, Lodolo EJ, Axcell BC. 1996. Mitochondrial relevance to yeast fermentative performance: a review. *J Inst Brewing* 102:19–25. <https://doi.org/10.1002/j.2050-0416.1996.tb00890.x>.
75. Kitagaki H, Takagi H. 2014. Mitochondrial metabolism and stress response of yeast: applications in fermentation technologies. *J Biosci Bioeng* 117:383–393. <https://doi.org/10.1016/j.jbiosc.2013.09.011>.
76. Zara G, van Vuuren HJ, Mannazzu I, Zara S, Budroni M. 2019. Transcriptomic response of *Saccharomyces cerevisiae* during fermentation under oleic acid and ergosterol depletion. *Fermentation* 5:57. <https://doi.org/10.3390/fermentation5030057>.
77. Zeng J, Smith KE, Chong P. 1993. Effects of alcohol-induced lipid interdigitation on proton permeability in L- α -dipalmitoylphosphatidylcholine vesicles. *Biophys J* 65:1404–1414. [https://doi.org/10.1016/S0006-3495\(93\)81204-9](https://doi.org/10.1016/S0006-3495(93)81204-9).
78. Landolfo S, Politi H, Angelozzi D, Mannazzu I. 2008. ROS accumulation and oxidative damage to cell structures in *Saccharomyces cerevisiae* wine strains during fermentation of high-sugar-containing medium. *Biochim Biophys Acta* 1780:892–898. <https://doi.org/10.1016/j.bbagen.2008.03.008>.
79. Molenaar D, Van Berlo R, De Ridder D, Teusink B. 2009. Shifts in growth strategies reflect tradeoffs in cellular economics. *Mol Syst Biol* 5:323. <https://doi.org/10.1038/msb.2009.82>.
80. Wimpenny J. 1969. The effect of Eh on regulatory processes in facultative anaerobes. *Biotechnol Bioeng* 11:623–629. <https://doi.org/10.1002/bit.260110409>.
81. Somlo M, Fukuhara H. 1965. On the necessity of molecular oxygen for the synthesis of respiratory enzymes in yeast. *Biochem Biophys Res Commun* 19:587–591. [https://doi.org/10.1016/0006-291X\(65\)90379-7](https://doi.org/10.1016/0006-291X(65)90379-7).
82. Pammer M, Briza P, Ellinger A, Schuster T, Stucka R, Feldmann H, Breitenbach M. 1992. *DIT101* (*CSD2*, *CAL1*), a cell cycle-regulated yeast gene required for synthesis of chitin in cell walls and chitosan in spore walls. *Yeast* 8:1089–1099. <https://doi.org/10.1002/yea.320081211>.
83. Argüello-Miranda O, Liu Y, Wood NE, Kositangool P, Donic A. 2018. Integration of multiple metabolic signals determines cell fate prior to commitment. *Mol Cell* 71:733–744. <https://doi.org/10.1016/j.molcel.2018.07.041>.
84. Zhao H, Wang Q, Liu C, Shang Y, Wen F, Wang F, Liu W, Xiao W, Li W. 2018. A role for the respiratory chain in regulating meiosis initiation in *Saccharomyces cerevisiae*. *Genetics* 208:1181–1194. <https://doi.org/10.1534/genetics.118.300689>.
85. Sipiczki M. 2011. Diversity, variability and fast adaptive evolution of the wine yeast (*Saccharomyces cerevisiae*) genome: a review. *Ann Microbiol* 61:85–93. <https://doi.org/10.1007/s13213-010-0086-4>.
86. Reiner S, Micolod D, Zellnig G, Schneiter R. 2006. A genomewide screen reveals a role of mitochondria in anaerobic uptake of sterols in yeast. *Mol Biol Cell* 17:90–103. <https://doi.org/10.1091/mbc.e05-06-0515>.
87. Perez-Gallardo RV, Briones LS, Diaz-Pérez AL, Gutiérrez S, Rodríguez-Zavala JS, Campos-García J. 2013. Reactive oxygen species production induced by ethanol in *Saccharomyces cerevisiae* increases because of a dysfunctional mitochondrial iron-sulfur cluster assembly system. *FEMS Yeast Res* 13:804–819. <https://doi.org/10.1111/1567-1364.12090>.
88. Carmel-Harel O, Storz G. 2000. Roles of the glutathione- and thioredoxin-dependent reduction systems in the *Escherichia coli* and *Saccharomyces cerevisiae* responses to oxidative stress. *Annu Rev Microbiol* 54:439–461. <https://doi.org/10.1146/annurev.micro.54.1.439>.

89. Toledano MB, Delaunay-Moisan A, Outten CE, Igbaria A. 2013. Functions and cellular compartmentation of the thioredoxin and glutathione pathways in yeast. *Antioxid Redox Signal* 18:1699–1711. <https://doi.org/10.1089/ars.2012.5033>.
90. Bridi R, González A, Bordeu E, López-Alarcón C, Aspée A, Diethelm B, Lissi E, Parpinello GP, Versari A. 2015. Monitoring peroxides generation during model wine fermentation by FOX-1 assay. *Food Chem* 175:25–28. <https://doi.org/10.1016/j.foodchem.2014.11.126>.
91. Maslanka R, Zadrag-Tecza R, Kwolek-Mirek M. 2020. Linkage between carbon metabolism, redox status and cellular physiology in the yeast *Saccharomyces cerevisiae* devoid of *SOD1* or *SOD2* gene. *Genes* 11:780. <https://doi.org/10.3390/genes11070780>.
92. Matsufuji Y, Yamamoto K, Yamauchi K, Mitsunaga T, Hayakawa T, Nakagawa T. 2013. Novel physiological roles for glutathione in sequestering acetaldehyde to confer acetaldehyde tolerance in *Saccharomyces cerevisiae*. *Appl Microbiol Biotechnol* 97:297–303. <https://doi.org/10.1007/s00253-012-4147-4>.
93. Koc A, Mathews CK, Wheeler LJ, Gross MK, Merrill GF. 2006. Thioredoxin is required for deoxyribonucleotide pool maintenance during S phase. *J Biol Chem* 281:15058–15063. <https://doi.org/10.1074/jbc.M601968200>.
94. Muller E. 1991. Thioredoxin deficiency in yeast prolongs S phase and shortens the G₁ interval of the cell cycle. *J Biol Chem* 266:9194–9202. [https://doi.org/10.1016/S0021-9258\(18\)31570-9](https://doi.org/10.1016/S0021-9258(18)31570-9).
95. Shenton D, Grant CM. 2003. Protein S-thiolation targets glycolysis and protein synthesis in response to oxidative stress in the yeast *Saccharomyces cerevisiae*. *Biochem J* 374:513–519. <https://doi.org/10.1042/BJ20030414>.
96. Raiser M, Wamelink MM, Kowald A, Gerisch B, Heeren G, Struys EA, Klipp E, Jakobs C, Breitenbach M, Lehrach H, Krobitsch S. 2007. Dynamic rerouting of the carbohydrate flux is key to counteracting oxidative stress. *J Biol* 6:10. <https://doi.org/10.1186/jbiol61>.
97. Deponte M. 2017. The incomplete glutathione puzzle: just guessing at numbers and figures? *Antioxid Redox Signal* 27:1130–1161. <https://doi.org/10.1089/ars.2017.7123>.
98. Saint-Prix F, Bönquist L, Dequin S. 2004. Functional analysis of the *ALD* gene family of *Saccharomyces cerevisiae* during anaerobic growth on glucose: the NADP⁺-dependent Ald6p and Ald5p isoforms play a major role in acetate formation. *Microbiology (Reading)* 150:2209–2220. <https://doi.org/10.1099/mic.0.26999-0>.
99. Greetham D, Vickerstaff J, Shenton D, Perrone GG, Dawes IW, Grant CM. 2010. Thioredoxins function as deglutathionylase enzymes in the yeast *Saccharomyces cerevisiae*. *BMC Biochem* 11:3. <https://doi.org/10.1186/1471-2091-11-3>.
100. Collinson EJ, Wheeler GL, Garrido EO, Avery AM, Avery SV, Grant CM. 2002. The yeast glutaredoxins are active as glutathione peroxidases. *J Biol Chem* 277:16712–16717. <https://doi.org/10.1074/jbc.M111686200>.
101. Ohdate T, Kita K, Inoue Y. 2010. Kinetics and redox regulation of Gpx1, an atypical 2-Cys peroxidoredoxin, in *Saccharomyces cerevisiae*. *FEMS Yeast Res* 10:787–790. <https://doi.org/10.1111/j.1567-1364.2010.00651.x>.
102. Ohdate T, Izawa S, Kita K, Inoue Y. 2010. Regulatory mechanism for expression of GPX1 in response to glucose starvation and Ca²⁺ in *Saccharomyces cerevisiae*: involvement of Snf1 and Ras/cAMP pathway in Ca²⁺ signaling. *Genes Cells* 15:59–75. <https://doi.org/10.1111/j.1365-2443.2009.01365.x>.
103. Greetham D, Grant CM. 2009. Antioxidant activity of the yeast mitochondrial one-Cys peroxidoredoxin is dependent on thioredoxin reductase and glutathione in vivo. *Mol Cell Biol* 29:3229–3240. <https://doi.org/10.1128/MCB.01918-08>.
104. Martins AMT, Cordeiro CAA, Freire AMJP. 2001. In situ analysis of methylglyoxal metabolism in *Saccharomyces cerevisiae*. *FEBS Lett* 499:41–44. [https://doi.org/10.1016/S0014-5793\(01\)02519-4](https://doi.org/10.1016/S0014-5793(01)02519-4).
105. Dhaoui M, Auchère F, Blaiseau P-L, Lesuisse E, Landoulsi A, Camadro J-M, Haguenaer-Tsapis R, Belgareh-Touzé N. 2011. Gex1 is a yeast glutathione exchanger that interferes with pH and redox homeostasis. *Mol Biol Cell* 22:2054–2067. <https://doi.org/10.1091/mbc.E10-11-0906>.
106. Cordente AG, Capone DL, Curtin CD. 2015. Unravelling glutathione conjugate catabolism in *Saccharomyces cerevisiae*: the role of glutathione/dipeptide transporters and vacuolar function in the release of volatile sulfur compounds 3-mercaptohexan-1-ol and 4-mercapto-4-methylpentan-2-one. *Appl Microbiol Biotechnol* 99:9709–9722. <https://doi.org/10.1007/s00253-015-6833-5>.
107. Kumar A, Tikoo S, Maity S, Sengupta S, Sengupta S, Kaur A, Kumar Bachhawat A. 2012. Mammalian proapoptotic factor ChaC1 and its homologues function as γ -glutamyl cyclotransferases acting specifically on glutathione. *EMBO Rep* 13:1095–1101. <https://doi.org/10.1038/embor.2012.156>.
108. Chen X, Li S, Liu L. 2014. Engineering redox balance through cofactor systems. *Trends Biotechnol* 32:337–343. <https://doi.org/10.1016/j.tibtech.2014.04.003>.
109. Fariña L, Medina K, Urruty M, Boido E, Dellacassa E, Carrau F. 2012. Redox effect on volatile compound formation in wine during fermentation by *Saccharomyces cerevisiae*. *Food Chem* 134:933–939. <https://doi.org/10.1016/j.foodchem.2012.02.209>.
110. Bloem A, Sanchez I, Dequin S, Camarasa C. 2016. Metabolic impact of redox cofactor perturbations on the formation of aroma compounds in *Saccharomyces cerevisiae*. *Appl Environ Microbiol* 82:174–183. <https://doi.org/10.1128/AEM.02429-15>.
111. Xu X, Bao M, Niu C, Wang J, Liu C, Zheng F, Li Y, Li Q. 2019. Engineering the cytosolic NADH availability in lager yeast to improve the aroma profile of beer. *Biotechnol Lett* 41:363–369. <https://doi.org/10.1007/s10529-019-02653-x>.
112. Xu X, Song Y, Guo L, Cheng W, Niu C, Wang J, Liu C, Zheng F, Zhou Y, Li X, Mu Y, Li Q. 2020. Higher NADH availability of lager yeast increases the flavor stability of beer. *J Agric Food Chem* 68:584–590. <https://doi.org/10.1021/acs.jafc.9b05812>.
113. Causton HC, Ren B, Koh SS, Harbison CT, Kanin E, Jennings EG, Lee TI, True HL, Lander ES, Young RA. 2001. Remodeling of yeast genome expression in response to environmental changes. *Mol Biol Cell* 12:323–337. <https://doi.org/10.1091/mbc.12.2.323>.
114. Mbuyane LL, de Kock M, Bauer FF, Divol B. 2018. *Torulaspota delbrueckii* produces high levels of C₅ and C₆ polyols during wine fermentations. *FEMS Yeast Res* 18:foy084. <https://doi.org/10.1093/femsyr/foy084>.
115. Endo A, Futagawa-Endo Y, Dicks LM. 2009. Isolation and characterization of fructophilic lactic acid bacteria from fructose-rich niches. *Syst Appl Microbiol* 32:593–600. <https://doi.org/10.1016/j.syapm.2009.08.002>.
116. Du Toit M, Pretorius IS. 2000. Microbial spoilage and preservation of wine: using weapons for nature's own arsenal. *S Afr J Enol Viticult* 21:74–96. <https://doi.org/10.21548/21-1-3559>.
117. Quain DE, Boulton CA. 1987. Growth and metabolism of mannitol by strains of *Saccharomyces cerevisiae*. *J Gen Microbiol* 133:1675–1684. <https://doi.org/10.1099/00221287-133-7-1675>.
118. Jordan P, Choe J-Y, Boles E, Oreb M. 2016. Hxt13, Hxt15, Hxt16 and Hxt17 from *Saccharomyces cerevisiae* represent a novel type of polyol transporters. *Sci Rep* 6:23502. <https://doi.org/10.1038/srep23502>.
119. Ramakrishnan V, Walker GA, Fan Q, Ogawa M, Luo Y, Luong P, Joseph C, Bisson LF. 2016. Inter-kingdom modification of metabolic behavior: [GAR⁺] prion induction in *Saccharomyces cerevisiae* mediated by wine ecosystem bacteria. *Front Ecol Evol* 4:137. <https://doi.org/10.3389/fevo.2016.00137>.
120. Brown JC, Lindquist S. 2009. A heritable switch in carbon source utilization driven by an unusual yeast prion. *Genes Dev* 23:2320–2332. <https://doi.org/10.1101/gad.1839109>.
121. Jarosz DF, Brown JCS, Walker GA, Datta MS, Ung WL, Lancaster AK, Rotem A, Chang A, Newby GA, Weitz DA, Bisson LF, Lindquist S. 2014. Cross-kingdom chemical communication drives a heritable, mutually beneficial prion-based transformation of metabolism. *Cell* 158:1083–1093. <https://doi.org/10.1016/j.cell.2014.07.025>.
122. Nurgel C, Pickering G. 2005. Contribution of glycerol, ethanol and sugar to the perception of viscosity and density elicited by model white wines. *J Texture Stud* 36:303–323. <https://doi.org/10.1111/j.1745-4603.2005.00018.x>.
123. Ansell R, Granath K, Hohmann S, Thevelein JM, Adler L. 1997. The two isoenzymes for yeast NAD⁺-dependent glycerol 3-phosphate dehydrogenase encoded by GPD1 and GPD2 have distinct roles in osmoadaptation and redox regulation. *EMBO J* 16:2179–2187. <https://doi.org/10.1093/emboj/16.9.2179>.
124. Remize F, Barnavon L, Dequin S. 2001. Glycerol export and glycerol-3-phosphate dehydrogenase, but not glycerol phosphatase, are rate limiting for glycerol production in *Saccharomyces cerevisiae*. *Metab Eng* 3:301–312. <https://doi.org/10.1006/mben.2001.0197>.
125. Smith KD, Gordon PB, Rivetta A, Allen KE, Berbasova T, Slayman C, Strobel SA. 2015. Yeast Fex1p is a constitutively expressed fluoride channel with functional asymmetry of its two homologous domains. *J Biol Chem* 290:19874–19887. <https://doi.org/10.1074/jbc.M115.651976>.
126. Clayton MG. 1997. Fluoride inhibition of wine yeasts: a thesis presented in partial fulfilment of the requirements for the degree of Master of Science in Microbiology at Massey University. Master's thesis. Massey University, Palmerston North, New Zealand.
127. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 29:15–21. <https://doi.org/10.1093/bioinformatics/bts635>.

128. Smith T, Heger A, Sudbery I. 2017. UMI-tools: modeling sequencing errors in Unique Molecular Identifiers to improve quantification accuracy. *Genome Res* 27:491–499. <https://doi.org/10.1101/gr.209601.116>.
129. Anders S, Pyl PT, Huber W. 2015. HTSeq: a Python framework to work with high-throughput sequencing data. *Bioinformatics* 31:166–169. <https://doi.org/10.1093/bioinformatics/btu638>.
130. Robinson MD, McCarthy DJ, Smyth GK. 2010. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* 26:139–140. <https://doi.org/10.1093/bioinformatics/btp616>.
131. Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK. 2015. limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res* 43:e47. <https://doi.org/10.1093/nar/gkv007>.
132. Liebermeister W, Noor E, Flamholz A, Davidi D, Bernhardt J, Milo R. 2014. Visual account of protein investment in cellular functions. *Proc Natl Acad Sci U S A* 111:8488–8493. <https://doi.org/10.1073/pnas.1314810111>.
133. Yu G, Wang L-G, Han Y, He Q-Y. 2012. clusterProfiler: an R package for comparing biological themes among gene clusters. *Omic* 16:284–287. <https://doi.org/10.1089/omi.2011.0118>.
134. Rorbach J, Bobrowicz A, Pearce S, Minczuk M. 2014. Polyadenylation in bacteria and organelles. *Methods Mol Biol* 1125:211–227. https://doi.org/10.1007/978-1-62703-971-0_18.
135. Brown CT, Irber L. 2016. sourmash: a library for MinHash sketching of DNA. *J Open Source Software* 1:27. <https://doi.org/10.21105/joss.00027>.
136. Pierce NT, Irber L, Reiter T, Brooks P, Brown CT. 2019. Large-scale sequence comparisons with sourmash. *F1000Res* 8:1006. <https://doi.org/10.12688/f1000research.19675.1>.
137. Li H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *arXiv* 1303.3997.
138. Ciriacy M. 1975. Genetics of alcohol dehydrogenase in *Saccharomyces cerevisiae*. *Mol Gen Genet* 138:157–164. <https://doi.org/10.1007/BF02428119>.
139. Zuzuarregui A, Monteoliva L, Gil C, del Olmo M. I. 2006. Transcriptomic and proteomic approach for understanding the molecular basis of adaptation of *Saccharomyces cerevisiae* to wine fermentation. *Appl Environ Microbiol* 72:836–847. <https://doi.org/10.1128/AEM.72.1.836-847.2006>.
140. Curiel JA, Salvadó Z, Tronchoni J, Morales P, Rodrigues AJ, Quirós M, Gonzalez R. 2016. Identification of target genes to control acetate yield during aerobic fermentation with *Saccharomyces cerevisiae*. *Microb Cell Fact* 15:156. <https://doi.org/10.1186/s12934-016-0555-y>.
141. Piper P, Mahé Y, Thompson S, Pandjaitan R, Holyoak C, Egner R, Mühlbauer M, Coote P, Kuchler K. 1998. The Pdr12 ABC transporter is required for the development of weak organic acid resistance in yeast. *EMBO J* 17:4257–4265. <https://doi.org/10.1093/emboj/17.15.4257>.
142. Caro LHP, Smits GJ, van Egmond P, Chapman JW, Klis FM. 1998. Transcription of multiple cell wall protein-encoding genes in *Saccharomyces cerevisiae* is differentially regulated during the cell cycle. *FEMS Microbiol Lett* 161:345–349. <https://doi.org/10.1111/j.1574-6968.1998.tb12967.x>.
143. Crépin L, Truong NM, Bloem A, Sanchez I, Dequin S, Camarasa C. 2017. Management of multiple nitrogen sources during wine fermentation by *Saccharomyces cerevisiae*. *Appl Environ Microbiol* 83:e02617-16. <https://doi.org/10.1128/AEM.02617-16>.
144. Hazelwood LA, Daran J-M, Van Maris AJ, Pronk JT, Dickinson JR. 2008. The Ehrlich pathway for fusel alcohol production: a century of research on *Saccharomyces cerevisiae* metabolism. *Appl Environ Microbiol* 74:2259–2266. <https://doi.org/10.1128/AEM.02625-07>.
145. Cebollero E, Reggiori F. 2009. Regulation of autophagy in yeast *Saccharomyces cerevisiae*. *Biochim Biophys Acta* 1793:1413–1421. <https://doi.org/10.1016/j.bbamcr.2009.01.008>.
146. Alexandre H, Ansanay-Galeote V, Dequin S, Blondin B. 2001. Global gene expression during short-term ethanol stress in *Saccharomyces cerevisiae*. *FEBS Lett* 498:98–103. [https://doi.org/10.1016/S0014-5793\(01\)02503-0](https://doi.org/10.1016/S0014-5793(01)02503-0).
147. Lillie SH, Pringle JR. 1980. Reserve carbohydrate metabolism in *Saccharomyces cerevisiae*: responses to nutrient limitation. *J Bacteriol* 143:1384–1394. <https://doi.org/10.1128/JB.143.3.1384-1394.1980>.
148. François J, Parrou JL. 2001. Reserve carbohydrates metabolism in the yeast *Saccharomyces cerevisiae*. *FEMS Microbiol Rev* 25:125–145. <https://doi.org/10.1111/j.1574-6976.2001.tb00574.x>.
149. Singer MA, Lindquist S. 1998. Multiple effects of trehalose on protein folding in vitro and in vivo. *Mol Cell* 1:639–648. [https://doi.org/10.1016/S1097-2765\(00\)80064-7](https://doi.org/10.1016/S1097-2765(00)80064-7).
150. Parrou JL, Teste M-A, Francois J. 1997. Effects of various types of stress on the metabolism of reserve carbohydrates in *Saccharomyces cerevisiae*: genetic evidence for a stress-induced recycling of glycogen and trehalose. *Microbiology* 143:1891–1900. <https://doi.org/10.1099/00221287-143-6-1891>.
151. Udom N, Chansongkrow P, Charoensawan V, Auesukaree C. 2019. Coordination of the cell wall integrity and high-osmolarity glycerol pathways in response to ethanol stress in *Saccharomyces cerevisiae*. *Appl Environ Microbiol* 85:e00551-19. <https://doi.org/10.1128/AEM.00551-19>.