

Lawrence Berkeley National Laboratory

Recent Work

Title

OBSERVATION OF SHARP AND INTENSE STRUCTURES IN THE EXCITATION SPECTRUM OF NO₂

Permalink

<https://escholarship.org/uc/item/8vm2v1vh>

Authors

Robrish, P.
Rosen, H.

Publication Date

1975

0 0 0 0 4 2 0 2 5 7 2

Submitted to Journal of Chemical
Physics

LBL-3233
Preprint c.1

OBSERVATION OF SHARP AND INTENSE STRUCTURES IN
THE EXCITATION SPECTRUM OF NO₂

P. Robrish and H. Rosen

RECEIVED
LAWRENCE
RADIATION LABORATORY

January 1975


MAR 3 1975

LIBRARY AND
DOCUMENTS SECTION

Prepared for the U. S. Atomic Energy Commission
under Contract W-7405-ENG-48

For Reference

Not to be taken from this room



LBL-3233
c.1

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

0 0 0 0 4 2 0 2 5 7 3

LBL-3233

OBSERVATION OF SHARP AND INTENSE STRUCTURES
IN THE EXCITATION SPECTRUM OF NO₂*

P. Robrish and H. Rosen

Lawrence Berkeley Laboratory
Energy & Environment Division
University of California
Berkeley, CA 94720

January 1975

*Research supported by National Science Foundation from RANN Division
and U.S. Atomic Energy Commission

The visible absorption spectrum of NO_2 extends from about 4000-7000 \AA ¹ and seems to consist of numerous overlapping lines forming a quasi-continuum, with a few recognizable bands² superimposed upon it. Although several limited regions, usually near fixed frequency laser wavelengths, have been successfully analyzed using laser induced fluorescence³ and microwave-optical double resonance^{4,5} techniques, there is still considerable uncertainty in the interpretation of the overall absorption spectrum, including the sharp line structures found by Douglas and Huber.² There is also a continuing problem of reconciling the short excited state lifetimes deduced from the absorption spectrum⁶ with the long measured lifetimes,^{6,7} although several workers have reported evidence of short lived intermediate states.^{5,8} We are reporting on a technique which may further help to unravel the complex spectra of simple polyatomic molecules like NO_2 .

We have measured the excitation spectrum for the ν_2 mode of NO_2 using a narrow band ($.1\text{\AA}$) nitrogen-laser-pumped dye laser.⁹ This technique allows one not only to selectively excite the NO_2 molecule, but also to monitor a particular re-emission mode. We tuned the laser through most of the visible absorption spectrum of NO_2 (4200-6000 \AA) and observed the fluorescence through a spectrometer with a bandwidth of 25 \AA . As the laser was tuned, the spectrometer was advanced so as to keep its bandpass centered on the ν_2 mode located $\sim 750 \text{ cm}^{-1}$ from the laser frequency. Our excitation spectra show several narrow regions where there are large enhancements in the intensity of the ν_2

fluorescence mode. These narrow structures, which consist of sharp and intense lines, are shown in Figure 1. For a given dye, other structures were observed, but they were either of lower intensity or did not consist of resolved sharp lines. The positions of five of the structures in Figure 1, located at 18215, 19523, 21099, 21996, and 22989 cm^{-1} , are given to within 10 cm^{-1} by the following formula:

$$\nu = 18215 + 576.22N + 36.38N^2$$

with N respectively equal to 0, 2, 4, 5, 6. The frequencies corresponding to N=1 and N=3 occur at gaps between the dyes and we speculate that there are similar structures in these two regions with all seven forming a smooth progression. We are uncertain of the explanation for this progression, but if it is due to vibrational spacing in the upper state, then it has a peculiar anharmonicity since the spacing between members of the progression increases with frequency. It should be noted that, of the six observed structures, only the one near 4545 $\overset{\circ}{\text{A}}$ (21996 cm^{-1}) corresponds to any of the bands reported by Douglas and Huber². For this band, however, there is a one to one correspondence between the lines in the excitation spectrum and those in the absorption spectrum. A more detailed analysis of this band will be presented elsewhere.¹⁰

When the laser is narrowed to a bandwidth of .01 $\overset{\circ}{\text{A}}$ and tuned to the peak of an intense line within one of these structures, the ν_2 fluorescence cross-section is found to be about two orders of magnitude larger than those typically obtained using fixed frequency lasers.¹¹ These large cross-sections are particularly surprising given the fact that the absorption cross-sections we have measured at lines near 4545 $\overset{\circ}{\text{A}}$ are comparable to that of the background absorption.

These sharp and intense structures may be due to transitions to intermediate states which are not perturbed by high lying vibrational levels of the ground state^{2,12}. Such an explanation is consistent with the large enhancement we have measured. One test of this hypothesis would be to measure the fluorescence lifetime within these structures to see if they are in better agreement with the short lifetimes deduced from the absorption spectrum.

ACKNOWLEDGEMENT

We would like to thank Owen Chamberlain for his encouragement and helpful discussions throughout the course of this work.


REFERENCES

- 1) T.C. Hall Jr. and F.E. Blacet, J.Chem. Phys. 20, 1745 (1952)
- 2) A.E. Douglas and K.P. Huber, Can. J. Phys. 43, 74 (1965)
- 3) K. Abe, Fred Myers, T.K. McCubbin and S.R. Polo, J. Mol. Spect. 38
552 (1971)
Koichi Abe, J. Mol Spect. 48, 395 (1973)
K. Abe, Fred Myers, T.K. McCubbin, Jr. and S.R. Polo, J. Mol. Spect.
50, 413 (1974)
J.C.D. Brand, J.L. Hardwick, R.J. Pirkle and C.J. Seliskar, Can. J.
Phys. 51, 2184 (1973)
W. Demtröder, F. Paech and R. Schmiedl, Chem. Phys. Letts 26, 381
(1974)
C.G. Stevens, M.W. Swagel, R. Wallace and R.N. Zare, Chem. Phys.
Letts. 18, 465 (1973)
R.E. Smalley, B.L. Ramakrishna, D.H. Levy and L. Wharton, J. Chem.
Phys. 61, 4363 (1974)
- 4) Richard Solarz and Donald H. Levy, J. Chem. Phys. 58, 4026 (1973)
Richard Solarz, Donald H. Levy, Koichi Abe and R.F. Curl, J. Chem.
Phys. 60, 1158 (1974)
Takehiko Tanaka, Alan D. English, Robert W. Field, Donald A. Jennings
and David O. Harris, J. Chem. Phys. 59, 5217 (1973)
Takehiko Tanaka, Koichi Abe, and R.F. Curl, J. Mol. Spect. 49, 310 (1974)
- 5) Richard Solarz and Donald H. Levy, J. Chem. Phys. 60, 842 (1974)
- 6) Dan Neuberger and A.B.F. Duncan, J. Chem. Phys. 22, 1693 (1954)
- 7) G.H. Myers, D.M. Silver and F. Kaufman, J. Chem. Phys. 44, 718 (1966)
K. Sakurai and H.P. Broida, J. Chem. Phys. 50, 2404 (1969)
Philip B. Sackett and James T. Yardley, Chem. Phys. Letts. 6, 323 (1970)

- L.F. Keyser, S.Z. Levine and F. Kaufman, J. Chem. Phys. 54, 355 (1971)
- Philip B. Sackett and James T. Yardley, J. Chem. Phys. 57, 152 (1972)
- 8) P.B. Sackett and J.T. Yardley, Chem. Phys. Letts. 9, 612 (1971)
- 9) T.W. Hänsch, Applied Optics 11, 895 (1972)
- 10) P. Robrish and H. Rosen (to be published)
- 11) D.G. Fouche, A. Herzenberg and R.K. Chang, J. Appl. Phys. 43, 3846 (1972)
- C.M. Penny, W.W. Morey, R.L. St. Peters, S.D. Silverstein, M. Lapp and D.R. White, N.A.S.A. Report number NASA-CR-132363.
- 12) A.E. Douglas, J. Chem. Phys. 45, 1007 (1966)

FIGURE CAPTIONS

- 1) Intensity in the ν_2 mode as a function of laser wavelength near certain selected wavelength regions where sharp and intense structures were observed. The bandwidth of the laser was $\approx .1\text{\AA}$ and the spectrometer bandwidth was set to 25\AA . The absolute intensities in the different scans can not be compared directly, due to different laser powers and scale factors. The spectra were taken from a cell containing 1 mm of NO_2 .



XBL 7412-8438

FIGURE 1

LEGAL NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

TECHNICAL INFORMATION DIVISION
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720