UC San Diego

UC San Diego Previously Published Works

Title

Evolutionary relationship between 5+5 and 7+7 inverted repeat folds within the amino acid-polyamine-organocation superfamily.

Permalink

<https://escholarship.org/uc/item/8ws8m6j0>

Journal

Proteins, 82(2)

ISSN

0887-3585

Authors

Västermark, Åke
Saier, Milton H

Publication Date

2014-02-01

DOI

10.1002/prot.24401

Peer reviewed

Evolutionary relationship between 5+5 and 7+7 inverted repeat folds within the amino acid-polyamine-organocation superfamily

Åke Västermark and Milton H. Saier Jr*

Department of Molecular Biology, University of California at San Diego, La Jolla, California 92093-0116

ABSTRACT

Evidence has been presented that 5+5 TMS and 7+7 TMS inverted repeat fold transporters are members of a single superfamily named the Amino acid-Polyamine-organoCation (APC) superfamily. However, the evolutionary relationship between the 5+5 and the 7+7 topological types has not been established. We have identified a common fold, consisting of a spiny membrane helix/sheet, followed by a U-like structure and a V-like structure that is recurrent between domain duplicated units of 5+5 and 7+7 inverted repeat folds. This fold is found in the following protein structures: AdiC, ApcT, LeuT, Mhp1, BetP, CaiT, and SglT (all 5+5 TMS repeats), as well as UraA and SulP (7+7 TMS repeats). AdiC, LeuT and Mhp1 have two extra TMSs after the second duplicated domain, SglT has four extra C-terminal TMSs, and BetP has two extra TMSs before the first duplicated domain. UraA and SulP on the other hand have two extra TMSs at the N-terminus of each duplicated domain unit. These observations imply that multiple hairpin and domain duplication events occurred during the evolution of the APC superfamily. We suggest that the five TMS architecture was primordial and that families gained two TMSs on either side of this basic structure via dissimilar hairpin duplications either before or after intragenic duplication. Evidence for homology between TMSs 1–2 of AdiC and TMSs 1–2 and 3–4 of UraA suggests that the 7+7 topology arose via an internal duplication of the N-terminal hairpin loop within the five TMS repeat unit followed by duplication of the 7 TMS domain.

Proteins 2014; 82:336–346.
© 2013 Wiley Periodicals, Inc.

Key words: AdiC; UraA; amino acid-polyamine-organocation; APC; superfamily; hairpin; duplication.

INTRODUCTION

Hydrophilic molecules such as ions, sugars, and amino acids are transported across biological membranes via proteinaceous transport systems. Solute transporters are classified into channels, primary and secondary active transporters and group translocators. While primary active transporters use ATP or another primary source of energy such as light, electron flow or a metabolic conversion reaction to drive transport, secondary transporters use electrochemical gradients derived by the pumping activities of primary active transporters.^{1,2} Further, while primary active transporters are often multicomponent systems, secondary carriers most frequently consist of a single polypeptide chain.

Most secondary carriers exhibit pseudo-twofold symmetry, consisting of two repeat units of four to seven transmembrane α -helical segments (TMSs). When these proteins have four to seven TMSs, the functional unit is usually a homo- or heterodimer, but when the protein has 8–14 TMSs, the fundamental unit is usually

a monomer with two internal repeat units. These observations suggest that a basic characteristic of most secondary carriers is the presence of two symmetrical or pseudosymmetrical domains both required for function. These conclusions are consistent with high resolution data as well as biochemical mechanistic information.³

The amino acid-polyamine-organocation (APC) superfamily represents the second largest recognized superfamily of secondary carriers,⁴ being second in size only to the major facilitator superfamily (MFS).⁵ Protein members of the former superfamily usually have repeat units of five TMS while those of the latter always have repeat

Additional Supporting Information may be found in the online version of this article.

Grant sponsor: NIH; grant numbers: GM077402-05A1; GM 094610-01.

*Correspondence to: Professor Milton H. Saier, Jr., Department of Molecular Biology, University of California at San Diego, La Jolla, California 92093-0116.
E-mail: msaier@ucsd.edu

Received 3 June 2013; Revised 19 July 2013; Accepted 15 August 2013

Published online 28 August 2013 in Wiley Online Library (wileyonlinelibrary.com). DOI: 10.1002/prot.24401

units of six TMS. While the topologies differ, an evolutionary relationship between the APC superfamily and the MFS has been proposed.⁵

The APC superfamily was first defined and described by Jack *et al.* in 2000.⁶ At that time, it included 10 subfamilies as members of the APC family (TC# 2.A.3), with weaker relationships to the Amino Acid/Auxin Permease (AAP) family (TC# 2.A.18) and the Hydroxy/Aromatic Amino Acid Permease (HAAAP) family (TC# 2.A.42). These investigators reported that most proteins in the APC family display 12 TMSs, except the spore germination proteins which had 10 TMSs, and eukaryotic members of the CAT family which had 14 TMSs. Most AAP and HAAAP family members had 11 TMSs, having lost TMS12 at their C-terminal end relative to 12 TMS APC family members.

In 2012, Wong *et al.* expanded the APC superfamily to its present state, including 11 families, tabulated in Table I.⁴ The Alanine or Glycine:Cation Symporter (AGCS) family (TC# 2.A.25) and the Cation-Chloride Cotransporter (CCC) family (TC# 2.A.30) had been identified as members of the APC superfamily prior to the work of Wong *et al.*, 2012 (unpublished results). The new 5+5 topology additions were: The Betaine/Carnitine/Choline Transporter (BCCT) family (TC# 2.A.15), the Solute:Sodium Symporter (SSS) family (TC# 2.A.21), the Neurotransmitter:Sodium Symporter (NSS) family (TC# 2.A.22) and the Nucleobase:Cation Symporter-1 (NCS1) family (TC# 2.A.39). While the five previously recognized families mainly transported amino acids and amino acid derivatives, the new members, which proved to be more sequence divergent, displayed a more varied substrate repertoire. Two topologically distinct families were also included: the Sulfate Permease (SulP) family (TC# 2.A.53), and the Nucleobase:Cation Symporter-2 (NCS2) family (TC# 2.A.40), both of which are believed to have a 7+7 TMS repeat unit topology.⁷

Since the first publication of the APC superfamily,⁶ high resolution three-dimensional X-ray structures of several members of the currently recognized APC superfamily have been published. These include: ApCT,⁸ LeuT,⁹ Mhp1,¹⁰ BetP,¹¹ CaiT,¹² and SglT¹³ representing constituent families displaying a common five TMS repeat unit fold with different constellations of “extra” TMSs, and UraA with a less common seven TMS repeat. In this report, we focus on AdiC and UraA, to understand the differences in their topologies. AdiC is an arginine:agmatine antiporter in the APC family while UraA is a uracil permease of the NCS2 family. The 2.8 Å resolution crystal structure of UraA⁷ revealed the 7 TMS inverted repeat with a “spiny” secondary structural element located near the substrate translocation site in equivalent positions of each repeat unit. The 3 Å resolution crystal structure of AdiC¹⁴ revealed structural modifications in TMS6 in the 5+5 TMS inverted repeat structure.

We here report a shared feature, the spiny secondary structural element found in UraA and the symmetry related TMSs 1 and 6 in AdiC. We establish a nomenclature to facilitate the identification, description, and comprehension of a common fold shared by these structures and compared our approaches and results with those of others. AdiC has two extra TMSs at the C-terminus, whereas UraA has two extra TMSs at the N-terminus of each domain duplicated unit. Other members of the APC superfamily may have two extra N-terminal TMSs (BCCT; TC# 2.A.15) or four extra C-terminal TMSs (SSS; TC# 2.A.21). We postulate that multiple hairpin and domain duplication events were responsible for the variations of extra TMSs, and we show that the basic fold is recurrent between all members of the 5+5 and 7+7 TMS architectures.

METHODS

RMSD measurements

The RMSD values reported here were obtained using Chimera 1.7, using the RMSD map function, part of the “morph” function. The values were confirmed using SuperPose 1.0 (<http://wishart.biology.ualberta.ca/SuperPose/>). They always refer to the RMSD, in units of Å, of all α -carbons present in the comparison. In some cases, when such values are tabulated in PDB under the 3D similarity tab, RMSD values for domain comparisons using only a subset of well-aligned residues are given as well.

Topology prediction, modeling

For topology prediction, the best current methods, TOPCONS and SPOCTOPUS were used.¹⁵ When results from TOPCONS are presented, we present an aggregation of any positive predictions, rather than the consensus, so that if even only one method found a TMS, it is considered as a positive finding. Homology modeling was done using MODELLER 9.11, 2012/08/29, r8834, using default parameters. We used the realign with MAC option.

Secondary structure matching

SSM (secondary structure matching)¹⁶ was used as implemented in the software PDBe Fold v2.55 to compare the model of rat prestin with vSGLT (chain A).

HMM comparisons

HHsuite (hhsuite-2.0.16) was used for HMM:HMM comparisons, and HHMAKE (HHmake version 2.0.15) was used to train models representing the first and second halves of the APC and SulP families. We then used the -M 50 flag, for “FASTA columns with fewer than X%

Table 1
Proteins and Families Within the APC Superfamily Considered in This Report.

Family: TC#	Protein Name	Protein: TC#	Topology	Fold	SLC#	Abbreviation	SulP	SLC26	7+7	TOPCONS prediction for 2.A.42.1.1: 11 TMS	2.A.53.1.1 2.A.42.1.1	sulfate permease tyrosine permease	SulP TyrP	c	c	Resolution
2.A.53 2.A.42	Sulfate Permease Hydroxy/Aromatic Amino Acid Permease				b	HAAAP		b		2.A.42.1.1: 11 TMS				c	c	
2.A.40 2.A.39	Nucleobase: Cation Symporter-2 Nucleobase: Cation Symporter-1				b	NCS2 NCS1		b	7+7 5+5	2.A.40.1.1 2.A.39.3.6		uracil permease benzyl-hydantoin: cation symporter-1	UraA Mhp1	3OE7 2JLN	2.78 Å 2.85 Å	
2.A.30	Cation-Chloride Cotransporter					CCC				prediction for 2.A.30.1.1: 13 TMS	2.A.30.1.1	NaCl/KCl symporter	NKCC2	c	c	
2.A.25	Alanine or Glycine: Cation Symporter				b	AGCS				prediction for 2.A.25.1.1: 11 TMS	2.A.25.1.1	Ala/Gly:Na ⁺ symporter	DagA	c	c	
2.A.22	Neuro-transmitter: Sodium Symporter					NSS			5+5	5+5+2	2.A.22.4.2	amino acid (leucine):2 Na ⁺ symporter	LeuT	2072	1.70 Å	
2.A.21 2.A.18	Solute-Sodium Symporter Amino Acid/Auxin Permease					SSS AAAP			5+5	5+5+2+2 prediction for 2.A.18.1.1: 12 TMS	2.A.21.3.2 2.A.18.1.1	galactose: Na ⁺ symporter Auxin:H ⁺ symporter	SgtT Aux-1	2XQ2, 3DH4 c	2.70 Å c	
2.A.15	Betaine/Carnitine/ Choline Transporter				b	BCCT			5+5	2+5+5	2.A.15.1.10, 2.A.15.2.1	glycine betaine transporter, carnitine:γ- butyrobetaine antiporter	BetP, CaiT	4AIN, 2WSW	3.10 Å A,2.29 Å	
2.A.3a	Amino Acid-Polyamine- Organocation					APC			5+5	5+5+2	2.A.3.2.5	arginine: agmatine antiporter	AdiC	3L1L	3 Å	
Family: TC#	Protein Name	Protein: TC#	Topology	Fold	SLC#	Abbreviation	SulP	SLC#						PDB#		Resolution

^aFor the APC family (2.A.3), the structures of GadC and ApcT are also available.

^bNo SLC number has been assigned for these families since they are not represented in humans.

^cNot applicable. An x-ray structure is not available.

gaps” that are match states, and HHsearch (2.0.15), to compare the domain halves of the APC and SulP families using a feature file from the TMHMM Server 2.0 and Jalview 2.8. For each comparison, we recorded the percentage probability of homology. Using the same programs, we trained HMMs on MAFFT v7.023b alignments of SulP and APC (as well as the other 5+5 families) and compared them against each other.

AlignMe

Using AlignMe¹⁷ (<http://www.bioinfo.mpg.de/AlignMe/>), LeuT-like folds were compared employing alignments that weigh in hydropathy profiles. Because the methodology is based on the Needleman-Wunch algorithm, there may be a bias towards aligning the ends of sequences if they have different lengths. We took AdiC (TMSs 1–5) and UraA (TMSs 1–7) and used AlignMe with the PST settings for distantly related proteins. The program generates secondary structure predictions using PSIPRED 3.2, PSSMs, a transmembrane topology using OCTUPUS and the alignment. This exercise was repeated for AdiC (TMSs 6–10) and UraA (TMSs 8–14).

InterCompare

InterCompare¹⁸ was used to compare alignments of the APC and NCS2 families, comparing aligned positions 300–400 (corresponding to TMSs 1–2) with 300–400 (corresponding to TMSs 3–4), respectively.

RESULTS

Relationship between 5+5 and 7+7 TMS inverted repeat folds

We downloaded the uracil permease (UraA; 3QE7) structure of the NCS2 family and the arginine/agmatine antiporter (AdiC; 3L1L) structure of the APC family from the Protein Databank (PDB). UraA, with 14 transmembrane α -helices (TMSs), can be subdivided into two clear halves (from residues #1–200 and 201–410). These halves are superimposable (global α -C RMSD 9.04 Å), showing that these two halves share the same fold, containing seven TMSs each, with no “extra” TMSs. The third TMS is discontinuous in both halves, appearing incomplete. In TOPCONS,¹⁵ for the first half, there is a clear prediction of seven TMSs. For the second half, a weak prediction for the 2nd TMS was obtained, and there appeared to be only six TMSs due to a fused prediction of TMSs 9 and 10.

The structure of AdiC can also be divided in two halves: #7–175 and 176–380 with a 5+5 TMS basic structure and two extra C-terminal TMSs (381–440), possibly due to duplication of a terminal hairpin structure. TMSs 1 and 6 are discontinuous and annotated as incomplete in the PDB file, even though they span the

membrane, physically similar to TMS3 in UraA. We observed a small helix located between TMSs 2 and 3, similar to a helix located between TMSs 4 and 5 in UraA. We introduced a simple nomenclature, where the symbol “I” represents an incomplete helix (or β sheet in the case of UraA¹⁹) that spans the membrane; the symbol “V” represents two helices at an angle to each other with a short connecting loop; and the symbol “U” represents two parallel TM helices separated by an extramembraneous α -helix, holding them apart. The I corresponds to the first half of the “bundle” motif, the U corresponds to second half of the “bundle” and the first half of the “hash,” and the V corresponds to the second half of the “hash” and the “arm” motif, described by Perez and Ziegler (2013).¹⁹ We also use the symbol V’ (V prime) to indicate a pair of N-terminal helices, with features similar to V but probably not derived from V. We also use a short dash to indicate space between two secondary structural elements that are not part of the same domain duplicated unit. Using this makeshift nomenclature, we could describe the domain duplicated structure of UraA as two consecutive units of V’IUUV: V’ - I - U - V || V’ - I - U - V. Similarly, AdiC would be I - U - V || I - U - V - V, having an extra V after the domain duplicated unit. These features are conserved in all of the proteins for which three-dimensional structures are available.

When we excised the first two TMSs of UraA, focusing on residues #65–200 (TMSs 3–7), and compared this with residues #7–175 of AdiC (TMSs 1–5), we recognized the same fold of I - U - V. The RMSD between 24 atom pairs was 0.93 Å, while the listed UraA-AdiC domain distance in PDB using the jFATCAT-rigid algorithm was 6.90 Å, using selected atoms. The 5th and corresponding 10th TMSs pointed in opposite directions relative to each other from the rest of the structure, but otherwise, they showed the same fold. What speaks for this interpretation is not only the overall RMSD, but also the matching up of irregularities, including firstly the spiny TMS1/ β 1-sheet, and secondly, the “U,” including the small non-membrane helix that separates the two parts of the “U.” There are no TMS irregularities that do not match up. As a control, we compared region I - U - V || V’ in UraA with the end of AdiC, I - U - V - V. UraA has a long membrane-parallel arm connecting residues #202 and 224. A comparison of #65–280 in UraA to the second half of AdiC (untruncated, with the two terminal extra TMSs present) resulted in a poor superposition (15.18 Å), rejecting this latter possibility.

AdiC, residues #7–175, could be compared with UraA, residues #65–200. This scored an RMSD of 9.46 Å. AdiC residues #176–380 and UraA residues #65–200, resulted in a similar matching, displaying an RMSD of 9.37 Å. On the other hand, the first two TMSs from AdiC (residues #19–67) and UraA (residues #5–63), a V-formed pair of helices, could not be superimposed; they pointed in different directions, casting doubt on the possibility

Figure 1

Locations of the domain duplicated unit of IUV in AdiC, LeuT, Mhp1, BetP, SglT, and UraA. This figure shows the parts of the structures that we have used to create the nomenclature of I, U, and V (corresponding to the first half of the bundle (I), the second half of the bundle and the first half of the hash (U), and the second half of the hash and the arm (V), respectively, where the terms hash, arm and bundle refer to the designations described in Ref. 19). In the second repeat unit, the I U V are inverted to illustrate the relative orientation of these two inverted repeats. The overall structure of AdiC, LeuT and Mhp1 is: I – U – V – I – U – V – V; UraA: V' – I – U – V – V' – I – U – V, and BetP: V' – I – U – V – I – U – V. vSGLT. (i) – I – U – V – I – U – V – V – V.

that TMSs 1 and 2 are equivalent between UraA and AdiC. However, UraA residues #113–200 could be superimposed on AdiC residues #68–170, that is, the last three TMSs in the seven and five TMS units including half of the “U” and all of the “V.” This is consistent with a domain duplication model. Furthermore, superimposing the “U” in AdiC (TMSs 2–3; residues #41–113) onto TMSs 4–7 in UraA (U – V; residues #40–153), resulted in matching between the U-like structures (i.e., TMS 4–5 in UraA).

The RMSD between the first and second halves of AdiC, removing the terminal two helices, resulted in an overall RMSD for α -carbons of 7.67 Å. We hypothesize that residues #65–200 of UraA and residues #7–175 of AdiC represent the domain duplicated unit of I – U – V in both structures. This comparison resulted in an overall α -carbon RMSD of 9.03 Å (the listed distance in PDB is 6.90 Å) for five TMSs, indicating a common fold. This possibility was confirmed by corresponding locations of TMS irregularities between the structures, including an incomplete helical prediction/ β segment for the first transmembrane secondary structural element, and the equivalent positioning of a small helix located between TMSs 2 and 3. The interpretation of these comparisons between UraA and AdiC are illustrated in Figures 1 and 2 and Supporting Information Figure S1.

Structures of other 5+5 architecture inverted repeat folds

To compare AdiC to other 5+5 fold members in the APC superfamily, members, we considered the outward-facing conformation of LeuT (2Q72) of the NSS family.²¹ Residue #230 was used as the arbitrary point subdividing the structure into the domain duplicated halves. The last two helices (the “extras”) were removed, taking the sequence for the second domain up to #430 (#230–

430). The basic I – U – V fold found in AdiC was present. TMS1 was divided into two halves (TMS1a and TMS1b), forming the “spiny” helix. However, the pairwise RMSD between the halves was ~ 15 Å. What affected the RMSD negatively was the presence of a small, non-TMS helix in a loop region between TMSs 3 and 4 which was only present in the first of the two domain duplicated units. In the second five TMS repeat, the connector between the two arms of the “U” was split into two small helices. Furthermore, the arm connecting the two domain duplicated units contained a small helix that was not found in AdiC. These features added up to a higher RMSD since these anomalies did not match between the two copies. As the length of the new helical sequence was smaller in the second duplicated domain, it scored better against residues #7–175 of AdiC (displaying an RMSD of 9.07 Å), comparable to the score obtained between UraA and AdiC.

The outward facing conformation of Mhp1 (2JLN) of the NCS1 family¹⁰ contains an intradomain linker peptide (residues #191–209). The structure is AdiC-like, starting with the I – U – V fold, followed by the peptide linker, a second set of I – U – V and a terminal V. To remove the two helices, we truncated the sequence at residue #385. A peptide tail at the N-terminus was removed as well, and hence domain #1 starts at residue 26. The listed distance of Mhp1 and UraA in PDB using the jFATCAT-rigid algorithm is 7.78 Å. The two units of the I – U – V fold produced a self-superimposition, scoring an RMSD of 6.34 Å, supporting the conclusion of the five TMS duplication.

The first half of Mhp1 superimposed on the first half of AdiC both contain an identical broken helix in TMS1, and U- and V-like structures of the same length and shape. For the second halves of both proteins superimposed on each other, two specific features are conserved, showing the close similarity: (1) the helical structure of

Figure 2

Self- and cross-superimposition of the domain duplicated unit I – U – V, found in both AdiC (3L1L) and UraA (3QE7). **A:** AdiC onto itself (the two halves); **B:** UraA onto itself (the two halves); **C:** UraA 65–200 onto AdiC 7–175; **D:** UraA 65–200 onto AdiC 176–380. The color scale for the five TMSs (using TMSs 3–7 in the UraA numbering) is: orange, yellow, green, forest green, cyan. UraA contains TMSs 1–2 (the initial V not included here), the third helix (the “I,” containing a small beta sheet), then TMSs 4 and 5 interspersed by a small extramembrane helix (the “U”), then TMSs 6–7 (the “V”), and finally the long arm to the second copy of the domain (not included here). UraA can be cut in half giving #1–215, and #216–410. In the model of AdiC, instead of such an arm, a cut in the structure (dashed lines can be seen). In AdiC, the first helices are orange-colored, and the spiny TMS contains helix parts, not sheets. In the 7+7 fold (UraA), the initial extra helices are close to the substrate binding capillary. The superimposition scores were: AdiC 7–175 onto AdiC 176–380 displayed RMSD of 7.67 Å; 1–200 onto 201–410 in UraA (3QE7) displayed RMSD of 9.04 Å. The cross-comparisons (C, D) gave: C: superimposition of UraA 65–200 onto AdiC 7–175 displayed an RMSD of 9.03 Å; D: superimposition of UraA #65–200 and AdiC #176–380 displayed an RMSD of 9.37 Å.

the second leg of the U is briefly interrupted in the equivalent position, and (2) the V-like structures have a tiny insert between the arms, making it slightly U-like (see Table II for details).

For the intermediate conformation of BetP (4AIN) of the BCCT family,^{22–24} we used chain A. Sequences #57–273 and 274–546 were used, removing helical segments at the beginning and end of the full sequence which were outside the membrane. The sequence was truncated at position #490, assuming (based on the AdiC configuration) that the last two TMSs were the additional ones. However, we noted that in both domain duplicated units, the first helices were not “spiny” as expected from AdiC, that the U was unexpectedly wide, that its second leg appeared interrupted, and that the supposed V was U-like. These halves were superimposable on each other, giving an RMSD of 9.88 Å, but they were not superimposable on the two halves of AdiC. Assuming instead that the N-terminal V was the irregularity and removing it, in that case starting at position #130 and proceeding to position #272 (past the U), and then adding the V up to position #324, the RMSD was 7.85 Å for the self-comparison with these portions. Beginning at position #324, the structure resembled the intra-domain arm seen in AdiC. Consequently, residues #130–324 (I – U – V) were chosen as were #360–546. We also compared residues #130–324 (the I – U – V fold) against residues #7–

175 of AdiC, and obtained an RMSD of 9.16 Å. The same structures have a listed RMSD of 3.96 in PDB using selected atoms. Thus, BetP can be described, using our nomenclature, as: V' – I – U – V – I – U – V. The fact that the two halves of BetP are more similar to each other than they are to AdiC suggests that duplication events occurred multiple times during the evolution of the superfamily (see Discussion).

In CaiT (2WSW) of the BCCT family, which is very similar to BetP, the intra-domain connector was identified, and the structure was divided into two halves, #1–280 and 281–500. For the first half, the elements V' – I – U – V were clearly visible. We removed #280–300, the domain linker, from the second half. Otherwise, the second half contained only the I – U – V fold. Like BetP, CaiT had two extra TMSs at its N-terminus. We removed residues #1–85 (the initial V) to get comparable structures and determined similarity for the comparison of residues #80–280 and 300–500, displaying an RMSD of 8.82 Å. In addition, CaiT residues #80–280 superimposed on residues #7–175 of AdiC displayed an RMSD of 9.00 Å, confirming that these two domains had the same fold. In PDB, the listed CaiT-UraA distance using the jFATCAT-rigid algorithm is 7.47 Å.

For the inward-facing conformation of SglT (vSGLT; 2XQ2) of the SSS family, we first selected chain A. What was immediately striking was that there were missing

Table II
Pairwise RMSD Comparisons Considered in This Report

[illegible]

N/A = The secondary structure elements of the common fold did not superpose with the current residue selection.

backbone atoms in several places and that there were more extra TMSs than seen for other 5+5 TMS-structured homologues. The structure started with a membrane helix, which appeared to be external to the domain duplicated unit (possibly external to the membrane), followed by a nick in the determined backbone trace. The “spiny I” started at position #50; after that, there was a normal “U,” but also a cut in the backbone trace in the V. Then, there was an uninterrupted domain linker peptide, and finally it continued directly with the spiny “I” in the next domain duplicated unit, ending at residue position #240. The second domain’s U contained a break, but the U-like structure was clearly recognizable followed by a V. We compared residues #50–212 (prior to the domain linker) and residues #240–450. After #450, two more “V” structures were observed that were not part of the domain duplicated unit. In total, we found 1 extra N-terminal helix, the I – U – V - I – U – V repeat fold, and a C-terminal VV. The self-superimposition of the two halves displayed an RMSD of 9.17 Å. The listed UraA-SgIT distance in PDB using the jFATCAT-rigid algorithm is 7.83 Å.

Homology modeling of rat prestin

Rat prestin (SLC26A5; TC# 2.A.53.2.5), a member of the SulP family, which functions as an anion transporter in the human ear,²⁵ has 744 residues and contains a region (#80–510) that can be homology-modeled on UraA's 7+7 inverted repeat fold. SPOCTOPUS and TOPCONS predicted that it had only 13 TMSs, but a homology model using MODELLER, based on UraA as the template, generated a model of rat prestin which displayed an RMSD of 3.75 Å to UraA (henceforth referred to as our "SulP model"), where all 14 TMSs matched up.

This model was subsequently compared with a closely related representative of the 5+5 architecture, vSGLT, using chain A of PDB entry 3DH4, not 2XQ2 as previously discussed. It appeared that the additional helix, prior to the “spiny” structural element, but partly located in an undetermined region, was in fact part of the spiny α -helical segment. The alignment of the SSS family revealed that the two extra C-terminal Vs are not well conserved between SSS family members, most of which display only a single pair of terminal Vs.

The SSM results showed that TMSs 1–2 of rat prestin matches TMSs 1–2 of vSGLT, and that the same is true for UraA and AdiC. TMSs 3–4 in the 7+7 structure are unmatched. The final three TMSs in both the 5+5 and 7+7 topologies match. In the second domain duplicated unit, however, it is TMSs 3–4 of the 7+7 topology that correlate with TMSs 1–2 of the 5+5 topology, implying asymmetry between the domain duplicated units. These results are consistent with our suggestion that the first two hairpin structures in each of the seven TMS repeat units resulted from intragenic duplication of the N-

terminal hairpin structure in each 5+5 repeat and are homologous (see next sections and Discussion).

HMM comparisons

To resolve the differences between the superimposition- and SSM-based results which suggested that TMSs 1–2 in vSGLT were equivalent to TMSs 1–2 in SulP/rat prestin in the first domain duplicated unit, we used HMM:HMM comparisons of AdiC and UraA to determine if sequence similarity could be found to support the SSM-based model of the TMS correspondences between the homology model of rat prestin and vSGLT.

The first and second domain duplicated halves of the APC alignment scored a 0.8% chance of homology to each other, despite the common fold. However, the first and second domain duplicated halves of the SulP alignment scored a significant 41.7% chance of homology to each other, possibly due in part to the fact that SulP is a smaller and consequently less diverse family than APC.

For the comparison between UraA and AdiC, we obtained an even more significant 61% chance of homology between TMS2 (in the 5+5 topology) and TMS2 (in the 7+7 topology), the primary similarity between these divergent but related families. The matching segments were residues #46–81 in 2.A.40.3.2 of the SulP family, containing a single TMS, and the segment from #45–79 in 2.A.3.11.1 of the APC family, containing a single TMS.

One explanation is that a hairpin loop duplicated at the N-terminus, meaning that TMSs 1–2 and 3–4 are homologous. It is possible that the sequence similarity between TMSs 2 in AdiC and UraA (as detected by HMMs) is partly independent of the structures, but that it still biased the SSM result for the whole structure comparison.

Use of AlignMe

The AlignMe results are shown in Supporting Information Figure S2A. The first two TMSs of UraA in both domain duplicated units are unmatched, indicating that they could be considered the “extra” TMSs, contradicting the SSM results above.

InterCompare results

We compared TMS 1–2 of the APC family (TC# 2.A.3) with either TMS 1–2 or 3–4 of the NCS2 family (TC# 2.A.40). The best scoring comparison was TMSs 1–2 of 2.A.3.4.5 (Tpo5p) and TMSs 3–4 of 2.A.40.4.4 (YbbY), scoring 65 (far better than the comparison of TMSs 1–2 in the two families, which gave 11% identity and a maximal comparison score of 3 standard deviations, S.D.). The exact aligning regions were: (302 GSIV-MLGF 373) in APC and (330 GTIL-LIGF 395) in SulP. The GSAT¹⁸ binary comparison shown in Supporting

Information Figure S2B gave a 37.5% identity, and a Z-score 8 S.D., a far better score than obtained when TMSs 1–2 was compared with TMSs 1–2 in the two proteins.

InterCompare was used to compare TMSs 1–2 and 3–4 within NCS2, comparing aligned positions 180–300 and 300–400. The best scoring comparison was between 2.A.40.7.6 (180–300) and 2.A.40.4.2 (300–400). We then used GSAT to make a binary comparison of these, giving a Z-score of 5 S.D. In summary, the trend we observed with the InterCompare scores, the %-identity in the binary comparisons, and the Z-scores, was that the highest ranked comparison is for TMSs 1–2 in the APC family and TMSs 3–4 in the NCS2 family.

Further analyses with InterCompare revealed that TMSs 9–10 (residues 1120–1200) and TMSs 11–12 (residues 1400–1480) of the APC family scored ~55 in InterCompare, the best pair of 2.A.3.9.2 and 2.A.3.7.3 displaying a Z-score of 6 S.D., constituting the strongest evidence for a hairpin loop duplication in the 2.A.3 family. TMSs 9–10 and 11–12 (residues 510–600 and 605–670) of 2.A.21 scored up to 45 in InterCompare, TMSs 1–2 and 3–4 in 2.A.15 (residues 60–120 and 140–220) scored only up to 40 in InterCompare, while TMSs 8–9 and 10–11 (630–680 and 680–720) of 2.A.40 scored up to 39, unlikely to and in the cases we checked, not translating to high comparison scores.

DISCUSSION

Membrane proteins constitute <1% of the known high resolution 3D protein structures, but restraints imposed by the membrane compress the fold space for these proteins considerably.²⁶ Contact prediction methods²⁷ suggest that the fold space becomes small for membrane proteins having 8+ TMSs. This consideration restricts the potential conformations a transmembrane protein can assume and renders structural properties of transmembrane transport proteins more similar to each other than is observed for cytoplasmic proteins which lack these restrictions.

Lu *et al.* (2011) expressed uncertainty about whether the UraA fold conforms to the LeuT fold. Four subsequent relevant structural studies since the publication of the UraA structure have appeared, but the authors of these papers did not specifically investigate the fold similarities and differences between UraA and 5+5 TMS type APC family homologues.^{28–31} One of these studies²⁸ related UraA/XanQ to NCS1 (Mhp1) and other nucleobase transporters, including UPS/NBUT (TC# 2.A.7.19) and PUP/POP (TC# 2.A.7.14) which are members of the drug/metabolite transporter (DMT) superfamily,³² and AzgA of the NCS2 family (TC# 2.A.40.7). Of these, only the NCS2 family is represented in humans, and these proteins are not known to transport nucleobases. The shared substrate specificities of NCS1 and NCS2 family

members could reflect a common fold, but the suggestion of Frillingos (2012) that these proteins are related to the DMT superfamily has not been substantiated. Indeed, the 3D structures of known proteins in these two superfamilies are strikingly dissimilar (2ENK of SLC30A9).

A multitude of studies of LeuT-like transporters have supported the “rocking bundle” transport paradigm. The model is supported by the observation that different states of the structurally related transporters are similar. Two alternate states may not be required for transport.³ The fold has been described with the terms “bundle” (TMSs 1–2 and 6–7), “hash” (TMSs 3–4 and 8–9) and “arm” (TMSs 5 and 10).³⁴ Conserved sodium binding sites are important in some LeuT fold homologues.²⁰ However, whereas animal NCS2 transporters function by symporting sodium, the bacterial NCS2 transporters use protons instead of sodium.⁷ One paper suggests that the LeuT fold may have its evolutionary origins in DedA,¹⁷ but the evidence is weak.

Both LeuT and UraA share a domain duplicated inverted repeat structure, a feature shared with many transporters.³⁵ Discontinuous (spiny) secondary structural elements that are important for transport have been described in several different types of transporters, including LeuT-like fold porters¹⁹ and UraA.⁷ It is known that TMSs 1 and 6 in the LeuT fold are important for function,³⁶ and even though the fold is different, there could be an evolutionary relationship.³⁷

Wong *et al.* (2012) established that TMSs 1–8 in LeuT correspond to TMSs 1–8 in AdiC/ApcT, in agreement with the structural information presented in this report. Furthermore, this same report established that TMSs 4–7 in BetP of the BCCT family are equivalent to TMSs 2–5 in Mhp1 of the NCS1 family. This is explicable in light of the structural findings that BetP/CaiT differ from AdiC/ApcT, Mhp1 and LeuT, in having a V' - I - U - V - I - U - V structure, rather than an I - U - V - I - U - V - V structure. UraA has a V' - I - U - V - V' - I - U - V structure.

The data presented by Wong *et al.* suggested that the APC family is closely related to the NSS family, consistent with members of both families having the same TMS configuration and displaying a comparison score of 16.3 S.D. for their best pair of homologues. The NSS family, in turn, proved to be closely related to the SSS family, which, in turn, appeared to be fairly closely related to members of the 7+7 topology SulP family for which only preliminary structural information exists (<http://dx.doi.org/10.1016/j.bpj.2012.11.648>). The 5+5 topology proteins also proved to be related to NCS2 family members which have the 7+7 structure. A weaker relationship exists between NSS and NCS1, although both display the same arrangement of I - U - V - I - U - V - V (comparison score of 13.4 S.D.). vSGLT of the SSS family displays the I - U - V - I - U - V - V - V structure, while

Figure 3

APC superfamily overview, adapted from Wong *et al.* Homology was established using the GAP and GSAT programs based on the superfamily principle. Previously established APC superfamily proteins and their homologues were used to establish homology between the six other families. The TC Number for each family is listed under the family abbreviation in parentheses (see also Table I). GAP and GSAT comparison scores are expressed in S.D.

BetP of the BCCT family displays the V' - I - U - V - I - U - V arrangement, and may be more closely related to SSS family members than to NCS1 family members (Table I). Given that several members of the APC, NSS and NCS1 families display an I - U - V - I - U - V - V architecture, that at least two members of the BCCT family display a V' - I - U - V - I - U - V structure, and that at least one member of the NCS2 family displays a V' - I - U - V - V' - I - U - V architecture, an evolutionary model of TMS gains and losses can be proposed. In this model, the 5+5 architecture was the original pattern, and members of certain families have gained two TMSs on one or the other side of this basic structure (Fig. 3).

The first part of the analysis revealed that all of these homologues contain a common repeat unit, located inside the 7+7 architecture, comprising TMSs 3–7 and TMSs 10–14. Although other results complicated this picture, rat prestin of the SulP family was compared to its closest pseudo- 5+5 neighbor, vSGLT of the SSS family using secondary structure matching (SSM).¹⁶ The result was that TMSs 1–2 in the 7+7 topology matched TMSs 1–2 in the 5+5 topology, but for the second repeat unit, TMSs 10–11 (not TMSs 8–9) in the 7+7 topological architecture matched TMSs 6–7 in the 5+5 topology. For several reasons, our interpretation of this observation is that there may have been a hairpin loop duplication, giving rise to TMSs 1–2 and 3–4 in the 7+7 proteins. Results from AlignMe¹⁷ supported the conclusion that the first two TMSs in both copies of the

Figure 4

Summary of the topologies of all representative APC superfamily proteins and the proposed evolutionary pathways from the basic 5+5 architecture. The schematic two dimensional structures have been drawn using TMRPres2D, University of Athens.³⁹ The schema shows the paths of gains and losses of TMSs from AdiC. The original 5+5 repeat units are shown in red and purple, respectively; extra TMSs at the C-terminus are shown in green, and at the N-terminus or internal in yellow and gold, respectively.

domain duplicated unit in the 7+7 architecture are the “extra” TMSs.

To objectively identify regions of strong sequence similarity between AdiC and UraA, we resorted to alignment comparisons using HMM:HMM.³⁸ This approach identified a region of strong similarity between UraA and AdiC in the first pair of TMSs in both alignments, agreeing with the initial results obtained with SSM. Wong *et al.* established that the APC superfamily includes both SulP and UraA as well as all of the 5+5 transporters discussed here. To objectively test if a hairpin loop duplica-

tion had occurred, we used InterCompare, which revealed that the best comparison scores were obtained when TMSs 1 and 2 of 5+5 were compared with TMSs 3 and 4 of 7+7 (8 S.D.), although the value of 3 S.D. was obtained when TMSs 1 and 2 of 7+7 were compared with TMSs 1 and 2 of the 5+5 architecture. Although 8 S.D. is not sufficient to establish homology, this huge difference demonstrates a higher degree of similarity between TMSs 3 and 4 of 7+7 and TMSs 1 and 2 of 5+5. In fact, when TMSs 1 and 2 were compared with TMSs 3 and 4 of 7+7, a value of 5 S.D. was obtained, substantially better than 3 S.D. These observations can best be reconciled by assuming that TMSs 1 and 2 in the 7+7 topology arose by duplication of TMSs 1 and 2 in the 5+5 topology or TMSs 3–4 in the 7+7 topology (Fig. 4).

In summary, we believe that a five TMS unit as observed in APC family proteins duplicated to create a 10 TMS protein. Additional hairpin loop duplications occurred on either side of the 10 TMS unit, creating the topological diversity of that in AdiC, LeuT, Mhp1, and BetP (Table I, Fig. 1). In the case of SglT, a second hairpin loop duplication may have occurred at the C-terminus, creating 4 “extra” TMSs after the core unit. SulP and UraA arose from duplication of a 7 TMS (2+5 TMSs) unit. We propose that the N-terminal hairpin duplicated first, and that the resultant 7 TMS unit then duplicated, forming the 7+7 TMS topology. Compared to the possibly related Major Facilitator Superfamily (MFS), where most proteins have the 6+6 architecture, but some have a 6+2+6 architecture, there is similar evidence for an internal hairpin loop duplication. This intragenic duplication of 2 TMSs appears to have occurred multiple times in different families during the evolution of the MFS.⁵ We therefore propose that during the evolution of either the APC or the MFS superfamily, hairpin and entire 5, 6, or 7 TMS bundles duplicated multiple times. Further experimentation will be required to fully establish this possibility.

ACKNOWLEDGMENTS

We thank Roberto Battistutta, Mattia Sturlese, University of Padua, and Dominik Oliver, University of Marburg for valuable discussions and insightful considerations.

REFERENCES

- Forrest LR, Kramer R, Ziegler C. The structural basis of secondary active transport mechanisms. *Biochim Biophys Acta* 2011;1807:167–188.
- Saier MH, Jr. Families of transmembrane transporters selective for amino acids and their derivatives. *Microbiology* 2000;146 (Part 8): 1775–1795.
- Forrest LR, Rudnick G. The rocking bundle: a mechanism for ion-coupled solute flux by symmetrical transporters. *Physiology (Bethesda)* 2009;24:377–386.

4. Wong FH, Chen JS, Reddy V, Day JL, Shlykov MA, Wakabayashi ST, Saier MH, Jr. The amino acid-polyamine-organocation superfamily. *J Mol Microbiol Biotechnol* 2012;22:105–113.
5. Reddy VS, Shlykov MA, Castillo R, Sun EI, Saier MH, Jr. The major facilitator superfamily (MFS) revisited. *FEBS J* 2012;279:2022–2035.
6. Jack DL, Paulsen IT, Saier MH. The amino acid/polyamine/organocation (APC) superfamily of transporters specific for amino acids, polyamines and organocations. *Microbiology* 2000;146 (Part 8):1797–1814.
7. Lu F, Li S, Jiang Y, Jiang J, Fan H, Lu G, Deng D, Dang S, Zhang X, Wang J, Yan N. Structure and mechanism of the uracil transporter UraA. *Nature* 2011;472:243–246.
8. Shaffer PL, Goehring A, Shankaranarayanan A, Gouaux E. Structure and mechanism of a Na⁺-independent amino acid transporter. *Science* 2009;325:1010–1014.
9. Singh SK, Yamashita A, Gouaux E. Antidepressant binding site in a bacterial homologue of neurotransmitter transporters. *Nature* 2007;448:952–956.
10. Weyand S, Shimamura T, Yajima S, Suzuki S, Mirza O, Krusong K, Carpenter EP, Rutherford NG, Hadden JM, O'Reilly J, Ma P, Saidijam M, Patching SG, Hope RJ, Norbertczak HT, Roach PC, Iwata S, Henderson PJ, Cameron AD. Structure and molecular mechanism of a nucleobase-cation-symport-1 family transporter. *Science* 2008;322:709–713.
11. Perez C, Koshy C, Yildiz O, Ziegler C. Alternating-access mechanism in conformationally asymmetric trimers of the betaine transporter BetP. *Nature* 2012;490:126–130.
12. Schulze S, Koster S, Geldmacher U, Terwisscha van Scheltinga AC, Kuhlbrandt W. Structural basis of Na⁺-independent and cooperative substrate/product antiport in CaiT. *Nature* 2010;467:233–236.
13. Watanabe A, Choe S, Chaptal V, Rosenberg JM, Wright EM, Grabe M, Abramson J. The mechanism of sodium and substrate release from the binding pocket of vSGLT. *Nature* 2010;468:988–991.
14. Gao X, Lu F, Zhou L, Dang S, Sun L, Li X, Wang J, Shi Y. Structure and mechanism of an amino acid antiporter. *Science* 2009;324:1565–1568.
15. Bernsel A, Viklund H, Hennerdal A, Elofsson A. TOPCONS: consensus prediction of membrane protein topology. *Nucleic Acids Res* 2009;37(Web Server issue):W465–W468.
16. Krissinel E, Henrick K. Secondary-structure matching (SSM), a new tool for fast protein structure alignment in three dimensions. *Acta Crystallogr D Biol Crystallogr* 2004;60(Part 12, Part 1):2256–2268.
17. Khafizov K, Staritzbichler R, Stamm M, Forrest LR. A study of the evolution of inverted-topology repeats from LeuT-fold transporters using AlignMe. *Biochemistry* 2010;49:10702–10713.
18. Screpanti E, Hunte C. Discontinuous membrane helices in transport proteins and their correlation with function. *J Struct Biol* 2007;159:261–267.
19. Perez C, Ziegler C. Mechanistic aspects of sodium-binding sites in LeuT-like fold symporters. *Biol Chem* 2013;394:641–648.
20. Yamashita A, Singh SK, Kawate T, Jin Y, Gouaux E. Crystal structure of a bacterial homologue of Na⁺/Cl⁻-dependent neurotransmitter transporters. *Nature* 2005;437:215–223.
21. Perez C, Khafizov K, Forrest LR, Kramer R, Ziegler C. The role of trimerization in the osmoregulated betaine transporter BetP. *EMBO Rep* 2011;12:804–810.
22. Tsai CJ, Khafizov K, Hakulinen J, Forrest LR, Kramer R, Kuhlbrandt W, Ziegler C. Structural asymmetry in a trimeric Na⁺/betaine symporter, BetP, from *Corynebacterium glutamicum*. *J Mol Biol* 2011;407:368–381.
23. Ressel S, Terwisscha van Scheltinga AC, Vornrhein C, Ott V, Ziegler C. Molecular basis of transport and regulation in the Na⁺/betaine symporter BetP. *Nature* 2009;458:47–52.
24. Schanzler M, Fahlke C. Anion transport by the cochlear motor protein prestin. *J Physiol* 2012;590 (Part 2):259–272.
25. Reddy VS, Saier MH, Jr. BioV Suite—a collection of programs for the study of transport protein evolution. *FEBS J* 2012;279:2036–2046.
26. Ebejer JP, Hill JR, Kelm S, Shi J, Deane CM. Memoir: template-based structure prediction for membrane proteins. *Nucleic Acids Res* 2013;41 (Web Server issue):W379–W383.
27. Neumann S, Hartmann H, Martin-Galiano AJ, Fuchs A, Frishman D. Camps 2.0: exploring the sequence and structure space of prokaryotic, eukaryotic, and viral membrane proteins. *Proteins* 2012;80:839–857.
28. Frillingos S. Insights to the evolution of nucleobase-ascorbate transporters (NAT/NCS2 family) from the Cys-scanning analysis of xanthine permease XanQ. *Int J Biochem Mol Biol* 2012;3:250–272.
29. Kosti V, Lambrinidis G, Myrianthopoulos V, Dhallinas G, Mikros E. Identification of the substrate recognition and transport pathway in a eukaryotic member of the nucleobase-ascorbate transporter (NAT) family. *PLoS One* 2012;7:e41939.
30. Papakostas K, Frillingos S. Substrate selectivity of YgfU, a uric acid transporter from *Escherichia coli*. *J Biol Chem* 2012;287:15684–15695.
31. Karena E, Frillingos S. The role of transmembrane segment TM3 in the xanthine permease XanQ of *Escherichia coli*. *J Biol Chem* 2011;286:39595–39605.
32. Jack DL, Yang NM, Saier MH, Jr. The drug/metabolite transporter superfamily. *Eur J Biochem* 2001;268:3620–3639.
33. Li J, Shaikh SA, Enkavi G, Wen PC, Huang Z, Tajkhorshid E. Transient formation of water-conducting states in membrane transporters. *Proc Natl Acad Sci USA* 2013;110:7696–7701.
34. Jeschke G. A comparative study of structures and structural transitions of secondary transporters with the LeuT fold. *Eur Biophys J* 2013;42:181–197.
35. Schweikhard ES, Ziegler CM. Amino acid secondary transporters: toward a common transport mechanism. *Curr Top Membr* 2012;70:1–28.
36. Courville P, Urbankova E, Rensing C, Chaloupka R, Quick M, Cellier MF. Solute carrier 11 cation symport requires distinct residues in transmembrane helices 1 and 6. *J Biol Chem* 2008;283:9651–9658.
37. Grishin NV. Fold change in evolution of protein structures. *J Struct Biol* 2001;134:167–185.
38. Soding J. Protein homology detection by HMM-HMM comparison. *Bioinformatics* 2005;21:951–960.
39. Spyropoulos IC, Liakopoulos TD, Bagos PG, Hamodrakas SJ. TMRPres2D: high quality visual representation of transmembrane protein models. *Bioinformatics* 2004;20:3258–3260.