

Lawrence Berkeley National Laboratory

Recent Work

Title

ON THE BRANCHING RATIO OF THE n^+ MESON

Permalink

<https://escholarship.org/uc/item/8x02w15n>

Author

Smith, Frances M.

Publication Date

1951-01-04

UNCLASSIFIED

UCRL-

1074

UNIVERSITY OF CALIFORNIA - BERKELEY

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

RADIATION LABORATORY

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UNIVERSITY OF CALIFORNIA

Radiation Laboratory

Contract No. W-7405-eng-48

ON THE BRANCHING RATIO OF THE π^+ MESON

Frances M. Smith

January 4, 1951

Berkeley, California

ON THE BRANCHING RATIO OF THE π^+ MESON

Frances M. Smith

Radiation Laboratory, Department of Physics
University of California, Berkeley, California

January 4, 1951

Early experiments on π^+ mesons, using photographic emulsions as detectors, have seemed to show⁽¹⁾ that some of the π^+ mesons, upon stopping in matter, do not decay into μ^+ mesons.

These studies were concerned with mesons of fairly low energy so that the emulsion would be the only stopping material. Ilford C₂ and C₃ emulsions were used in order to facilitate the identification of the mesons. In many cases the processed emulsions showed an apparent non-uniformity in sensitivity and since the μ -meson track is rather tenuous in the region of the terminus of the π track, there is a chance of missing the decay. For a more intensive study of the decay scheme, a much more sensitive emulsion is required. Ilford G₅ and Eastman NTB₃ emulsions were chosen for the present study.

The apparatus used in this study consisted of a brass chamber for holding the plates and the target, as shown in the figure. The target was 0.036-inch carbon. This assembly was mounted on a probe and inserted into the vacuum chamber of the 184-inch cyclotron. The circulating beam of 345 Mev protons irradiated the target. Mesons emitted in the backward direction entered a channel cut into the brass holder. This channel was of such dimensions that π^+ mesons from the target with energies between 6 and 8 Mev only will enter the emulsion after a turn of 180°. No μ mesons from decay of the π stopping in

¹ J. Burfening, E. Gardner, and C.M.G. Lattes, Phys. Rev. 75, 383-4 (1949)

the target can get into the plate chamber. μ mesons from decay in flight of the π mesons could get into the emulsion only if they were emitted in a narrow cone in the forward or backward direction. These would not be confusable with π mesons from the target as their ranges in the emulsion would be too great or too small to have the correct energy.

The plates were studied using a high power microscope. Only those mesons which stopped in the emulsion at a distance greater than 10 microns from either surface of the undeveloped emulsion were counted.

Meson scattering from the channel walls gave a background fairly uniformly distributed with respect to range in the emulsion.

Analysis of results consisted of calculating the number of background μ mesons expected to fall in the main distribution. This number was subtracted from the number of mesons showing no decay found in the main distribution.

A preliminary estimate of the percentage of π mesons from the target which do not decay into μ mesons is: $R = 0.3 \pm 0.4\%$. This indicates that the branching ratio of the π^+ mesons is less than 1 percent and probably zero. A more complete account of this work will be published at a later date.

I wish to thank Dr. L. W. Alvarez for his many helpful suggestions in this study. I wish also to thank J. Vale and the cyclotron crew for their help in the use of the cyclotron and J. Willat for microscope work.

This work was performed under the auspices of the Atomic Energy Commission.

Information Division
1/8/51 cg




FIG. 1