UC Davis

UC Davis Previously Published Works

Title

Bridging tumor genomics to patient outcomes through an integrated patient-derived xenograft platform.

Permalink

<https://escholarship.org/uc/item/8x80h4g0>

Journal

Clinical lung cancer, 16(3)

ISSN

1525-7304

Authors

Gandara, David R
Mack, Philip C
Bult, Carol
et al.

Publication Date

2015-05-01

DOI

10.1016/j.clcc.2015.03.001

Peer reviewed

Bridging Tumor Genomics to Patient Outcomes Through an Integrated Patient-Derived Xenograft Platform

David R. Gandara,¹ Philip C. Mack,¹ Carol Bult,² Tianhong Li,¹
 Primo N. Lara, Jr,¹ Jonathan W. Riess,¹ Stephanie H. Astrow,³
 Regina Gandour-Edwards,¹ David T. Cooke,¹ Ken Y. Yoneda,¹
 Elizabeth H. Moore,¹ Chong-xian Pan,¹ Rebekah A. Burich,¹ Elizabeth A. David,¹
 James G. Keck,² Susan Airhart,² Neal Goodwin,² Ralph W. de Vere White,¹
 Edison T. Liu²

Abstract

New approaches to optimization of cancer drug development in the laboratory and the clinic will be required to fully achieve the goal of individualized, precision cancer therapy. Improved preclinical models that more closely reflect the now recognized genomic complexity of human cancers are needed. Here we describe a collaborative research project that integrates core resources of The Jackson Laboratory Basic Science Cancer Center with genomics and clinical research facilities at the UC Davis Comprehensive Cancer Center to establish a clinically and genomically annotated patient-derived xenograft (PDX) platform designed to enhance new drug development and strategies for targeted therapies. Advanced stage non-small-cell lung cancer (NSCLC) was selected for initial studies because of emergence of a number of “druggable” molecular targets, and recent recognition of substantial inter- and inpatient tumor heterogeneity. Additionally, clonal evolution after targeted therapy interventions make this tumor type ideal for investigation of this platform. Using the immunodeficient NOD *scid* gamma mouse, > 200 NSCLC tumor biopsies have been xenotransplanted. During the annotation process, patient tumors and subsequent PDXs are compared at multiple levels, including histomorphology, clinically applicable molecular biomarkers, global gene expression patterns, gene copy number variations, and DNA/chromosomal alterations. NSCLC PDXs are grouped into panels of interest according to oncogene subtype and/or histologic subtype. Multiregimen drug testing, paired with next-generation sequencing before and after therapy and timed tumor pharmacodynamics enables determination of efficacy, signaling pathway alterations, and mechanisms of sensitivity-resistance in individual models. This approach should facilitate derivation of new therapeutic strategies and the transition to individualized therapy.

Clinical Lung Cancer, Vol. 16, No. 3, 165-72 © 2015 Elsevier Inc. All rights reserved.

Keywords: Clinical trials, Lung cancer, Mouse model, Patient derived xenograft

Current address for Neal Goodwin: Champions Oncology, Hackensack, NJ.

¹University of California, Davis Comprehensive Cancer Center, Sacramento, CA

²The Jackson Laboratory, Bar Harbor, ME and Sacramento, CA

³Response Genetics, Inc, Los Angeles, CA

Submitted: Jan 29, 2015; Revised: Mar 10, 2015; Accepted: Mar 10, 2015; Epub: Mar 18, 2015

Address for correspondence: David R. Gandara, MD, Division of Hematology and Oncology, Suite 3016, UC Davis Comprehensive Cancer Center, 4501 X Street, Suite 3016, Sacramento, CA 95817

Fax: 916-734-7946; e-mail contact: david.gandara@ucdmc.ucdavis.edu

Introduction

Substantial advances have been made in understanding the molecular biology that drives carcinogenesis and cancer-associated proliferative and antiapoptotic signaling pathways. A wide variety of potentially “druggable” molecular targets for cancer therapy have emerged from these studies. Although a large number of molecular targeted agents have subsequently been tested, most of which showed substantial activity in available preclinical models, relatively few have been successful

Tumor Genomics to Patient Outcomes Through PDX Platform

Figure 1 Algorithm of Patient-Derived Xenograft (PDX) Creation. Candidate Patients (PTs) Were Identified and Provided Consent for Tumor Collection. After Biopsy, Pleural Effusion Fluid Collection, or Surgical Resection, Viable Portions of the Fresh Specimens Were Rapidly Transported to the Jackson Laboratories-West Facilities for Implantation Into the NSG Mouse Model. Concurrently, Remaining Portions of the PT Specimen Were Fixed and Subsequently Characterized and Molecularly Profiled. A PDX Model Was Considered “Established” After Demonstrating Growth in Passage 1, After Successful Implantation, Development and Transplantation From Passage 0. Histomorphologic Evaluation and Molecular Profiling of the PDX Model Was Conducted and Results Compared With the Contributing Human Specimen. When PDX Models Reach Passage 2, Cohorts Can Be Prepared for Growth Inhibition and Tumor Pharmacodynamic Studies

Abbreviations: CNV = copy number variation; FFPE = Formalin-fixed, paraffin-embedded; IHC = immunohistochemistry; NGS = next-generation sequencing; NSCLC = non-small-cell lung cancer; SNP = single-nucleotide polymorphism.

Figure 2 Panel of Non-Small-Cell Lung Cancer (NSCLC) Patient-Derived Xenograft (PDX) Models That Harbor Epidermal Growth Factor Receptor (EGFR)-Activating Mutations. Models Are Organized According to Patient Clinical Status at the Time of the Originating Biopsy. “Erlotinib Sensitive-Naïve” Models Were Acquired Before Treatment With Any EGFR Inhibitor; “Primary Resistance” Models Were Derived From Tumors That Showed No Clinical Benefit From Erlotinib; “Acquired Resistance” Models Were Derived From Tumors That Initially Responded to Erlotinib But Had Progressed at the Time of Biopsy. Models Were Also Subdivided According To Those With EGFR Gene Amplification (Upper Circle). Additional Information Is Provided in Each Box, Indicating Oncogenic Abnormalities of Interest

Abbreviations: AMP = amplification; mut = mutation.

Figure 3 Case Example of Clinical Annotation of Patient-Derived Xenograft (PDX) Models. A Large Amount of Anonymized, Annotated Patient Information Is Collected From Each Consenting Donor on PDX Implantation. This Information Includes Age, Sex, Race, Diagnosis, Course of Disease, and Previous Treatment, Known Mutation Status, Confounding Medical Issues, Family History of Cancer, Timing of the Establishment of PDX Relative to Treatment, and Therapeutic Outcomes. The Figure Shows Estimated Disease Burden of the Patient Who Contributed the LG0703 model. As Indicated, the PDX Was Established From a Biopsy at the Time of Disease Progression After a Successful Course of Erlotinib. The Patient Subsequently Went on to Receive the Combination of Afatinib With Cetuximab, With a Durable Response

Abbreviations: Chemo = chemotherapy; P0-3 = Passage 0-3.

in the clinic.¹⁻⁵ Thus, there remains an unmet need in drug development for new approaches toward testing patient-relevant treatment models in the laboratory, before or concurrent with initiation of clinical trials. The present preclinical evaluation process for new anticancer agents, based predominantly on human cancer cell lines or cell line-based xenograft models, has proven largely ineffective at predicting therapeutic potential in patients.^{6,7} Despite promising preclinical results, only 5% of cancer drugs under development are eventually approved for use. Most fail because of lack of efficacy in phase III clinical trials, as exemplified by recent trials for non-small-cell lung cancer (NSCLC).⁸ Chief among the many obstacles to development of effective anticancer regimens is the complexity of tumor signal transduction networks, with parallel pathways, crosstalk, compensatory feedback mechanisms, and extensive interactions between tumor cells and their microenvironment, which cannot be replicated in conventional preclinical models taken in isolation. Adding to these complexities are the inter- and inpatient tumor heterogeneity characteristic of human cancers and the innate adaptability of tumor mutator phenotypes, resulting in rapid development of resistance mechanisms. Taken together, these challenges necessitate new ways of thinking regarding the role of preclinical models in cancer drug development.⁹

Addressing an Unmet Need for Improving Drug Development Strategies in NSCLC

This need for improved preclinical platforms is particularly relevant to NSCLC, a worldwide health care epidemic for which most systemic therapies offer only modest benefit. For most patients with advanced NSCLC, the therapeutic decision-making process has remained largely empiric, based on factors such as historical treatment precedent, individual patient characteristics, and physician or patient preferences. Although the feasibility of selecting treatment for individual cancer patients based on tumor molecular profiles (personalized therapy or precision medicine) is already being explored in NSCLC, these efforts have been hampered considerably by tumor heterogeneity and the complexity of the underlying biologic pathways, and suboptimal preclinical models in which individualized therapeutic strategies can be tested.^{1,3,7} Further, improved strategies to identify and overcome mechanisms of de novo and acquired resistance to treatment are essential to increasing survival and cure rates.¹⁰⁻¹³

Already, transgenic or knock-in preclinical systems such as genetically engineered mouse models (GEMMs) have proven to be invaluable tools for understanding carcinogenesis, tumor biology, and target validation for therapeutics in ways not addressable with other modeling approaches.¹⁴⁻¹⁸ Nevertheless,

Tumor Genomics to Patient Outcomes Through PDX Platform

Table 1 Fidelity Between PT and PDX

	Mutational Status		
	EML4-ALK Fusion Transcript	K-RAS Mutation	EGFR Mutation
LG-0476			
PT	No translocation	Wild type	Wild type
PDX	No translocation	Wild type	Wild type
LG-0481			
PT	No translocation	Gly12Cys	Wild type
PDX	No translocation	Gly12Cys	Wild type
LG-0703			
PT	No translocation	Wild type	EGFR L858R
PDX	No translocation	Wild type	EGFR L858R
LG-1193			
PT	No translocation	Wild type	EGFR E19del
PDX	No translocation	Wild type	EGFR E19del
LG-0812			
PT	Fusion-positive	Wild type	Wild type
PDX	Fusion-positive	Wild type	Wild type
LG-0552			
PT	No translocation	Gly12Ala	Wild type
PDX	No translocation	Gly12Ala	Wild type
LG-0567			
PT	No translocation	Gly12Cys	Wild type
PDX	No translocation	Gly12Cys	Wild type

Abbreviations: PDX = patient-derived xenograft; PT = patient tumor.

GEMMs represent nonhuman cancers lacking the range of tumor heterogeneity and complexity of biologic pathways inherent to human cancers and present within patient-derived xenografts (PDXs). Although preliminary studies have already demonstrated that chemotherapeutic agents and biologic therapies can be administered and assessed in a systematic fashion using in vivo xenotransplantation PDX models, systematic development, clinical-genomic annotation, and drug testing paradigms have yet to be comprehensively evaluated.¹⁹⁻²³

Development of the Integrated University of California Davis Comprehensive Cancer Center-The Jackson Laboratory PDX Resource in NSCLC

Herein we describe development of a clinically and genomically annotated PDX resource in NSCLC and initial pilot projects that focus on epidermal growth factor receptor (EGFR) pathways and EGFR-directed therapies, to directly address and overcome these limitations. This NSCLC PDX research platform integrates core components of The Jackson Laboratory (JAX) National Cancer Institute (NCI)-designated Basic Science Cancer Center and the JAX In Vivo Pharmacology and Clinical Lab Services together with laboratory resources and the clinical research program of

the NCI-designated University of California Davis Comprehensive Cancer Center (UCD). The platform is designed to dynamically engage external institutions, investigators, additional organizations, and pharmaceutical partners as appropriate on project-specific bases. Altogether, > 25 medical centers and other partners have participated in the Primary Human Tumor Consortium. Since the launching of the program in 2009, > 1700 tumor specimens from individual cancer patients, including > 200 lung tumors, have been xenotransplanted into immunodeficient NOD *scid* gamma (NSG; NOD.Cg-Prkdc^{scid} Il2rg^{tm1Wjl}/SzJ) mice. Information and data about the PDX models are publicly accessible from the PDX portal at the Mouse Tumor Biology database (MTB; <http://tumor.informatics.jax.org/mtbwil/pdxSearch.do>).²⁴

The Jackson Laboratory NSG Model for PDX

The JAX NSG model is of particular interest for PDX development. Lacking functional B and T cells and natural killer cell activity, the NSG mouse is the most immunodeficient yet physiologically durable murine model available for consistent xenograftment of human primary tumors.²⁴ These attributes of the NSG mouse model facilitate engraftment of a wide variety of human cancers, with excellent correlation of histomorphological and molecular features between PDX tumors and the original human cancers as described herein. Ongoing studies at JAX are focusing on optimization of the engraftment algorithm, such as use of small biopsy samples acquired using core needle biopsy (minimum of 1 mm³ of tissue) and “transportability” of viable patient tumor specimens from other institutions. For example, engraftment rates appear similar between specimens acquired locally in Sacramento versus those express-shipped overnight from sites across the country, overall exceeding 40% for NSCLC specimens from advanced-stage disease.

Integrated Clinical and Genomic Annotation of Patient Tumor and Corresponding PDX

The algorithm for PDX creation and analysis after patient tumor biopsy is shown in Figure 1. In a synchronized evaluation process, patient tumors (PTs) and subsequent PDXs undergo histomorphologic assessment by a single reference pathologist, biomarker testing at the Clinical Laboratory Improvement Amendments (CLIA) laboratory at Response Genetics, Inc, and genomic analysis using multiple platforms including gene expression arrays and next-generation sequencing. PDXs of interest, based on molecular subtyping (eg, EGFR-mutated, Kirsten rat sarcoma or KRAS-mutated, squamous, etc) are then grouped into panels for subsequent drug testing. Figure 2 demonstrates how a panel (eg, EGFR-mutated) is subgrouped according to mutation type, resistance mechanism (if known), and clinical annotation (sensitive to EGFR tyrosine kinase inhibitor, de novo resistance, or acquired resistance). Patient clinical annotation is tracked (case example as shown in Figure 3) to account for all pertinent patient demographic characteristics, including smoking status, treatment records, and timing of PDX

Figure 4 Diagram of Experimental Approach to Testing Drug Efficacy in Patient-Derived Xenograft (PDX) Models. Tumor Fragments From a Single Passage 2 (P2) Mouse Were Implanted Simultaneously Into a Treatment Cohort of NSG Mice. As Tumors Reach 300 mm³, They Are Randomized Into Arms and Receive Treatment for the Indicated Number of Cycles. All Treated Mice Are Further Observed for an Extended Period of Time or Until the Tumor Reached a Terminal Size and Mice Were Euthanized. In Parallel, Pharmacodynamic Studies Were Conducted, With Mice Treated for Short Periods (For Instance, 6 or 24 Hours) to Examine the Immediate Molecular Effects of Treatment Such as Degree of Target Inhibition and Effects on Signal Transduction. Tumors Were Harvested Using a Rapid Resection Protocol Designed to Minimize Tumor Ischemia

Abbreviation: PK = protein kinase.

creation. Mutational fidelity is demonstrated in Table 1 for a representative group of PTs and PDXs characterized for EGFR and KRAS mutations and anaplastic lymphoma kinase gene rearrangements in a CLIA laboratory environment (Response Genetics, Inc).^{25,26}

Drug Testing Paradigm

A variety of projects are ongoing to test PDX panels of interest for new drug treatment strategies, and to determine mechanisms of resistance and how to overcome them. A combination of tumor growth inhibition studies and real-time tumor pharmacodynamics are incorporated into each drug testing project (Figure 4). An advantage of this testing paradigm in these is that degrees of responsiveness can be quantitatively measured to provide a more nuanced assessment of the pharmacologic effect of any drug or drug combination. Moreover, the availability of an annotated panel of xenografts for a particular genetic mutation or histology should increase the likelihood of discovery of new genomic signatures of drug sensitivity and resistance. The use of these models in coclinical trials is of particular interest.

An ongoing pilot study to investigate mechanisms of acquired drug resistance in EGFR-mutated lung cancer, for example, is using a panel of EGFR-mutated PDXs to investigate a drug regimen of afatinib with or without cetuximab, using erlotinib and

vehicle as controls. Results in 2 PDX models, one in which the host patient proved to subsequently be responsive to the afatinib-cetuximab combination and the other patient was unresponsive, mimicked the clinical outcomes, as shown in Figure 5. Associated tumor pharmacodynamics (Figure 6) illustrates abrogation of multiple signal transduction pathways globally mediated by maximal EGFR inhibition in the sensitive model (LG0703), in contrast to incomplete EGFR inhibition coupled with compensatory upregulation of survival pathways in the resistant model (LG1049). These data suggest that PDX models can recapitulate drug treatment outcomes in patients from whom the PDXs were derived, and that investigations can be designed to increase understanding of underlying biologic pathways and to devise strategies to improve effectiveness of therapy. This hypothesis will be tested in a future Southwestern Oncology Group phase II/III clinical trial (S1403) that will compare afatinib with or without cetuximab in the first-line therapy of patients with EGFR-mutated lung cancer, in which selected patients will undergo repeat tumor biopsy at the time of progressive disease, for genomic assessment and for PDX creation and drug testing.

Summary

The integrated preclinical-clinical modeling strategy described herein, that uses a large annotated NSG mouse resource of PDX

Figure 5 Activity of Epidermal Growth Factor Receptor (EGFR) Inhibitors in EGFR-Mutant Patient-Derived Xenograft (PDX) Models. The Relative Tumor Growth Inhibition Induced By Erlotinib, Afatinib, Cetuximab or the Combination of Afatinib With Cetuximab Was Tested in 2 PDX Models Derived From Erlotinib-Refractory Patients Who Subsequently (After PDX Biopsy) Were Treated With the Combination of Afatinib With Cetuximab. (A) The Patient From Whom the PDX LG0703 Was Derived (See Figure 3) Was Responsive to Subsequent Therapy With Afatinib With Cetuximab. This Combination Proved to Be Effective in The PDX Model, Recapitulating the Clinical Results. (B) In Contrast, PDX LG1049, Derived From a T790M-Positive Cancer Patient Who Did Not Respond to Subsequent Cetuximab With Afatinib Therapy, Exhibited No Significant Response to Afatinib With Cetuximab. Again, Clinical Results Were Recapitulated in the Related PDX Models

A Effects of Afatinib, Erlotinib and Cetuximab on LG0703 PDX Model

B Effects of Afatinib, Erlotinib and Cetuximab on LG1049 PDX Model

Abbreviations: ANOVA = Analysis of Variance; p.o. = orally; q.d. = once per day.

models, provides a novel research tool for drug development and for understanding the genomic complexity of lung cancer. Prospective development of PDXs from mutation-specific cohorts of patients treated in a homogeneous fashion, such as the planned PDX project associated with the S1403 clinical trial, should provide a unique resource for future study.

It is anticipated that availability of such an annotated resource will assist in optimization of therapeutic strategies and lead to improved patient outcomes.

Acknowledgments

Research reported in this publication was partially supported by the NCI award to UCD (P30CA034196) by JAX Director's Innovation Fund, the Addario Lung Cancer Foundation, and the HOPE Foundation. The MTB database is supported by the National Institutes of Health (CA089713).

Disclosure

The authors have stated that they have no conflicts of interest.

Figure 6 Comparison of Tumor Pharmacodynamic Effects After Afatinib/Cetuximab in Two Erlotinib-Resistant Patient-Derived Xenograft (PDX) Models LG0703 (T790M-Negative) and LG1049 (T790M-Positive). (A) LG0703 Untreated, Revealing Intense Signaling Through ERK and p38. (B) LG0703 After Treatment With Afatinib/Cetuximab for 24 Hours, Showing Substantially Diminished MAPK and PI3K-AKT Signaling Concurrent With Loss of Epidermal Growth Factor Receptor (EGFR) Phosphorylation. (C) LG1049 Untreated, Showing Enhanced AKT Activity With Moderate ERK Activity at Baseline. (D) LG1049 After Treatment With Afatinib/Cetuximab Showing Minimal Diminishment of EGFR Phosphorylation and Signal Transduction, With Compensatory Upregulation of p38. Phosphorylation States of Receptors and Signaling Intermediaries Were Measured Using A Combination of immunoblot and Kinase Arrays (Data Not Shown, Publication In Preparation)

Abbreviations: ERK = extracellular signal-regulated kinase; MAPK = Mitogen-activated protein kinase; mTor = mammalian target of rapamycin; Mut = mutant; PI3K = phosphatidylinositol-3-kinase.

References

- Collins I, Workman P. New approaches to molecular cancer therapeutics. *Nat Chem Biol* 2006; 2:689-700.
- Workman P. Genomics and the second golden era of cancer drug development. *Mol Biosyst* 2005; 1:17-26.
- Daigo Y, Takano A, Teramoto K, Chung S, Nakamura Y. A systematic approach to the development of novel therapeutics for lung cancer using genomic analyses. *Clin Pharmacol Ther* 2013; 94:218-23.
- Paul SM, Mytelka DS, Dunwiddie CT, et al. How to improve R&D productivity: the pharmaceutical industry's grand challenge. *Nat Rev Drug Discov* 2010; 9:203-14.
- Hutchinson L, Kirk R. High drug attrition rates—where are we going wrong? *Nat Rev Clin Oncol* 2011; 8:189-90.
- Rubin EH, Gilliland DG. Drug development and clinical trials—the path to an approved cancer drug. *Nat Rev Clin Oncol* 2012; 9:215-22.
- Li T, Kung HJ, Mack PC, Gandara DR. Genotyping and genomic profiling of non-small-cell lung cancer: implications for current and future therapies. *J Clin Oncol* 2013; 31:1039-49.
- Gandara DR, Li T, Lara PN Jr, et al. Algorithm for codevelopment of new drug-predictive biomarker combinations: accounting for inter- and inpatient tumor heterogeneity. *Clin Lung Cancer* 2012; 13:321-5.
- Collins FS. Reengineering translational science: the time is right. *Sci Transl Med* 2011; 3:90cm17.
- Rosell R, Karachaliou N, Morales-Espinoso D, et al. Adaptive resistance to targeted therapies in cancer. *Transl Lung Cancer Res* 2013; 2:152-9.
- Gandara DR, Li T, Lara PN, et al. Acquired resistance to targeted therapies against oncogene-driven non-small-cell lung cancer: approach to subtyping progressive disease and clinical implications. *Clin Lung Cancer* 2014; 15:1-6.
- Gower A, Wang Y, Giaccone G. Oncogenic drivers, targeted therapies, and acquired resistance in non-small-cell lung cancer. *J Mol Med (Berl)* 2014; 92:697-707.
- Iwasa Y, Michor F. Evolutionary dynamics of intratumor heterogeneity. *PLoS One* 2011; 6:e17866.
- Politi K, Pao W. How genetically engineered mouse tumor models provide insights into human cancers. *J Clin Oncol* 2011; 29:2273-81.
- Carver BS, Pandolfi PP. Mouse modeling in oncologic preclinical and translational research. *Clin Cancer Res* 2006; 12:5305-11.
- Sharpless NE, Depinho RA. The mighty mouse: genetically engineered mouse models in cancer drug development. *Nat Rev Drug Discov* 2006; 5:741-54.
- Regales L, Balak MN, Gong Y, et al. Development of new mouse lung tumor models expressing EGFR T790M mutants associated with clinical resistance to kinase inhibitors. *PLoS One* 2007; 2:e810.
- Weaver Z, Difilippantonio S, Carretero J, et al. Temporal molecular and biological assessment of an erlotinib-resistant lung adenocarcinoma model

Tumor Genomics to Patient Outcomes Through PDX Platform

- reveals markers of tumor progression and treatment response. *Cancer Res* 2012; 72:5921-33.
19. Tentler JJ, Tan AC, Weekes CD, et al. Patient-derived tumour xenografts as models for oncology drug development. *Nat Rev Clin Oncol* 2012; 9: 338-50.
 20. Shultz LD, Ishikawa F, Greiner DL. Humanized mice in translational biomedical research. *Nat Rev Immunol* 2007; 7:118-30.
 21. Olive KP, Tuveson DA. The use of targeted mouse models for preclinical testing of novel cancer therapeutics. *Clin Cancer Res* 2006; 12:5277-87.
 22. Hidalgo M, Bruckheimer E, Rajeshkumar NV, et al. A pilot clinical study of treatment guided by personalized tumorgrafts in patients with advanced cancer. *Mol Cancer Ther* 2011; 10:1311-6.
 23. John T, Kohler D, Pintilie M, et al. The ability to form primary tumor xenografts is predictive of increased risk of disease recurrence in early-stage non-small-cell lung cancer. *Clin Cancer Res* 2011; 17:134-41.
 24. Bult CJ, Krupke DM, Begley DA, et al. Mouse Tumor Biology (MTB): a database of mouse models for human cancer. *Nucleic Acids Res* 2015; 43(Database issue): D818-24.
 25. Gandara DR, Grimminger P, Mack PC, et al. Association of epidermal growth factor receptor activating mutations with low ERCC1 gene expression in non-small-cell lung cancer. *J Thorac Oncol* 2010; 5:1933-8.
 26. Maus MK, Grimminger PP, Mack PC, et al. KRAS mutations in non-small-cell lung cancer and colorectal cancer: implications for EGFR-targeted therapies. *Lung Cancer* 2014; 83:163-7.