

UC Irvine

UC Irvine Previously Published Works

Title

Glaucoma Surgery Calculator: Limited Additive Effect of Phacoemulsification on Intraocular Pressure in Ab Interno Trabeculectomy

Permalink

<https://escholarship.org/uc/item/8xs96230>

Journal

PLOS ONE, 11(4)

ISSN

1932-6203

Authors

Neiweem, Ashley E
Bussel, Igor I
Schuman, Joel S
[et al.](#)

Publication Date

2016

DOI

10.1371/journal.pone.0153585

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

RESEARCH ARTICLE

Glaucoma Surgery Calculator: Limited Additive Effect of Phacoemulsification on Intraocular Pressure in Ab Interno Trabeculectomy

Ashley E. Neiveem¹, Igor I. Bussel², Joel S. Schuman², Eric N. Brown^{2,3}, Nils A. Loewen^{2*}

1 Rosalind Franklin University of Medicine and Science, Chicago Medical School, Chicago, United States of America, **2** Department of Ophthalmology, School of Medicine, University of Pittsburgh, Pittsburgh, United States of America, **3** Department of Ophthalmology, School of Medicine, Vanderbilt University, Nashville, Tennessee, United States of America

These authors contributed equally to this work.

* Loewen.nils@gmail.com



OPEN ACCESS

Citation: Neiveem AE, Bussel II, Schuman JS, Brown EN, Loewen NA (2016) Glaucoma Surgery Calculator: Limited Additive Effect of Phacoemulsification on Intraocular Pressure in Ab Interno Trabeculectomy. PLoS ONE 11(4): e0153585. doi:10.1371/journal.pone.0153585

Editor: Pedro Gonzalez, Duke University, UNITED STATES

Received: October 23, 2015

Accepted: March 31, 2016

Published: April 14, 2016

Copyright: © 2016 Neiveem et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper.

Funding: This work was supported by Research to Prevent Blindness Departmental Grant and Individual Grant from Department of Ophthalmology, University of Pittsburgh. The sponsor or funding organization had no role in the design or conduct of this research. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Abstract

Purpose

To compare intraocular pressure (IOP) reduction and to develop a predictive surgery calculator based on the results between trabectome-mediated ab interno trabeculectomy in pseudophakic patients versus phacoemulsification combined with trabectome-mediated ab interno trabeculectomy in phakic patients.

Methods

This observational surgical cohort study analyzed pseudophakic patients who received trabectome-mediated ab interno trabeculectomy (AIT) or phacoemulsification combined with AIT (phaco-AIT). Follow up for less than 12 months or neovascular glaucoma led to exclusion. Missing data was imputed by generating 5 similar but non-identical datasets. Groups were matched using Coarsened Exact Matching based on age, gender, type of glaucoma, race, preoperative number of glaucoma medications and baseline intraocular pressure (IOP). Linear regression was used to examine the outcome measures consisting of IOP and medications.

Results

Of 949 cases, 587 were included consisting of 235 AIT and 352 phaco-AIT. Baseline IOP between groups was statistically significant ($p \leq 0.01$) in linear regression models and was minimized after Coarsened Exact Matching. An increment of 1 mmHg in baseline IOP was associated with a 0.73 ± 0.03 mmHg IOP reduction. Phaco-AIT had an IOP reduction that was only 0.73 ± 0.32 mmHg greater than that of AIT. The resulting calculator to determine IOP reduction consisted of the formula $-13.54 + 0.73 \times (\text{phacoemulsification yes:1,}$

Competing Interests: The authors have read the journal's policy and have the following competing interests: Nils Loewen has received honoraria for wet labs and lectures by Neomedix, Inc. The other authors have declared that no competing interests exist. This does not alter the authors' adherence to PLOS ONE policies on sharing data and materials.

$\text{no:0} + 0.73 \times (\text{baseline IOP}) + 0.59 \times (\text{secondary open angle glaucoma yes:1, no:0}) + 0.03 \times (\text{age}) + 0.09 \times (\text{medications}).$

Conclusions

This predictive calculator for minimally invasive glaucoma surgery can assist clinical decision making. Only a small additional IOP reduction was observed when phacoemulsification was added to AIT. Patients with a higher baseline IOP had a greater IOP reduction.

Introduction

Cataract surgery is often associated with a moderate intraocular pressure (IOP) reduction of 1.5–3 mmHg in patient with ocular hypertension or glaucoma [1–3]. Minimally invasive glaucoma surgery (MIGS) allows to combine IOP lowering with vision improvement from cataract surgery in an age group often affected by both. Such a practice pattern has become more common because it is standardized, safe [4] and also cost effective [5]. The first randomized controlled trials comparing phacoemulsification alone to phacoemulsification combined with implantation of trabecular bypass microstents, a form of MIGS, (iStent, Glaukos, Laguna Hills, CA), showed a relatively small additional effect of these implants on IOP reduction [6]. Ab interno trabeculectomy with the trabectome (Neomedix Corp; Tustin, CA), another MIGS modality, lowers IOP by plasma-mediated ionization and ablation of trabecular meshwork (TM) of up to 180° thereby increasing aqueous outflow in eyes with an intact downstream drainage system [4]. Both ab interno trabeculectomy (AIT) and phacoemulsification combined with ab interno trabeculectomy (phaco-AIT) can be used in patients with different angle opening [7] and surgical status [8,9]. The purpose of the comparison in this study was to assess reduction of IOP after AIT performed in pseudophakic patients versus phaco-AIT in phakic patients using a *Coarsened Exact Matched* cohort. Based on results with microstents [10], we hypothesized that in this matched comparison the benefit of adding phacoemulsification to AIT would be associated with a greater reduction in IOP and medications during 12 month follow-up. The resulting calculator can help clinicians to predict the IOP reduction.

Methods

Participants

Data for this study were collected with approval by the Institutional Review Board of the University of Pittsburgh, in accordance with the Declaration of Helsinki and the Health Insurance Portability and Accountability Act. No informed consent was necessary for this retrospective, observational cohort study. Patient records were anonymized and de-identified prior to analysis. Subjects were divided into pseudophakic patients who received AIT and phakic patients who received phaco-AIT. Outcomes were determined for all patients with a diagnosis of glaucoma with or without a visually significant cataract, who had 12 months of follow-up. The specific target IOP was set on a case-by-case basis by the individual treating physician and was the maximum IOP estimated to prevent further nerve damage. Patients who were followed for less than 12 months or diagnosed with neovascular glaucoma were excluded. Indications for AIT consisted of worsening glaucoma on maximally tolerated topical therapy while indications for phaco-AIT were the same or stable glaucoma with desire to reduce medications plus a visually significant cataract with visual brightness acuity testing equal or worse than least 0.4 logMAR

(20/50 Snellen). The postoperative medications consisted of 1% pilocarpine four times per day for 1 month, then three times per day for 1 month, 1% prednisolone acetate four times per day for 1 week to be tapered by one drop each week, and a third or fourth generation fluoroquinolone four times per day for 1 week. Glaucoma medications could be continued as deemed necessary to achieve target pressures. Visual field status of all patients was categorized as early, moderate, or advanced by individual glaucoma specialists based on the most recent Humphrey visual field exams (Zeiss, Jena, Germany). All patients had a comprehensive slit lamp, gonioscopy and dilated ophthalmoscopy exam prior to surgery.

Statistics

Demographics were compared by Mann-Whitney U test and chi-squared test for continuous and categorical variables, respectively. To avoid eliminating data with missing values multiple imputation was used. Missing values of the incomplete dataset were imputed $m > 1$ times, thus creating m completed datasets. Second, each of the m completed datasets were independently analyzed. Finally, the results from each of the m analysis were pooled into a final result. Missing data such as age, gender and race were imputed by generating 5 similar but non-identical datasets. Groups were then matched by utilizing *Coarsened Exact Matching* [11–13] based on age, gender, type of glaucoma, race, preoperative number of glaucoma medications and baseline IOP. Univariate linear regression was performed first and those variables that were statistically significant were included in the final multivariate regression model. A p-value of less than 0.05 was considered statistically significant. Continuous variables were expressed as mean \pm SD. All analyses were performed using R [14].

Results

Baseline Demographics

After applying exclusion criteria and matching, a total of 587 patients were included in the study consisting of 235 AIT and 352 phaco-AIT (Fig 1). Baseline demographics are shown in Table 1.

ACG (angle closure glaucoma); POAG (primary open angle glaucoma); SOAG (secondary open angle glaucoma).

Primary open angle glaucoma (POAG) comprised 86% and 89% of AIT and phaco-AIT, respectively. Secondary open angle glaucoma (SOAG) included 14% and 11% of AIT and phaco-AIT, respectively. From the matched subjects, 75% in AIT and also in phaco-AIT were Caucasian, followed by Asian, African American, and others. These values were not statistically significant. Additionally, age, gender, baseline number of glaucoma medications, and baseline IOP were found to be statistically different ($p < 0.01$) between groups (Table 1). Following *Coarsened Exact Matching*, these preoperative differences between treatment groups were minimized (Table 2).

Multiple Imputation and Coarsened Exact Matching

Missing data in each category are recorded. Data missing from baseline number of medications, type of glaucoma, and IOP were 0% for both groups. Conversely, age, gender, and race had missing data among both groups. Six percent of AIT had an unknown age, 4% were without defined gender and 8% without defined race. Two percent of phaco-AIT had an unknown age, 2% were without defined gender, and 11% without race.




Fig 1. Trabectome Surgeries. Cases analyzed after accounting for exclusion criteria (AIT, ab interno trabeculectomy; phaco-AIT, AIT combined with phacoemulsification; IOP, intraocular pressure; POAG, primary open angle glaucoma; SOAG, secondary open angle glaucoma).

doi:10.1371/journal.pone.0153585.g001

Table 1. Raw data demographics of study population. Demographics for AIT-only and phaco-AIT show significant difference ($p < 0.05$) for age, gender, baseline IOP, and baseline number of medications in unmatched data.

	AIT (n = 368)	Phaco-AIT (n = 581)	p-value
Age			<0.01
Mean±SD	75±10	74±9	
Range	(50, 96)	(51, 94)	
Gender			0.01
Male	130 (35%)	248 (43%)	
Female	223 (61%)	323 (56%)	
Types of Glaucoma			0.19
ACG	2 (1%)	11 (2%)	
POAG	289 (78%)	442 (76%)	
SOAG	77 (21%)	128 (22%)	
Race			0.34
African Americans	19 (5%)	29 (5%)	
Asians	92 (25%)	166 (29%)	
Caucasians	216 (59%)	303 (52%)	
Others	10 (3%)	20 (3%)	
Baseline IOP			<0.01
Mean±SD	24.1±7.1	20.6±6.6	
Range	(10, 51)	(10, 59)	
Baseline Number of Glaucoma Medications			<0.01
Mean±SD	2.9±1.1	2.4±1.0	
Range	(1, 6)	(1, 5)	

ACG (angle closure glaucoma); POAG (primary open angle glaucoma); SOAG (secondary open angle glaucoma).

doi:10.1371/journal.pone.0153585.t001

Linear Regression Models

Linear regression of the multiple imputed, matched data was used to identify the influence of the parameters above on the IOP lowering effect of surgery. Univariate linear regression was performed first (Table 3) using the variables phacoemulsification, baseline IOP, SOAG, age, number of medications at baseline, race and gender (male).

Of these variables, phacoemulsification, baseline IOP, SOAG, age and number of medications at baseline were found to be statistically significant and included in the final multivariate regression model (Table 4). Only baseline IOP and phacoemulsification were statistically significant in both models ($p = 0.02$ and $p < 0.01$, respectively).

Each increment of 1 mmHg in baseline IOP was associated with an IOP reduction of 0.73 ± 0.03 mmHg ($p < 0.01$). After adjusting for baseline IOP, age, baseline number of glaucoma medications, and type of glaucoma, phacoemulsification conferred an additional IOP reduction of 0.73 ± 0.32 mmHg IOP. Preoperative IOP was 22.6 ± 6.4 mmHg in AIT and 19.9 ± 5.8 mmHg in phaco-AIT with 2.8 ± 1.1 medications in AIT and 2.4 ± 1.1 in phaco-AIT. At one year, IOP in AIT was reduced to 16.9 ± 4.5 mmHg (mean±SD) and in phaco-AIT to 15.4 ± 3.6 mmHg ($p < 0.01$), while medications in AIT declined to 2.3 ± 1.3 and in phaco-AIT to 1.7 ± 1.2 ($p < 0.01$). Postoperative AIT and phaco-AIT were significantly different at all time points for both IOP and medications due to the large sample sizes with a narrow confidence interval and small standard error (Fig 2).

Table 2. Matched data demographics of study population. Matched data is shown for both AIT-only and phaco-AIT group demographics.

	AIT (n = 235)	phaco-AIT (n = 352)	p-value
Age			0.04
Mean±SD	76±9	75±8	
Range	(51,96)	(56,94)	
Gender			0.15
Female	156 (66%)	212 (60%)	
Male	79 (34%)	140 (40%)	
Types of Glaucoma			0.34
ACG	0 (0%)	0 (0%)	
POAG	202 (86%)	313 (89%)	
SOAG	33 (14%)	39 (11%)	
Race			0.77
African Americans	7 (3%)	7 (2%)	
Asians	48 (20%)	78 (22%)	
Caucasians	176 (75%)	263 (75%)	
Others	4 (2%)	4 (1%)	
Baseline IOP			<0.01
Mean±SD	22.6±6.4	19.9±5.9	
Range	(10, 46)	(10, 42)	
Baseline Number of Glaucoma Medications			<0.01
Mean±SD	2.8±1.1	2.4±1.1	
Range	(1,6)	(1,5)	

doi:10.1371/journal.pone.0153585.t002

Glaucoma Surgery IOP Reduction Calculator

The calculator predicting the IOP reduction had the formula: $-13.54 + 0.73 \times (\text{phaco}; \text{yes}:1, \text{no}:0) + 0.73 \times (\text{baseline IOP}) + 0.59 \times (\text{SOAG}; \text{yes}:1, \text{no}:0) + 0.03 \times (\text{age}) + 0.09 \times (\text{medications})$.

For example, a 75-year-old pseudophakic patient with POAG with a baseline IOP of 21 and 2 different medications receiving AIT alone would be expressed as: $13.54+0.73(0)+0.73 \times (21) + 0.59(0)+0.03 \times (75)+0.09 \times (2) = 4.22$ mmHg reduction in IOP, with a resulting postoperative IOP of 16.78 mmHg.

Table 3. Univariate linear regression of patient parameters of study. A p-value of <0.05 is considered statistically significant.

	Coefficient	Standard Error	p-value
Phaco	-1.35	0.50	<0.01
Baseline IOP	0.74	0.03	<0.01
SOAG	4.17	0.74	<0.01
Age	0.10	0.03	<0.01
Baseline # of medications	0.56	0.22	<0.01
Race			
Asian	1.10	2.30	0.64
Caucasian	1.94	2.19	0.39
Other	4.84	3.47	0.18
Male	-0.75	0.50	0.14

Phaco (phacoemulsification); IOP (intraocular pressure); SOAG (secondary open angle glaucoma).

doi:10.1371/journal.pone.0153585.t003

Table 4. Multivariate linear regression of patient parameters that were found to be statistically significant ($p < 0.05$) in univariate linear regression.

	Coefficient	Standard Error	p-value
Intercept	-13.54	1.67	<0.01
Phaco	0.73	0.32	0.02
Baseline IOP	0.73	0.03	<0.01
SOAG	0.59	0.50	0.24
Age	0.03	0.02	0.10
Baseline # of Medications	0.09	0.14	0.55

Phaco (phacoemulsification); IOP (intraocular pressure); SOAG (secondary open angle glaucoma).

doi:10.1371/journal.pone.0153585.t004

Conversely, a 75-year-old phakic patient with POAG with the same baseline IOP and medications receiving phaco-AIT in a combined approach would have an IOP reduction of $-13.54 + 0.73(1) + 0.73 \times (21) + 0.59(0) + 0.03(75) + 0.09(2) = 4.95$ mmHg, thus a postoperative IOP of 16.05 mmHg.

The linear relationship of pre- and postoperative IOP can be seen in the scattergrams that show every single data point (Fig 3).

Discussion

We created a glaucoma surgery calculator to determine the postoperative IOP based on the preoperative IOP, type of open angle glaucoma, age, medications, and type of surgery. This first calculator for minimally invasive glaucoma surgery (MIGS) can help clinicians to estimate outcomes and anticipate the need for postoperative glaucoma medications. Using *Coarsened Exact Matching*, we found only a small additional contribution of phacoemulsification to the considerable IOP reduction from AIT. The impact of baseline IOP on total pressure reduction





Fig 2. IOP and medication plots. Preoperative and postoperative intraocular pressure (IOP; left) and number of glaucoma medications (Meds; right) over the 12 month follow-up for both groups. Represented as mean \pm standard error. Statistically significantly different at all time points for both plots ($p < 0.05$).

doi:10.1371/journal.pone.0153585.g002

AIT Scattergrams



Phaco-AIT Scattergrams




Fig 3. Phaco-AIT and AIT Scattergrams. Scattergrams of AIT and phaco-AIT after 1 month (left) and 12 months (right). Baseline IOP plotted against IOP at 1 month and 12 months with $x = y$ line. Red line represents linear fit.

doi:10.1371/journal.pone.0153585.g003

was substantial, adding 0.73 mmHg IOP reduction per 1 mmHg higher baseline IOP. Both groups achieved a significant decline in medications.

Compared to the limited IOP effect of only 0.73 ± 0.32 mmHg by phacoemulsification in our study, a more significant IOP reduction can sometimes be seen after phacoemulsification [1–3]. It has been hypothesized to be caused by TM and Schlemm's canal distension that increase the outflow facility [15,16], activation of a TM stress response pathway from ultrasound and

fluids [17], a trabeculoplasty-like effect [18,19] or resolution of relative pupillary block [15]. Our results are consistent with the concept that this may be mediated by the remaining temporal TM [16,20]. Different from the open angle glaucoma patients examined in this study, angle closure glaucoma patients often experience a profound IOP reduction from cataract surgery alone, both in the chronic [21] and the acute form [22].

We caution against use of phacoemulsification alone for the purpose of IOP reduction as recently advocated [15]. Phacoemulsification on its own does not always lower IOP reliably in glaucoma patients because of a relatively more diseased TM [23] and can cause potentially dangerous IOP spikes during the postoperative course [23,24]. This can be prevented by combining AIT with phacoemulsification [24]. A higher than normal IOP before surgery is a risk factor for IOP spikes even in patients without glaucoma [25]. Eyes with a more decreased outflow facility may be more prone to this but also experience a larger IOP reduction from AIT [26]. The impact of phacoemulsification on IOP in patients with same session TM bypass microstents may be relatively higher compared to ab interno trabeculectomy because of more remaining TM [4] and fewer drainage segments accessed in the former [27].

The raw baseline age differences and higher number of medications between AIT and phaco-AIT groups match the increased incidence of cataracts and glaucoma with age [28]. Following coarsened exact matching, these differences were minimized and allowed a statistically valid comparison with linear regression [29]. It was previously assumed that IOP reduction following AIT is relatively independent of preoperative IOP [7,8] and only limited by episcleral venous pressure and other downstream elements [30]. The linear correlation between pre- and postoperative IOPs seen here suggests that patients with higher baseline IOP may have both a higher TM-mediated outflow resistance and a somewhat higher outflow resistance that is downstream of the TM.

This study had limitations. Instead of a randomized controlled trial, we applied newer statistical methods, *Coarsened Exact Matching* and *Multiple Imputation*, that belong to the class of *Monotonic Imbalance Bounding (MIB)* and make no assumptions about the data generation process [12,31]. This accounts for missing demographic values and avoids losing data thereby increasing the validity and sample size of the study. Additionally, the 12 month follow-up is still a relatively short-term follow-up endpoint for patients with good life expectancy and ongoing ocular disease. The calculator presented here may not apply to the pediatric or juvenile glaucoma population. Interpretations of IOP outcomes presented here have to take into account that phaco-AIT patients in reality have a mixed indication of vision improvement (cataract surgery) with often optional IOP or medication reduction (AIT). As a result, IOPs can be considerably lower if the second of the two average glaucoma medication is not eliminated.

In conclusion, this first glaucoma surgery calculator advises clinicians on IOP after trabeculectomy-mediated ab interno trabeculectomy. Phacoemulsification has only a small additional IOP lowering effect when combined with trabeculectomy surgery. Patients with higher baseline IOP are expected to have a greater IOP reduction.

Author Contributions

Conceived and designed the experiments: NAL ENB IIB. Performed the experiments: AN ENB IIB. Analyzed the data: AN ENB IIB NAL. Contributed reagents/materials/analysis tools: JSS NAL. Wrote the paper: AN IIB JSS NAL.

References

1. Mansberger SL, Gordon MO, Jampel H, Bhorade A, Brandt JD, Wilson B, et al. Reduction in intraocular pressure after cataract extraction: the Ocular Hypertension Treatment Study. *Ophthalmology*. 2012; 119: 1826–1831. doi: [10.1016/j.ophtha.2012.02.050](https://doi.org/10.1016/j.ophtha.2012.02.050) PMID: [22608478](https://pubmed.ncbi.nlm.nih.gov/22608478/)

2. Yang HS, Lee J, Choi S. Ocular biometric parameters associated with intraocular pressure reduction after cataract surgery in normal eyes. *Am J Ophthalmol.* 2013; 156: 89–94.e1. doi: [10.1016/j.ajo.2013.02.003](https://doi.org/10.1016/j.ajo.2013.02.003) PMID: [23628350](https://pubmed.ncbi.nlm.nih.gov/23628350/)
3. Shingleton BJ, Pasternack JJ, Hung JW, O'Donoghue MW. Three and five year changes in intraocular pressures after clear corneal phacoemulsification in open angle glaucoma patients, glaucoma suspects, and normal patients. *J Glaucoma.* 2006; 15: 494–498. PMID: [17106361](https://pubmed.ncbi.nlm.nih.gov/17106361/)
4. Kaplowitz K, Schuman JS, Loewen NA. Techniques and outcomes of minimally invasive trabecular ablation and bypass surgery. *Br J Ophthalmol.* 2014; 98: 579–585. doi: [10.1136/bjophthalmol-2013-304256](https://doi.org/10.1136/bjophthalmol-2013-304256) PMID: [24338085](https://pubmed.ncbi.nlm.nih.gov/24338085/)
5. Iordanous Y, Kent JS, Hutnik CML, Malvankar-Mehta MS. Projected cost comparison of Trabectome, iStent, and endoscopic cyclophotocoagulation versus glaucoma medication in the Ontario Health Insurance Plan. *J Glaucoma.* 2014; 23: e112–8. PMID: [23807348](https://pubmed.ncbi.nlm.nih.gov/23807348/)
6. Samuelson TW, Katz LJ, Wells JM, Duh Y-J, Giamporcaro JE, US iStent Study Group. Randomized evaluation of the trabecular micro-bypass stent with phacoemulsification in patients with glaucoma and cataract. *Ophthalmology.* 2011; 118: 459–467. doi: [10.1016/j.ophtha.2010.07.007](https://doi.org/10.1016/j.ophtha.2010.07.007) PMID: [20828829](https://pubmed.ncbi.nlm.nih.gov/20828829/)
7. Bussell II, Kaplowitz K, Schuman JS, Loewen NA, Trabectome Study Group. Outcomes of ab interno trabeculectomy with the trabectome by degree of angle opening. *Br J Ophthalmol.* 2015; 99: 914–919. doi: [10.1136/bjophthalmol-2014-305577](https://doi.org/10.1136/bjophthalmol-2014-305577) PMID: [25336577](https://pubmed.ncbi.nlm.nih.gov/25336577/)
8. Bussell II, Kaplowitz K, Schuman JS, Loewen NA, Trabectome Study Group. Outcomes of ab interno trabeculectomy with the trabectome after failed trabeculectomy. *Br J Ophthalmol.* 2014; doi: [10.1136/bjophthalmol-2013-304717](https://doi.org/10.1136/bjophthalmol-2013-304717)
9. Mosaed S, Chak G, Haider A, Lin KY, Minckler DS. Results of Trabectome Surgery Following Failed Glaucoma Tube Shunt Implantation: Cohort Study. *Medicine.* 2015; 94: e1045. PMID: [26222842](https://pubmed.ncbi.nlm.nih.gov/26222842/)
10. Samuelson TW, Katz LJ, Wells JM, Duh Y-J, Giamporcaro JE, US iStent Study Group. Randomized evaluation of the trabecular micro-bypass stent with phacoemulsification in patients with glaucoma and cataract. *Ophthalmology.* Elsevier; 2011; 118: 459–467.
11. Iacus SM, King G, Porro G. Causal Inference without Balance Checking: Coarsened Exact Matching. *Polit Anal.* 2011; doi: [10.1093/pan/mpr013](https://doi.org/10.1093/pan/mpr013)
12. Iacus SM, King G, Porro G. Multivariate Matching Methods That Are Monotonic Imbalance Bounding. *J Am Stat Assoc.* 2011; 106: 345–361.
13. Parikh HA, Bussell II, Schuman JS, Brown EN, Loewen NA. Coarsened Exact Matching of Phaco-Trabectome to Trabectome in Phakic Patients: Lack of Additional Pressure Reduction from Phacoemulsification. *PLoS One.* 2016; 11: e0149384. doi: [10.1371/journal.pone.0149384](https://doi.org/10.1371/journal.pone.0149384) PMID: [26895293](https://pubmed.ncbi.nlm.nih.gov/26895293/)
14. R Core Team. R: A Language and Environment for Statistical Computing [Internet]. Vienna, Austria: R Foundation for Statistical Computing; 2015. Available: <http://www.R-project.org/>
15. Poley BJ, Lindstrom RL, Samuelson TW, Schulze R Jr. Intraocular pressure reduction after phacoemulsification with intraocular lens implantation in glaucomatous and nonglaucomatous eyes: evaluation of a causal relationship between the natural lens and open-angle glaucoma. *J Cataract Refract Surg.* 2009; 35: 1946–1955. doi: [10.1016/j.jcrs.2009.05.061](https://doi.org/10.1016/j.jcrs.2009.05.061) PMID: [19878828](https://pubmed.ncbi.nlm.nih.gov/19878828/)
16. Kee C, Moon SH. Effect of cataract extraction and posterior chamber lens implantation on outflow facility and its response to pilocarpine in Korean subjects. *Br J Ophthalmol.* 2000; 84: 987–989. PMID: [10966950](https://pubmed.ncbi.nlm.nih.gov/10966950/)
17. Wang N, Chintala SK, Fini ME, Schuman JS. Ultrasound activates the TM ELAM-1/IL-1/NF-kappaB response: a potential mechanism for intraocular pressure reduction after phacoemulsification. *Invest Ophthalmol Vis Sci.* 2003; 44: 1977–1981. PMID: [12714632](https://pubmed.ncbi.nlm.nih.gov/12714632/)
18. Shazly TA, Latina MA, Dagianis JJ, Chitturi S. Effect of prior cataract surgery on the long-term outcome of selective laser trabeculoplasty. *Clin Ophthalmol.* 2011; 5: 377–380. doi: [10.2147/OPHTH.S17237](https://doi.org/10.2147/OPHTH.S17237) PMID: [21468349](https://pubmed.ncbi.nlm.nih.gov/21468349/)
19. Damji KF, Konstas AGP, Liebmann JM, Hodge WG, Ziakas NG, Giannikakis S, et al. Intraocular pressure following phacoemulsification in patients with and without exfoliation syndrome: a 2 year prospective study. *Br J Ophthalmol.* 2006; 90: 1014–1018. PMID: [16672324](https://pubmed.ncbi.nlm.nih.gov/16672324/)
20. Meyer MA, Savitt ML, Kopitas E. The effect of phacoemulsification on aqueous outflow facility. *Ophthalmology.* 1997; 104: 1221–1227. PMID: [9261307](https://pubmed.ncbi.nlm.nih.gov/9261307/)
21. Tham CCY, Kwong YYY, Leung DY, Lam SW, Li FCH, Chiu TYH, et al. Phacoemulsification versus combined phacotrabeculectomy in medically uncontrolled chronic angle closure glaucoma with cataracts. *Ophthalmology.* Elsevier; 2009; 116: 725–31, 731.e1–3.
22. Jacobi PC, Dietlein TS, Lücke C, Engels B, Krieglstein GK. Primary phacoemulsification and intraocular lens implantation for acute angle-closure glaucoma. *Ophthalmology.* Elsevier; 2002; 109: 1597–1603.

23. Slabaugh MA, Bojjikian KD, Moore DB, Chen PP. Risk factors for acute postoperative intraocular pressure elevation after phacoemulsification in glaucoma patients. *J Cataract Refract Surg.* 2014; 40: 538–544. doi: [10.1016/j.jcrs.2013.08.048](https://doi.org/10.1016/j.jcrs.2013.08.048) PMID: [24440104](https://pubmed.ncbi.nlm.nih.gov/24440104/)
24. Weiner Y, Severson ML, Weiner A. Intraocular pressure 3 to 4 hours and 20 hours after cataract surgery with and without ab interno trabeculectomy. *J Cataract Refract Surg.* Elsevier; 2015; 41: 2081–2091.
25. O'Brien PD, Ho SL, Fitzpatrick P, Power W. Risk factors for a postoperative intraocular pressure spike after phacoemulsification. *Can J Ophthalmol.* Elsevier; 2007; 42: 51–55.
26. Loewen R, Roy P, Parikh H, Dang Y, Schuman J, Loewen N. Impact of a Glaucoma Severity Index on Results of Trabectome Surgery: Larger Pressure Reduction in More Severe Glaucoma. *PLoS One.* 2016; In press.
27. Rosenquist R, Epstein D, Melamed S, Johnson M, Grant WM. Outflow resistance of enucleated human eyes at two different perfusion pressures and different extents of trabeculectomy. *Curr Eye Res.* 1989; 8: 1233–1240. PMID: [2627793](https://pubmed.ncbi.nlm.nih.gov/2627793/)
28. Klaver CC, Wolfs RC, Vingerling JR, Hofman A, de Jong PT. Age-specific prevalence and causes of blindness and visual impairment in an older population: the Rotterdam Study. *Arch Ophthalmol.* 1998; 116: 653–658. PMID: [9596502](https://pubmed.ncbi.nlm.nih.gov/9596502/)
29. Honaker J, King G, Blackwell M, Others. Amelia II: A program for missing data. *J Stat Softw.* 2011; 45: 1–47.
30. Schuman JS, Chang W, Wang N, de Kater AW, Allingham RR. Excimer laser effects on outflow facility and outflow pathway morphology. *Invest Ophthalmol Vis Sci.* 1999; 40: 1676–1680. PMID: [10393035](https://pubmed.ncbi.nlm.nih.gov/10393035/)
31. Iacus SM, King G, Porro G. Causal Inference without Balance Checking: Coarsened Exact Matching. *Polit Anal.* 2012; 20: 1–24.