

UCSF

UC San Francisco Previously Published Works

Title

Throwing Injury Prevention Strategies with a Whole Kinetic Chain-Focused Approach

Permalink

<https://escholarship.org/uc/item/8xv6d0mg>

Journal

Current Reviews in Musculoskeletal Medicine, 15(2)

ISSN

1935-973X

Authors

Mayes, Michael
Salesky, Madeleine
Lansdown, Drew A

Publication Date

2022-04-01

DOI

10.1007/s12178-022-09744-9

Peer reviewed



Throwing Injury Prevention Strategies with a Whole Kinetic Chain-Focused Approach

Michael Mayes¹ · Madeleine Salesky¹ · Drew A. Lansdown¹

Accepted: 17 December 2021 / Published online: 7 April 2022

© The Author(s) 2022

Abstract

Purpose of Review This review examines the relationship between the baseball pitching motion and the kinetic chain. The goal was to determine the underlying causes of a deficiency in throwing mechanics throughout a throwing motion, and to provide an evidence-based approach on how to prevent injuries caused by a lack of proper mechanics. In doing so, we sought to provide a warm-up strategy that can be added to every baseball player's daily on-field routine that is tailored to each phase of the throwing motion.

Recent Findings To help minimize the risk of injury to overhead throwing athletes, a thorough understanding of the throwing motion is critical. Throwing a ball places extreme stress on the body, notably the shoulder and elbow joints. With a clear understanding of the biomechanics of throwing, we can develop an injury prevention routine to minimize unnecessary stresses throughout the kinetic chain.

Summary The throwing cycle is a complex motion that places various stresses throughout the thrower's body, from the ankle to the core, and from the back to the shoulder and elbow. A thorough understanding of the mechanics of this motion, along with specific exercises to target the specific actions of each phase, may allow for throwers, regardless of their age and experience, to minimize injury risk.

Keywords Baseball · Throwing motion · Pitching mechanics · Kinetic chain · Shoulder · Injury prevention · Warm-up

Introduction

Baseball and softball are among the most common sports for children and adolescents, with over 5 million children participating annually in the USA [1]. Throwing a ball places extreme stress on the shoulder and elbow joints [2–5]. The throwing motion, however, also requires careful coordination and strength throughout the entire kinetic chain [6]. With baseball increasingly becoming a year-round sport, overuse injuries are becoming more common, and youth pitchers who throw more than 100 innings per calendar year have a

3.5 times greater risk of sustaining a serious sports-related injury [7]. While USA Baseball has published pitching safety guidelines and regulations, sports-related injury among overhead throwing athletes is still common. Hamstring, lumbar paraspinal, and oblique muscle strains represent three of the five most common injuries in a report on Major and Minor League Baseball Players with data from the Major League Baseball (MLB) Health and Injury Tracking System (HITS) [8]. This potential for injury throughout the kinetic chain emphasizes the importance of a comprehensive approach to injury prevention in addition to pitch count restrictions to allow for players to continue to safely participate in throwing sports. The majority of shoulder injuries in sports are caused by repetitive overhead motion leading to overuse injuries [9, 10]. Injuries in the throwing athlete often lead to missed time for treatment and decreased performance capabilities, highlighting the need for effective injury prevention programs [11, 12].

A clear understanding of the biomechanics of throwing is essential for sports medicine specialists, including athletic trainers, physical therapists, and physicians, to effectively care

This article is part of the Topical Collection on *Sports Injuries and Rehabilitation: Getting Athletes Back to Play*

✉ Drew A. Lansdown
Drew.Lansdown@ucsf.edu

¹ Department of Orthopedic Surgery, Sports Medicine & Shoulder Surgery, University of California, San Francisco, 1500 Owens Street, San Francisco, CA 94158, USA

for throwing athletes. The six phases of throwing include windup, stride, arm cocking, acceleration, deceleration, and follow-through [3, 13]. Deficiencies at any point in this cycle may transfer increased stress to the thrower's shoulder and elbow or may stem from problems with total shoulder rotation, external rotation weakness, and scapular dyskinesis. Understanding the dynamic phases of throwing is imperative to understanding the development of overuse injuries in the overhead athlete [14]. Each phase localizes force to different parts of the body, leading to potential injury in that phase. Given these differential stresses in each phase, there is also an opportunity to target specific exercises for the muscle groups active during a given phase of the throwing cycle.

The purpose of this article is to provide an evidence-based review of injury prevention strategies based on each phase of the throwing cycle that may be incorporated by healthy throwing athletes. The objectives of this paper are to describe the complex motion and muscular activation in each phase of throwing, to demonstrate how variability in throwing mechanics can contribute to sports-related injury, and to identify exercises to target each phase of throwing to limit injury risk.

Windup

The windup phase begins with the pitcher's first movement from the static position of facing the batter with both feet on the mound and is completed when the lead leg reaches maximum knee height [15]. Muscular activations will be seen in the iliopsoas, rectus femoris, pectineus, and sartorius to elevate the stride leg [13]. The final moment during the windup when the knee is at maximum height is referred to as the "balance point" [15]. Subsequently, the pitcher starts to remove the ball from the glove in order to begin the next phase.

The risk of injury during this time is relatively low, but the pitcher is setting the timing and tone for the remainder of the pitching motion [15]. While in this "balance point" position, we should see the pitcher's shoulders aligned between home plate and second base demonstrating a stable center of gravity (COG) [15]. When observing the throwing motion, special attention should be focused on stability at the balance point, as alterations at this early point may result in unnecessarily increased stress through the upper extremity. The primary contributors to successfully completing the windup phase arise from strength and proprioception in the lower extremity.

First, the ankle joint contributes to forming a stable foundation for the stance leg. Athletes with chronic ankle instability demonstrate increased postural sway, which can result in loss of control during the windup phase [16–19]. The primary mechanical impairments from an unstable ankle are increased anterior joint laxity [20], reduced posterior talar glide [21], and reduced range of motion, most notably in decreased dorsiflexion range of motion [22–24]. Restrictions in the anterior-

posterior glide of the talus on the tibia have been well documented in those with lateral ankle instability [25]. These restrictions can be seen in those who have only experienced one incidence of lateral ankle sprain, or those who suffer from chronic ankle instability [25]. Arthokinematic restrictions may not allow the thrower to achieve the necessary dorsiflexion range of motion, which in turn can affect their COG within the windup phase. Dorsiflexion range of motion has been shown to have a significant influence of dynamic balance regardless of a history of lateral ankle sprains [26]. An ankle box stretch (Figure 1) should be a part of every pitcher's warm-up routine to maintain necessary flexibility.

Additionally, the tensor fascia latae, gluteus medius, and gluteus minimus must isometrically contract to maintain a stable pelvis [13]. If the COG is positioned too far posteriorly or anteriorly, torque will transfer to the upper extremity, thus predisposing the shoulder and elbow to injury [15]. Factors that can contribute to faulty windup mechanics include poor balance at maximum knee height secondary to reduced lower extremity strength, poor trunk control, and tilting of the COG [15].

As the hips develop power throughout the windup phase, it is transferred through the lumbopelvic region and finally to the throwing arm [27]. The lumbopelvic muscles show significant activity during the pitching motion [28, 29]. Poor lumbopelvic control has been associated with increased shoulder horizontal abduction torque and elbow valgus torque, both of which may result in increased injury risk and decreased performance [30]. Additionally, there is increased compressive force in the glenohumeral joint with increased pelvic tilt toward the throwing side, increased pelvic axial rotation velocity, and a decreased stride length [31]. From an athlete performance standpoint, Chaudhari et al. [32] found that pitchers with less lumbopelvic control produced more walks and hits per inning when compared to those with more lumbopelvic control. Likewise, pitchers with decreased lumbopelvic control have been shown to have an increased likelihood of injury and spending more time on the disabled list [33].

Hip and core strength are directly linked to dynamic balance [34–39]. Evaluation of stability can be done through tests like the Star Excursion Balance Test (SEBT) and the Y Balance test [40]. Deficiencies in balance are treatable through short courses of training programs. For instance, prior studies have demonstrated significant improvement in SEBT with a range of 5- to 12-week core training protocols [39, 41, 42]. Exercises like banded side-walks ("lateral monster walks") for gluteus medius strength and activation [43] and bird dogs [44] for core stability and strength are easy to incorporate into on-field practice routines without excessive equipment. Forward lunges and lateral lunges are also excellent exercises to activate and strengthen the gluteus medius [43] (Figure 1).






<p>Ankle Box Stretch</p> 	<p>Exercise Description: Place your foot on a bench and use your body weight and arms to force your knee over your toes. It is important to keep your heel in contact with the bench, and to keep your knee straight over your toes. You should feel a stretch in the back of your ankle.</p> <p>Duration: Hold this stretched position for 1 minute</p> <p>Goal: Increase ankle dorsiflexion</p> <p>Phase: Wind-up</p>
<p>Lateral Monster Walk</p> 	<p>Exercise Description: Use a looped rubber band. Secure the band just above your knee. Squat to approximately 45 degrees. While staying in this squat position step one leg out to the side, widening the space between your feet. Then step the other leg in, narrowing the space between your feet. Be sure to maintain a constant tension in the band and not bring your feet too close together.</p> <p>Duration: Perform 2 sets of 30 steps in each direction. Maintain a good squat position throughout the exercise.</p> <p>Goal: Gluteus Medius muscle activation</p> <p>Phase: Wind-up</p>
<p>Forward Lunge</p> 	<p>Exercise Description: Stand with your feet shoulder width apart. Take a long step forward and slowly lower the back knee toward the ground. The front knee should bend to approximately 90 degrees and the patella should track with the second toe. Push off the front leg to return to stranding position. Alternate legs.</p> <p>Duration: Perform 2 sets of 10 repetitions on each leg</p> <p>Goal: Gluteus Medius muscle activation</p> <p>Phase: Wind-up</p>
<p>Lateral Lunge</p> 	<p>Exercise Description: Stand with your feet shoulder width apart. Take a big step to one side, push your hips back while bending your stride knee. Lower your body until your stride knee is bent 90 degrees. Push back to the starting position. Alternate legs.</p> <p>Duration: Perform 2 sets of 10 repetitions on each leg.</p> <p>Goal: Gluteus Medius muscle activation</p> <p>Phase: Wind-up</p>
<p>Bird Dog</p> 	<p>Exercise Description: Begin on all fours in your Table Top Position. Brace your core and then slowly lift and extend your right arm and left leg until they are parallel to the floor. You should squeeze your shoulder blade down and back on the right arm and flex your foot and press your heel rearward to engage your glutes on the left leg. Hold this position for 10 seconds. Return to table top position without touching the ground and repeat. Do all the reps on one side and then switch.</p> <p>Duration: Hold position for 10 seconds. Perform 10 repetitions on each side.</p> <p>Goal: Core muscle activation</p> <p>Phase: Wind-up</p>

Figure 1 Specific stretches and exercises to help assist with flexibility, hip strength, and core strength

Stride

The stride phase begins with the throwing shoulder horizontally abducting to approximately 90° and ends with the front foot striking the ground [13]. From the ground up, the athlete will be engaging their ankles, legs, pelvis, core, thoracic spine, and shoulder [15, 45, 46]. Again, especially with sufficient

injury history, a clinician may begin to see breakdowns within the throwing motion and kinetic chain already. Areas of common concern at this point include core strength, lumbar extension, and thoracic spine rotation. Deficiencies in the kinetic chain in these early portions of a throwing motion can change mechanics later on through the “catch-up phenomenon” and result in an altered delivery and possible injury [15, 45, 46]. Low back pain (LBP) and/or injury may develop as a consequence to poor core muscle activation/stability, which in turn may cause deficiencies later within the throwing motion [47].

LBP is a significant issue for baseball players, as it can lead to missed participation time and early career termination [48]. The prevalence of LBP in active baseball players ranges between 3 and 15% [49–53]. This pain can present itself in forms of stress reactions, stress fractures, vertebral disc degeneration, and mechanical LBP involving surrounding musculature. Each is products of repetitive, high-velocity spinal movement, and loading. For instance, Hangai et al. found that 59.7% of the tested population of Japanese baseball players had radiographic disc degeneration at one or more levels [54]. Improper mechanics may cause forces to concentrate in these regions and subsequently place throwers at risk for injury. Toyoshima et al. reported that the trunk contributes to as much as 50% of the kinetic energy and force production during the entire throwing motion [55]. Wasser et al. [56] discussed the importance of maintaining a neutral spine by avoiding excessive lumbar extension and rotation during the stride and early cocking phases.

To maintain a neutral spine through the throwing cycle, throwers must maintain lumbar stability and thoracic mobility. Additionally, the risk of LBP may be limited by combining core stability with thoracic mobility exercises better than using core exercises alone [57–60]. This approach effectively reduces the stress caused by excessive movement of segments from lumbar instability by mobilizing segments above, as found by Yang et al., Kaltenborn et al., and Sung YB et al. [57, 59, 60]. During the stride and cocking phase, instead of motion and force coming from the lumbar region, causing excessive lumbar extension and rotation, thoracic flexibility may allow for force distribution and decrease stress concentration.

Specific exercises that allow for proper lumbar spine stability and thoracic spine mobility include bird dogs [44], pallof presses, lifts, and chops, in addition to thoracic spine rotation mobilization, and thoracic spine extension mobilization (Figure 2).

Cocking

The cocking phase begins with the front foot striking the ground and ends with maximum shoulder external rotation at 150 to 180° [61, 62]. This stage of throwing can be further




Figure 2 Specific exercises that allow for proper lumbar spine stability and thoracic spine mobility

divided into early and late cocking phases. Potential energy is accumulated in the early cocking phase and transferred to the throwing arm in the late cocking phase to prepare for acceleration and ball release [63].

Early Cocking

The early cocking phase begins with lead foot contact. The quadriceps of the lead leg contracts to stabilize a fulcrum point [61, 64]. The pelvis then rotates toward home plate. Trunk rotation and extension lag behind pelvic rotation, transferring energy from the pelvis to the upper torso [3, 65]. During rotation, the abdominal and oblique musculature is activated to stabilize the trunk through the delay between pelvic and upper torso rotation [61].

Significant shoulder muscle activity is required to stabilize the throwing arm as the trunk rotates and extends. During this phase, the deltoid muscle activates to maintain 90° abduction of the throwing arm with the elbow flexed at 90 to 100° [62, 64]. The rotator cuff muscles achieve high activity to resist the compressive force generated by the trunk [62]. The shoulder girdle muscles (levator scapulae, serratus anterior, trapezius, rhomboids, and pectoralis minor) are activated to stabilize the scapula and glenoid for subsequent humeral head external rotation [61]. Importantly, the scapula must protract and rotate upwards to ensure that the humeral head is positioned in the “safe zone” on the glenoid as rotation of the throwing arm lags behind the torso [66].

Scapular dyskinesia describes a disruption of normal scapular kinematics and is a major cause of injury during the early cocking phase. At rest, excessive scapulothoracic protraction and upward rotation lead to misalignment between the scapula and glenoid [66]. While throwing, abnormal scapular movement results in a loss of coordination between the glenohumeral and scapulothoracic joints. Scapular dyskinesia has been associated with shoulder pain, shoulder impingement syndrome, rotator cuff tendinopathy, and rotational deficits which disturb the scapulohumeral rhythm [67, 68]. A meta-analysis found that asymptomatic athletes with shoulder dyskinesia had a 43% higher chance of developing shoulder pain than athletes without scapular abnormalities [69].

Scapular stabilization exercises are effective both to treat and to prevent scapular dyskinesia and secondary shoulder injuries in throwing athletes. For athletes with scapular dyskinesia, exercises to counteract the abnormal protraction, depression, and rotation of the scapula are indicated [70, 71]. In healthy athletes, strengthening exercises for scapular stabilizers including the upper and lower trapezius and serratus anterior muscles promote normal scapular movement. The low row, inferior glide, lawnmower, and robbery exercises

have been shown to effectively activate and strengthen these muscle groups in asymptomatic and symptomatic patients [68, 71, 72]. Incorporating these exercises into routine on-field training will promote appropriate scapular stabilization and glenoid positioning during the early cocking phase without disrupting the normal scapulohumeral rhythm (Figure 3).

Late Cocking

During the late cocking phase, potential energy from the trunk is transferred to the throwing arm as it externally rotates and horizontally adducts. The late cocking phase prepares the arm

for forward acceleration and subsequent ball release. The degree of external rotation achieved in this phase determines the range of forward movement in the following stages with greater external rotation leading to increased ball velocity [61]. The late cocking phase ends when the throwing arm is maximally externally rotated to 150 to 180° and horizontally adducted to approximately 20° [15, 62].

The infraspinatus and teres minor muscles contract concentrically to externally rotate the shoulder. As the arm externally rotates, the shoulder internal rotators (subscapularis, teres major, pectoralis major) contract eccentrically to control the speed of rotation. The posterior rotator cuff muscles (infraspinatus and teres minor) and latissimus dorsi contract to generate a posterior force which resists anterior humeral head translation and supports the anterior capsule [62]. The pectoralis major and anterior deltoid muscles also contract concentrically to horizontally adduct the throwing arm with peak angular velocity of 600° per second [15, 62].

Athletes often complain of shoulder pain in the late cocking position, where the throwing arm is abducted to 90°, the elbow is flexed to 90°, and the arm is maximally externally rotated in preparation for forward acceleration. In this position, the throwing arm holds its maximum potential energy [64]. Shoulder pain in this case can limit the athlete’s range of motion in external rotation and lead to decreased ball velocity during the acceleration phase [73]. Among professional pitchers, preseason deficits in external rotation are associated with in-season injury [74]. Pitchers with insufficient external rotation are more likely to be placed on the disabled list for a shoulder injury and require shoulder surgery than pitchers with deficits in internal or total shoulder rotation [75].

Having greater flexibility in external rotation can benefit throwing performance, but excessive stretching can exacerbate capsule laxity and lead to shoulder instability. Wilk refers to these competing interests as the “thrower’s paradox” [70, 76]. Athletes with inadequate external rotation may benefit from stretching exercises to promote increased range of motion, but repetitive throwing motions with extreme external rotation can lead to capsular instability and injury to the labrum or rotator cuff muscles [61]. For throwers with excessive laxity, one strategy to combat this paradox is increasing range of motion elsewhere in the chain. Increasing thoracic spine mobility in extension and rotation toward the pitching side may eliminate the need for excessive glenohumeral external rotation range of motion and help avoid resultant anterior capsule laxity and additional stress on the long head of the biceps tendon (Figure 3).

A proper warm-up routine to prevent pain with external rotation and capsular laxity during the late cocking phase should encompass thoracic spine mobility as well as facilitate activation of shoulder stabilizers including the middle and








<p>Low Row</p> 	<p>Exercise Description: Stand with good posture in front of a surface. Place your hand on the surface with the palm facing backward and elbow extended. Apply maximal force to the surface while pulling your scapula down and back. You should feel the lower to mid trapezius and rhomboids activated. Hold this position for three seconds, then slowly release.</p> <p>Duration: Perform 10 repetitions of maximal force.</p> <p>Goal: Activation of posterior deltoid, serratus anterior, scapular stabilizers muscular activation.</p> <p>Phase: Cocking</p>
<p>Inferior Glide</p> 	<p>Exercise Description: Sit upright with one arm extended to the side with fist supported on a surface at approximately chest height. Apply maximal downward force to the surface while pulling the scapula down. Focus on activating your lower trapezius muscles. Hold for 3 seconds, then slowly release.</p> <p>Duration: Perform 10 repetitions of maximal force.</p> <p>Goal: Serratus anterior, lower trapezius, and scapular stabilizers muscular activation</p> <p>Phase: Cocking</p>
<p>Lean Mower</p> 	<p>Exercise Description: Stand with trunk flexed, knees bent, and hand positioned at the contralateral thigh/knee. Rotate and extend the trunk while raising the arm and retracting the scapula. Focus on keeping the movement fluid and controlled. Hold at maximal retraction for three seconds, then slowly release.</p> <p>Duration: Perform 10 repetitions</p> <p>Goal: Serratus anterior, lower trapezius, posterior deltoid, scapular stabilizer, and core musculature activation</p> <p>Phase: Cocking</p>
<p>Rubber</p> 	<p>Exercise Description: Attach exercise band at chest height and stand with good posture with your core engaged. Arms out from the side at 40 degrees. Elbows remain bent at 90. Pinch shoulder blades together, rotating arms out, keeping elbows bent. Hold lower and middle trapezius contraction for 5 seconds</p> <p>Duration: Perform 1 set of 10 repetitions</p> <p>Goal: Posterior deltoid, posterior rotator cuff, lower and middle trapezius, and scapular stabilizers muscular activation</p> <p>Phase: Cocking</p>

Figure 3 Incorporating these exercises into routine on-field training will promote appropriate scapular stabilization and glenoid positioning

Figure 4 Stretches used to isolate the posterior aspect of the shoulder and triceps

<p style="text-align: center;"><u>Sleeper Stretch</u></p> 	<p>Exercise Description: Lay on your side with shoulder flexed in front of your body. Bend elbow to 90 degrees. Slowly apply downward pressure with opposite arm.</p> <p>Duration: Perform for 3 sets of 60 seconds</p> <p>Goal: Stretching of the posterior capsule</p> <p>Phase: Acceleration</p>
<p style="text-align: center;"><u>Cross Body Stretch</u></p> 	<p>Exercise Description: Grab one arm above your elbow with your opposite hand, and pull it across your body toward your chest until you feel a stretch in your shoulder. Make sure to keep your shoulder blade retracted by engaging your lower/middle trapezius</p> <p>Duration: Hold for 1 minute</p> <p>Goal: Stretching of the posterior capsule and posterior deltoid</p> <p>Phase: Acceleration</p>
<p style="text-align: center;"><u>Overhead Triceps Stretch</u></p> 	<p>Exercise Description: Standing up straight with a tight core, extend your arm straight into the air. Keep the elbow up as you bend your arm behind your head. Take the opposite hand and gently pull your elbow down and towards the opposite side.</p> <p>Duration: Hold stretch for 1 minute</p> <p>Goal: Stretching of the triceps brachii muscle</p> <p>Phase: Acceleration</p>

lower trapezius, rotator cuff musculature, biceps, and pec minor. With this in mind, we can address potential pain with external rotation and capsular laxity through the aforementioned thoracic spine rotation mobilization and thoracic spine extension mobilization in order to increase thoracic spine mobility toward the pitching side. In addition, the low row, inferior glide, lawn mower, and robber exercises can facilitate activation of the scapular stabilizing musculature [68, 71, 72].


Acceleration

The acceleration phase takes place from maximal external rotation to the moment of ball release [47]. The potential energy accumulated by the throwing arm is utilized to accelerate the ball to its maximal velocity. The trunk flexes forward to neutral position as the throwing arm rotates internally and the elbow extends. Arm rotation lags behind elbow extension,

with maximal elbow velocity occurring halfway through the acceleration phase and maximal internal rotation velocity occurring at ball release [3, 61, 77]. This delay reduces the inertia of the shoulder to increase the torque and angular velocity of the throwing arm [3]. The mean angular velocity of the arm at ball release is approximately 7,000° per second, making this one of the fastest human movements [3, 10, 47].

Muscle recruitment in this phase facilitates rapid rotation and extension of the arm. The quadriceps of the lead leg contracts concentrically to extend the leg. The trunk flexors (rectus abdominis, obliques) tilt the trunk forward to allow the throwing arm to accelerate through a greater distance [47, 61]. Internal rotation of the shoulder to 90 or 100° is achieved via concentric activation of the internal rotators (latissimus dorsi, pectoralis). The rotator cuff muscles, trapezius, serratus anterior, and levator scapulae remain active to stabilize the scapula and glenohumeral joint [15, 61]. Of note, rotator cuff and biceps activation

Figure 5 A banded reverse throw allows for activation of the scapular stabilizers, warming up the posterior deltoid and rotator cuff, and preparing the shoulder to decelerate a throw by eccentrically resisting a band



is up to three times higher in amateur pitchers than in professional pitchers during this phase, potentially contributing to overuse injuries in young athletes [15]. Extension of the elbow during ball acceleration is achieved via the centrifugal force generated by the trunk and concentric activation of elbow extensors. Deceleration of elbow extension at ball release is achieved via eccentric activation of elbow flexors (biceps brachii, brachialis, and brachioradialis) [47, 61]. Wrist flexor muscles (flexor carpi radialis, flexor carpi ulnaris, and flexor digitorum) shift the wrist from hyperextension to neutral position at ball release [10, 47, 61].

Most elbow injuries among overhead athletes result from the substantial stresses applied to the elbow during the late cocking and acceleration phases. Injury to the ulnar collateral

ligament (UCL) is common because the valgus stress applied to the medial elbow during ball acceleration exceeds the tensile strength of the UCL at 64 N·m [10, 78, 79]. Sidarm tracking due to decreased trunk flexion contributes to UCL trauma by increasing the force applied to the medial elbow [15]. UCL injury results in problems including elbow pain, decreased throwing velocity and control, joint instability, and muscle weakness [7, 10, 80]. Laxity of the UCL can also lead to additional ligamentous tears and ulnar nerve injury [79].

The medial shear force of 300 N and compressive force of 500 N applied to the elbow can also result in valgus extension overload syndrome (VEO) with impingement of the posteromedial elbow [10, 79]. This syndrome is characterized by olecranon tip osteophytes, loose bodies, and damage to the posteromedial trochlea [79, 81]. Athletes with VEO often

experience posterior elbow pain with elbow extension. VEO is common among baseball players who undergo elbow surgery with 65% of patients diagnosed with posterolateral olecranon osteophytes [82].

Elbow injury in overhand throwers often occurs secondary to adaptive changes in the shoulder which result in excessive force on the elbow. Pitchers often demonstrate an increase in external rotation, loss of internal rotation, and decrease in total range of motion of the throwing shoulder [83–87]. While these deficits are often described chronically, changes in shoulder range of motion have also been observed acutely after pitching. Reinold et al. described significant changes in shoulder and elbow range of motion within 30 min of pitching [85]. Kilber et al. similarly found significant and sustained loss of internal rotation up to 72 h after throwing [88]. Garrison and colleagues further demonstrated that deficits in total shoulder rotation range of motion are associated with UCL tears in high school and collegiate baseball players [84]. The acute time course of these adaptations suggests that stretching before and after a pitching session may be effective to prevent long-term changes.

Drills to prevent elbow injury in overhead athletes act by maintaining 180° total shoulder range of motion. The sleeper stretch is used to isolate the posterior aspect of the shoulder (posterior capsule, deltoid, and latissimus dorsi) and has been shown to effectively recover internal range of motion after pitching [89] (Figure 4). Other techniques including the cross-body stretch and overhead triceps stretch can also prevent glenohumeral internal rotation deficit (GIRD) [90] (Figure 4). A prevention program involving the cross-body stretch, overhead triceps stretch, and sleeper stretch among other exercises

was found to significantly reduce the incidence of medial elbow injury and shoulder injury while improving hip and thoracic flexibility in youth pitchers [12, 91]. A separate study found that in a cohort of twenty pitchers, the two-out drill was effective to restore internal, external, and total range of motion of the throwing shoulder after a 40-pitch session, which can be incorporated in-game, between innings [92].

Deceleration

The deceleration phase takes place between the time of ball release and maximal humeral head internal rotation with elbow extension [47]. During this last point of contact prior to completing the pitch, there are 0° of glenohumeral rotation, 100° of shoulder abduction, and 35° of horizontal adduction [93]. The teres minor, infraspinatus, and posterior deltoid are responsible for slowing the shoulder down and dissipating compressive forces across the joint [47]. Requiring a large eccentric contraction, the posterior musculature and posterior capsule are repeatedly placed in situations of potential injury [94]. In a clinical setting, these areas facing repeated trauma, coupled with continued anterior capsule stretching with the external rotation during the late cocking phase, pose a risk for glenohumeral internal rotation deficit (GIRD).

The biceps brachii and brachialis are also active during deceleration by contracting eccentrically to slow down elbow extension and forearm pronation [3]. Other structures to take note of are the trapezius, rhomboids, and serratus anterior, as they all assist in the deceleration phase and help the thrower stabilize

Table 1 A thorough understanding of the mechanics of this motion, along with specific exercises to target the specific actions of each phase as described here, may allow for throwers to minimize injury risk

Phase	Notable active musculature	Potential concerns	Warm-up exercises
Windup	Iliopsoas, rectus femoris, pectineus, sartorius, tensor fascia latae, gluteus medius, gluteus minimus, core	Center of gravity, ankle dorsiflexion, lumbopelvic control	Ankle box stretch, lateral monster walk, forward lunge, lateral lunge, bird dog
Stride	Tensor fascia latae, gluteus medius, gluteus minimus, core	Lumbar hypermobility, Thoracic hypomobility	Pallof press, lift, chop, thoracic spine extension mobilization, thoracic spine rotation mobilization
Cocking	Deltoid, rotator cuff, levator scapulae, serratus anterior, trapezius, rhomboid, pectoralis minor	Scapular dyskinesis, glenohumeral capsular laxity	Low row, inferior glide, lawn mower, robber
Acceleration	Latissimus dorsi, pectoralis, rotator cuff, trapezius, serratus anterior, levator scapulae, biceps brachii	Glenohumeral internal rotation deficit	Sleeper stretch, cross body stretch, overhead triceps stretch
Deceleration	Teres minor, infraspinatus, posterior deltoid, biceps brachii, brachialis, trapezius, rhomboid, serratus anterior	Glenohumeral internal rotation deficit, subacromial impingement, biceps brachii tendinitis, posterior rotator cuff tendinitis, posterior capsule tightening	Reverse throw
Follow-through	Culmination of kinetic chain delivering the pitch. Low risk of injury during this phase		

their scapula throughout movement [47]. Without proper kinematics, muscle activation, and form, there will be a higher risk of injury during this time. A comprehensive warm-up routine is one avenue to combat this increased injury risk.

A focus on scapular stability and posterior deltoid strength is paramount for a successful deceleration phase in order to combat the extreme rotational and distraction forces being placed on the shoulder. GIRD, along with subacromial impingement, biceps brachii tendinitis, posterior rotator cuff tendinitis, and posterior capsule tightening are just some of the injuries that can be seen within a baseball throwing shoulder as a result of overuse, and stress placed during this phase.

The low row, inferior glide, lawnmower, and robbery exercises as used in the early cocking phases are helping prime the shoulder for the violent motion of a pitch and activate the musculature stated above. Additionally, a banded reverse throw allows for activation of the scapular stabilizers, warming up the posterior deltoid and rotator cuff, and preparing the shoulder to decelerate a throw by eccentrically resisting a band (Figure 5).

Follow-through

The follow-through phase is the culmination of the kinetic chain: linking together, generating forces, and delivering the pitch. Due to decreased joint loading during this phase, the risk of injury is reduced relative to other phases. The pitcher will simply continue to move forward toward the catcher until arm motion has ceased, and then become a fielder for the remainder of the play.

Conclusion

The throwing cycle is a complex motion that places various stresses throughout the thrower's body, from the ankle to the core, and from the back to the shoulder and elbow. A thorough understanding of the mechanics of this motion, along with specific exercises to target the specific actions of each phase as described here, may allow for throwers to minimize injury risk (Table 1).

Availability of Data and Material All data was obtained from published articles online.

Code Availability Not applicable

Author Contribution Michael Mayes—conceptualization, methodology, literature search, writing—original draft preparation

Madeleine Salesky—conceptualization, methodology, literature search, writing—original draft preparation

Drew Lansdown—conceptualization, methodology, literature search, writing—reviewing and editing

Declarations

Conflict of Interest Michael Mayes and Madeleine Salesky declare that they have no conflicts of interest.

Drew Lansdown has the following disclosures: Received fellowship-related research/education funding from Arthrex, Inc. and Smith & Nephew; educational support from Medwest; hospitality payments from Wright Medical; Arthroscopy Association of North America: Board or committee member, American Orthopedic Society for Sports Medicine: Board or committee member.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

Papers of particular interest, published recently, have been highlighted as:

- Of importance

1. Norton R, Honstad C, Joshi R, Silvis M, Chinchilli V, Dhawan A. Risk Factors for Elbow and Shoulder Injuries in Adolescent Baseball Players: A Systematic Review. *Am J Sports Med.* 2019;47(4):982–990. <https://doi.org/10.1177/0363546518760573>.
2. Conte SA, Fleisig GS, Dines JS, Wilk KE, Aune KT, Patterson-Flynn N, et al. Prevalence of Ulnar Collateral Ligament Surgery in Professional Baseball Players. *Am J Sports Med.* 2015;43(7):1764–9. <https://doi.org/10.1177/0363546515580792>.
3. Dillman CJ, Fleisig GS, Andrews JR. Biomechanics of pitching with emphasis upon shoulder kinematics. *J Orthop Sports Phys Ther.* 1993;18(2):402–8. <https://doi.org/10.2519/jospt.1993.18.2.402>.
4. Lyman S, Fleisig GS, Waterbor JW, Funkhouser EM, Pulley L, Andrews JR, et al. Longitudinal study of elbow and shoulder pain in youth baseball pitchers. *Med Sci Sports Exerc.* 2001;33(11):1803–10.
5. Olsen SJ 2nd, Fleisig GS, Dun S, Loftice J, Andrews JR. Risk factors for shoulder and elbow injuries in adolescent baseball pitchers. *Am J Sports Med.* 2006;34(6):905–12. <https://doi.org/10.1177/0363546505284188>.
6. Fleisig GS, Andrews JR, Dillman CJ, Escamilla RF. Kinetics of baseball pitching with implications about injury mechanisms. *Am J Sports Med.* 1995;23(2):233–9.
7. Bruce JR, Andrews JR. Ulnar collateral ligament injuries in the throwing athlete. *J Am Acad Orthop Surg.* 2014;22(5):315–25. <https://doi.org/10.5435/JAAOS-22-05-315>.
8. Camp CL, Dines JS, van der List JP, Conte S, Conway J, Altchek DW, et al. Summative Report on Time Out of Play for Major and Minor League Baseball: An Analysis of 49,955 Injuries From 2011 Through 2016. *Am J Sports Med.* 2018;46(7):1727–32. <https://doi.org/10.1177/0363546518765158>.

9. Anderson MW, Alford BA. Overhead throwing injuries of the shoulder and elbow. *Radiol Clin North Am.* 2010;48(6):1137–54. <https://doi.org/10.1016/j.rcl.2010.07.002>.
10. Patel NB, Thomas S, Lazarus ML. Throwing injuries of the upper extremity. *Radiol Clin North Am.* 2013;51(2):257–77. <https://doi.org/10.1016/j.rcl.2012.09.016>.
11. Melugin HP, Leafblad ND, Camp CL, Conte S. Injury prevention in baseball: from youth to the pros. *Curr Rev Musculoskelet Med.* 2018;11(1):26–34 **Melugin et al. explain in detail how injuries in the throwing athlete often lead to missed time for treatment and decreased performance capabilities, which highlights the need for effective injury prevention programs.**
12. Sakata J, Nakamura E, Suzuki T, Suzukawa M, Akaie A, Shimizu K, et al. Efficacy of a prevention program for medial elbow injuries in youth baseball players. *Am J Sports Med.* 2018;46(2):460–9.
13. Weber AE, Kontaxis A, O'Brien SJ, Bedi A. The biomechanics of throwing: simplified and cogent. *Sports Med Arthrosc Rev.* 2014;22(2):72–9. <https://doi.org/10.1097/JSA.000000000000019>.
14. Lin DJ, Wong TT, Kazam JK. Shoulder Injuries in the Overhead-Throwing Athlete: Epidemiology, Mechanisms of Injury, and Imaging Findings. *Radiology.* 2018;286(2):370–87. <https://doi.org/10.1148/radiol.2017170481> **Lin et al. detail how understanding the dynamic phases of throwing is imperative to understanding the development of overuse injuries in the overhead athlete.**
15. Calabrese GJ. Pitching mechanics, revisited. *Int J Sports Phys Ther.* 2013;8(5):652–60.
16. Fu AS, Hui-Chan CW. Ankle joint proprioception and postural control in basketball players with bilateral ankle sprains. *Am J Sports Med.* 2005;33(8):1174–82. <https://doi.org/10.1177/0363546504271976>.
17. Wikstrom EA, Hubbard TJ. Talar positional fault in persons with chronic ankle instability. *Arch Phys Med Rehabil.* 2010;91(8):1267–71. <https://doi.org/10.1016/j.apmr.2010.04.022>.
18. Nyska M, Shabat S, Simkin A, Neeb M, Matan Y, Mann G. Dynamic force distribution during level walking under the feet of patients with chronic ankle instability. *Br J Sports Med.* 2003;37(6):495–7. <https://doi.org/10.1136/bjism.37.6.495>.
19. Morrison KE, Hudson DJ, Davis IS, Richards JG, Royer TD, Dierks TA, et al. Plantar pressure during running in subjects with chronic ankle instability. *Foot Ankle Int.* 2010;31(11):994–1000. <https://doi.org/10.3113/FAI.2010.0994>.
20. Hubbard TJ, Kaminski TW, Vander Griend RA, Kovalski JE. Quantitative assessment of mechanical laxity in the functionally unstable ankle. *Med Sci Sports Exerc.* 2004;36(5):760–6. <https://doi.org/10.1249/01.mss.0000126604.85429.29>.
21. Denegar CR, Hertel J, Fonseca J. The effect of lateral ankle sprain on dorsiflexion range of motion, posterior talar glide, and joint laxity. *J Orthop Sports Phys Ther.* 2002;32(4):166–73. <https://doi.org/10.2519/jospt.2002.32.4.166>.
22. Drewes LK, McKeon PO, Kerrigan DC, Hertel J. Dorsiflexion deficit during jogging with chronic ankle instability. *J Sci Med Sport.* 2009;12(6):685–7. <https://doi.org/10.1016/j.jsams.2008.07.003>.
23. Youdas JW, McLean TJ, Krause DA, Hollman JH. Changes in active ankle dorsiflexion range of motion after acute inversion ankle sprain. *J Sport Rehabil.* 2009;18(3):358–74. <https://doi.org/10.1123/jsr.18.3.358>.
24. Kim SG, Kim WS. Effect of Ankle Range of Motion (ROM) and Lower-Extremity Muscle Strength on Static Balance Control Ability in Young Adults: A Regression Analysis. *Med Sci Monit.* 2018;24:3168–75. <https://doi.org/10.12659/MSM.908260>.
25. Hertel J, Corbett RO. An Updated Model of Chronic Ankle Instability. *J Athl Train.* 2019;54(6):572–88. <https://doi.org/10.4085/1062-6050-344-18>.
26. Hoch MC, Staton GS, McKeon PO. Dorsiflexion range of motion significantly influences dynamic balance. *J Sci Med Sport.* 2011;14(1):90–2. <https://doi.org/10.1016/j.jsams.2010.08.001>.
27. Hirashima M, Yamane K, Nakamura Y, Ohtsuki T. Kinetic chain of overarm throwing in terms of joint rotations revealed by induced acceleration analysis. *J Biomech.* 2008;41(13):2874–83. <https://doi.org/10.1016/j.jbiomech.2008.06.014>.
28. Oliver GD, Keeley DW. Gluteal muscle group activation and its relationship with pelvis and torso kinematics in high-school baseball pitchers. *J Strength Cond Res.* 2010;24(11):3015–22. <https://doi.org/10.1519/JSC.0b013e3181c865ce>.
29. Oliver GD, Weimar WH, Plummer HA. Gluteus medius and scapula muscle activations in youth baseball pitchers. *J Strength Cond Res.* 2015;29(6):1494–9. <https://doi.org/10.1519/JSC.0000000000000797>.
30. Laudner KG, Wong R, Meister K. The influence of lumbopelvic control on shoulder and elbow kinetics in elite baseball pitchers. *J Shoulder Elbow Surg.* 2019;28(2):330–4. <https://doi.org/10.1016/j.jse.2018.07.015> **Laudner et al. demonstrate the importance of addressing the entire kinetic chain when developing injury prevention strategies. With increased lumbopelvic control, they found a decreased risk of injury at the shoulder and elbow.**
31. Keeley DW, Oliver GD, Dougherty CP, Tory MR. Lower body predictors of glenohumeral compressive force in high school baseball pitchers. *J Appl Biomech.* 2015;31(3):181–8. <https://doi.org/10.1123/jab.2011-0229>.
32. Chaudhari AM, McKenzie CS, Borchers JR, Best TM. Lumbopelvic control and pitching performance of professional baseball pitchers. *J Strength Cond Res.* 2011;25(8):2127–32. <https://doi.org/10.1519/JSC.0b013e31820f5075>.
33. Chaudhari AM, McKenzie CS, Pan X, Onate JA. Lumbopelvic control and days missed because of injury in professional baseball pitchers. *Am J Sports Med.* 2014;42(11):2734–40. <https://doi.org/10.1177/0363546514545861>.
34. Ambegaonkar JP, Mettinger LM, Caswell SV, Burt A, Cortes N. Relationships between core endurance, hip strength, and balance in collegiate female athletes. *Int J Sports Phys Ther.* 2014;9(5):604–16.
35. Haruyama K, Kawakami M, Otsuka T. Effect of Core Stability Training on Trunk Function, Standing Balance, and Mobility in Stroke Patients. *Neurorehabil Neural Repair.* 2017;31(3):240–9. <https://doi.org/10.1177/1545968316675431>.
36. Hodges PW, Richardson CA. Contraction of the abdominal muscles associated with movement of the lower limb. *Phys Ther.* 1997;77(2):132–42; discussion 42–4. <https://doi.org/10.1093/ptj/77.2.132>.
37. Zazulak BT, Hewett TE, Reeves NP, Goldberg B, Cholewicki J. Deficits in neuromuscular control of the trunk predict knee injury risk: a prospective biomechanical-epidemiologic study. *Am J Sports Med.* 2007;35(7):1123–30. <https://doi.org/10.1177/0363546507301585>.
38. López-Valenciano A, Ayala F, De Ste CM, Barbado D, Vera-García FJ. Different neuromuscular parameters influence dynamic balance in male and female football players. *Knee Surg Sports Traumatol Arthrosc.* 2019;27(3):962–70. <https://doi.org/10.1007/s00167-018-5088-y>.
39. Sandrey MA, Mitzel JG. Improvement in dynamic balance and core endurance after a 6-week core-stability-training program in high school track and field athletes. *J Sport Rehabil.* 2013;22(4):264–71. <https://doi.org/10.1123/jsr.22.4.264>.
40. Powden CJ, Dodds TK, Gabriel EH. THE RELIABILITY OF THE STAR EXCURSION BALANCE TEST AND LOWER QUARTER Y-BALANCE TEST IN HEALTHY ADULTS: A SYSTEMATIC REVIEW. *Int J Sports Phys Ther.* 2019;14(5):683–94.

41. Bashir SF, Nuhmani S, Dhall R, Muaidi QI. Effect of core training on dynamic balance and agility among Indian junior tennis players. *J Back Musculoskelet Rehabil.* 2019;32(2):245–52. <https://doi.org/10.3233/BMR-170853>.
42. Imai A, Kaneoka K, Okubo Y, Shiraki H. Effects of two types of trunk exercises on balance and athletic performance in youth soccer players. *Int J Sports Phys Ther.* 2014;9(1):47–57.
43. Ebert JR, Edwards PK, Fick DP, Janes GC. A Systematic Review of Rehabilitation Exercises to Progressively Load the Gluteus Medius. *J Sport Rehabil.* 2017;26(5):418–36. <https://doi.org/10.1123/jsr.2016-0088>.
44. McGill SM, Karpowicz A. Exercises for spine stabilization: motion/motor patterns, stability progressions, and clinical technique. *Arch Phys Med Rehabil.* 2009;90(1):118–26. <https://doi.org/10.1016/j.apmr.2008.06.026>.
45. van der Hoeven H, Kibler WB. Shoulder injuries in tennis players. *Br J Sports Med.* 2006;40(5):435–40; discussion 40. <https://doi.org/10.1136/bjsm.2005.023218>.
46. Kibler WB, Wilkes T, Sciascia A. Mechanics and pathomechanics in the overhead athlete. *Clin Sports Med.* 2013;32(4):637–51. <https://doi.org/10.1016/j.csm.2013.07.003>.
47. Seroyer ST, Nho SJ, Bach BR, Bush-Joseph CA, Nicholson GP, Romeo AA. The kinetic chain in overhand pitching: its potential role for performance enhancement and injury prevention. *Sports Health.* 2010;2(2):135–46. <https://doi.org/10.1177/1941738110362656>.
48. Dick R, Sauers EL, Agel J, Keuter G, Marshall SW, McCarty K, et al. Descriptive epidemiology of collegiate men's baseball injuries: National Collegiate Athletic Association Injury Surveillance System, 1988-1989 through 2003-2004. *J Athl Train.* 2007;42(2):183–93.
49. Bono CM. Low-back pain in athletes. *J Bone Joint Surg Am.* 2004;86(2):382–96. <https://doi.org/10.2106/00004623-200402000-00027>.
50. d'Hemecourt PA, Gerbino PG 2nd, Micheli LJ. Back injuries in the young athlete. *Clin Sports Med.* 2000;19(4):663–79. [https://doi.org/10.1016/s0278-5919\(05\)70231-3](https://doi.org/10.1016/s0278-5919(05)70231-3).
51. Dick R, Agel J, Marshall SW. National Collegiate Athletic Association Injury Surveillance System commentaries: introduction and methods. *J Athl Train.* 2007;42(2):173–82.
52. Posner M, Cameron KL, Wolf JM, Belmont PJ Jr, Owens BD. Epidemiology of Major League Baseball injuries. *Am J Sports Med.* 2011;39(8):1676–80. <https://doi.org/10.1177/0363546511411700>.
53. Wasser JG, Zaremski JL, Herman DC, Vincent HK. Prevalence and proposed mechanisms of chronic low back pain in baseball: part I. *Res Sports Med.* 2017;25(2):219–30. <https://doi.org/10.1080/15438627.2017.1282361>.
54. Hangai M, Kaneoka K, Hinotsu S, Shimizu K, Okubo Y, Miyakawa S, et al. Lumbar intervertebral disk degeneration in athletes. *Am J Sports Med.* 2009;37(1):149–55. <https://doi.org/10.1177/0363546508323252>.
55. Toyoshima S, Hoshikawa, T, Miyashita, M, and Oguri T. The contribution of body parts to throwing performance. In: Nelson, R.C., Morehouse, C.A. (eds) *Biomechanics IV. International Series on Sport Sciences.* Palgrave, London. 1974:169-174. https://doi.org/10.1007/978-1-349-02612-8_24.
56. Wasser JG, Zaremski JL, Herman DC, Vincent HK. Assessment and rehabilitation of chronic low back pain in baseball: part II. *Res Sports Med.* 2017;25(2):231–43. <https://doi.org/10.1080/15438627.2017.1282362>.
57. Yang SR, Kim K, Park SJ, Kim K. The effect of thoracic spine mobilization and stabilization exercise on the muscular strength and flexibility of the trunk of chronic low back pain patients. *J Phys Ther Sci.* 2015;27(12):3851–4. <https://doi.org/10.1589/jpts.27.3851>.
58. Heo MY, Kim K, Hur BY, Nam CW. The effect of lumbar stabilization exercises and thoracic mobilization and exercises on chronic low back pain patients. *J Phys Ther Sci.* 2015;27(12):3843–6. <https://doi.org/10.1589/jpts.27.3843>.
59. Kaltenborn FM. *Manual mobilization of the extremity joint: basic evaluation and treatment techniques.* Oslo: Olaf norisbokhandel; 1989. p. 23–48.
60. Sung YB, Lee JH, Park YH. Effects of thoracic mobilization and manipulation on function and mental state in chronic lower back pain. *J Phys Ther Sci.* 2014;26(11):1711–4. <https://doi.org/10.1589/jpts.26.1711>.
61. Fleisig GS, Barrentine SW, Escamilla RF, Andrews JR. Biomechanics of Overhand Throwing with Implications for Injuries. *Sports Medicine.* 1996;21(6):421–37. <https://doi.org/10.2165/00007256-199621060-00004>.
62. Escamilla RF, Andrews JR. Shoulder muscle recruitment patterns and related biomechanics during upper extremity sports. *Sports Med.* 2009;39(7):569–90. <https://doi.org/10.2165/00007256-200939070-00004>.
63. Dououguih WA, Dolce DL, Lincoln AE. Early Cocking Phase Mechanics and Upper Extremity Surgery Risk in Starting Professional Baseball Pitchers. *Orthop J Sports Med.* 2015;22(4). <https://doi.org/10.1177/2325967115581594>.
64. Chorley J, Eccles RE, Scurfield A. Care of Shoulder Pain in the Overhead Athlete. *Pediatr Ann.* 2017;46(3):e112–e3. <https://doi.org/10.3928/19382359-20170216-01>.
65. Chalmers PN, Wimmer MA, Verma NN, Cole BJ, Romeo AA, Cvetanovich GL, et al. The Relationship Between Pitching Mechanics and Injury: A Review of Current Concepts. *Sports Health.* 2017;9(3):216–21. <https://doi.org/10.1177/1941738116686545>.
66. Konda S, Yanai T, Sakurai S. Configuration of the Shoulder Complex During the Arm-Cocking Phase in Baseball Pitching. *Am J Sports Med.* 2015;43(10):2445–51. <https://doi.org/10.1177/0363546515594379>.
67. Kibler WB, Ludewig PM, McClure PW, Michener LA, Bak K, Sciascia AD. Clinical implications of scapular dyskinesis in shoulder injury: the 2013 consensus statement from the 'Scapular Summit'. *Br J Sports Med.* 2013;47(14):877–85. <https://doi.org/10.1136/bjsports-2013-092425>.
68. Panagiotopoulos AC, Crowther IM. Scapular Dyskinesia, the forgotten culprit of shoulder pain and how to rehabilitate. *SICOT J.* 2019;5:29. <https://doi.org/10.1051/sicotj/2019029>.
69. Hickey D, Solvig V, Cavalheri V, Harrold M, McKenna L. Scapular dyskinesis increases the risk of future shoulder pain by 43% in asymptomatic athletes: a systematic review and meta-analysis. *Br J Sports Med.* 2018;52(2):102–10. <https://doi.org/10.1136/bjsports-2017-097559>.
70. Kennedy DJ, Visco CJ, Press J. Current concepts for shoulder training in the overhead athlete. *Curr Sports Med Rep.* 2009;8(3):154–60. <https://doi.org/10.1249/JSR.0b013e3181a64607>.
71. Kibler WB, McMullen J, Uhl T. Shoulder rehabilitation strategies, guidelines, and practice. *Orthop Clin North Am.* 2001;32(3):527–38. [https://doi.org/10.1016/s0030-5898\(05\)70222-4](https://doi.org/10.1016/s0030-5898(05)70222-4).
72. Kibler WB, Sciascia AD, Uhl TL, Tambay N, Cunningham T. Electromyographic analysis of specific exercises for scapular control in early phases of shoulder rehabilitation. *Am J Sports Med.* 2008;36(9):1789–98. <https://doi.org/10.1177/0363546508316281>.
73. Wilk KE, Obma P, Simpson CD, Cain EL, Dugas JR, Andrews JR. Shoulder injuries in the overhead athlete. *J Orthop Sports Phys Ther.* 2009;39(2):38–54. <https://doi.org/10.2519/jospt.2009.2929>.
74. Camp CL, Zajac JM, Pearson DB, Sinatra AM, Spiker AM, Werner BC, et al. Decreased Shoulder External Rotation and Flexion Are Greater Predictors of Injury Than Internal Rotation Deficits: Analysis of 132 Pitcher-Seasons in Professional Baseball.

- Arthroscopy. 2017;33(9):1629–36. <https://doi.org/10.1016/j.arthro.2017.03.025>.
75. Wilk KE, Macrina LC, Fleisig GS, Aune KT, Porterfield RA, Harker P, et al. Deficits in Glenohumeral Passive Range of Motion Increase Risk of Shoulder Injury in Professional Baseball Pitchers: A Prospective Study. *Am J Sports Med.* 2015;43(10):2379–85. <https://doi.org/10.1177/0363546515594380>.
 76. Wilk KE, Meister K, Andrews JR. Current concepts in the rehabilitation of the overhead throwing athlete. *Am J Sports Med.* 2002;30(1):136–51. <https://doi.org/10.1177/03635465020300011201>.
 77. Fortenbaugh D, Fleisig GS, Andrews JR. Baseball pitching biomechanics in relation to injury risk and performance. *Sports Health.* 2009;1(4):314–20. <https://doi.org/10.1177/1941738109338546>.
 78. Paulino FE, Villacis DC, Ahmad CS. Valgus Extension Overload in Baseball Players. *Am J Orthop (Belle Mead NJ).* 2016;45(3):144–51.
 79. Patel RM, Lynch TS, Amin NH, Calabrese G, Gryzlo SM, Schickendantz MS. The thrower's elbow. *Orthop Clin North Am.* 2014;45(3):355–76. <https://doi.org/10.1016/j.jocl.2014.03.007>.
 80. Ellenbecker TS, Wilk KE, Altchek DW, Andrews JR. Current concepts in rehabilitation following ulnar collateral ligament reconstruction. *Sports Health.* 2009;1(4):301–13. <https://doi.org/10.1177/1941738109338553>.
 81. Fleisig GS, Weber A, Hassell N, Andrews JR. Prevention of elbow injuries in youth baseball pitchers. *Curr Sports Med Rep.* 2009;8(5):250–4. <https://doi.org/10.1249/JSR.0b013e3181b7ee5f>.
 82. Andrews JR, Timmerman LA. Outcome of elbow surgery in professional baseball players. *Am J Sports Med.* 1995;23(4):407–13. <https://doi.org/10.1177/036354659502300406>.
 83. Lee BJ, Garrison JC, Conway JE, Pollard K, Aryal S. The Relationship Between Humeral Retrotorsion and Shoulder Range of Motion in Baseball Players With an Ulnar Collateral Ligament Tear. *Orthop J Sports Med.* 2016;4(10):2325967116667497. <https://doi.org/10.1177/2325967116667497>.
 84. Garrison JC, Cole MA, Conway JE, Macko MJ, Thigpen C, Shanley E. Shoulder range of motion deficits in baseball players with an ulnar collateral ligament tear. *Am J Sports Med.* 2012;40(11):2597–603. <https://doi.org/10.1177/0363546512459175>.
 85. Reinold MM, Wilk KE, Macrina LC, Sheheane C, Dun S, Fleisig GS, et al. Changes in shoulder and elbow passive range of motion after pitching in professional baseball players. *Am J Sports Med.* 2008;36(3):523–7. <https://doi.org/10.1177/0363546507308935>.
 86. Borsa PA, Laudner KG, Sauers EL. Mobility and stability adaptations in the shoulder of the overhead athlete: a theoretical and evidence-based perspective. *Sports Med.* 2008;38(1):17–36. <https://doi.org/10.2165/00007256-200838010-00003>.
 87. Borsa PA, Wilk KE, Jacobson JA, Scibek JS, Dover GC, Reinold MM, et al. Correlation of range of motion and glenohumeral translation in professional baseball pitchers. *Am J Sports Med.* 2005;33(9):1392–9. <https://doi.org/10.1177/0363546504273490>.
 88. Kibler WB, Sciascia A, Moore S. An acute throwing episode decreases shoulder internal rotation. *Clin Orthop Relat Res.* 2012;470(6):1545–51. <https://doi.org/10.1007/s11999-011-2217-z>.
 89. Reuther KE, Larsen R, Kuhn PD, Kelly JD, Thomas SJ. Sleeper stretch accelerates recovery of glenohumeral internal rotation after pitching. *J Shoulder Elbow Surg.* 2016;25(12):1925–9. <https://doi.org/10.1016/j.jse.2016.07.075>.
 90. Mine K, Nakayama T, Milanese S, Grimmer K. Effectiveness of Stretching on Posterior Shoulder Tightness and Glenohumeral Internal-Rotation Deficit: A Systematic Review of Randomized Controlled Trials. *J Sport Rehabil.* 2017;26(4):294–305. <https://doi.org/10.1123/jsr.2015-0172>.
 91. Sakata J, Nakamura E, Suzuki T, Suzukawa M, Aakeda M, Yamazaki T, et al. Throwing Injuries in Youth Baseball Players: Can a Prevention Program Help? A Randomized Controlled Trial. *Am J Sports Med.* 2019;47(11):2709–16. <https://doi.org/10.1177/0363546519861378>.
 92. Escamilla RF, Yamashiro K, Mikla T, Collins J, Lieppman K, Andrews JR. Effects of a Short-Duration Stretching Drill After Pitching on Elbow and Shoulder Range of Motion in Professional Baseball Pitchers. *Am J Sports Med.* 2017;45(3):692–700. <https://doi.org/10.1177/0363546516671943>.
 93. Meister K. Injuries to the shoulder in the throwing athlete. Part one: Biomechanics/pathophysiology/classification of injury. *Am J Sports Med.* 2000;28(2):265–75. <https://doi.org/10.1177/03635465000280022301>.
 94. Burkhart SS, Morgan CD, Kibler WB. The disabled throwing shoulder: spectrum of pathology Part III: The SICK scapula, scapular dyskinesis, the kinetic chain, and rehabilitation. *Arthroscopy.* 2003;19(6):641–61. [https://doi.org/10.1016/s0749-8063\(03\)00389-x](https://doi.org/10.1016/s0749-8063(03)00389-x).

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.