UC Davis

UC Davis Previously Published Works

Title

Adult ADHD and emerging models of maladaptive personality: a meta-analytic review.

Permalink

<https://escholarship.org/uc/item/90d76515>

Journal

BMC psychiatry, 21(1)

ISSN

1471-244X

Authors

Jacobsson, Peter
Hopwood, Christopher J
Söderpalm, Bo
et al.

Publication Date

2021-06-01

DOI

10.1186/s12888-021-03284-1

Peer reviewed

RESEARCH

Open Access

Adult ADHD and emerging models of maladaptive personality: a meta-analytic review

Peter Jacobsson^{1,2*}, Christopher J. Hopwood^{2,3}, Bo Söderpalm¹ and Thomas Nilsson^{2,4}

Abstract

Background: ADHD is a highly consequential disorder that is estimated to affect 2.5% of the adult population. Emerging models of psychopathology posit that disorders like ADHD can be usefully situated within general models of individual differences in personality, such as those recently implemented in the DSM and ICD for the diagnosis of personality disorder. Previous research and systematic reviews have linked adult ADHD to the personality traits Conscientious Inhibition and Negative Emotionality. However, there have been some inconsistencies in the literature and research embedding ADHD-personality connections in the DSM-5 and ICD-11 personality disorder models has been limited. The goal of this paper was to systematically review associations between adult ADHD and personality traits, organized within a maladaptive five factor framework.

Method: A comprehensive literature search yielded 13 papers whose effects were meta-analyzed.

Results: Results supported associations between ADHD and low Conscientious Inhibition and high Negative Emotionality. However, interesting patterns of variability were observed, potentially related to issues such as instrumentation and facet variation.

Conclusion: Results support the clinical application of personality assessment for suggesting risk for ADHD symptoms, and point to important directions for further research.

Keywords: Personality, Personality disorder, ADHD, Personality traits, Clinical assessment, Diagnosis

Attention Deficit Hyperactivity Disorder (ADHD) is a prevalent psychiatric disorder [1–3]. Although it is commonly conceptualized as a neurodevelopmental condition, it also includes features that resemble basic personality traits, such as Neuroticism and Impulsivity [4, 5]. Similar to broad personality traits, ADHD tends to co-occur with a wide range of other disorders, including mood disorders, anxiety disorders, personality disorders and substance disorders [6–9]. Moreover, deficits in personality traits such as emotion-regulation, distractibility, irresponsibility, risk-

taking and impulsivity are thought to be at the core of ADHD symptoms [10, 11].

Most research on personality and psychopathology organizes individual differences in the form of the Five Factor Model [4], which includes the traits Neuroticism, Extraversion, Openness, Agreeableness, and Conscientiousness. Notably, several comparable models have gained attention in mental health research and practice recently, including the Research Domain Criteria (RDoC) [12]; the Hierarchical Taxonomy of Psychopathology (HiTOP) [13]; the Alternative Model of Personality Disorder (AMPD) [14]; and the ICD-11 proposal for personality disorder [15]. All of these models have in common the proposal that a few broad traits underlie

* Correspondence: peter.jacobsson@regionhalland.se

¹The Sahlgrenska Academy at University of Gothenburg, Gothenburg, Sweden

²Psychiatry Halland, Region Halland, Sweden

Full list of author information is available at the end of the article

© The Author(s). 2021 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

functioning in a wide range of areas, including dysfunction related to inattention, impulsivity and hyperactivity. Associations between personality and psychopathology are particularly strong when maladaptive trait measures, or instruments that focus on personality-related problems, are used [16].

Several previous reviews have documented associations between FFM personality traits and ADHD [17–19]. These reviews indicate that ADHD as a unified construct is most consistently related to low Conscientiousness, low Agreeableness, and high Neuroticism. This research also suggests some specificity in associations between inattentive and hyperactive/impulsive aspects of ADHD and certain traits [20, 21]. Specifically, Inattention is positively correlated to Neuroticism and negatively to Conscientiousness; whereas Hyperactivity/Impulsivity relates negatively to Agreeableness and positively to Extraversion [17]. As such, there is considerable potential in using basic traits like those of the FFM, RDoC, HiTOP, AMPD, and ICD-11 to understand how individuals diagnosed with ADHD differ from individuals with other diagnoses in terms of underlying personality traits. The goal of the present study was to use meta-analysis to summarize associations between adult ADHD and create a bridge from non-clinical to maladaptive personality traits.

The integrated five-factor model

The FFM consists of the following five factors: *Neuroticism* which refers to the extent of negative emotions, i.e., sadness, fear, hostility, and emotional lability that the individual experiences [22]. Individuals high in this domain may be at risk for many different psychiatric and physical disorders, high comorbidity, lower quality of life, shorter lifespan and more extensive use of health care [23]. *Extraversion* concerns how outgoing or talkative the individual is in most situations. Core traits are sociability, assertiveness, positive affect, and activity level [22]. Low Extraversion, or introversion, may include social withdrawal, social detachment, intimacy avoidance, restricted affectivity and anhedonia [24–26] and extremely high Extraversion may represent personality pathology in sexual promiscuity, emotional intrusiveness, excessive self-disclosure and thrill-seeking behavior [27]. *Openness to experience* describes the depth and breadth of an individual's intellectual, artistic, and experiential life, with key facets such as aesthetic sensitivity, intellectual interests, and imagination [22]. Although open individuals generally tend to have greater psychological well-being, maladaptive Openness can be found in distinguishing between major depression and bipolar disorder as well as different variants of schizotypy [28]. *Agreeableness* concerns the extent of the individual's motivation for prosocial behavior and pleasant interpersonal

relationships. Important traits are compassion, trust, and politeness [22]. Individuals low in Agreeableness tend to be critical, skeptical, try to push limits, express hostility and being condescending [28]. Agreeableness is also inversely linked to psychopathy and aggressive behavior [29, 30]. Maladaptive variants of extreme Agreeableness can result in gullibility, submissiveness, clinging, subservience, servility etc. [28]. *Conscientiousness* refers to the individual's level of organization, ability to complete tasks, and persistence in achieving long-term goals. Key concepts are orderliness, self-discipline, and reliability [22]. However, both high and low Conscientiousness is associated with decreased functioning. Low Conscientiousness is characterized by disinhibition, irresponsibility, negligence, and rashness [31, 32]; and inflexible high Conscientiousness can result in perfectionism, fastidiousness, punctiliousness, workaholism and other facets of compulsivity [33–35].

Research has shown that multiple models of both normal-range and maladaptive personality traits can be integrated into the structure of the FFM [36–38]. For instance, the trait Neuroticism is empirically and conceptually similar to the trait Negative Affectivity in the AMPD and ICD-11, Emotion Dysregulation in Livesley's Dimensional Assessment of Personality Pathology (DAPP [39]); and the traits Harm Avoidance and (low) Self-Transcendence in Cloninger's Temperament and Character Inventory (TCI [40]). Similarly, FFM Conscientiousness is similar to Disinhibition in the AMPD/ICD-11, Compulsivity on the DAPP-BQ, and low Novelty Seeking and Persistence in the TCI. As such, in this study, we will conceptualize personality traits in terms of an Integrative Five Factor Model (IFFM) inclusive of instruments that are explicitly conceptualized as measures of the FFM, as well as those, such as the DAPP and TCI, that have commensurate scales (See Table 1). We will use domain labels that integrate normal-range and maladaptive aspects of each trait [37]: Negative Emotionality, Positive Emotionality, Openness, Conscientious Inhibition, and Agreeable Inhibition.

Conceptualizing ADHD within an integrative individual differences framework

Conceptualizing ADHD as a dimension or set of dimensions that can be placed within a general, evidence-based model of maladaptive personality traits has a number of potential clinical advantages [41–44]. For instance, this model may help provide a principled and efficient means for conceptualizing heterogeneity in the presentation of individuals with ADHD diagnosis. Specifically, previous research suggests that Negative Emotionality and low Conscientious Inhibition are most strongly related to inattention, whereas Agreeable Inhibition and Positive Emotionality maybe more strongly

Table 1 A translation of how personality traits from different assessment instruments are related to the integrated five-factor model traits

The Integrated Five-Factor Model ^a	Five-Factor Model	DAPP-BQ	Cloninger TCI	AMPD/ICD-11 ^b
<i>Negative Emotionality</i>	Neuroticism	Emotional dysregulation	Harm avoidance; Self-directedness (–)	Negative affectivity
<i>Positive Emotionality</i>	Extraversion	Inhibitedness (–)	Harm avoidance (–); Reward dependence	Detachment (–)
<i>Openness</i>	Openness to experience		Self-transcendence	Psychoticism (AMPD only)
<i>Conscientious Inhibition</i>	Conscientiousness	Compulsivity	Novelty seeking (–); Persistence	Disinhibition (AMPD) (–); Anankastia (ICD-11)
<i>Agreeable Inhibition</i>	Agreeableness	Dissocial behavior (–)	Cooperativeness	Antagonism (AMPD) (–); Dissociality (ICD-11) (–)

^aAdapted from Markon et al (2005) [37] and Gomez & Corr (2014). Note: Negative sign indicates that the scale loaded negatively

^bEmerging personality models AMPD and ICD-11 factors are not represented in included studies of the current review

related to hyperactivity and impulsivity [17, 45–47]. Evidence-based models of individual differences also help explain comorbidity. Insofar as many disorders share underlying propensities for high Negative emotionality and low Conscientious and Agreeable Inhibition [48], these disorders should be expected to co-occur with inattentive, impulsive and hyperactive symptoms. Conceptualizing ADHD within a multidimensional personality model that can help provide explanations and order to patterns of heterogeneity and comorbidity may thus pave the way for more targeted treatments and research. As such, individual differences models such as the IFFM have the potential to improve the efficiency, validity, and utility of ADHD diagnoses [43].

Four previous studies summarize associations between personality traits and adult ADHD [17–19, 47]. This study provides additional information not included in those studies in the following ways. First, this unlike some reviews [18], uses formal meta-analytic procedures to quantify associations between personality traits and ADHD. Second, six not previously reviewed studies are included in the current meta-analytic review [46, 49–53]. Third, whereas the cited earlier reviews have included both child and adult studies with varying approaches to the assessment of ADHD [17–19, 47], the present review has only included papers with adult individuals and excluded studies without a clear description of a formal ADHD diagnosis using a stringent clinical assessment. Fourth, this study conceptualizes personality using various models that can be integrated within an IFFM framework.

In summary, the aim of this study was to quantify associations between ADHD and the traits of an Integrated FFM to better understand how ADHD fits into evidence-based models of personality and psychopathology, and ultimately enable improved clinical practice

and research. Given the existing body of research in adult populations, we focused on traits from the FFM, DAPP, HEXACO [54], and TCI. Given that relatively few studies have distinguished between ADHD subtypes, we conceptualized ADHD as unitary construct.

Method

Literature searches were conducted in November 2019 via Cochrane, PubMed, PsychInfo and SCOPUS using the following keywords: (ADHD OR ADD OR ADDH OR Attention Deficit Disorder with Hyperactivity) AND (personality traits OR personality dimensions OR NEO OR five-factor model OR FFM OR BFI OR Temperament and Character Inventory OR TCI). The reference sections of each article retrieved was searched manually for any potentially relevant studies. Included studies were a) written in English; b) published after January 1, 2000; c) contained adult cases with a diagnosis of ADHD according to DSM-IV or DSM-5; d) used validated measures to assess adult ADHD; e) used established dimensional personality measures (FFM, DAPP, HEXACO, or TCI); and f) presented results in a format that allowed for straightforward conversion to the FFM i.e. into the Integrated Five-Factor Model (IFFM [37]; (Fig. 1).

The first author screened titles and abstracts and excluded papers that did not meet inclusion criteria. The remaining articles were assessed, discussed, and consented upon by the first author and two of the coauthors (TN and BS). This process resulted in 14 papers that sampled adult patients with ADHD [17, 46, 49–53, 55–61]. Six studies used an explicit FFM measure [17, 46, 49, 50, 52, 53], one used the HEXACO [61], six used the Temperament and Character Inventory (TCI [55–60]), and Jacob et al. [51] included both FFM and TCI measures. A study by Koerting et al. [52] used measures from the DAPP-BQ in addition to an FFM instrument.

As explained above, and illustrated in Table 1, traits from these measures were categorized as falling within the IFFM domains of *Negative Emotionality*, *Positive Emotionality*, *Openness*, *Conscientious Inhibition*, or *Agreeable Inhibition*. Within the IFFM framework, the DAPP-BQ traits *Emotional dysregulation* and *Compulsivity* relate to *Negative Emotionality* and *Conscientious Inhibition*, respectively; *Inhibitedness* loads negatively on *Positive Emotionality*; and *Dissocial behavior* has a negative association with *Agreeable Inhibition*. No specific DAPP-BQ trait is associated with *Openness* in the IFFM framework. The TCI traits *Harm avoidance*, *Reward dependence*, *Self-transcendence*, *Persistence*, and *Cooperativeness* are associated with the respective IFFM domains *Negative Emotionality*, *Positive Emotionality*, *Openness*, *Conscientious Inhibition*, and *Agreeable Inhibition*. *Self-directedness* loads negatively on *Negative Emotionality*, *Harm avoidance* is negatively related to *Positive Emotionality*, and *Novelty seeking* is negatively related to *Conscientious Inhibition*.

All formal ADHD diagnoses were based on the DSM criteria, although different instruments were used to assess current and historical symptoms and impairments. All data included in this analysis examined group differences as they relate to personality traits. Control groups included community samples, non-clinical parents of children with ADHD, blood donors, clinical groups without ADHD, and population norms extracted from manuals (Table 2). In five of the papers [17, 46, 51, 58, 62] correlations with the ADHD symptom clusters: Combined (C), Inattentive (I), Hyperactive/Impulsive (HI), were described. Only the Jacob [51] 2016 study presented individual data according to specific subtype of ADHD. Thus, results on individual ADHD type or presentation could not be aggregated and analyzed separately. However, indicative heuristic patterns are presented and discussed.

Meta-analysis was used to aggregate associations of personality traits and ADHD in Meta-Essentials [63, 64]. Group differences were converted to the Cohen's d

Table 2 A compilation of included studies presenting study characteristics, data about personality scales (*M*, *SD*) for ADHD and control subjects, and correlations between personality traits and ADHD

Author	ADHD N	Male (%)	Age <i>M</i> (<i>SD</i>)	Clinical recruitment	Control N	Male (%)	Age <i>M</i> (<i>SD</i>)	Control Group	Personality scale	ADHD <i>M</i> (<i>SD</i>)	Control <i>M</i> (<i>SD</i>)	Correlation trait - ADHD dx
He & Antschel (2019)	184	51	36.3 (10.8)	Psychiatric clinics	118	46	29.8 (8.7)	Community	TCI Harm Avoidance	60.0 (13.5)	52.78 (9, 9)	.29
									TCI Self- directedness	61.4 (13.9)	80.3 (9.3)	-.62
									TCI Reward Dependence	67.2 (11.2)	72.0 (9.4)	-.23
									TCI Self- transcendence	52.4 (16.5)	48.8 (16, 3)	.11
									TCI Novelty seeking	64.0 (9, 7)	53.0 (7.2)	.54
									TCI Persistence	63.3 (11.8)	72.9 (9.9)	-.40
									TCI Cooperativeness	80.3 (9, 2)	74.5 (11, 0)	-.27
Wallace (2016) [49]	217	56	23.6 (4.7)	Psychiatric clinics and advertisements	114	50	22.2 (4.7)	Community	NEO FFI Neuroticism	28.5 (7, 9)	21.50 (7, 8)	.40
									NEO-FFI Extraversion	28.8 (7, 7)	29.5 (7.4)	-.05
									NEO-FFI Openness to experience	33.6 (7, 1)	31.9 (6.5)	.12
									NEO-FFI Conscientiousness	16.1 (4, 8)	20.4 (4.6)	-.42
									NEO-FFI Agreeableness	31 (6.9)	32.9 (6.6)	-.14
Di Nicola (2014) [50]	16	No information	No information	Affective disorders unit	102	38 ^a	47.4 (13.2)	BD	NEO-PI-R Neuroticism	140.5 (23.5)	112.0 (21.4)	.54
	8	No information	No information	Affective disorders unit	106	40 ^a	46.1 (14.3)	MDD		133.4 (22.9)	111.3 (22.8)	.44
								BD	NEO-PI-R Extraversion	93.4 (25.0)	94.8 (21, 5)	-.03;
								MDD		82.3 (20.3)	93.1 (20, 9)	-.25
								BD	NEO-PI-R Openness	104.7 (21.3)	105.8 (20.7)	-.03
								MDD		107.4 (21.1)	101.4 (14.5)	.16
								BD	NEO-PI-R	88.7	106.0	-.41

Table 2 A compilation of included studies presenting study characteristics, data about personality scales (*M*, *SD*) for ADHD and control subjects, and correlations between personality traits and ADHD (*Continued*)

Author	ADHD N	Male (%)	Age <i>M</i> (<i>SD</i>)	Clinical recruitment	Control N	Male (%)	Age <i>M</i> (<i>SD</i>)	Control Group	Personality scale	ADHD <i>M</i> (<i>SD</i>)	Control <i>M</i> (<i>SD</i>)	Correlation trait - ADHD dx
Faraone et al. (2009) [56]	127	53	36.1 (10.8)	Psychiatric clinics and advertisements	123	46	29.9 (9.0)	Community	Conscientiousness	(17.9)	(21.0)	
										94.4 (22.7)	108.6 (21.8)	-.30
									NEO-PI-R Agreeableness	118.5 (18.3)	119.0 (18.9)	-.01
										124.0 (15.6)	119.2 (15.4)	.15
									TCI Harm Avoidance	59.5 (13.5)	52.7 (9, 95)	.26,
									TCI Self- directedness	61.3 (13.5)	80.2 (94)	-.57
									TCI Reward Dependence	67.4 (10.7)	72.0 (9.5)	-.20
									TCI Self- transcendence	52.4 (16.8)	48.6 (16, 2)	.12
									TCI Novelty seeking	64.2 (10.6)	53.2 (7.2)	.53
									TCI Persistence	63.1 (12.2)	72.9 (10, 0)	-.33
Jacob et al. (2016) [51]	Male 441	50	34.5 (10.2)	Psychiatric clinic	80	No information	No information	Reference values male	TCI Cooperativeness	75.6 (10.0)	80.2 (9.2)	-.23
									TPQ Harm Avoidance	18.2 (7, 1)	14.7 (6.4)	.25
	Female 440	50	34.5 (10.2)	Psychiatric clinic	80	No information	No information	Reference values female	TPQ Reward Dependence	20.9 (6, 7)	16.3 (6)	.34
										16.1 (4, 5)	14.8 (4.7)	.14
										18.6 (4, 2)	17.5 (4.7)	.12
									TPQ Novelty seeking	19.5 (5, 7)	12.8 (5.8)	.50
										19 (5.9)	15.1 (4.6)	.34
	Male 426		34.5 (10.2) Total	Psychiatric clinic	4219	No information	No information	Reference values male	NEO-PI-R Neuroticism	108.7 (24.6)	84.2 (22, 5)	.46
										122.6 (25.1)	95 (23.3)	.50
	Female 432		34.5 (10.2) Total	Psychiatric clinic	7505	No information	No information	Reference values female	NEO-PI-R	100.1	109.6	-.22

Table 2 A compilation of included studies presenting study characteristics, data about personality scales (*M*, *SD*) for ADHD and control subjects, and correlations between personality traits and ADHD (*Continued*)

Author	ADHD N	Male (%)	Age <i>M</i> (<i>SD</i>)	Clinical recruitment	Control N	Male (%)	Age <i>M</i> (<i>SD</i>)	Control Group	Personality scale	ADHD <i>M</i> (<i>SD</i>)	Control <i>M</i> (<i>SD</i>)	Correlation trait - ADHD <i>dx</i>
Koerting (2016) [52]	30	43	33.5 (8.8)	Psychiatric clinic	30	43	28.2 (7.0)	Community	Extraversion	(21.6)	(20.4)	
										100.3 (23.7)	111 (19, 5)	-.24
									NEO-PI-R Openness	109.4 (19.1)	120.4 (20.1)	-.27
										114.2 (21.2)	125.7 (18.6)	-.28
									NEO-PI-R Conscientiousness	92.3 (21.2)	114.1 (20.7)	-.46
										89.9 (22.8)	113.8 (19.7)	-.49
									NEO-PI-R Agreeableness	107.7 (16.9)	108.6 (17.3)	-.03
										116 (16, 1)	114.9 (16.4)	.03
									NEO-PI-R Neuroticism;	2.4 (0.4)	1.8 (0.4)	.58
									NEO-PI-R Extraversion	2.1 (0.4)	2.3 (0.3)	-.046
									NEO-PI-R Openness	2.5 (0.4)	2.5 (0.3)	-.030
									NEO-PI-R Conscientiousness	1.6 (0.5)	2.5 (0.4)	-.001
									NEO-PI-R Agreeableness	2.3 (0.3)	2.4 (0.3)	0.11
									DAPP-BQ Emotional Dysregulation	2.7 (0.5)	2.1 (0.4)	.52
									DAPP-BQ Inhibitedness	2.5 (0.4)	2.0 (0.3)	.56
Nigg et al. (2002) [17]	88	38	21.6 (3.9)	University disability office and advertisements	26	No information	No information	Community	DAPP-BQ Compulsivity	2.6 (0.7)	3.0 (0.6)	-.31
									DAPP-BQ DissocialBehavior	2.5 (0.4)	2.3 (0.4)	.32
									NEO-FFI Neuroticism	26.4 (8, 7)	19.2 (7.3)	.41
									NEO-FFI	32 (7.2)	31.3 (4.7)	.06

Table 2 A compilation of included studies presenting study characteristics, data about personality scales (*M*, *SD*) for ADHD and control subjects, and correlations between personality traits and ADHD (*Continued*)

Author	ADHD N	Male (%)	Age <i>M</i> (<i>SD</i>)	Clinical recruitment	Control N	Male (%)	Age <i>M</i> (<i>SD</i>)	Control Group	Personality scale	ADHD <i>M</i> (<i>SD</i>)	Control <i>M</i> (<i>SD</i>)	Correlation trait - ADHD <i>dx</i>
Knouse (2013) [46]	117	50	42.8 (11.0)	Hospital	1539	No information	No information	Reference values	Extraversion			
									NEO-FFI Openness to experience	31.6 (5, 8)	30.7 (5.9)	.08
									NEO-FFI Conscientiousness	22.3 (7, 6)	34.6 (5.2)	-.69
									NEO-FFI Agreeableness	29.7 (6)	33.3 (5.9)	-.29
									NEO-FFI Neuroticism	26.3 (8, 2)	19.1 (7.7)	.41
									NEO-PI-R Extraversion	28.5 (6, 4)	27.7 (5.8)	.07
									NEO-PI-R Openness	31.3 (6, 5)	27.0 (5.8)	.33
									NEO-PI-R Conscientiousness	19.7 (6, 7)	34.6 (5.9)	-.76
									NEO-PI-R Agreeableness	31.7 (6, 5)	32.8 (5.0)	-.10
									VTCI Harm Avoidance	55.2 (11.4)	50 (10)	.24
Sizoo et al. (2009) [57]	53	68	31.4 (9.9)	Center developmental disorders	657	No information	No information	Expected population norms (T-score)	VTCI Self-directedness	34.8 (14.8)	50.0 (10, 0)	-.52
									VTCI Reward Dependence	50.3 (10.9)	50.0 (10, 0)	.14
									VTCI Self-transcendence	53.7 (10.8)	50.0 (10, 0)	.18
									VTCI Novelty seeking	61.3 (9, 6)	50.0 (10, 0)	.50
									VTCI Persistence	50.5 (10.7)	50.0 (10, 0)	.02
Steinhausen et al (2013) [53]	30	No information	No information	Parents with ADHD of children with ADHD	34	50	44.7 (5.4)	Parents of children with ADHD	VTCI Cooperativeness	46.1 (13.4)	50.0 (10, 0)	.16
									NEO-FFI Neuroticism	2.2 (0.8)	1.5 (0.6)	.44
									NEO-FFI Extraversion	2.2 (0.7)	2.4 (0.5)	-0.16
									NEO-FFI Openness to experience	2.3 (0.5)	2.4 (0.4)	-0.11

Table 2 A compilation of included studies presenting study characteristics, data about personality scales (*M*, *SD*) for ADHD and control subjects, and correlations between personality traits and ADHD (*Continued*)

Author	ADHD N	Male (%)	Age <i>M</i> (<i>SD</i>)	Clinical recruitment	Control N	Male (%)	Age <i>M</i> (<i>SD</i>)	Control Group	Personality scale	ADHD <i>M</i> (<i>SD</i>)	Control <i>M</i> (<i>SD</i>)	Correlation trait - ADHD <i>dx</i>
Perroud (2016) [58]	119	69	37.3 (11.0)	ADHD center	403	63	45.5 (12.7)	Blood donors	NEO-FFI Conscientiousness	2.4 (0.6)	2.9 (0.5)	−0.41
									NEO-FFI Agreeableness	2.5 (0.5)	2.6 (0.4)	−0.11
									TCI Harm Avoidance	18.3 (8, 1)	12.7 (6.2)	.36
									TCI Self- directedness	21.9 (8, 5)	34.5 (5.8)	−.65
									TCI Reward Dependence	14.2 (4, 0)	14.6 (3.8)	−.05
									TCI Self- transcendence	15.3 (6, 8)	13.5 (6.3)	.14
									TCI Novelty seeking	23 (5.9)	18.8 (5.6)	.34
									TCI Persistence	4.8 (2.0)	4.8 (2.0)	.00
									TCI Cooperativeness	30.5 (7, 4)	33.5 (5)	−.23
									VTCI Harm Avoidance	55.7 (11.4)	50.0 (10, 0)	.26
Sizoo et al. (2015) [59]	53	66	32.1 (11.4)	Center developmental disorders	399	No information	No information	Expected population norms (T-scores)	VTCI Self- directedness	34.4 (14.7)	50.0 (10, 0)	−.53
									VTCI Reward Dependence	50 (10, 7)	50.0 (10, 0)	.00
									VTCI Self- transcendence	53.8 (10.8)	50.0 (10, 0)	.18
									VTCI Novelty seeking	60.9 (9, 7)	50.0 (10, 0)	.48
									VTCI Persistence	50.6 (10.7)	50.0 (10, 0)	.03
									VTCI Cooperativeness	45.9 (13.2)	50.0 (10, 0)	−.17
									TCI Harm Avoidance	63.4 (12.1)	50.0 (10, 0)	.51
									TCI Self- directedness	29 (12, 5)	50.0 (10, 0)	−.68
Anckarsater et al. (2006)	100	No information	No information		1300	No information	No information	Expected population norms (T-scores)	TCI Reward Dependence	46.2 (12.5)	50.0 (10, 0)	−.16

Table 2 A compilation of included studies presenting study characteristics, data about personality scales (*M*, *SD*) for ADHD and control subjects, and correlations between personality traits and ADHD (*Continued*)

Author	ADHD N	Male (%)	Age <i>M</i> (<i>SD</i>)	Clinical recruitment	Control N	Male (%)	Age <i>M</i> (<i>SD</i>)	Control Group	Personality scale	ADHD <i>M</i> (<i>SD</i>)	Control <i>M</i> (<i>SD</i>)	Correlation trait - ADHD <i>dx</i>
									TCI Self-transcendence	55,1 (12,9)	50,0 (10, 0)	.22
									TCI Novelty seeking	59,2 (10,9)	50,0 (10, 0)	.40
									TCI Persistence	50,2 (10,1)	50,0 (10, 0)	.01
									TCI Cooperativeness	35,2 (17,5)	50,0 (10, 0)	-.46

Note: TCI Temperament and Character Inventory, NEO FFI NEO Five Factor Inventory, NEO-PI-R The Revised NEO Personality Inventory, TPQ Tridimensional Personality Questionnaire, DAPP-BQ Dimensional Assessment of Personality Pathology, Brief Questionnaire, VTCI The brief Dutch version of the Temperament and Character Inventory, BD Bipolar Disorder, MDD Major Depressive Disorder

^aTotal number including ADHD

metric, weighted by sample size, and analyzed separately for each personality dimension. Effect sizes, 95% confidence intervals and prediction intervals with a random effects model were calculated and illustrated in forest plots. Heterogeneity of effects was also calculated to estimate the robustness of results [65]. I^2 was used to evaluate the heterogeneity of mean effect sizes. An I^2 value above 50% can be interpreted as meaningful heterogeneity around the mean effect size [66].

Quality assessment of papers

The 14 papers were ranked for scientific quality according to an adaptation of QUADAS [67]. QUADAS was developed as a tool to systematically assess scientific papers that concern diagnostic tests. The following QUADAS items: a) design, b) sample size, c) relevant population, d) assessment method, e) relevant assessment instrument, f) sources of recruited cases, g) blinding, h) conflict of interest, and i) drop-out, were assessed by the first author (PJ) and two co-authors (TN and BS). The papers were first assessed independently and then discussed to reach consensus. Each aspect was scored from 0 to 2: unsatisfactory = 0; fair = 1; good = 2. The scores for the aspects were summarized for each paper to generate total scores between 0 and 18. The 14 papers ranged from 5 to 13. The paper with lowest score [5] was removed after quality assessment and subsequent preliminary analyses due to deviating both in quality assessment and being identified as an outlier in subsequent analyses, thus resulting in 13 papers that were included in the final analysis. The excluded paper did not meet the quality criteria (unsatisfactory = 0) for five of the nine assessed criteria: small sample, limited number of sources from which participants were recruited, no blinding, no description of conflict of interest, no description of

drop-out. High drop-out rate or not specifying drop-out and insufficient blinding were the most common quality deficits in the other papers. Assessment methods and relevant populations were generally study strengths. The range of the included papers was 7 to 13 (median 10).

Results

Thirteen original articles consisting of 2023 unique adult subjects diagnosed with ADHD and 16,835 control participants, were included in this meta-analysis. Mean age ranged from 21.6 to 42.8 years; 54% of the included population identified as male. Most of the clinical cases of ADHD were recruited through psychiatric clinics, either general or specialized centers (see Table 2). A few clinical cases were recruited through advertisements.

The meta-analytic analyses used in this review indicated a range of combined effect sizes from $d = 0.15$ (negligible) for Openness to $d = 1.11$ (large) for Negative Emotionality [68, 69] (Fig. 2). The effect size for Conscientious Inhibition was also large ($d = -0.89$). Effect sizes for Agreeable Inhibition and Positive Emotionality were more modest, $d = -0.39$ and -0.43 , respectively. All associations with the exception of Openness were significant. *Negative Emotionality* and *Conscientious Inhibition* are thus the personality dimensions that were consistently and most strongly elevated in the ADHD samples in this review.

Even though the combined effect sizes for Negative Emotionality and Conscientious Inhibition were large, there was significant variability in the ADHD populations within these personality domains. Effect sizes for Negative Emotionality ranged from $d = 0.50$ (medium to large) to $d = 2.06$ (very large) (Fig. 3). An even greater range was observed for Conscientious Inhibition: $d = 0.06$ (negligible) for the Cloninger TCI facet Persistence

Combined effect sizes (Cohen's d) from 16 to 30 datasets derived from the FFM family, TCI related measures, one DAPP-Q study.

Fig. 2 Personality traits in ADHD

to $d = -2.51$ (very large) for the Five-Factor domain Conscientiousness (Fig. 4). Combined confidence intervals (CI) and prediction intervals (PI) were largest for Conscientious Inhibition (CI = -1.15 to -0.63 ; PI = -2.09 to 0.30) and smallest for Positive Emotionality (CI = -0.57 to -0.28 ; PI = -1.23 to 0.38).

The I^2 is a measure for the proportion of observed variance that reflects real differences in effect size. It is expressed as a percentage with a range from 0 to 100% [70]. All effect sizes for the various personality domains in the ADHD samples ranged from 85 to 97%. Heterogeneity decreased when FFM measures were used exclusively. The I^2 for FFM Negative Emotionality was 47% and I^2 for FFM Agreeable Inhibition was 64%.

Discussion

The results of this systematic, quantitative review suggest a robust connection between adult ADHD and the IFFM personality domains *Negative Emotionality* and *low Conscientious Inhibition*. *Positive Emotionality* and *Agreeable Inhibition* have somewhat smaller, although significant, associations with adult ADHD. These results are consistent with the review from Gomez et al. [47], with the exception that Gomez et al. observed a larger

effect size for Agreeable Inhibition ($d = -.64$) than we did ($d = -.39$).

Low *Conscientious Inhibition* consists of features that are also implicated in the core symptoms of ADHD. These features are conceptualized in ADHD as difficulties in executive functions or self-regulatory processes [71, 72]. Barkley [73] proposed that *behavioral inhibition*, consisting of: a) inhibition of pre-potent responses, b) stopping ongoing responses, and c) interference control, can be seen as a superordinate factor that affects several cognitive and behavioral modalities, eventually manifesting themselves in specific symptoms commonly found in ADHD. This construct is largely commensurate with Conscientious Inhibition as conceptualized in trait measures, thus providing a bridge between neuropsychiatric and quantitative trait approaches to conceptualizing underlying deficits in ADHD [14, 48, 74].

Negative Emotionality is elevated across many psychiatric disorders and is highly predictive of psychological and physical comorbidity as well as general quality of life [23, 48]. Although symptoms of Negative Emotionality are not central to ADHD criteria, this review and other evidence clearly shows that emotional distress and dysregulation are common among individuals with ADHD

diagnoses [75]. There is also evidence that high Negative Emotionality also predicts worse outcomes in individuals with ADHD [76–78].

Significant variability was observed in effect sizes between and within samples in the current review, as well as between instruments. For example, the high percentage in the I^2 measure, described above, indicates that there is a high degree of heterogeneity between studied populations within the different personality domains in the meta-analysis. Ideally, subgroup or moderator analyses would be performed to explain this variation. However, the number of studies available for this meta-analysis was too small to test moderation. Nevertheless, some potentially reliable patterns were observed.

First, when specific measures of FFM were extracted for separate analysis, heterogeneity decreased. This suggests more consistency when personality traits were measured with FFM instruments. One particularly important source of variability may have to do with differing facet models across those instruments. Facet variability is complicated by the interstitial nature of some traits, which leads different models to place similar facets on different domains. For instance, we conceptualized TCI Self-transcendence as an aspect of Negative Emotionality, although it also shares some features with Conscientious Inhibition. Future work exploring a broader range of traits, and conceptualizing personality

at the level of facets, would shed significant light on this issue.

Second, stronger effects with psychopathology can generally be expected with maladaptive as opposed to normal range personality measures [16]. As maladaptive range measures are more likely to be used in clinical settings, associations with these measures are potentially both stronger, and more informative for clinical practice. As such, future research examining links between adult ADHD and maladaptive range personality measures would be particularly useful.

A third source of variability involves ADHD presentation, given that traits may differentially relate to inattentive, impulsive and hyperactive features [17, 45, 47, 51, 58]. Unfortunately, our relatively small sample size precluded moderator analyses of ADHD subtypes. Future studies with larger samples and symptom or subtype-level assessments of ADHD are needed to further explore the heterogeneity of effects we observed here.

Clinical implications

Overall, these findings support the emerging view that personality, personality disorders, and neurodevelopmental disorders can be conceptualized within an integrated and evidence-based dimensional trait system. This supports the clinical application of personality trait assessment models according to the DSM and ICD

diagnostic systems to broaden the clinical description of adult ADHD, as well as research on the shared etiological factors between personality variables and ADHD.

Categorical psychiatric diagnoses are fraught with high comorbidity, complicating the clinical picture and reducing treatment specificity. Dimensional models provide an opportunity to integrate comorbidity into a more comprehensive and individualized clinical description, which makes treatment planning more in line with current clinical praxis. The current review suggests that the features of ADHD can be economically situated within evidence-based models of personality and psychopathology. As such, the clinical use of personality trait measures and the emerging dimensional personality models included in the AMPD and the ICD-11 proposal provide opportunities to implement a more individualized approach for patients with problems related to hyperactivity, impulsivity, and inattention. Specifically, multidimensional trait profiles may help clinicians estimate long-term risk for psychosocial consequences of ADHD, as well as the likelihood of other comorbid conditions. They also help provide explanations for heterogeneous clinical presentations and point to appropriate pathways for intervention [41–43, 79].

In clinical practice it is useful to rely on heuristics to make efficient clinical decisions [80]. For example, these findings indicate that individuals with scores on personality measures that are more than one standard deviation higher on Negative Emotionality (NE) constructs or lower on Conscientious Inhibition (CI), in relation to non-clinical populations, may be at risk for ADHD or other disorders. Such heuristics can be helpful in screening because they both indicate risk for specific kinds of problems, while also efficiently providing a comprehensive portrait of the individual's strengths and weakness, which has significant value for making prognostic predictions and guiding treatment [17, 43–45].

Limitations

The literature on associations between adult ADHD and personality traits is scarce, thus limiting the degree to which the current results can provide robust information about these links. The existing literature, while sufficient to quantitatively summarize connections between adult ADHD and IFFM trait domains, limited our ability to test moderators such as ADHD subtype, personality measure, or various sample features, and it necessitated inclusion of measures with varying connection to the FFM or DSM/ICD personality disorder proposals. Some of our conclusions were based on assumptions about how to organize facets into maladaptive five factor domain [37, 47] that have not been fully tested empirically. All personality measures were self-report. This is particularly important because some aspects of personality

(developmental, variability, neurocognitive, reflective functioning) may be better assessed using other methods (e.g., informant reports, repeated reporting, implicit tests, neuropsychological testing). Also, the initial screening was performed only by the primary author, which might have risked possible exclusion of relevant studies. Subsequent multiple rater inclusion process was based on consensus; however, reliability statistics for how consensus was reached, were not calculated. Finally, the samples used in this study were WEIRD (i.e. *White Educated Industrialized Rich and Democratic*). Greater diversity in samples is needed to add confidence to these findings and enhance generalizability.

Conclusion

Psychiatric nosology is in a transition from legacy diagnostic categories to evidence-based models of individual differences that closely resemble personality traits [14, 81]. Research on how ADHD fits into these models is relatively nascent, particularly in adult samples. This study shows that existing research on personality associations with ADHD suggests that ADHD is strongly related to Negative Emotionality and low Conscientious Inhibition, and moderately related to low Positive Emotionality and low Agreeable Inhibition. These results support the clinical assessment of personality traits in ADHD diagnosis, clinical care, and research, while also pointing to the need for further research to more specifically delineate how ADHD can be fit into the personality/psychopathology hierarchy.

Acknowledgements

We wish to acknowledge FOU Halland for consulting on analyses.

Authors' contributions

PJ, CH, BS and TN conceptualized and designed the study, PJ performed all the analyses and drafted the manuscript. All authors reviewed and consented to the final version of the manuscript. The authors read and approved the final manuscript.

Funding

PJ was supported by the Region Halland and the study project received research grants from Region Halland (Dnr: 836292) and from Sparbanken Varberg (Dnr: 939031). Open Access funding provided by University of Gothenburg.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

Approved by the Ethics Review Board in Lund, Sweden, Dnr: 2018/53.

Consent for publication

Not applicable.

Competing interests

All authors declare that they have no competing interests.

Author details

¹The Sahlgrenska Academy at University of Gothenburg, Gothenburg, Sweden. ²Psychiatry Halland, Region Halland, Sweden. ³University of California, Davis, USA. ⁴Department of Forensic Psychiatry, National Board of Forensic Medicine, The Sahlgrenska Academy at University of Gothenburg, Gothenburg, Sweden.

Received: 29 January 2021 Accepted: 13 May 2021

Published online: 01 June 2021

References

- Simon V, Czobor P, Balint S, Meszaros A, Bitter I. Prevalence and correlates of adult attention-deficit hyperactivity disorder: meta-analysis. *Br J Psychiatry*. 2009 Mar;194(3):204–11. <https://doi.org/10.1192/bjp.bp.107.048827>.
- Nylander L, Holmqvist M, Gustafson L, Gillberg C. ADHD in adult psychiatry. Minimum rates and clinical presentation in general psychiatry outpatients. *Nordic J Psychiatr*. 2009;63(1):64–71.
- Deberdt W, Thome J, Lebec J, Kraemer S, Fregenal I, Ramos-Quiroga JA, et al. Prevalence of ADHD in nonpsychotic adult psychiatric care (ADPSYC): A multinational cross-sectional study in Europe. *BMC Psychiatr*. 2015;15:242–015–0624-5.
- Costa PT Jr, McCrae RR. Domains and facets: hierarchical personality assessment using the revised NEO personality inventory. *J Pers Assess*. 1995 Feb;64(1):21–50. https://doi.org/10.1176/s15327752jpa6401_2.
- Nigg JT, Karalunas SL, Feczko E, Fair DA. Toward a revised nosology for attention-deficit/hyperactivity disorder heterogeneity. *Biol Psychiatry Cogn Neurosci Neuroimaging*. 2020 Aug;5(8):726–37. <https://doi.org/10.1016/j.bpsc.2020.02.005>.
- Kessler RC, Adler L, Barkley R, Biederman J, Conners CK, Demler O, et al. The prevalence and correlates of adult ADHD in the United States: results from the National Comorbidity Survey Replication. *Am J Psychiatry*. 2006;163(4):716–23. <https://doi.org/10.1176/ajp.2006.163.4.716>.
- Martel MM. Research review: a new perspective on attention-deficit/hyperactivity disorder: emotion dysregulation and trait models. *J Child Psychol Psychiatry*. 2009 Sep;50(9):1042–51. <https://doi.org/10.1111/j.1469-7610.2009.02105.x>.
- Reimherr FW, Marchant BK, Gift TE, Steans TA, Wender PH. Types of adult attention-deficit hyperactivity disorder (ADHD): baseline characteristics, initial response, and long-term response to treatment with methylphenidate. *Atten Defic Hyperact Disord*. 2015 Jun;7(2):115–28. <https://doi.org/10.1007/s12402-015-0176-z>.
- Matthies SD, Philipsen A. Common ground in Attention Deficit Hyperactivity Disorder (ADHD) and Borderline Personality Disorder (BPD)-review of recent findings. *Borderline Personal Disord Emot Dysregul*. 2014;1:3 6673-1-3. eCollection 2014.
- Reimherr FW, Marchant BK, Strong RE, Hedges DW, Adler L, Spencer TJ, et al. Emotional dysregulation in adult ADHD and response to atomoxetine. *Biol Psychiatry*. 2005;58(2):125–31. <https://doi.org/10.1016/j.biopsych.2005.04.040>.
- Bo Bach MS, Huxley E. Related personality disorders located within an elaborated externalizing psychopathology Spectrum. *The Wiley Handbook of Disruptive and Impulse-Control Disorders*. 2017;2021/01:103–23.
- Insel T, Cuthbert B, Garvey M, Heinssen R, Pine DS, Quinn K, et al. Research domain criteria (RDoC): toward a new classification framework for research on mental disorders. *Am J Psychiatry*. 2010;167(7):748–51. <https://doi.org/10.1176/appi.ajp.2010.09091379>.
- Kotov R, Krueger RF, Watson D, Achenbach TM, Althoff RR, Bagby RM, et al. The hierarchical taxonomy of psychopathology (HiTOP): a dimensional alternative to traditional nosologies. *J Abnorm Psychol*. 2017;126(4):454–77. <https://doi.org/10.1037/abn0000258>.
- American Psychiatric Association. American Psychiatric Association, DSM-5 Task Force. Diagnostic and statistical manual of mental disorders : DSM-5. Arlington, VA: American Psychiatric Association; 2013.
- World Health Organization (WHO). The ICD-10 classification of mental and behavioural disorders: Diagnostic criteria for research. Genève: World Health Organization; 1993.
- Morey LC, Hopwood CJ, Markowitz JC, Gunderson JG, Grilo CM, McGlashan TH, et al. Comparison of alternative models for personality disorders, II: 6-, 8- and 10-year follow-up. *Psychol Med*. 2012;42(8):1705–13. <https://doi.org/10.1017/S0033291711002601>.
- Nigg JT, John OP, Blaskey LG, Huang-Pollock C, Willcutt EG, Hinshaw SP, et al. Big five dimensions and ADHD symptoms: links between personality traits and clinical symptoms. *J Pers Soc Psychol*. 2002;83(2):451–69. <https://doi.org/10.1037/0022-3514.83.2.451>.
- Pinzone V, De Rossi P, Trabucchi G, Lester D, Girardi P, Pompili M. Temperament correlates in adult ADHD: a systematic review (bigstar bigstar). *J Affect Disord*. 2019;252:394–403. <https://doi.org/10.1016/j.jad.2019.04.006>.
- Gomez R, Van Doorn G, Watson S, Gomez A, Stavropoulos V. Cloninger's personality dimensions and ADHD: A meta-analytic review. *Personality and Individual Differences*. 2017;107:219–27.
- Connor DF, Ford JD. Comorbid symptom severity in attention-deficit/hyperactivity disorder: a clinical study. *J Clin Psychiatry*. 2012;73(5):711–7. <https://doi.org/10.4088/JCP.11m07099>.
- Achenbach TM. The classification of children's psychiatric symptoms: a factor-analytic study. *Psychol Monogr*. 1966;80(7):1–37. <https://doi.org/10.1037/h0093906>.
- McCrae RR, Costa PT Jr. NEO inventories, Professional Manual: PAR; 2010.
- Lahey BB. Public health significance of neuroticism. *Am Psychol*. 2009;64(4):241–56. <https://doi.org/10.1037/a0015309>.
- Gore WL, Widiger TA. The DSM-5 dimensional trait model and five-factor models of general personality. *J Abnorm Psychol*. 2013;122(3):816–21. <https://doi.org/10.1037/a0032822>.
- Watson D, Stasik SM, Ro E, Clark LA. Integrating normal and pathological personality: relating the DSM-5 trait-dimensional model to general traits of personality. *Assessment*. 2013;20(3):312–26. <https://doi.org/10.1177/1073191113485810>.
- Skodol AE, Clark LA, Bender DS, Krueger RF, Morey LC, Verheul R, et al. Proposed changes in personality and personality disorder assessment and diagnosis for DSM-5 part I: description and rationale. *Personal Disord*. 2011;2(1):4–22. <https://doi.org/10.1037/a0021891>.
- McCrae RR, Löckenhoff CE, Costa PT Jr. A step toward DSM-V: cataloguing personality-related problems in living. *Eur J Pers*. 2020;19(4):269–86 2005 06/01.
- Widiger TA. The Oxford handbook of the five factor model; 2017.
- Decuyper M, De Pauw S, De Fruyt F, De Bolle M, De Clercq BJ. A meta-analysis of psychopathy-, antisocial PD- and FFM associations. *Eur J Pers*. 2020;23(7):531–65 2009 11/01.
- Jones SE, Miller JD, Lynam DR. Personality, antisocial behavior, and aggression: a meta-analytic review. *J Crim Just*. 2011;39(4):329–37. <https://doi.org/10.1016/j.jcrimjus.2011.03.004>.
- De Fruyt F, De Clercq B, De Bolle M, Wille B, Markon K, Krueger RF. General and maladaptive traits in a five-factor framework for DSM-5 in a university student sample. *Assessment*. 2013;20(3):295–307. <https://doi.org/10.1177/1073191113475808>.
- Griffin SA, Samuel DB. A closer look at the lower-order structure of the personality inventory for DSM-5: comparison with the five-factor model. *Personal Disord*. 2014;5(4):406–12. <https://doi.org/10.1037/per0000074>.
- Livesley WJ, Jang KL, Vernon PA. Phenotypic and genetic structure of traits delineating personality disorder. *Arch Gen Psychiatry*. 1998;55(10):941–8. <https://doi.org/10.1001/archpsyc.55.10.941>.
- Crego C, Samuel DB, Widiger TA. The FFOCI and other measures and models of OCPD. *Assessment*. 2015 Apr;22(2):135–51. <https://doi.org/10.1177/1073191114539382>.
- Samuel DB, Gore WL. Maladaptive Variants of Conscientiousness and Agreeableness. *J Pers*. 2020;80(6):1669–96 2012 12/01.
- O'Connor BP. A quantitative review of the comprehensiveness of the five-factor model in relation to popular personality inventories. *Assessment*. 2002;9(2):188–203. <https://doi.org/10.1177/1079102009002010>.
- Markon KE, Krueger RF, Watson D. Delineating the structure of normal and abnormal personality: an integrative hierarchical approach. *J Pers Soc Psychol*. 2005;88(1):139–57. <https://doi.org/10.1037/0022-3514.88.1.139>.
- John O, Naumann L, Soto C. Paradigm shift to the integrative big five trait taxonomy: History, measurement, and conceptual issues; 2008. p. 114–58.
- Livesley WJ, Jackson DN. Manual for the Dimensional Assessment Of Personality Pathology — Basic Questionnaire. Port Huron: Sigma Press; 2009.
- Cloninger CR, Svrakic DM, Przybeck TR. A psychobiological model of temperament and character. *Arch Gen Psychiatry*. 1993 Dec;50(12):975–90. <https://doi.org/10.1001/archpsyc.1993.01820240059008>.
- Ruggero CJ, Kotov R, Hopwood CJ, First M, Clark LA, Skodol AE, et al. Integrating the hierarchical taxonomy of psychopathology (HiTOP) into

- clinical practice. *J Consult Clin Psychol.* 2019;87(12):1069–84. <https://doi.org/10.1037/ccp0000452>.
42. Conway CC, Forbes MK, Forbush KT, Fried EI, Hallquist MN, Kotov R, et al. A Hierarchical Taxonomy of Psychopathology Can Transform Mental Health Research. *Perspect Psychol Sci* 2019 05/01. 2020;14(3):419–36.
43. Hopwood C, Bagby R, Gralnick T, Ro E, Ruggero C, Mullins-Sweatt S, et al. Integrating Psychotherapy With the Hierarchical Taxonomy of Psychopathology (HiTOP). *J Psychotherapy Integration.* 2020;30:477–97.
44. Hopwood CJ. A framework for treating DSM-5 alternative model for personality disorder features. *Personality and Mental Health* 2018 05/01. 2020;12(2):107–25.
45. Nigg JT. Temperament and developmental psychopathology. *J Child Psychol Psychiatry.* 2006;47(3–4):395–422. <https://doi.org/10.1111/j.1469-7610.2006.01612.x>.
46. Knouse LE, Traeger L, O'Leirigh C, Safren SA. Adult attention deficit hyperactivity disorder symptoms and five-factor model traits in a clinical sample: a structural equation modeling approach. *J Nerv Ment Dis.* 2013; 201(10):848–54. <https://doi.org/10.1097/NMD.0b013e3182a5bf33>.
47. Gomez R, Corr PJ. ADHD and personality: a meta-analytic review. *Clin Psychol Rev.* 2014;34(5):376–88. <https://doi.org/10.1016/j.cpr.2014.05.002>.
48. Kotov R, Gamez W, Schmidt F, Watson D. Linking "big" personality traits to anxiety, depressive, and substance use disorders: a meta-analysis. *Psychol Bull.* 2010;136(5):768–821. <https://doi.org/10.1037/a0020327>.
49. Wallace A, Ullsperger JM, Nikolas MA. Do personality traits explain the association between childhood attention-deficit hyperactivity disorder symptoms and substance use and problems in young adults? *Personality and Individual Differences.* 2016;92:22–8.
50. Di Nicola M, Sala L, Romo L, Catalano V, Even C, Dubertret C, et al. Adult attention-deficit/hyperactivity disorder in major depressed and bipolar subjects: role of personality traits and clinical implications. *Eur Arch Psychiatry Clin Neurosci.* 2014;264(5):391–400. <https://doi.org/10.1007/s00406-013-0456-6>.
51. Jacob CP, Gross-Lesch S, Reichert S, Geissler J, Jans T, Kittel-Schneider S, et al. Sex- and Subtype-Related Differences of Personality Disorders (Axis II) and Personality Traits in Persistent ADHD. *J Attention Disord.* 2016;20(12): 1056–65.
52. Koerting J, Pukrop R, Klein P, Ritter K, Knowles M, Banzhaf A, et al. Comparing Dimensional Models Assessing Personality Traits and Personality Pathology Among Adult ADHD and Borderline Personality Disorder. *J Attention Disord.* 2016;20(8):715–24.
53. Steinhausen HC, Gollner J, Brandeis D, Muller UC, Valko L, Drechsler R. Psychopathology and personality in parents of children with ADHD. *J Atten Disord.* 2013;17(1):38–46. <https://doi.org/10.1177/1087054711427562>.
54. Ashton MC, Lee K, Perugini M, Szarota P, de Vries RE, Di Blas L, et al. A six-factor structure of personality-descriptive adjectives: solutions from psycholexical studies in seven languages. *J Pers Soc Psychol.* 2004;86(2): 356–66. <https://doi.org/10.1037/0022-3514.86.2.356>.
55. He JA, Antshel KM, Biederman J, Faraone SV. Do personality traits predict functional impairment and quality of life in adult ADHD? A controlled study. *J Atten Disord.* 2019;23(1):12–21. <https://doi.org/10.1177/1087054715613440>.
56. Faraone SV, Kunwar A, Adamson J, Biederman J. Personality traits among ADHD adults: implications of late-onset and subthreshold diagnoses. *Psychol Med.* 2009;39(4):685–93. <https://doi.org/10.1017/S0033291708003917>.
57. Sizoo B, van den Brink W, van Eenige MG, van der Gaag RJ. Personality characteristics of adults with autism spectrum disorders or attention deficit hyperactivity disorder with and without substance use disorders. *J Nerv Ment Dis.* 2009;197(6):450–4. <https://doi.org/10.1097/NMD.0b013e3181a61dd0>.
58. Perroud N, Hasler R, Golay N, Zimmermann J, Prada P, Nicastro R, et al. Personality profiles in adults with attention deficit hyperactivity disorder (ADHD). *BMC Psychiatr.* 2016;16:199.
59. Sizoo BB, van der Gaag RJ, van den Brink W. Temperament and character as endophenotype in adults with autism spectrum disorders or attention deficit/hyperactivity disorder. *Autism : the international journal of research and practice.* 2015;19(4):400–8. <https://doi.org/10.1177/1362361314522352>.
60. Anckarsäter H, Stahlberg O, Larson T, Hakansson C, Jutblad S, Niklasson L, et al. The impact of ADHD and autism spectrum disorders on temperament, character, and personality development. *Am J Psychiatry.* 2006;163(7):1239–44.
61. Mesrobian SK, Villa AEP, Bader M, Gotte L, Lintas A. Event-related potentials during a gambling task in young adults with attention-deficit/hyperactivity disorder. *Front Hum Neurosci.* 2018;12:79.
62. Salgado CA, Bau CH, Grevet EH, Fischer AG, Victor MM, Kalil KL, et al. Inattention and hyperactivity dimensions of ADHD are associated with different personality profiles. *Psychopathology.* 2009;42(2):108–12. <https://doi.org/10.1159/000203343>.
63. Suurmond R, van Rhee H, Hak T. Introduction, comparison, and validation of Meta-Essentials: A free and simple tool for meta-analysis. *Res Syn Meth* 2017 12/01. 2020;8(4):537–53.
64. Van Rhee HJ, Suurmond R, Hak T. User manual for meta-essentials: workbooks for meta-analysis (version 1.4) Rotterdam, The Netherlands: Erasmus Research Institute of Management. Retrieved from www.erim.eur.nl/research-support/meta-essentials. 2015.
65. Hak T, van Rhee H, Suurmond R. How to Interpret Results of Meta-Analysis. Available at SSRN: <https://ssrn.com/abstract=3241367>. <http://dx.doi.org/https://doi.org/10.2139/ssrn.3241367>. 2016.
66. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ.* 2003;327(7414):557–60. <https://doi.org/10.1136/bmj.327.7414.557>.
67. Whiting P, Rutjes AWS, Reitsma JB, Bossuyt PMM, Kleijnen J. The development of QUADAS: a tool for the quality assessment of studies of diagnostic accuracy included in systematic reviews. *BMC Med Res Methodol.* 2003;3(1):25.
68. Cohen J. Statistical Power Analysis. *Curr Dir Psychol Sci* 1992 06/01. 2020; 1(3):98–101.
69. Sawilowsky S. New Effect Size Rules of Thumb. *J Modern Appl Stat Methods.* 2009;8:597–9.
70. Subgroup Analyses. *Introduction to Meta-Analysis* 2009 03/13; 2020. p. 149–86.
71. Barkley RA. Attention-deficit hyperactivity disorder: a handbook for diagnosis and treatment. 4th ed. New York; New York: The Guilford Press; The Guilford Press; 2015.
72. Brown TE. ADHD comorbidities: handbook for ADHD complications in children and adults. Washington, DC: American Psychiatric Pub; 2009.
73. Barkley RA. Behavioral inhibition, sustained attention, and executive functions: constructing a unifying theory of ADHD. *Psychol Bull.* 1997;121(1): 65–94. <https://doi.org/10.1037/0033-2909.121.1.65>.
74. Smith TE, Samuel DB. A multi-method examination of the links between ADHD and personality disorder. *J Personal Disord.* 2017;31(1):26–48. https://doi.org/10.1521/pedi_2016_30_236.
75. Hopwood CJ, Morey LC. Emotional Problems Suppress Disorder/ Performance Associations in Adult ADHD Assessment. *J Psychopathol Behav Assessment.* 2008;30(3):204–10.
76. Karalunas SL, Fair D, Musser ED, Aykes K, Iyer SP, Nigg JT. Subtyping attention-deficit/hyperactivity disorder using temperament dimensions: toward biologically based nosologic criteria. *JAMA Psychiatry.* 2014;71(9): 1015–24. <https://doi.org/10.1001/jamapsychiatry.2014.763>.
77. Karalunas SL, Nigg JT. Heterogeneity and subtyping in attention-deficit/hyperactivity disorder-considerations for emerging research using person-centered computational approaches. *Biol Psychiatry.* 2020;88(1):103–10. <https://doi.org/10.1016/j.biopsych.2019.11.002>.
78. Smith TE, Martel MM. Trait-Based Profiles of ADHD in Adolescents and Young Adults. *J Clin Child Adolescent Psychol.* 2018;53:1–15.
79. Bach B, Markon K, Simonsen E, Krueger RF. Clinical utility of the DSM-5 alternative model of personality disorders: six cases from practice. *J Psychiatr Pract.* 2015;21(1):3–25. <https://doi.org/10.1097/01.pra.0000460618.02805.ef>.
80. Bodemer N, Hanoch Y, Katsikopoulos KV. Heuristics: foundations for a novel approach to medical decision making. *Internal and Emergency Med.* 2015; 10(2):195–203.
81. Reed GM, First MB, Kogan CS, Hyman SE, Gureje O, Gaebel W, et al. Innovations and changes in the ICD-11 classification of mental, behavioural and neurodevelopmental disorders. *World Psychiatry.* 2019;18(1):3–19. <https://doi.org/10.1002/wps.20611>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.