

Lawrence Berkeley National Laboratory

Recent Work

Title

SUMMARY OF THE RESEARCH PROGRESS MEETING OF JAN. 18, 1951.

Permalink

<https://escholarship.org/uc/item/90n3863p>

Author

Folden, Margaret Foss.

Publication Date

1951-06-26

UNIVERSITY OF CALIFORNIA - BERKELEY

UCRL- 1245

UNCLASSIFIED

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

RADIATION LABORATORY

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UNIVERSITY OF CALIFORNIA

Radiation Laboratory

Contract No. W-7405-eng-48

SUMMARY OF THE RESEARCH PROGRESS MEETING OF JANUARY 18, 1951

Margaret Foss Folden

June 26, 1951

"Some of the results reported in this document may be of a preliminary or incomplete nature. It is the request of the Radiation Laboratory that the document not be circulated off the project nor the results quoted without permission."

Berkeley, California

SUMMARY OF THE RESEARCH PROGRESS MEETING OF JANUARY 18, 1951

Margaret Foss Folden

Radiation Laboratory, Department of Physics
University of California, Berkeley, California

June 26, 1951

Physics at Duke University - H. W. Newson.

Studies at Duke University on level separations in compound nuclei after the absorption of a neutron were described. The rising levels of excitation measured by time of flight studies were described. Bethe's results and those at Duke were compared. Their results indicate with a surprising constancy that the spacing from neutron 10 to neutron 50 is about 5×10^5 volts. Studies are being made in the region of $N = 28$ to 45 with Sc^{81} , Cu^{65} and As^{75} have been investigated.

Measurement of Liver Circulation - Ernest L. Dobson.

The circulatory system was first described in general.

In these experiments the quartz rod transillumination technique was used to obtain information on liver circulation, and injections of india ink made observations more certain. The phagocytes interlaced among the sinusoids take up certain colloidal material; this fact was used to study the liver circulation. P^{32} , labeled chromic phosphate colloid, was injected into the blood. A short mixing time of about two minutes for large animals is required. The logarithm disappearance equation obtained from this treatment is given by:

$$\text{concentration} = ce^{-(K)t}.$$

The disappearance rate is measured to determine K. Assume that all the blood going to the liver is cleared of the colloid, then a measure of the rate of blood flow through the liver is obtained. The spleen takes out from 1 to 20 percent of the colloid, but since these two organs are in series it does not matter.

The lymph nodes do not take out the colloid. Bone marrow takes up a small amount of the chromic phosphate, so the equation must be written

$$C = ce^{-(m_l k_l + n_{bm} k_{bm})t}$$

where l = liver and bm = bone marrow. The actual disappearance curve is shown in Fig. 1. The three components shown can not be explained on the basis of liver efficiency, taking out the large particles and letting the small ones go through. In the samples of blood early in the curve centrifugation has removed the colloidal material; however, late in the curve material was not removed. The colloidal particles varied from .01 to 5 microns.

The blood tissue perfusion factor is the factor most constant from one animal to another. It has been found that in man the factor is 1.2 volume blood per volume liver per unit of time. In chickens the factor is 5.0; in dogs 1.2; in the anesthetized mouse 1.7; in the normal mouse 1.4; in the normal rabbit 0.74; and in the anesthetized rabbit 1.12.

Photographic Fourier Summations - David H. Templeton.

Studies have been made on the density of electrons using the equation:

$$\rho(xy) = \sum \sum_{hko} F_{hk} \cos(hx + ky) 2\pi$$

where $\rho(xy)$ = projection on the xy plane

F = structure factor

$|F|^2$ kI = intensity of Laue spots.

This method is a photographic way of representing the theory and makes use of photographic masks. The light transmitted is proportional to

$$1 + \cos(hx + ky) 2\pi.$$

The masks are exposed for a length of time proportional to F . When the film is developed the dark spots give the location of the atoms; that is, the method

determines h and k by the process of approximating the Laue pattern by a succession of superposed patterns with known h and k. Figure 1 gives examples of the patterns used corresponding to typical terms in the Fourier series. Thorium boride, yttrium fluoride and rhenium bromide are among the compounds investigated.

Information Division
6/26/51 nw




Fig. 1