

UC-70
LBL-23636
c.1



Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

EARTH SCIENCES DIVISION

RECEIVED
LAWRENCE
BERKELEY LABORATORY

AUG 21 1987

LIBRARY AND
DOCUMENTS SECTION


To be presented at the American Nuclear Society
Winter Meeting, Los Angeles, CA, November 15-19, 1987

PRESSURE-INDUCED BRINE MIGRATION INTO AN OPEN BOREHOLE IN A SALT REPOSITORY

Y. Hwang, P.L. Chambré, W.W.-L. Lee, and T.H. Pigford

June 1987

For Reference
Not to be taken from this room



LBL-23636
c.1

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Pressure-Induced Brine Migration into an Open Borehole in a Salt Repository*

Y. Hwang, P. L. Chambré, W. W.-L. Lee and T. H. Pigford
Department of Nuclear Engineering & Lawrence Berkeley Laboratory
University of California, Berkeley, CA 94720

In the *Environmental Assessment*¹ for a potential nuclear waste repository in salt, estimates of release rates of radionuclides from the engineered barrier system were made by multiplying the brine flow rate into an assumed open borehole at 300 years by the solubilities of radioelements in the waste package. The borehole was conservatively assumed to remain open, at atmospheric pressure, for hundreds and thousands of years. Brine was assumed to migrate into the borehole at a rate given by the Jenks equation² for temperature-gradient induced movement of brine inclusions within individual salt crystals. However, it is now recognized that inclusions that reach grain boundaries are likely to move along grain boundaries under the influence of a pressure gradient, along with brine originally present in the grain boundaries. Here we present estimates of the rate of brine accumulation in an assumed open borehole, based on grain-boundary migration theory.

Within a few years after emplacement, brine inclusions inside salt crystals within a few meters of a hot waste package will have migrated to the grain boundaries. Pressure gradients to cause grain-boundary migration result from the difference between the far-field hydrostatic pressure of brine in the undisturbed salt and the assumed atmospheric pressure in the borehole, and from the pressure gradients resulting from thermal expansion of the salt and brine in the time-dependent temperature field. The formulation of brine movement with salt as a thermoelastic porous medium, in the context of the continuum theory of mixtures, was published by McTigue.³ Chambré obtained the analytic solutions to the governing equations for a spherical-equivalent waste form and to the coupled radionuclide transport problem, driven by thermoelastic effects.⁴ In numerical calculations we found that the thermoelastic effects are minute for this case and the McTigue equations reduce to those of standard time-dependent pressure-driven porous media flow.

We present some numerical illustrations of the solution for a spherical-equivalent waste package embedded in an infinite salt medium, using a permeability of 10^{-21} m², a value suggested by McTigue³. We assume

*Work supported in part by U. S. Department of Energy via Contract DE-AC03-76SF00098

a far-field lithostatic pressure of 16.3 MPa and obtain the predicted velocity of brine flow into the open borehole shown in Figure 1. The velocity is small and rapidly approaches steady state, in contrast to the analysis in the *Environmental Assessment* where steady state is reached only after several hundred years. Figure 2 is a comparison of the cumulative brine flow volumes predicted by various models. In Figure 2 the *Environmental Assessment* prediction of cumulative brine inflow, primarily thermally-driven, is shown along with our results for pressure gradient flow. Also shown are cumulative brine flows calculated using McTigue's model equations. For the parameter values adopted herein, grain-boundary migration into an open borehole is driven almost entirely by the large difference between pressures in the far field and in the borehole. On the long time scales of Figure 2, brine is calculated to accumulate linearly with time. It does not level off after about 1,000 years as is predicted in the *Environmental Assessment*:

We have presented selected results from a detailed model for brine movement in the first few years after emplacement. Companion papers deal with brine migration after the borehole has closed, and possible simplification in the analysis.^{5,6}

References

1. U.S. Department of Energy, 1986, *Environmental Assessment, Deaf Smith County Site, Texas*, DOE/RW-0069.
2. G. H. Jenks and H. C. Claiborne, 1981, *Brine Migration in Salt and its Implications in the Geologic Disposal of Nuclear Waste*, ORNL-5818
3. D. T. McTigue, 1986, "Thermoelastic Response of Fluid-Saturated, Porous Rock," *J. Geophys. Res.*, 91, B9, 9533
4. P. L. Chambré, To be published
5. Y. Hwang, P. L. Chambré, W. W.-L. Lee and T. H. Pigford, "Pressure-Induced Brine Migration in Consolidated Salt in a Repository," UCB-NE-4101, Paper submitted for the 1987 Winter Meeting, American Nuclear Society
6. P. L. Chambré, Y. Hwang, W. W.-L. Lee and T. H. Pigford, "Release Rates in Waste Packages in a Salt Repository," UCB-NE-4102, Paper submitted for the 1987 Winter Meeting, American Nuclear Society




Figure 1. Brine Migration Velocity into an Open Borehole Due to Pressure Difference




Figure 2. Brine Accumulation in an Open Borehole Calculated by Three Methods

- Note 1 Primarily thermally-driven
- Note 2 Primarily pressure-difference driven

*LAWRENCE BERKELEY LABORATORY
TECHNICAL INFORMATION DEPARTMENT
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720*