UC San Diego

UC San Diego Previously Published Works

Title

Cyclohexeno[3,4]cyclodec-1,5-diyne-3-ene: A Convenient Enediyne.

Permalink

<https://escholarship.org/uc/item/9213x2w2>

Journal

Organic letters, 23(17)

ISSN

1523-7060

Authors

Shrinidhi, Annadka

Perrin, Charles L

Publication Date

2021-09-01

DOI

10.1021/acs.orglett.1c02459

Peer reviewed

Cyclohexeno[3,4]cyclodec-1,5-diyne-3-ene: A Convenient Enediyne

Annadka Shrinidhi and Charles L. Perrin

Department of Chemistry, University of California—San Diego, La Jolla, California 92093-0358, USA
Supporting Information Placeholder

ABSTRACT: Enediynes are widely studied to understand their cycloaromatization and the trapping of the resulting *p*-dehydrobenzene diradical. However, few model substrates are known, and they are hard to synthesize and difficult to handle. Herein we report cyclohexeno[3,4]cyclodec-1,5-diyne-3-ene as a convenient model for studying reactivity of enediynes. It can be easily synthesized from 1,2-diethynylcyclohexene and 1,4-diiodobutane. It is a solid, stable at room temperature. In solution the *p*-dehydrobenzene diradical derived from its cycloaromatization can be trapped by nucleophiles. The rate-limiting step is the cyclization, slightly slower than parent cyclodec-1,5-diyne-3-ene but faster than its benzo analog, consistent with the distances between the reacting carbon atoms.

INTRODUCTION

Enediynes are one of the common scaffolds in medically important organic compounds.¹ However, the lability of their enediyne core limits their medicinal use because of cytotoxicity and nonspecificity, especially as anticancer antibiotics.² This behavior triggered studies toward understanding the reactivity of enediynes, particularly their cycloaromatization to a *p*-dehydrobenzene ("p-benzyne", a term discouraged by IUPAC) diradical, a reaction known as the Bergman cyclization.³ Many researchers have explored the effects on cyclization of ring strain,⁴ conformational and electronic effects,⁵ and metal-ion coordination,⁶ in order to design enediynes for medicinal purposes.^{1f,7} Most of those studies used model enediynes, either cyclodec-1,5-diyne-3-ene (**1**) or its homologues, or its derivatives, such as 3,4-benzocyclodec-3-ene-1,5-diyne (**2**).^{4a,8}

Some earlier studies explored the reaction of enediyne **1** with nucleophiles. The reaction of **1** with halides follows kinetics first-order in enediyne and zero-order in both halide and carboxylic acid.^{8b} Therefore the rate-limiting step is the cycloaromatization, and the nucleophilic addition to the *p*-dehydrobenzene diradical is fast,^{8c} as is protonation of the resulting "naked" aryl anion, by water, carboxylic acid, or deuterated solvent.^{8c} This mechanism is shown in Scheme 1 for a more general series of enediynes. The novel feature is the nucleophilic trapping of the diradical, in contrast to its expected reaction by radical pathways.

Scheme 1. Reactions of enediynes 1 (R = H), 2 (RR = (CH)₄), and 3 (RR = (CH₂)₄).

However, detailed studies of enediyne cycloaromatization and the trapping of the resulting *p*-dehydrobenzene diradical, as well as the reactivity and selectivity of the aryl anion, are challenging. One of the obstacles to thorough studies of enediyne **1** has been its low yield in synthesis. Jones et al. reported an isolated yield of 95% using an exceptional HMPA analog "TEP" as an additive whose identity was never revealed.⁹ In contrast, the synthesis of **1** using the more usual HMPA as additive consistently produced a yield of only 5-10%.¹⁰ An alternative but more circuitous procedure produced a 12% overall yield.^{4b} Furthermore, handling of **1** is challenging. It is a volatile and irritating liquid, difficult to isolate. Because it decomposes within a day at room temperature it must be stored in solution at -35°C or below.

Therefore alternative enediynes continue to be developed. Basak et al. studied the regioselectivity of nucleophilic addition to unsymmetrical diradicals derived from azaenediynes.¹¹ Many researchers studied enediyne **2**, for which a simple synthesis provides a 48% yield.¹² However, its disappearance does not follow simple first-order kinetics, because it was observed that the efficiency of trapping of its *p*-dehydrobenzene diradical depends on the concentration of 1,4-cyclohexadiene.¹² Nor are the kinetics zero-order in nucleophile.¹⁰ Therefore it can be concluded that the initial cycloaromatization must be reversible.

Another enediyne is desirable. An ideal candidate is cyclohexeno[3,4]cyclodec-1,5-diyne-3-ene (**3**, IUPAC name bicyclo[8.4.0]tetradeca-1,7-diyne-13-ene), a cyclohexene-fused cyclodec-1,5-diyne-3-ene. The additional alkylation increases the stabilities of both the enediyne and its cycloaromatization product. Herein we report the robust synthesis of **3**, explore its cycloaromatization, study the reaction of nucleophiles with diradicals formed from **3**, and determine the kinetics of disappearance of **3**. We also report that cyclopentenoenediyne **4** has been too elusive.

Enediyne **3** has the further feature that reversal of the initial cycloaromatization can produce not only the original enediyne but also an identical isomer ("automer"), as shown in Scheme 2. In contrast, reversal of the cycloaromatization of

enediyne **2** would produce a dienediyne, although no evidence was found for that complication.¹² Even if this rearrangement occurs for **3**, it cannot complicate the kinetics.

Scheme 2. Degenerate rearrangement of

enediyne **3**.

EXPERIMENTAL

Synthesis. Cyclodec-1,5-diyne-3-ene (**1**) was prepared by cyclization of 1,10-dibromo-2,8-octadiyne via carbenoid coupling-elimination according to the modified procedure,¹⁰ derived from the reported literature.⁹ 3,4-Benzocyclodeca-1,5-diyne-3-ene (**2**) was prepared by cyclization of 1,2-diethynylbenzene with 1,4-diiodobutane according to the literature procedure.¹² Cyclohexeno[3,4]cyclodec-1,5-diyne-3-ene (**3**) was prepared by cyclization of 1,2-diethynylcyclohexene with 1,4-diiodobutane. Details are provided in Supporting Information. Enediyne **3** was found to be an off-white solid of mp 105-106°C. It is stable under ambient conditions for more than a month and can be stored in powder form at -10°C for more than a year with no apparent decomposition.

Kinetics. Enediyne was added to a solution of nucleophile X⁻ and pivalic acid (or with added 1,4-cyclohexadiene for enediyne **2**) in DMSO-*d*₆ containing 1,3,5-trichlorobenzene as internal standard. The sample was heated in an oil bath. Disappearance of starting material was monitored every 1.0 h by removing the sample and analyzing it by ¹H NMR. The first-order rate constant *k* was obtained by least-squares analysis of a logarithmic plot of [enediyne] versus time (Eq. 1). Rate constants *k* and free energies of activation (Δ*G*[‡]) for the disappearance of enediynes **1-3** in the presence of LiX and pivalic acid in DMSO-*d*₆ are given in Table S1. Details are provided in Supporting Information.

$$\ln ([\mathbf{1-3}]_t) = -kt + \ln ([\mathbf{1-3}]_0) \quad (1)$$

RESULTS

In order to assign the rate-limiting step for the transformation of enediyne **3** plus nucleophile X⁻ to 9-X-1,2,3,4,5,6,7,8-octahydroanthracene (**5X**), we have measured the kinetics of these reactions. The initial concentration of **3** was kept low while the concentration of nucleophile or acid was varied. Table 1 lists rate constants and yields of **5X** for reactions of enediyne **3** with MX in DMSO-*d*₆ containing pivalic acid. The first-order

rate constants for these reactions at 75°C lie between 5.1 and 6.2 x 10⁻⁵ s⁻¹.

Table 1. Rate constants and yields of 5X for reactions of enediyne 3 (5 mM) with MX and pivalic acid (HA) in DMSO-d₆ at 75°C.

MX	[X] ⁻ ₀ (mM)	[HA] ₀ (mM)	10 ⁵ k (s ⁻¹)	5X, Yield ^a (%)
Lil	500	50	5.23	>99
Lil	250	25	5.21	>99
Bu ₄ NI	250	25	5.16	>99
LiBr	500	50	5.63	82
LiBr	250	25	5.60	83
Bu ₄ NBr	250	25	5.57	85
NaCN	500	50	5.78	78
NaCN	250	25	5.75	79
Bu ₄ NCN	250	25	5.70	81
KSCN	500	50	6.16	66
KSCN	250	25	6.12	68
Bu ₄ NSC N	250	25	6.03	69
none	0	50	8.43	43 ^b

^aAverage of ¹H NMR and GC-FID yields. ^bProduct is 5H.

The rate is independent of the concentration of nucleophile or of acid, but not entirely independent of nucleophile. It is slightly lower than the rate of decomposition of enediyne **3** in the absence of nucleophiles. That is an indication that the variability is due to the incursion of polymerization, which occurs when the *p*-dehydrobenzene diradical reacts with enediyne. Nucleophilic capture of the diradical prevents polymerization. According to computations,^{8c} that capture is encounter-controlled, with no activation energy other than that required for desolvation of the nucleophile. Consequently smaller ions, which are more strongly solvated, have a higher activation energy and are less effective in preventing polymerization. Therefore we conclude that the most reliable rate constant for cycloaromatization of **3** is (5.20±0.025) x 10⁻⁵ s⁻¹, as observed with I⁻ as nucleophile, which most effectively intercepts the diradical before it can induce polymerization. Further studies with a mixture of competing nucleophiles will assess the role of solvation.

The data show that the observed rate constant for disappearance of enediyne **3** is independent of the concentration of nucleophile. Therefore the kinetics are simply first-order in [**3**] and zero-order in [I⁻]. This is the same result as for **1**,^{8c} for which it was concluded that the rate-limiting step is the initial cyclization to a *p*-dehydrobenzene diradical, which is attacked by nucleophile in a fast second step, followed by a rapid protonation of the "naked" aryl anion (Scheme 1).^{8d}

The free energies of activation for the conversion of enediynes **1-3** to their diradicals were evaluated from the rate constants of enediyne decomposition in DMSO-d₆ in the presence of Lil and pivalic acid. The values are tabulated in Table 2.

Table 2. Free energy of activation for the conversion of enediynes (1-3) to the corresponding *p*-dehydrobenzene diradicals

Enediyne	1	2	3
ΔG [‡] , kcal- mol ⁻¹	25.3±0.0	28.6±0.1	27.2±0.1
Lit.	24.8 ^{4b,27} , 4 ¹³	29.5 ¹²	---

For comparison with activation energies, calculated 1,6 distances between the two carbon atoms that form the new C-C bond during *p*-dehydrobenzene formation are given in Table 3.

Table 3. C1-C6 Distances (Å) in enediynes

Enediyne	1	2	3
<i>d</i>	3.26	3.46	3.32
Lit.	3.25 ^{4b}	---	---

DISCUSSION

By following Semmelhack's protocol,¹² macrocyclization of 1,2-diethynylcyclohexene with 1,4-diiodobutane produces enediyne **3** as a major product (45%). The fact that enediyne **3** is solid and stable under ambient conditions allows easy handling and accurate measurement of reactivity and kinetics.

Reaction of enediyne **3** with nucleophiles.

On heating, the *p*-dehydrobenzene diradical generated from enediyne **1** reacts with nucleophiles such as I⁻, SCN⁻, Br⁻, NC⁻, and N₃⁻.^{8c} Similarly, the *p*-dehydrobenzene diradical generated from enediyne **2** reacts with I⁻, SCN⁻, and Br⁻.¹⁰ In order to study cycloaromatization and nucleophilic addition to the resulting *p*-dehydrobenzene diradical, enediyne **3** was heated with nucleophiles. With iodide, **3** reacts to form 9-iodo-1,2,3,4,5,6,7,8-octahydroanthracene **5I** in >99% isolated yield. It was found necessary to use a dilute solution of enediyne to reduce polymerization, and 5 mM was found optimal. Heating enediyne **3** without any nucleophile produced 1,2,3,4,5,6,7,8-octahydroanthracene **5H** in 40% yield, whereas no **5H** was formed in the presence of I⁻. Consequently an excess of nucleophile was used during the reaction to promote the capture of the diradical. Reaction of **3** with SCN⁻, Br⁻, NC⁻, or NO₂⁻ produced moderate to good yields of 9-X-1,2,3,4,5,6,7,8-octahydroanthracene **5X** (Table 1). Lower yields of **5X** compared to **5I** are due to the lower nucleophilicity of X⁻, leading to increased

conversion to **5H** and other unidentified compounds (Fig. S8).

The significant result in Fig. S10-11 is that the rate of disappearance of enediyne **3** is effectively independent of the concentration of nucleophile or acid and regardless of which nucleophile is present. Thus, the rate law can be written simply as a first-order equation, $-d[\mathbf{3}]/dt = k[\mathbf{3}]$. It follows that the rate-limiting step is the cycloaromatization of **3** to form the *p*-dehydrobenzene diradical, which then reacts rapidly with nucleophile to form product **5X**.

According to Table S2 the rates of disappearance of enediynes **1**, **2**, and **3** in the presence of LiI and pivalic acid decrease in the order **1** > **3** > **2**, and this trend remains the same even on varying $[I^-]$ or temperature. Fig. 1 shows a plot of the free energy of activation for cyclization of enediynes **1**, **2**, and **3** versus the calculated distances between carbon atoms C1 and C6, between which the new bond is formed (Tables 2 and 3). For **2** $\Delta G^\ddagger$ was taken as 30 kcal·mol⁻¹, from data (Table S1) in the presence of Cl⁻, which is least likely to add directly to the enediyne **2**. The least-squares slope is 23 kcal·mol⁻¹·Å⁻¹, and the correlation coefficient R^2 is 0.99. Therefore an 0.1-Å reduction in the distances between carbon atoms C1 and C6 is associated with a 2.3 kcal·mol⁻¹ reduction in activation energy.

Figure 1. Free energy of activation (kcal·mol⁻¹) for conversion of enediynes **1-3** to the corresponding *p*-dehydrobenzene diradicals versus C1-C6 distance (Å).

The behavior of enediyne **2** is more complicated. According to the data in Table S1, the rate of disappearance of **2** increases on doubling the concentration of LiX, but by less than twofold. More significantly, the rates of disappearance of enediyne **2** are lower with the less nucleophilic LiBr and even lower with LiCl. No such variations are seen with **1** or **3**. We attribute these deviations from zero- or first-order dependence on halide to a direct nucleophilic addition of halide to the C≡C of **2** as well as to its *p*-dehydrobenzene diradical. Moreover, **2** is slower to cyclize because cyclization produces a dehydronaphthalene diradical, which gains less aromatic stabilization than for **1** and **3**, which produce a dehydrobenzene from a nonaromatic precursor. Therefore the plot in Fig. 1 is not

necessarily linear, because the aromatic stabilization is not constant, and the point for **2** may be higher if **2** reacts directly with Cl⁻ or if capture of dehydronaphthalene diradical by I⁻ is competitive with its recyclization. Nevertheless, the plot is treated as linear for simplicity.

It is not surprising that the rate of cyclization correlates with the C(1)-C(6) distance. Nicolaou et al. reported that saturation of the double bond following intramolecular thiol addition to the α,β-unsaturated ketone in calicheamicin-γ₁ shortens the calculated distance between the alkyne carbons. Specifically, they reported that the shorter C1-C6 distance (3.16 Å) between alkyne carbons in the cyclic aglycone increases the chance of spontaneous cyclization, relative to acyclic aglycone (3.35 Å).^{4a} They also performed calculations on several other cyclic enediynes and proposed that a critical C1-C6 distance (*cd*) for room-temperature cyclization is 3.20-3.31 Å.^{4b} Later work suggested that this range may be extended to 2.9-3.4 Å.¹⁴ Furthermore, the synthesis of model cyclic enediynes with varying tendencies towards cycloaromatization due to varying C1-C6 distances provided evidence for this trend. Indeed, enediyne **3**, with *cd* = 3.32 Å, is stable at room temperature. It was further demonstrated that the critical factor for ease of cyclization is more accurately described as the difference between the strain energies of ground and transition states, in part because there is a repulsive contribution from the in-plane π orbitals.¹⁵

In summary, enediyne **3** undergoes cycloaromatization at a convenient 65-75 °C, slightly higher than the 45-55 °C for **1** but lower than the 85-95 °C for **2**, and these differences can be understood in terms of the structures of those enediynes.

CONCLUSIONS

We have demonstrated that cyclohexeno[3,4]cyclodec-1,5-diyne-3-ene (**3**) is a convenient model for the study of the reactivity of enediynes. It was synthesized in a relatively good yield and is stable for more than a month at room temperature. The first-order rate constant for its disappearance at 75 °C is $(5.4 \pm 0.1) \times 10^{-5} \text{ s}^{-1}$, independent of nucleophile concentration. Therefore, as with cyclodec-1,5-diyne-3-ene (**1**), the *p*-dehydrobenzene diradical derived from **3** reacts rapidly with nucleophiles. Comparison of activation energies for cyclization of enediynes **1**, **2**, and **3** shows that they parallel the critical C1-C6 distance across which the bond forms.

Enediyne **3** is a valuable addition to the suite of enediynes that probe not only their cycloaromatization but also the unusual nonradical reaction between nucleophiles and the *p*-dehydrobenzene diradical, as well as the reactivity and selectivity of the resulting aryl anion. These are reactions that can now be more

easily probed with this more readily available enediyne.

ASSOCIATED CONTENT

SUPPORTING INFORMATION

Details on materials, instrumentation, synthetic procedures and characterization, procedures for kinetics, free energy of activation, molecular mechanics, ^1H NMR analysis, GC-MS analysis, spectra, plots of $[3]_v/[3]_o$ and $\ln [3]_v/[3]_o$, first-order rate constants and free energies of activation for disappearance of enediynes **1-3**, and calculated distances, bond angles, and strain energies in their MM2 energy-minimized structures. Supporting Information File (SI.pdf)

AUTHOR INFORMATION

Corresponding Author

Charles L. Perrin – Department of Chemistry, University of California—San Diego, La Jolla, California 92093-0358, United States
Email: cperrin@ucsd.edu

Author(s)

Annadka Shrinidhi – Department of Chemistry, University of California—San Diego, La Jolla, California 92093-0358, United States

Author Contributions

A.S designed and performed all experiments. A.S and C.L.P wrote the manuscript. C.L.P supervised the work.

ACKNOWLEDGMENTS

This research was supported by grants CHE124B, CHEFFPE, CHE63CP, and CHEGDCP from UCSD to C.L.P. We sincerely thank Dr. Anthony Mrse of the Departmental NMR Facility and Dr. Yongxuan Su of the Departmental Molecular Mass Spectrometry Facility.

REFERENCES

- (1) (a) Golik, J.; Clardy, J.; Dubay, G.; Groenewold, G.; Kawaguchi, H.; Konishi, M.; Krishnan, B.; Ohkuma, H.; Saitoh, K.; Doyle, T. W. Esperamicins, a Novel Class of Potent Antitumor Antibiotics. 2. Structure of Esperamicin X. *J. Am. Chem. Soc.* **1987**, *109*, 3461-3462. (b) Nicolaou, K. C.; Skokotas, G.; Furuya, S.; Suemune, H.; Nicolaou, D. C. Golfomycin A, a Novel Designed Molecule with DNA-Cleaving Properties and Antitumor Activity. *Angew. Chem., Int. Ed.* **1990**, *29*, 1064-1067. (c) Ben, S.; Wen, L.; Koichi, N. Enediyne Natural Products: Biosynthesis and Prospect Towards Engineering Novel Antitumor Agents. *Curr. Med. Chem.* **2003**, *10*, 2317-2325. (d) Fenical, W.; Jensen, P. R. Developing a New Resource for Drug Discovery: Marine Actinomycete Bacteria. *Nat. Chem. Biol.* **2006**, *2*, 666-673. (e) Shen, B.; Hindra; Yan, X.; Huang, T.; Ge, H.; Yang, D.; Teng, Q.; Rudolf, J. D.; Lohman, J. R. Enediynes: Exploration of Microbial Genomics to Discover New Anticancer Drug Leads. *Bioorg. Med. Chem. Lett.* **2015**, *25*, 9-15. (f) Bhattacharya, P.; Basak, A.; Campbell, A.; Alabugin, I. V. Photochemical Activation of Enediyne Warheads: A Potential Tool for Targeted Antitumor Therapy. *Mol. Pharm.* **2018**, *15*, 768-797. (g) Cohen, D. R.; Townsend, C. A. A dual role for a polyketide synthase in dynemicin enediyne and anthraquinone biosynthesis. *Nature Chem.* **2018**, *10*, 231-236. (h) Yan, X.; Chen, J.-J.; Adhikari, A.; Teijaro, C. N.; Ge, H.; Crnovcic, I.; Chang, C.-Y.; Annaval, T.; Yang, D.; Rader, C.; Shen, B. Comparative Studies of the Biosynthetic Gene Clusters for Anthraquinone-Fused Enediynes Shedding Light into the Tailoring Steps of Tancimycin Biosynthesis. *Org. Lett.* **2018**, *20*, 5918-5921. (i) Nicolaou, K. C.; Li, R.; Chen, Q.; Lu, Z.; Pitsinos, E. N.; Schammel, A.; Lin, B.; Gu, C.; Sarvaiya, H.; Tchelepi, R.; Valdiosera, A.; Clubb, J.; Barbour, N.; Sisodiya, V.; Sandoval, J.; Lee, C.; Aujay, M.; Gavriluk, J. Synthesis and Biological Evaluation of Shishijimicin A-Type Linker-Drugs and Antibody-Drug Conjugates. *J. Am. Chem. Soc.* **2020**, *142*, 12890-12899. (j) Nicolaou, K. C.; Das, D.; Lu, Y.; Rout, S.; Pitsinos, E. N.; Lyssikatos, J.; Schammel, A.; Sandoval, J.; Hammond, M.; Aujay, M.; Gavriluk, J. Total Synthesis and Biological Evaluation of Tancimycins A and B, Yangpumicin A, and Related Anthraquinone-Fused Enediyne Antitumor Antibiotics. *J. Am. Chem. Soc.* **2020**, *142*, 2549-2561. (k) Nicolaou, K. C.; Rigol, S.; Pitsinos, E. N.; Das, D.; Lu, Y.; Rout, S.; Schammel, A. W.; Holte, D.; Lin, B.; Gu, C.; Sarvaiya, H.; Trinidad, J.; Barbour, N.; Valdiosera, A. M.; Sandoval, J.; Lee, C.; Aujay, M.; Fernando, H.; Dhar, A.; Karsunky, H.; Taylor, N.; Pysz, M.; Gavriluk, J. Uncialamycin-based antibody-drug conjugates: Unique enediyne ADCs exhibiting bystander killing effect. *Proc. Natl. Acad. Sci.* **2021**, *118*, e2107042118.
- (2) (a) Nicolaou, K. C.; Smith, A. L.; Yue, E. W. Chemistry and Biology of Natural and Designed Enediynes. *Proc. Natl. Acad. Sci.* **1993**, *90*, 5881. (b) Liang, Z.-X. Complexity and Simplicity in the Biosynthesis of Enediyne Natural Products. *Nat. Prod. Rep.* **2010**, *27*, 499-528. (c) Adhikari, A.; Shen, B.; Rader, C. Challenges and opportunities to develop enediyne natural products as payloads for antibody-drug conjugates. *Antibody Therapeutics* **2021**, *4*, 1-15.
- (3) Jones, R. R.; Bergman, R. G. *p*-Benzyne. Generation as an Intermediate in a Thermal Isomerization Reaction and Trapping Evidence for the 1,4-Benzenediyl Structure. *J. Am. Chem. Soc.* **1972**, *94*, 660-661.
- (4) (a) Nicolaou, K. C.; Ogawa, Y.; Zuccarello, G.; Schweiger, E. J.; Kumazawa, T. Cyclic Conjugated Enediynes Related to Calicheamicins and Esperamicins: Calculations, Synthesis, and Properties. *J. Am. Chem. Soc.* **1988**, *110*, 4866-4868. (b) Nicolaou, K. C.; Zuccarello, G.; Riemer, C.; Estevez, V. A.; Dai, W. M. Design, Synthesis, and Study of Simple Monocyclic Conjugated Enediynes. The 10-Membered Ring Enediyne Moiety of the Enediyne Anticancer Antibiotics. *J. Am. Chem. Soc.* **1992**, *114*, 7360-7371.
- (5) (a) Nicolaou, K. C.; Dai, W. M.; Hong, Y. P.; Baldrige, K. K.; Siegel, J. S.; Tsay, S. C. Molecular Design, Chemical Synthesis, Kinetic Studies, Calculations, and Biological Studies of Novel Enediynes Equipped with Triggering, Detection, and Deactivating Devices. Model Dynemicin A Epoxide and cis-Diol Systems. *J. Am. Chem. Soc.* **1993**, *115*, 7944-7953. (b) Jones, G. B.; Plourde, G. W. Electronic Control of the Bergman Cycloaromatization: Synthesis and Chemistry of Chloroenediynes. *Org. Lett.* **2000**, *2*, 1757-1759. (c) Zeidan, T. A.; Manoharan, M.; Alabugin, I. V. Ortho Effect in the Bergman Cyclization: Interception of *p*-Benzyne Intermediate by Intramolecular Hydrogen Abstraction. *J. Org. Chem.* **2006**, *71*, 954-961. (d) Zeidan, T. A.; Kovalenko, S. V.; Manoharan, M.; Alabugin, I. V. Ortho Effect in the Bergman Cyclization: Comparison of Experimental Approaches and Dissection of Cycloaromatization Kinetics. *J. Org. Chem.* **2006**, *71*, 954-961. (e) Mohamed, R. K.; Paul W. Peterson, P. W.; Alabugin, I. V.

Concerted Reactions That Produce Diradicals and Zwitterions: Electronic, Steric, Conformational, and Kinetic Control of Cycloaromatization Processes. *Chem. Rev.* **2013**, *113*, 7089-7129. (f) Peterson, P. W.; Shevchenko, N.; Breiner, B.; Manoharan, M.; Lufti, F.; Delaune, J.; Kingsley, M.; Kovnir, K.; Alabugin, I. V. Orbital Crossings Activated through Electron Injection: Opening Communication between Orthogonal Orbitals in Anionic C1–C5 Cyclizations of Eneidyne. *J. Am. Chem. Soc.* **2016**, *138*, 15617-15628. (g) Hu, H.; Zhang, B.; Luxon, A.; Scott, T.; Wang, B.; Parish, C. A. An Extended Multireference Study of the Singlet and Triplet States of the 9,10-didehydroanthracene Diradical. *J. Phys. Chem. A* **2018**, *122*, 3688-3696.

(6) (a) König, B.; Hollnagel, H.; Ahrens, B.; Jones, P. G. Activation of Macrocyclic Biaryl-Eneidyne by Metal Ion Coordination. *Angew. Chem., Int. Ed.* **1995**, *34*, 2538-2540. (b) Warner, B. P.; Millar, S. P.; Broene, R. D.; Buchwald, S. L. Controlled Acceleration and Inhibition of Bergman Cyclization by Metal Chlorides. *Science* **1995**, *269*, 814. (c) Coalter, N. L.; Concolino, T. E.; Streib, W. E.; Hughes, C. G.; Rheingold, A. L.; Zaleski, J. M. Structure and Thermal Reactivity of a Novel Pd(0) Metalloenediynes. *J. Am. Chem. Soc.* **2000**, *122*, 3112-3117. (d) O'Connor, J. M.; Lee, L. I.; Gantzel, P.; Rheingold, A. L.; Lam, K.-C. Inhibition and Acceleration of the Bergman Cycloaromatization Reaction by the Pentamethylcyclopentadienyl Ruthenium Cation. *J. Am. Chem. Soc.* **2000**, *122*, 12057-12058. (e) O'Connor, J. M.; Friese, S. J.; Tichenor, M. Ruthenium-Mediated Cycloaromatization of Acyclic Eneidyne and Dienes at Ambient Temperature. *J. Am. Chem. Soc.* **2002**, *124*, 3506-3507. (f) Basak, A.; Mandal, S.; Bag, S. S. Chelation-Controlled Bergman Cyclization: Synthesis and Reactivity of Eneidyne Ligands. *Chem. Rev.* **2003**, *103*, 4077-4094. (g) O'Connor, J. M.; Friese, S. J.; Rodgers, B. L. A Transition-Metal-Catalyzed Eneidyne Cycloaromatization. *J. Am. Chem. Soc.* **2005**, *127*, 16342-16343. (h) Byers, P. M.; Rashid, J. I.; Mohamed, R. K.; Alabugin, I. V. Polyaromatic Ribbon/Benzofuran Fusion via Consecutive Endo Cyclizations of Eneidyne. *Org. Lett.* **2012**, *14*, 6032-6035. (i) Gomes, G. d. P.; Alabugin, I. V. Drawing Catalytic Power from Charge Separation: Stereoelectronic and Zwitterionic Assistance in the Au(I)-Catalyzed Bergman Cyclization. *J. Am. Chem. Soc.* **2017**, *139*, 3406-3416.

(7) Nicolaou, K. C.; Smith, A. L. Molecular Design, Chemical Synthesis, and Biological Action of Eneidyne. *Acc. Chem. Res.* **1992**, *25*, 497-503.

(8) (a) Semmelhack, M. F.; Neu, T.; Foubelo, F. Arene 1,4-Diradical Formation from o-Dialkynylarenes. *Tetrahedron Lett.* **1992**, *33*, 3277-3280. (b) Perrin, C. L.; Rodgers, B. L.; O'Connor, J. M. Nucleophilic Addition to a p-Benzyne Derived from an Eneidyne: A New Mechanism for Halide Incorporation into Biomolecules. *J. Am. Chem. Soc.* **2007**, *129*, 4795-4799. (c) Perrin, C. L.; Reyes-Rodríguez, G. J. Reactivity of Nucleophiles Toward a p-Benzyne Derived from an Eneidyne. *J. Phys. Org. Chem.* **2012**, *26*, 206-210. (d) Perrin, C. L.; Reyes-Rodríguez, G. J. Selectivity and Isotope Effects in Hydronation of a Naked Aryl Anion. *J. Am. Chem. Soc.* **2014**, *136*, 15263-15269.

(9) Jones, G. B.; Wright, J. M.; Plourde, G. W.; Hynd, G.; Huber, R. S.; Mathews, J. E. A Direct and Stereocontrolled Route to Conjugated Eneidyne. *J. Am. Chem. Soc.* **2000**, *122*, 1937-1944.

(10) Reyes-Rodríguez, G. J. Nucleophilic Additions to a para-Benzyne Derived from an Eneidyne: Exploring

the Non-Radical Reactivity of a Diradical. PhD Thesis, University of California: San Diego, 2013.

(11) (a) Das, E.; Basak, S.; Anoop, A.; Basak, A. Experiment and Computational Study on the Regioselectivity of Nucleophilic Addition to Unsymmetrical p-Benzynes Derived from Bergman Cyclization of Eneidyne. *J. Org. Chem.* **2018**, *83*, 7730-7740. (b) Das, E.; Basak, S.; Anoop, A.; Chand, S.; Basak, A. How To Achieve High Regioselectivity in Barrier-less Nucleophilic Addition to p-Benzynes Generated via Bergman Cyclization of Unsymmetrical Cyclic Azaenediynes? *J. Org. Chem.* **2019**, *84*, 2911-2921. (c) Das, E.; Basak, A. Regioselective Synthesis of Benzo-Fused Tetrahydroisoquinoline-Based Biaryls through a Tandem One-Pot Halogenation of p-Benzynes from Eneidyne and Suzuki-Miyaura Coupling. *J. Org. Chem.* **2020**, *85*, 2697-2703. (d) Singha, M.; Bhattacharya, P.; Ray, D.; Basak, A. Sterically Hindering the Trajectory of Nucleophilic Attack Towards p-Benzynes by a Properly Oriented Hydrogen Atom: an Approach to Achieve Regioselectivity. *Org. Biomol. Chem.* **2021**, *19*, 5148-5154.

(12) Semmelhack, M. F.; Neu, T.; Foubelo, F. Arene 1,4-Diradical Formation from o-Dialkynylarenes. *J. Org. Chem.* **1994**, *59*, 5038-5047.

(13) Lockhart, T. P.; Comita, P. B.; Bergman, R. G. Kinetic Evidence for the Formation of Discrete 1,4-Dehydrobenzene Intermediates. Trapping by Inter- and Intra-molecular Hydrogen Atom Transfer and Observation of High-Temperature CIDNP. *J. Am. Chem. Soc.* **1981**, *103*, 4082-4090.

(14) (a) Schreiner, P. R. Cyclic Eneidyne: Relationship between Ring Size, Alkyne Carbon Distance, and Cyclization Barrier. *Chem. Commun.* **1998**, 483-484. (b) Schreiner, P. R. Monocyclic Eneidyne: Relationships between Ring Sizes, Alkyne Carbon Distances, Cyclization Barriers, and Hydrogen Abstraction Reactions. Singlet–Triplet Separations of Methyl-Substituted p-Benzynes. *J. Am. Chem. Soc.* **1998**, *120*, 4184-4190.

(15) (a) Snyder, J. P. The Cyclization of Calicheamicin-Esperamicin Analogs: a Predictive Biradicaloid Transition State. *J. Am. Chem. Soc.* **1989**, *111*, 7630-7632. (b) Magnus, P.; Fortt, S.; Pittner, T.; Snyder, J. P. Synthetic and Mechanistic Studies on Esperamicin A1 and Calicheamicin γ . Molecular Strain rather than π -Bond Proximity Determines the Cycloaromatization Rates of Bicyclo[7.3.1]eneidyne. *J. Am. Chem. Soc.* **1990**, *112*, 4986-4987. (c) Snyder, J. P. Monocyclic Eneidyne Collapse to 1,4-diyl Biradicals: a Pathway Under Strain Control. *J. Am. Chem. Soc.* **1990**, *112*, 5367-5369. (d) Snyder, J. P.; Tipson, G. E. Proposal for Blending Classical and Biradical Mechanisms in Antitumor Antibiotics: Dynemicin A. *J. Am. Chem. Soc.* **1990**, *112*, 4040-4042. (e) Magnus, P.; Carter, P.; Elliott, J.; Lewis, R.; Harling, J.; Pittner, T.; Bauta, W. E.; Fortt, S. Synthetic and Mechanistic Studies on the Antitumor Antibiotics Esperamicin A1 and Calicheamicin γ 1: Synthesis of 2-Ketobicyclo[7.3.1]eneidyne and 13-Ketocyclo[7.3.1]eneidyne Cores Mediated by η^2 -Dicobalt Hexacarbonyl Alkyne Complexes. Cycloaromatization Rate Studies. *J. Am. Chem. Soc.* **1992**, *114*, 2544-2559. (f) Alabugin, I. V.; Manoharan, M. Reactant Destabilization in the Bergman Cyclization and Rational Design of Light- and pH-Activated Eneidyne. *J. Phys. Chem. A* **2003**, *107*, 3363-3371.