

UC Irvine

Western Journal of Emergency Medicine: Integrating Emergency Care with Population Health

Title

Emergency Nurses' Perceptions of Opioid Use Disorder and Its Treatment in the Emergency Department

Permalink

<https://escholarship.org/uc/item/9230q4fn>

Journal

Western Journal of Emergency Medicine: Integrating Emergency Care with Population Health, 23(5.1)

ISSN

1936-900X

Authors

Prucnal, Christiana
Williamson, Dawn
Samuels, Elizabeth
et al.

Publication Date

2022

DOI

10.5811/westjem.58866

Copyright Information

Copyright 2022 by the author(s). This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

asked how successful their efforts were to diversify their staff, 3 respondents reported that their efforts were very successful and 20 reported either partially or not very successful. There was an association between the type of location (suburban vs urban) and the number of physicians of color hired when looking at programs that hired 3 or more physicians of color.

3 Emergency Nurses' Perceptions of Opioid Use Disorder and Its Treatment in the Emergency Department

Christiana K. Prucnal; Dawn Williamson; Elizabeth A. Samuels; Kristina Monteiro; Margaret Samuels-Kalow

Objectives: To describe the knowledge and attitudes of emergency nurses regarding caring for patients with opioid use disorder in the emergency department.

Background: Many eligible patients with opioid use disorder do not receive available emergency department services for treatment and harm mitigation. While prior study examined contributing provider factors, little is known of nursing factors. This study describes knowledge and attitudes of emergency nurses regarding patients with opioid use disorder and their evidence-based treatment services in the emergency department setting.

Methods: Anonymous email surveys with novel and previously validated questions based on The Theory of Planned Behavior Framework were distributed to emergency department nurses at a large, urban tertiary-care hospital. Chi-Square and independent samples t-tests were used in analyses.

Results: More than one third of nurses completed the questionnaire (39%, 85/218). Most showed willingness and confidence screening for substance use disorder (95% and 88% respectively). Higher confidence providing buprenorphine and take-home naloxone was significantly associated with having worked fewer years (8.33 v. 15.62, $p=0.01$ and 7.38 v. 12.03, $p=0.03$ respectively). Confidence administering buprenorphine was significantly associated with receiving in-service training ($p=0.03$). Staff with knowledge of take-home naloxone, positive attitudes toward syringe service programs, and a belief in a biopsychosocial basis of addiction were significantly younger and had worked significantly fewer years than those not indicating these beliefs. Specific educational gaps were identified.

Conclusion: Emergency nurses display willingness to champion evidence-based care for patients with opioid use disorder. Younger age and having worked fewer years were significantly associated with positive attitudes towards recovery science, harm mitigation, and services knowledge. Having worked fewer years was significantly associated with greater confidence performing treatment and harm mitigation. In-service training was significantly associated with greater confidence administering buprenorphine. Further study

should support generalizability and determine which staff development measures generate improved outcomes.

4 Effect of the COVID-19 Pandemic on ED Adult Psychiatric Visits

Barnet Eskin; Crystal Bauman; John R. Allegra

Objectives: Our goal was to determine whether the proportion of ED visits for specific psychiatric conditions, namely anxiety disorders, depression, self-harm/suicidal thoughts, bipolar disorder, and psychotic disorders, changed after the arrival of COVID-19.

Background: In March 2020, the COVID-19 pandemic reached the New York tri-state area, which, at the time, was one of the regions in the United States (US) that the virus most severely affected. ED visits dramatically declined, likely due to social isolation mandates and fear of exposure to the virus. Quarantining at home, fear of becoming sick, and job disruptions caused the level of stress in the population to increase. In a previous US study, the proportion of ED visits for some psychiatric conditions increased.

Methods: Design: Retrospective cohort. Setting: EDs of 28 hospitals within 150 miles of New York City. Hospitals were teaching and non-teaching in rural, suburban and urban areas. Total annual ED volumes were 12,000 to 122,000. Population: Consecutive ED patients = 21 years old from March 1 to November 30 in 2019 and 2020. Data analysis: We tallied the number of patients in 2019 and 2020 with anxiety disorders, depression, self-harm/suicidal thoughts, bipolar disorder, and psychotic disorders, identified using International Classification of Disease codes (version 10). We calculated the proportion of these visits to total ED visits in 2019 and 2020. We report the changes in these proportions from 2019 to 2020, along with 95% CIs.

Results: Total ED visits decreased 27%, from 844,017 in 2019 to 618,195 in 2020. In 2019 and 2020 combined, the number of patients were: 13,151 with anxiety disorders, 6884 with depression, 8886 with suicidal ideation/self-harm, 3252 with bipolar disorder, and 7129 with psychotic disorders. The changes [with 95% CIs] in the proportion of visits from 2019 to 2020 were: anxiety disorders -1% [-4,+3%], depression -5% [-10,-1%], self-harm/suicidal thoughts +23% [+18,+29%], bipolar disorder +14% [+6, +22%], and psychotic disorders +23% [+18,+29%].

Conclusion: The proportion of adult ED visits for self-harm/suicidal thoughts, bipolar disorder, and psychotic disorders increased following the arrival of COVID-19, whereas the proportions for anxiety and depression changed minimally. These results are somewhat different from the findings in the previously reported study. Our study highlights the need for continued surveillance of the impact of