

UC Davis

UC Davis Previously Published Works

Title

Isavuconazole in the Treatment of Coccidioidal Meningitis.

Permalink

<https://escholarship.org/uc/item/92q3n6tw>

Journal

Antimicrobial Agents and Chemotherapy, 63(3)

ISSN

0066-4804

Authors

Heidari, Arash
Quinlan, Miriam
Benjamin, David J
et al.

Publication Date

2019-03-01

DOI

10.1128/aac.02232-18

Peer reviewed



Isavuconazole in the Treatment of Coccidioidal Meningitis

Arash Heidari,^{a,b} Miriam Quinlan,^a David J. Benjamin,^c Brett Laurence,^d Anandit Mu,^e Tiffany Ngai,^e Wes J. Hoffman,^f Stuart H. Cohen,^g  Ian McHardy,^h Royce Johnson,^{a,b} George R. Thompson III^{g,h}

^aKern County Medical Center, Bakersfield, California, USA

^bValley Fever Institute, Bakersfield, California, USA

^cDepartment of Internal Medicine, University of California Davis Medical Center, Sacramento, California, USA

^dSutter Medical Center, Sacramento, California, USA

^eDepartment of Internal Medicine, Division of Infectious Diseases, University of California San Francisco-Fresno, Fresno, California, USA

^fDepartment of Pharmacy, University of California Davis Medical Center, Sacramento, California, USA

^gDepartment of Internal Medicine, Division of Infectious Diseases, University of California Davis Medical Center, Sacramento, California, USA

^hDepartment of Medical Microbiology and Immunology, University of California Davis, Davis, California, USA

ABSTRACT Patients with coccidioidal meningitis require lifelong antifungal therapy. Cumulative toxicity and lack of antifungal efficacy require salvage therapy in the treatment of some patients. In a retrospective review of nine patients with coccidioidal meningitis treated with isavuconazole, successful therapy was seen in three patients and stable disease was confirmed in six patients. Isavuconazole may be a useful addition to the therapeutic choices currently available for coccidioidal meningitis.

KEYWORDS coccidioides, isavuconazole, meningitis

C*occidioides immitis* and *posadasii* are dimorphic fungi endemic to the southwestern United States (1). The spectrum of disease ranges from asymptomatic acquisition with resultant immunity to severe and life-threatening disseminated infection. Disseminated infection is uncommon, occurring in only 1% to 3% of infected individuals, and is more frequent in immunocompromised individuals, women during the third trimester of pregnancy or the immediate postpartum period, or individuals of Filipino or African ancestry. Dissemination may occur to any site, although the skin, musculoskeletal system, and central nervous system (CNS) are the most frequently encountered sites (2).

CNS involvement that causes meningitis is the most serious complication and carries significant morbidity and mortality (3). Current guidelines recommend high-dose fluconazole or itraconazole in the treatment of coccidioidal meningitis (1). Limited data are available describing the efficacy of other triazoles, including voriconazole and posaconazole (4, 5). Despite the availability of these agents and the utility of polyenes or combination therapy in select cases requiring salvage therapy (6, 7), toxicity and/or lack of efficacy requires investigation of alternative agents.

Isavuconazole is a novel triazole with a broad spectrum of antifungal activity. Administered as a water-soluble prodrug, i.e., isavuconazonium sulfate, it undergoes cleavage by plasma esterases to the active moiety isavuconazole (8). Clinical trials demonstrated efficacy in the treatment of invasive aspergillosis (9), candidiasis (10), and mucormycosis (11) and in the treatment of endemic mycoses without evidence of CNS involvement (12). However, efficacy in the treatment of coccidioidal meningitis has not previously been described.

Patients with coccidioidal meningitis who received isavuconazole were identified by review of the ICD-9 and ICD-10 codes cross-referenced with pharmacy databases. Abstracted data included demographic and clinical information; results of laboratory and radiographic studies; microbiologic, pathologic, and serologic results; outcomes;

Citation Heidari A, Quinlan M, Benjamin DJ, Laurence B, Mu A, Ngai T, Hoffman WJ, Cohen SH, McHardy I, Johnson R, Thompson GR, III. 2019. Isavuconazole in the treatment of coccidioidal meningitis. *Antimicrob Agents Chemother* 63:e02232-18. <https://doi.org/10.1128/AAC.02232-18>.

Copyright © 2019 American Society for Microbiology. All Rights Reserved.

Address correspondence to George R. Thompson III, grthompson@ucdavis.edu.

Received 22 October 2018

Accepted 10 December 2018

Accepted manuscript posted online 17 December 2018

Published 26 February 2019

TABLE 1 Clinical characteristics, treatment, and outcomes of patients with coccidioidal meningitis treated with isavuconazole

Patient by age (yr), sex, ethnicity, underlying condition	Prior antifungal ^a (days)	Reason for ISAV ^b treatment	Response to ISAV and change in MSG ^c score at 3 mo	Clinician assessment at 3 mo	Outcome, duration of ISAV therapy (days)
46, M, African American, none	FLU (734)→POS (2,920)→VOR (607)	FLU, failure; POS, nausea/vomiting; VOR, photopsia, headache confusion	Success (7→3)	Success	Alive, no relapses (621)
49, F, Caucasian, none	FLU (2,079)→VOR (1,455)	VOR, failure	Success (2→0)	Success	Alive, no relapses (441)
44, M, Hispanic, none	FLU (549)	FLU, failure	Success (7→5)	Success	Alive, no relapses (518)
33, M, Hispanic, none	FLU (3,205)→VOR (1,694)	VOR, photodermatitis	Stable (1→1)	Success	Alive, no relapses (637)
55, M, Indian, none	FLU (811)→VOR (148)	VOR, dyspepsia	Stable (8→7)	Success	Alive, no relapses ^d (138)
51, M, Hispanic, none	FLU (560)→VOR (1,560)	VOR, xerosis, photodermatitis	Stable (4→4)	Stable	Alive, no relapses (708)
41, M, Hispanic, none	FLU (915)→VOR (3,071)	VOR, photodermatitis	Stable (0→0)	Stable	Alive, no relapses (274)
43, M, Hispanic, none	FLU (361)→VOR (3,319)	VOR, squamous cell carcinoma	Stable (0→0)	Stable	Alive, no relapses (385)
59, M, Caucasian, none	FLU (1,252)→VOR (410)→FLU (941)	FLU, failure; VOR, hepatic injury; FLU, nausea/vomiting	Stable (10→8)	Stable	Alive, no relapses (810)

^aFLU, fluconazole; VOR, voriconazole.

^bISAV, isavuconazole.

^cMSG, Mycoses Study Group scoring system (13).

^dISAV discontinued because of worsening dyspepsia and transitioned back to voriconazole.

and prior treatment regimens. Responses to therapy were measured with a previously validated scoring system used in clinical trials of coccidioidal meningitis (13). This scoring system combines clinical, radiographic, and cerebrospinal fluid (CSF) data to evaluate a therapeutic response, defined as a $\geq 40\%$ reduction in abnormalities. Each patient was assigned a composite score at the time of isavuconazole initiation and after 3 months of therapy. The treating clinician's assessment was also abstracted from patient medical records, and defined as success, stable, or failure.

Nine patients met criteria for inclusion (Table 1). The median patient age was 46 years, and eight of the patients were male. No comorbidities were present in this patient cohort. All patients were initiated on fluconazole as primary therapy for coccidioidal meningitis, and eight were transitioned to voriconazole, although the reasons for this change were not always identifiable on chart review. Treatment failure was found in only one patient treated with voriconazole. All other patients were transitioned from voriconazole to isavuconazole secondary to adverse effects, including photodermatitis ($n = 3$), photopsia ($n = 1$), squamous cell carcinoma ($n = 1$), dyspepsia ($n = 1$), and hepatic injury ($n = 1$).

Patient outcomes assessed with Mycosis Study Group (MSG) scoring criteria showed successful therapy in three of nine patients and stable disease in six patients. Mean time on isavuconazole therapy at last assessment was 504 days (range, 274 to 810 days) per patient. The abstracting treating clinician assessment revealed success in five of nine patients and stable disease in four patients. Discrepancies between the assessment by the treating clinician and MSG scores were seen in two patients. These differences in patient assessments were secondary to MSG criteria requiring a decrease in CSF protein to < 100 mg/dL for a change in point assignment, whereas patient 4 had a decline in CSF protein from 599 to only 469 mg/dL. The second discrepant patient had a significant improvement in CSF glucose level (from 22 to 45 mg/dL) with only a moderate change in CSF white blood cell count (from 470 to 450 cells/mm³; protein values, from 2,325 to 2,006 mg/dL). No isavuconazole clinical failures were identified in this retrospective evaluation, although one patient was transitioned back to voriconazole because of worsening dyspepsia after a change to isavuconazole.

Morbidity and mortality associated with coccidioidal meningitis remain unacceptably high, and, in an attempt to prevent relapse(s), patients are maintained on lifelong antifungal therapy (14). The commitment to lifelong therapy subjects patients to cumulative toxicity, ranging from benign but irritating (e.g., xerosis, cheilitis, alopecia) (15) to more serious (e.g., fluorosis, cutaneous malignancy) (16–18) manifestations. These adverse reactions may require the use of alternative agents in attempts to improve patient compliance and tolerability.

Furthermore, despite medication compliance, relapse/breakthrough may occur (19, 20). The reasons for relapse remain unclear, and a data-driven approach to management decisions for these patients has not been presented. In cases of breakthrough fungal infection, a change in class or a change of agent is typically recommended, pending further investigation (e.g., assessment of compliance, determination of possible resistance) (21).

Cumulative toxicity and/or drug efficacy failure have necessitated the evaluation of newer antifungals. Isavuconazole is an attractive new triazole for coccidioidomycosis because of the availability of both oral and intravenous formulations, a favorable side-effect profile when directly compared with voriconazole in the treatment of invasive aspergillosis (9), and low isavuconazole MICs when evaluated against *Coccidioides* isolates (8, 12, 22). However, few patients with coccidioidomycosis have been treated with isavuconazole, and data have been limited to primary pulmonary infections, none with dissemination or severe manifestations of disease (12).

Despite the lack of demonstrated clinical efficacy in cases of disseminated or meningeal coccidioidomycosis, animal models of other mycoses have demonstrated the potential efficacy of isavuconazole in CNS infections (23), and early experience in human cryptococcal meningitis cases has been favorable (12). The current study establishes precedence for use in coccidioidomycosis meningitis cases with a history of treatment failure or adverse reactions to front-line antifungals.

We observed successful or stable disease in all patients included in this study, although the retrospective nature, limited duration of follow-up, and number of patients studied limit implications for wide-scale adoption. Evaluation of isavuconazole in the treatment of coccidioidomycosis would benefit from use of animal models and prospective clinical trials. The response rates observed were favorable, and isavuconazole may be a useful agent in the salvage setting for this morbid disease.

REFERENCES

- Galgiani JN, Ampel NM, Blair JE, Catanzaro A, Geertsma F, Hoover SE, Johnson RH, Kusne S, Lisse J, MacDonald JD, Meyerson SL, Raksin PB, Siever J, Stevens DA, Sunenshine R, Theodore N. 2016. 2016 Infectious Diseases Society of America (IDSA) Clinical Practice Guideline for the Treatment of Coccidioidomycosis. *Clin Infect Dis* 63:e112–e146. <https://doi.org/10.1093/cid/ciw360>.
- Stockamp NW, Thompson GR, III. 2016. Coccidioidomycosis. *Infect Dis Clin North Am* 30:229–246. <https://doi.org/10.1016/j.idc.2015.10.008>.
- Thompson G, III, Wang S, Bercovitch R, Bolaris M, Van Den Akker D, Taylor S, Lopez R, Catanzaro A, Cadena J, Chin-Hong P, Spellberg B. 2013. Routine CSF analysis in coccidioidomycosis is not required. *PLoS One* 8:e64249. <https://doi.org/10.1371/journal.pone.0064249>.
- Kim MM, Vikram HR, Kusne S, Seville MT, Blair JE. 2011. Treatment of refractory coccidioidomycosis with voriconazole or posaconazole. *Clin Infect Dis* 53:1060–1066. <https://doi.org/10.1093/cid/cir642>.
- Freifeld A, Proia L, Andes D, Baddour LM, Blair J, Spellberg B, Arnold S, Lentnek A, Wheat LJ. 2009. Voriconazole use for endemic fungal infections. *Antimicrob Agents Chemother* 53:1648–1651. <https://doi.org/10.1128/AAC.01148-07>.
- Stewart ER, Eldridge ML, McHardy I, Cohen SH, Thompson GR, III. 2018. Liposomal amphotericin B as monotherapy in relapsed coccidioidal meningitis. *Mycopathologia* 183:619–622. <https://doi.org/10.1007/s11046-017-0240-7>.
- Mathisen G, Shelub A, Truong J, Wigen C. 2010. Coccidioidal meningitis: clinical presentation and management in the fluconazole era. *Medicine (Baltimore)* 89:251–284. <https://doi.org/10.1097/MD.0b013e3181f378a8>.
- Thompson GR, III, Wiederhold NP. 2010. Isavuconazole: a comprehensive review of spectrum of activity of a new triazole. *Mycopathologia* 170: 291–313. <https://doi.org/10.1007/s11046-010-9324-3>.
- Maertens JA, Raad II, Marr KA, Patterson TF, Kontoyiannis DP, Cornely OA, Bow EJ, Rahav G, Neofytos D, Aoun M, Baddley JW, Giladi M, Heinz WJ, Herbrecht R, Hope W, Karthaus M, Lee D-G, Lortholary O, Morrison VA, Oren I, Selleslag D, Shoham S, Thompson GR, III, Lee M, Maher RM, Schmitt-Hoffmann A-H, Zeiher B, Ullmann AJ. 2016. Isavuconazole versus voriconazole for primary treatment of invasive mould disease caused by *Aspergillus* and other filamentous fungi (SECURE): a phase 3, randomised-controlled, non-inferiority trial. *Lancet* 387:760–769. [https://doi.org/10.1016/S0140-6736\(15\)01159-9](https://doi.org/10.1016/S0140-6736(15)01159-9).
- Kullberg BJ, Viscoli C, Pappas PG, Vazquez J, Ostrosky-Zeichner L, Rotstein C, Sobel JD, Herbrecht R, Rahav G, Jaruratanasirikul S, Chetchotisakd P, Van Wijngaerden E, De Waele J, Lademacher C, Engelhardt M, Kovanda LL, Croos-Dabrera R, Fredericks C, Thompson GR, III. 2018. Isavuconazole versus caspofungin in the treatment of candidemia and other invasive candida infections: the ACTIVE trial. *Clin Infect Dis* <https://doi.org/10.1093/cid/ciy827>.
- Marty FM, Ostrosky-Zeichner L, Cornely OA, Mullane KM, Perfect JR, Thompson GR, III, Alangaden GJ, Brown JM, Fredricks DN, Heinz WJ, Herbrecht R, Klimko N, Klyasova G, Maertens JA, Melinkeri SR, Oren I, Pappas PG, Racil Z, Rahav G, Santos R, Schwartz S, Vehreschild JJ, Young JH, Chetchotisakd P, Jaruratanasirikul S, Kanj SS, Engelhardt M, Kaufhold A, Ito M, Lee M, Sasse C, Maher RM, Zeiher B, Vehreschild M, VITAL and FungiScope Mucormycosis Investigators. 2016. Isavuconazole treatment for mucormycosis: a single-arm open-label trial and case-control analysis. *Lancet Infect Dis* 16:828–837. [https://doi.org/10.1016/S1473-3099\(16\)00071-2](https://doi.org/10.1016/S1473-3099(16)00071-2).
- Thompson GR, III, Rendon A, Ribeiro dos Santos R, Queiroz-Telles F, Ostrosky-Zeichner L, Azie N, Maher R, Lee M, Kovanda L, Engelhardt M, Vazquez JA, Cornely OA, Perfect JR. 2016. Isavuconazole treatment of cryptococcosis and dimorphic mycoses. *Clin Infect Dis* 63:356–362. <https://doi.org/10.1093/cid/ciw305>.
- Galgiani JN, Catanzaro A, Cloud GA, Higgs J, Friedman BA, Larsen RA, Graybill JR. 1993. Fluconazole therapy for coccidioidal meningitis: the NIAID-Mycoses Study Group. *Ann Intern Med* 119:28–35. <https://doi.org/10.7326/0003-4819-119-1-199307010-00005>.
- Dewsnup DH, Galgiani JN, Graybill JR, Diaz M, Rendon A, Cloud GA, Stevens DA. 1996. Is it ever safe to stop azole therapy for *Coccidioides immitis* meningitis? *Ann Intern Med* 124:305–310. <https://doi.org/10.7326/0003-4819-124-3-199602010-00004>.
- Pappas PG, Kauffman CA, Perfect J, Johnson PC, McKinsey DS, Bam-

- berger DM, Hamill R, Sharkey PK, Chapman SW, Sobel JD. 1995. Alopecia associated with fluconazole therapy. *Ann Intern Med* 123:354–357. <https://doi.org/10.7326/0003-4819-123-5-199509010-00006>.
16. Pfizer. 2018. Voriconazole package insert. Pfizer, New York, NY. http://www.pfizer.com/files/products/uspi_vfend.pdf.
 17. Thompson GR, III, Bays D, Cohen SH, Pappagianis D. 2012. Fluoride excess in coccidioidomycosis patients receiving long-term antifungal therapy: an assessment of currently available triazoles. *Antimicrob Agents Chemother* 56:563–564. <https://doi.org/10.1128/AAC.05275-11>.
 18. Epaulard O, Villier C, Ravaud P, Chosidow O, Blanche S, Mamzer-Bruneel MF, Thiebaut A, Leccia MT, Lortholary O. 2013. A multistep voriconazole-related phototoxic pathway may lead to skin carcinoma: results from a French nationwide study. *Clin Infect Dis* 57:e182. <https://doi.org/10.1093/cid/cit600>.
 19. Ho J, Fowler P, Heidari A, Johnson RH. 2017. Intrathecal amphotericin B: a 60-year experience in treating coccidioid meningitis. *Clin Infect Dis* 64:519–524. <https://doi.org/10.1093/cid/ciw794>.
 20. McHardy IH, Dinh BN, Waldman S, Stewart E, Bays D, Pappagianis D, Thompson GR, III. 2018. Coccidioidomycosis complement fixation titer trends in the age of antifungals. *J Clin Microbiol* 56:e01318-18. <https://doi.org/10.1128/jcm.01318-18>.
 21. Maschmeyer G, Patterson TF. 2014. Our 2014 approach to breakthrough invasive fungal infections. *Mycoses* 57:645–651. <https://doi.org/10.1111/myc.12213>.
 22. Gonzalez GM. 2009. In vitro activities of isavuconazole against opportunistic filamentous and dimorphic fungi. *Med Mycol* 47:71–76. <https://doi.org/10.1080/13693780802562969>.
 23. Wiederhold NP, Kovanda L, Najvar LK, Bocanegra R, Olivo M, Kirkpatrick WR, Patterson TF. 2016. Isavuconazole is effective for the treatment of experimental cryptococcal meningitis. *Antimicrob Agents Chemother* 60:5600–5603. <https://doi.org/10.1128/AAC.00229-16>.