

UC Berkeley

UC Berkeley Previously Published Works

Title

Protein Quality Control and Lipid Droplet Metabolism

Permalink

<https://escholarship.org/uc/item/92x2d45n>

Journal

Annual Review of Cell and Developmental Biology, 36(1)

ISSN

1081-0706

Authors

Roberts, Melissa A

Olzmann, James A

Publication Date

2020-10-06

DOI

10.1146/annurev-cellbio-031320-101827

Peer reviewed



Published in final edited form as:

Annu Rev Cell Dev Biol. 2020 October 06; 36: 115–139. doi:10.1146/annurev-cellbio-031320-101827.

Protein Quality Control and Lipid Droplet Metabolism

Melissa A. Roberts¹, James A. Olzmann^{1,2}

¹Department of Molecular and Cell Biology and Department of Nutritional Sciences and Toxicology, University of California, Berkeley, California 94720, USA

²Chan Zuckerberg Biohub, San Francisco, California 94158, USA

Abstract

Lipid droplets (LDs) are endoplasmic reticulum–derived organelles that consist of a core of neutral lipids encircled by a phospholipid monolayer decorated with proteins. As hubs of cellular lipid and energy metabolism, LDs are inherently involved in the etiology of prevalent metabolic diseases such as obesity and nonalcoholic fatty liver disease. The functions of LDs are regulated by a unique set of associated proteins, the LD proteome, which includes integral membrane and peripheral proteins. These proteins control key activities of LDs such as triacylglycerol synthesis and breakdown, nutrient sensing and signal integration, and interactions with other organelles. Here we review the mechanisms that regulate the composition of the LD proteome, such as pathways that mediate selective and bulk LD protein degradation and potential connections between LDs and cellular protein quality control.

Keywords

lipid droplet; endoplasmic reticulum; triacylglycerol; ubiquitin; proteasome; lipophagy; chaperone-mediated autophagy; metabolism; protein targeting

1. INTRODUCTION

Lipid droplets (LDs) are neutral lipid storage organelles that function as hubs of cellular lipid and energy metabolism (Olzmann & Carvalho 2019, Renne et al. 2020, Walther et al. 2017). LDs provide a dynamic source of stored lipids that can be rapidly mobilized, releasing fatty acids that can be broken down by β -oxidation for energy, employed as building blocks for membrane biosynthesis, and used as lipid signaling molecules (Olzmann & Carvalho 2019). By sequestering lipids, LDs also play important roles in suppressing lipotoxicity by preventing cell death, endoplasmic reticulum (ER) stress, and mitochondrial dysfunction caused by free fatty acids (Olzmann & Carvalho 2019). Beyond their roles in lipid metabolism, LDs have also been implicated in less conventional roles such as the regulation of gene expression (Gallardo-Montejano et al. 2016, Mejhert et al. 2020), histone sequestration (Cermelli et al. 2006), viral replication (Miyanari et al. 2007), and drug

olzmann@berkeley.edu.

DISCLOSURE STATEMENT

The authors are not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

activity (Dubey et al. 2020, Greenwood et al. 2019). The underproduction and overproduction of LDs have been implicated in the etiology of several human diseases (Krahmer et al. 2013a, S. Xu et al. 2018).

LDs have a unique architecture consisting of a core of neutral lipids encircled by a phospholipid (PL) monolayer and a host of resident proteins. LDs are essentially surfactant-stabilized oil droplets suspended in the aqueous cytosol that undergo dynamic cycles of growth and degradation. There are two main steps in LD biogenesis. The first step is the biosynthesis of neutral lipids [e.g., triacylglycerols (TAGs) and sterol esters (SEs)] and their deposition between the leaflets of the ER bilayer. When neutral lipids accumulate to a particular level, a neutral lipid lens forms in the membrane driven by a liquid–liquid phase separation process (Khandelia et al. 2010, Zoni et al. 2020). This process is modulated by membrane curvature (e.g., ER tubules versus sheets) and the protein and lipid composition at sites of LD biogenesis (Khandelia et al. 2010, Santinho et al. 2020, Zoni et al. 2020). The second step of LD biogenesis involves the budding of the nascent LD toward the cytosol. The directionality of LD budding is driven by asymmetrical surface tension that is mediated by differences in the luminal and cytosolic leaflets of the ER bilayer (Ben M’barek et al. 2017, Chorlay et al. 2019).

Proteins such as seipin and PEX30 localize to sites of LD formation and regulate LD biogenesis (Olzmann & Carvalho 2019, Renne et al. 2020). Seipin-associated factors, such as lipid droplet–associated factor 1 (LDAF1) (also known as promethin and TMEM159) in mammalian cells and lipid droplet organization proteins of 16- and 45-kDa (LDO16 and LDO45) in yeast, also contribute to organized LD biogenesis (Olzmann & Carvalho 2019, Renne et al. 2020). These integral membrane proteins may cooperate with recruited cytosolic factors such as the perilipin (PLIN) family of LD scaffold proteins (Olzmann & Carvalho 2019, Renne et al. 2020). Precisely how the emerging set of LD biogenesis factors influences neutral lipid phase separation and membrane surface tension to promote organized LD formation is incompletely understood. Mature LDs are subject to regulated degradation through two pathways: a lipolytic pathway that employs the consecutive actions of lipases recruited to the LD surface and a selective autophagy pathway termed lipophagy characterized by the autophagosome-mediated delivery of LDs to autolysosomes for breakdown (Schulze et al. 2017, Zechner et al. 2017).

The unique complement of proteins decorating the LD surface (i.e., the LD proteome) regulates the LD life cycle and its functions, including growth, degradation, nutrient signal integration, and the formation of membrane contact sites with other organelles. In this review, we focus on how the LD proteome is established, our emerging understanding of the mechanisms that mediate LD proteome remodeling, and the connections between LDs and cellular protein quality control pathways.

2. LIPID DROPLET STRUCTURE AND PROTEOME

2.1. Physical Properties and Lipid Composition of Lipid Droplets

LDs are an emulsion of neutral lipids stabilized by a shell of surfactants consisting of amphipathic lipids and proteins. The LD core is an amalgam of neutral lipids, predominantly

TAGs and SEs, that is devoid of proteins. LDs may also contain retinyl esters, ether lipids, acylceramides, squalene, fat-soluble vitamins, and waxes, with the precise composition differing between cell types and metabolic conditions. The diverse composition of the lipid core suggests that LDs impact many cellular processes beyond their canonical role in energy metabolism. Electron and cryo-electron microscopy studies indicate an additional layer of complexity, providing evidence that the neutral lipid core has an internal structure that depends on the cellular state (Cheng et al. 2009, Czabany et al. 2008, Mahamid et al. 2019, Tauchi-Sato et al. 2002). Although the internal structure of LDs normally consists of an amorphous emulsion of neutral lipids, during mitotic arrest and nutrient deprivation this structure reorganizes into a liquid crystalline shell (possibly composed of SEs) with an amorphous core (Fujimoto & Parton 2011, Mahamid et al. 2019). The impact of the altered internal structure of the LD on lipid access and metabolism, lipid exchange, and protein binding is as yet unknown.

The LD hydrophobic core is bounded by a monolayer of PLs that act as surfactants to maintain the structural integrity of the LD. Lipidomic analyses identify a complex composition of PLs (Bartz et al. 2007a) that is critical for LD biogenesis and the maintenance of LD morphology. The initial PLs surrounding LDs are derived from the outer leaflet of the ER, and continuous PL synthesis is essential during LD biogenesis both to maintain ER membrane homeostasis and to support directional LD emergence (Chorlay et al. 2019). Similarly, PL synthesis is required for the growth of mature LDs and for the prevention of LD coalescence (Krahmer et al. 2011). How the LD PL composition is maintained and modulated remains for now mostly unknown. As most PLs are synthesized in the ER, ER–LD membrane continuities and lipid-transfer proteins (LTPs) provide mechanisms for PL delivery. The oxysterol-binding protein (OSBP) and OSBP-related proteins (ORPs) are an important family of LTPs, and ORP2 and ORP5 have been observed at ER–LD contacts (Du et al. 2020, Hynynen et al. 2009). ORP5 mediates the trafficking of both phosphatidylserine from the ER to LDs and phosphatidylinositol 4-phosphate from LDs to the ER (Du et al. 2020). Several members of the chorein domain–containing family of LTPs also localize to ER–LD contacts, including ATG2, VPS13a, and VPS13c (Kumar et al. 2018, Valverde et al. 2019). This unique family of LTPs is distinguished by the ability to bind and transfer tens of glycerophospholipids at once via a large hydrophobic cavity (Kumar et al. 2018, Valverde et al. 2019), providing a mechanism for rapid PL delivery and the equilibration of membrane compositions. Future research is necessary to define the full complement of LD-associated LTPs and their functions at LD–organelle membrane contact sites.

2.2. Lipid Droplet Proteome Composition

A unique complement of integral and peripheral proteins associates with the LD PL monolayer (Bersuker & Olzmann 2017, Zhang & Liu 2019). LD-localized proteins were first observed in the early 1990s, when perilipin-1 (PLIN1) was found in adipocyte LD fractions and localized to the LD surface (Greenberg et al. 1991). More recently, proteomic analyses of LDs isolated by biochemical fractionation on sucrose gradients have enabled the characterization of the LD proteome composition in numerous species, cell types, and tissues (Bersuker & Olzmann 2017, Zhang & Liu 2019). The associated repertoire of LD

proteins differs between cell types and reflects the cell's metabolic state, and alterations are observed in cultured adipocytes following induction of lipolysis (Brasaemle et al. 2004) in mice on high-fat diets (Crunk et al. 2013, D'Aquila et al. 2019, Ding et al. 2012, Khan et al. 2015, Kraemer et al. 2018, Liu et al. 2017b) and in human liver and subcutaneous adipose tissue biopsies from patients with nonalcoholic fatty liver disease and insulin resistance, respectively (Su et al. 2014, Xie et al. 2016). These dynamic changes highlight the importance of LD proteome remodeling in modulating LD functions to match cellular energy demands.

LD-associated proteins constitute a variety of functional classes (Bersuker & Olzmann 2017, Zhang & Liu 2019). Many LD proteins are lipid metabolic enzymes that mediate LD growth and degradation. These proteins include enzymes involved in TAG synthesis, such as the acyl-CoA synthetase ACSL3 and the acyltransferases GPAT4 and DGAT2, as well as enzymes involved in TAG breakdown, such as the lipolysis rate-limiting enzyme adipose triglyceride lipase (ATGL) and its associated regulators CGI-58, HILPDA, and G0S2. The PLIN family of proteins (consisting of PLINs 1–5) are invariably found on LDs, with the expression of family members differing by tissue. PLINs are generally thought to function as scaffold proteins that maintain the structural integrity of the LD, but they have also been implicated as integrators of nutrient signaling pathways, regulators of lipolysis, tethering proteins at LD–organelle membrane contact sites, and direct mediators of lipid signaling pathways (Kimmel & Sztalryd 2016). Some cells express members of the cell death-inducing DNA fragmentation factor 45-like effector (CIDE) family of proteins, which mediate the regulated fusion of LDs. Beyond lipid metabolism, other functional classes of proteins are present on LDs, including proteins involved in trafficking (e.g., motor and cytoskeletal proteins) and many Rab guanosine triphosphatases (GTPases) (Bartz et al. 2007b; Liu et al. 2004, 2007). Some of the Rabs affect LD functions, such as Rab18 regulation of ER–LD contacts (D. Li et al. 2019, D. Xu et al. 2018) and Rab10 regulation of the autophagic clearance of LDs (Li et al. 2016, Z. Li et al. 2019), but the effects of most Rab family proteins on LDs is as yet unclear. Several proteins have been implicated in the formation of LD–organelle contact sites that promote fatty acid oxidation. For example, M1 spastin links LDs to peroxisomes through its interactions with peroxisomal ABCD1 and facilitates fatty acid transfer to the peroxisome at these sites by recruiting the membrane-shaping ESCRT-III proteins IST1 and CHMP1B (Chang et al. 2019). Other functional classes include ubiquitination, signaling, transcription, and redox, but their roles in LD biology are not yet well understood.

An ongoing challenge for LD proteomic approaches is distinguishing putative LD proteins from contaminants that copurify in LD-enriched buoyant fractions. Protein correlation profiling (Currie et al. 2014; Kraemer et al. 2013b, 2018) and proximity labeling proteomic analyses of LD proteins using the engineered ascorbate peroxidase APEX2 (Bersuker & Olzmann 2019, Bersuker et al. 2018) provide two methods that are useful for identifying high-confidence LD proteomes and avoid the identification of common, abundant contaminant proteins. These approaches identify approximately 100–150 LD proteins in mammalian cells and 30–40 LD proteins in yeast, some with exclusive LD localization and others found in multiple subcellular compartments. A second challenge is mapping the proteomes of heterogeneous subpopulations of LDs, which can differ in size, localization,

function, and organelle contacts. This heterogeneity is lost during bulk isolation of fractionated LDs. Further separation of the isolated LD-enriched buoyant fraction has been used to compare the proteomes of LDs with different sizes (Zhang et al. 2016). Proximity labeling proteomics approaches provide a promising method for profiling separate sets of LD proteomes by tagging differentially localized LD proteins with enzymes that mediate the biotinylation of neighboring proteins.

3. PROTEIN TARGETING TO LIPID DROPLETS

Targeting proteins to LDs presents several unique challenges. In contrast to most other organelles, neither protein insertion machinery (analogous to the ER translocon or peroxisomal importomer) nor specific lipids have been identified on LDs for protein recruitment and insertion. LD proteins also do not contain a signal or targeting sequence that can be predicted by informatic approaches. However, LD proteins exhibit some shared structural features because of the topological constraints imposed by the architecture of the LD monolayer, which requires LD proteins to adopt conformations that avoid exposing hydrophilic domains to the neutral lipid core of the LD. For this reason, integral LD proteins are monotopic proteins that insert into the LD monolayer and orient their soluble domains toward the cytosol. Bitopic and polytopic transmembrane proteins that typically associate with bilayers are excluded from LDs. A general classification system for integral LD proteins has been adopted based on their trafficking pathways: Class I LD proteins insert first into the ER and traffic to LDs, and Class II LD proteins insert directly into LDs from the cytosol (Kory et al. 2016) (Figure 1). Both Class I and Class II LD proteins can recruit soluble, peripheral proteins to LDs.

3.1. Class I Lipid Droplet Proteins: Endoplasmic Reticulum to Lipid Droplet Targeting

Class I LD proteins insert into the ER and laterally diffuse from the outer leaflet of the ER to the PL monolayer of nascent LDs (Figure 1). In the absence of LDs, Class I LD proteins typically exhibit an ER distribution. Examples of Class I LD proteins include the lipid biosynthesis enzymes GPAT4 and DGAT2, the acyl-CoA synthetase ACSL3, the ER-associated degradation (ERAD) factors AUP1 and UBXD8, the putative methyltransferase AAM-B, caveolin-1 and caveolin-2, HSD17B11, the hepatitis C virus (HCV) core protein, and ALDI.

Class I LD proteins are generally thought to adopt a hairpin conformation in which a hydrophobic domain embedded in the monolayer is flanked by hydrophilic regions that are exposed to the cytosol, forming a V shape extending from the membrane. These proteins lack the luminal loops or domains typically found in transmembrane proteins, allowing them to be accommodated in the outer leaflet of the ER and the monolayer of the LD. Crystal structures of Class I LD proteins are lacking, and the model of hairpin conformation for these proteins is primarily based on the bioinformatic prediction of hydrophobic regions, the biochemical characterization of membrane insertion, and the protease sensitivity of N- and C-terminal tags. The hairpin structure was first noted for plant structural proteins called oleosins, which contain a critical conserved proline knot buried in the hydrophobic sequence (Abell et al. 1997). Although a proline knot is required for LD insertion of oleosins (Abell et

al. 1997) and HCV core protein (Hope et al. 2002), mutation of central prolines in the hydrophobic domains of AAM-B, ALDI, and CYB5R3 (Zehmer et al. 2008) does not abolish LD localization, indicating that the proline knot is not a universal requirement for Class I LD protein targeting.

Most secreted and transmembrane proteins are inserted into the ER through well-characterized pathways, with proteins bearing an N-terminal signal sequence inserted cotranslationally via the Sec61 translocon and proteins bearing C-terminal transmembrane domains (i.e., tail-anchored proteins) inserted posttranslationally via the guided entry of tail-anchor (GET) pathway (Shao & Hegde 2011). The mechanism of ER insertion for Class I LD proteins has not been extensively studied. At least one Class I LD protein, UBXD8, relies on a distinct posttranslational insertion pathway mediated by the peroxisome biogenesis factors PEX19 and PEX3 (Schrul & Kopito 2016). In this pathway, the cytosolic receptor PEX19 binds to the hydrophobic segment of UBXD8 and mediates UBXD8 delivery to specific ER subdomains containing its binding partner PEX3, an ER-resident integral membrane protein (Schrul & Kopito 2016). Whether LD proteins other than UBXD8 use this insertion pathway is unknown.

The mechanisms that control Class I LD protein partitioning between the ER and LDs remain unresolved. The junction between the ER bilayer and the LD monolayer provides a diffusion barrier that is permissive for monotopic protein trafficking but excludes bitopic and polytopic membrane protein movement. However, this diffusion barrier does not explain why some Class I LD proteins concentrate in LDs. The distribution of Class I LD proteins between the ER and LDs can be impacted by selective ubiquitin-dependent degradation of their ER pools (Ruggiano et al. 2016) and by their tethering to protein-binding partners in the ER. ER–LD partitioning of these proteins may be driven by differences in their conformations or mobility in the ER bilayer and the LD monolayer that impact the proteins' stability, degradation, and residence time.

3.2. Class II Lipid Droplet Proteins: Cytosol to Lipid Droplet Targeting

Class II LD proteins are generally thought to be translated on cytosolic ribosomes and posttranslationally inserted into the LD directly from the cytosol (Figure 1). Examples of Class II proteins include the PLIN family, the rate-limiting enzyme of phosphatidylcholine (PC) synthesis CTP:phosphocholine cytidyltransferase (CCT), the LD–LD fusogenic protein CIDEA, and the antiviral protein viperin.

How LD proteins are selectively targeted to the LD monolayer as opposed to other membrane-bound compartments has been an enigma. Recent *in silico* molecular dynamics studies provide an answer, revealing that a PL monolayer encircling a neutral lipid core exhibits persistent and large PL packing defects at the interfacial region in which acyl chains of the bounding PLs and core TAGs are exposed to the cytosol (Bacle et al. 2017, Prévost et al. 2018). This unique feature of LD membranes is exploited by Class II LD proteins, which often contain amphipathic helices with large hydrophobic residues that detect and bind to the hydrophobic PL packing defects (Prévost et al. 2018). Amphipathic helices are not unique to LD proteins, and they are common bilayer-targeting domains found in many proteins that localize to other subcellular compartments. Indeed, promiscuous LD localization of

amphipathic helices from a range of proteins is observed when they are ectopically expressed (Prévost et al. 2018). The mechanisms that mediate the selective targeting of LD proteins and exclusion of non-LD proteins that contain amphipathic helices are unclear but may involve differences in their affinities for transiently exposed neutral lipids (Chorlay & Thiam 2020, Prévost et al. 2018), exclusion by macromolecular crowding effects on the LD surface (Kory et al. 2015), and regulation by dominant targeting information present in regions flanking the amphipathic helix (Ingelmo-Torres et al. 2009). The localization of Class II LD proteins may also be favored by a series of amphipathic helices that would increase avidity and binding. The PLIN family of LD proteins localize almost exclusively to LDs and exhibit a series of N-terminal 11-mer repeats that are predicted to form amphipathic helices (Ajjaji et al. 2019, opí et al. 2018, Rowe et al. 2016). The PLIN4 11-mer repeat region forms a particularly long amphipathic helix (>60 amino acids), binds directly to TAG, and acts as a potent surfactant to stabilize LDs during periods in which PLs are limiting (opí et al. 2018). PLINs 1–3 also contain a C-terminal amphipathic helix consisting of a four-helix bundle domain that can differentially stabilize LD binding and may underlie the observed hierarchical association of PLINs with the LD (Ajjaji et al. 2019). Not all Class II LD proteins contain amphipathic helices. Some LD proteins contain large hydrophobic residues important for insertion that could still exploit LD PL packing defects as a recruitment mechanism (Boeszoermyeni et al. 2015).

Fatty acid modifications are important for the localization of some Class II LD proteins. The conjugation of fatty acids (e.g., myristate and palmitate) to proteins provides a hydrophobic anchor that facilitates binding to PL membranes. Palmitoylation of the Arf-GTPase activating protein ELMOD2 (Suzuki et al. 2015) and the SNARE protein SNAP-23 (Boström et al. 2007) is required for their LD localization. Palmitoylation is a reversible fatty acid modification that provides a potential switchlike mechanism for regulating the LD association of palmitoylated proteins. Myristoylation is an irreversible fatty acid modification that was recently shown to be required for the LD and plasma membrane localization of FSP1, a Coenzyme Q10 oxidoreductase implicated in preventing oxidative lipid damage and ferroptosis (Bersuker et al. 2019, Doll et al. 2019) and in regulating NADH redox during thermogenesis in brown adipocytes (Nguyen et al. 2020). Additionally, modification by prenylation mediates LD targeting of the aldehyde dehydrogenase ALDH3B2 (Kitamura et al. 2015). It will be important to determine the extent to which other LD proteins are modified by fatty acids and the signals in flanking regions of the protein that contribute to their LD targeting and binding, as several hundred mammalian proteins are modified with fatty acids yet most are not LD localized.

4. UBIQUITIN-DEPENDENT LIPID DROPLET PROTEIN DEGRADATION

Cellular physiology and organelle functions are determined by the composition and integrity of the cellular proteome. Cells have evolved elaborate networks of protein-folding and degradation factors that monitor and maintain the integrity of the cellular proteome by recognizing misfolded proteins and mediating their refolding, sequestration, or degradation (i.e., protein quality control). These same pathways also regulate the levels of folded proteins, modulating the composition of the proteome and enabling cells to adapt to changes in cellular requirements (i.e., protein quantity control). Organelles contain specialized

pathways for protein quality and quantity control. For example, proteins in the early secretory pathway can be targeted for clearance either through ERAD (Olzmann et al. 2013a), which mediates the ubiquitin-dependent delivery of substrates to the proteasome, or through a selective ER autophagy pathway known as ER-phagy (Chino & Mizushima 2020), which mediates substrate delivery to the lysosome.

The mechanisms of LD protein quality and quantity control remain poorly understood. LD proteins traffic to and from different subcellular compartments (e.g., the ER and the cytosol), and multiple folding and degradation pathways could be involved in their regulation (Figure 2). Several LD proteins are known to be targeted for degradation by the ubiquitin-proteasome system (UPS) under various metabolic conditions, including PLIN1 (Xu et al. 2006), PLIN2 (Eastman et al. 2009, Masuda et al. 2006, Nguyen et al. 2019, Takahashi et al. 2016, Xu et al. 2005), CIDEA (Chan et al. 2007, Qi et al. 2008), CIDEA/FSP27 (Nian et al. 2010, Qian et al. 2017, Zhang et al. 2014), ATGL (Dai et al. 2013, Ghosh et al. 2016, Niyogi et al. 2019, Olzmann et al. 2013b), DGAT2 (Choi et al. 2014, Luo et al. 2018), CGI-58 (Patel et al. 2014), G0S2 (Heckmann et al. 2016, Kamikubo et al. 2019, Yang et al. 2011), PNPLA3 (BasuRay et al. 2017, 2019), and c18orf32 (Bersuker et al. 2018). In the UPS pathway, ubiquitin is conjugated to substrates by the consecutive actions of E1 activating enzymes, E2 conjugating enzymes, and E3 ligases. For most LD substrates, the ubiquitination machinery involved in their degradation is unknown and challenging to characterize, as the cell contains hundreds of E3 ligases and numerous adaptor proteins and accessory factors.

4.1. Endoplasmic Reticulum–Associated Degradation

The LD and ER membranes are, at least initially, contiguous, and they continue to be held in close contact by multiple ER–LD tethering proteins (Olzmann & Carvalho 2019). The levels of Class I LD proteins trafficking through the ER en route to LDs may be regulated by ERAD, a process in which a network of factors organized around membrane-embedded E3 ligases mediates substrate recognition, ubiquitination, membrane extraction by the AAA-ATPase p97/VCP (Cdc48 in yeast), and delivery to the proteasome (Figure 2) (Olzmann et al. 2013a).

In yeast, the ER-resident E3 ligase Doa10 targets several LD proteins (e.g., Pgc1, Dga1, and Yeh1) for proteasomal degradation (Ruggiano et al. 2016). The hydrophobic hairpins of certain Class I LD proteins act as Doa10 degrons (Ruggiano et al. 2016), possibly because the ER contains machinery capable of recognizing the hairpin as an aberrant structure and/or because the hairpin adopts a distinct, destabilized conformation in the ER bilayer in contrast to a more stable conformation in the LD monolayer. Doa10-mediated degradation of mislocalized LD proteins limits their levels in the ER and leads to their concentration in LDs (Ruggiano et al. 2016), indicating that ERAD is essential for the maintenance of ER and LD proteome compositions. The clearance of mislocalized LD proteins is reminiscent of other forms of spatial quality control such as the degradation of mislocalized ER proteins in the inner nuclear membrane (Foresti et al. 2014, Khmelinskii et al. 2014) and mislocalized tail-anchored proteins in the mitochondria (Chen et al. 2014, Okreglak & Walter 2014).

The process of Class I LD protein degradation by ERAD is conserved in more complex eukaryotes, as DGAT2 (Choi et al. 2014, Luo et al. 2018) and c18orf32 (Bersuker et al. 2018) are degraded by ERAD in human cell lines. DGAT2, and the evolutionarily unrelated DGAT1, are enzymes that mediate the conversion of diacylglycerol (DAG) into TAG for storage in LDs. In contrast to DGAT1, which is a polytopic ER-resident enzyme (McFie et al. 2010), DGAT2 consists of a membrane-embedded hairpin with its C-terminal active site oriented toward the cytosol (Stone et al. 2006). DGAT2 is present in the ER and traffics to LDs (Stone et al. 2009, Wilfling et al. 2013) or regions of ER tethered to LDs (McFie et al. 2018), where it mediates local TAG synthesis and LD expansion (Wilfling et al. 2013). Two ERAD pathways using distinct E3 ligases, gp78 (Choi et al. 2014) and Hrd1 (Luo et al. 2018), have been implicated in the ubiquitin-dependent proteasomal degradation of DGAT2. Whether gp78 directly recognizes DGAT2 or uses adaptor proteins to mediate their interaction is unclear. In the case of Hrd1-mediated DGAT2 degradation, Axin interactor dorsalization-associated protein (AIDA) has been implicated as an adaptor that couples Hrd1 and DGAT2 (Luo et al. 2018). AIDA is also required for the degradation of the ER-resident lipid metabolism enzymes MOGAT2 and GPAT3, suggesting a key role for AIDA in regulating the levels of multiple components of a TAG synthesis pathway (Luo et al. 2018).

Another Class I LD protein that is degraded via ERAD is c18orf32 (Bersuker et al. 2018). C18orf32 is a small protein with an N-terminal hydrophobic region that is required for its ER and LD targeting (Bersuker et al. 2018). Cells lacking c18orf32 exhibit changes in the lipid composition of their LDs (Bersuker et al. 2018), but the function of c18orf32 is unknown. The degradation of c18orf32 requires the E3 ligase gp78, the rhomboid pseudoprotease derlin-1, and VCP (Bersuker et al. 2018). Inhibition of gp78 increases the levels of c18orf32 on LDs (Bersuker et al. 2018), consistent with a role for ERAD in regulating the LD proteome composition by controlling the pool of Class I LD proteins available to traffic from the ER to LDs.

The features of DGAT2 and c18orf32 recognized by ERAD machinery are unclear. The data in yeast (Ruggiano et al. 2016) raise the possibility that the hairpin region of the proteins could act as the degron. The first putative transmembrane domain of DGAT2 has been proposed to function as a degron (Choi et al. 2014), but this finding is controversial (Brandt et al. 2016). The presence of a hairpin structure is not always sufficient to induce LD protein degradation in the ER. For example, the Class I LD protein UBXD8 contains a hairpin and is stable both in the ER and in LDs (Olzmann et al. 2013b). This may be due to the association of UBXD8 with ER-binding partners such as UBAC2 (Olzmann et al. 2013b) that could prevent recognition of a degron.

Although examples of ERAD-mediated clearance of Class I LD proteins have been reported, whether ERAD targets Class II LD proteins for degradation is less clear because this would require trafficking of the proteins from LDs to the ER. LDs and the ER form dynamic bridging contacts, fusions of the LD PL monolayer with the outer leaflet of the ER bilayer, which provide a mechanism for protein exchange (Wilfling et al. 2013) and could allow LD proteins to access ERAD machinery. In cells with low basal levels of LDs, uninserted PLIN2 is degraded through a UPS pathway that involves the ERAD E3 ligase MARCH6 (Nguyen et al. 2019). MARCH6 and its yeast ortholog Doa10 have been implicated in the N-degron

pathway (also referred to as the N-end rule pathway), which recognizes proteins with destabilizing N-terminal amino acids that act as degrons (Varshavsky 2019). Indeed, the initiator methionine of PLIN2 is cleaved, yielding an alanine that is acetylated (Nguyen et al. 2019, Takahashi et al. 2016). This N-terminal acetylated alanine facilitates the binding of MARCH6 and the ubiquitin-dependent degradation of PLIN2 (Nguyen et al. 2019). MARCH6 and Doa10 have also been reported to degrade cytosolic proteins, and, in some cases, the protein first associates with the outer leaflet of the ER through an amphipathic helix (Stefanovic-Barrett et al. 2018). Whether, in the absence of LDs, PLIN2 inserts into the ER to associate with MARCH6 or cytosolic PLIN2 associates with MARCH6 for clearance is unknown. Oleate-induced LD biogenesis prevents PLIN2 degradation (Masuda et al. 2006, Xu et al. 2005), indicating that PLIN2 LD insertion can impair recognition by the MARCH6-mediated N-degron pathway, despite the presence of the N-terminal destabilizing residue. Whether PLIN2 that has been inserted into LDs can be degraded by MARCH6 or another ERAD pathway during lipolytic degradation of LDs is not yet known.

4.2. Lipid Droplet–Associated Degradation

The association of ubiquitination machinery with LDs raises the possibility of an LD-associated degradation pathway (Figure 2). Several proteins implicated in ERAD traffic from the ER to LDs, including UBXD8 and UBXD2, which recruit p97/VCP to LDs through their UBX domains (Olzmann et al. 2013b, Suzuki et al. 2012), and AUP1, which recruits the E2 enzyme UBE2G2 to LDs through its C-terminal UBE2G2-binding motif (Klemm et al. 2011, Spandl et al. 2011). This set of ubiquitination and VCP-recruitment factors is suggestive of a complex that could extract ubiquitinated proteins for proteasomal degradation. However, an LD-associated E3 ligase that cooperates with these factors and substrates for this putative degradation pathway has yet to be identified. AIP4/5, which are recruited to LDs by spartin (Eastman et al. 2009, Edwards et al. 2009, Hooper et al. 2010, Tokunaga et al. 2013), and SMURF1 (Petrasek et al. 2019) have been proposed to function as LD-associated E3 ligases. These proteins have not been observed in most proteomic analyses of LD fractions, although their association may be lost during the biochemical isolation of LDs or be limited to specific cell types. RNF213 (also known as mysterin), which is mutated in the rare cardiovascular disorder moyamoya disease, also localizes to a subset of LDs (Sugihara et al. 2019). RNF213 is a large protein that contains two AAA-ATPase domains and a RING finger domain that confers E3 ligase activity (Liu et al. 2011, Morito et al. 2014). RNF213 competes for LD localization with ATGL, the rate-limiting enzyme in lipolysis, and thereby impacts LD abundance and turnover (Sugihara et al. 2019). The localization of RNF213, and hence its ability to displace ATGL, requires its RING finger and AAA-ATPase domains (Sugihara et al. 2019), but ATGL does not appear to be an RNF213 substrate. Whether RNF213 ubiquitinates any LD proteins is unknown at this time.

An LD-associated protein degradation pathway has also been proposed in plants (Deruyffelaere et al. 2018, Kretzschmar et al. 2018). Oleosins are plant LD scaffold proteins that stabilize LDs and are degraded by the UPS following seed germination. Studies in *Arabidopsis thaliana* indicate that the removal of ubiquitinated oleosins from LDs and their subsequent proteasomal degradation are dependent on PUX10, a UBXD8 ortholog that recruits the VCP ortholog Cdc48 to LDs via its C-terminal UBX domain (Deruyffelaere et

al. 2018, Kretzschmar et al. 2018). The signals that recruit PUX10 to the LD, as well as the machinery required for oleosin ubiquitination, remain to be determined.

4.3. Cytosolic Degradation Pathways

Class II LD proteins synthesized in the cytosol may be degraded via cytosolic degradation pathways prior to insertion (Figure 2). In addition, as LDs are degraded during lipolysis, they shrink, and protein crowding forces weakly bound proteins off of the LD into the cytosol (Kory et al. 2015). Whether these LD proteins employ cytosolic UPS pathways is mostly unexplored. Nonetheless, ATGL is reported to be targeted for degradation by the E3 ligase COP1 (Ghosh et al. 2016, Niyogi et al. 2019), and GOS2 is targeted for degradation by the E3 ligase RNF126 along with the chaperone Bag6 (Kamikubo et al. 2019). Bag6 and RNF126 play a well-characterized role as a cytosolic complex that recognizes hydrophobic regions of mislocalized secretory proteins that have failed to insert into the ER or have been retrotranslocated in certain ERAD pathways. Whether LD proteins that fail to insert into LDs and mislocalize to the cytosol engage this pathway remains an open question.

5. AUTOPHAGIC DEGRADATION OF LIPID DROPLET PROTEINS

5.1. Bulk Degradation by Lipophagy

Orchestrated by a host of autophagy-related (ATG) proteins, autophagy is an evolutionarily conserved process that mediates the delivery of cytoplasmic components to the lysosome for the catabolism and recycling of macromolecules (Mizushima 2018). Macroautophagy (hereafter referred to as autophagy) involves the sequestration of portions of cytoplasm within double-membrane-bounded vesicles (i.e., autophagosomes) that then fuse with lysosomes to enable the degradation of cargo by lysosomal enzymes. Autophagy can nonselectively degrade portions of cytoplasm, or it can be harnessed for the selective clearance of specific cargo such as protein aggregates and damaged organelles. A seminal study revealed that LDs in hepatocytes are broken down by a selective form of autophagy now known as lipophagy, resulting in the nonselective, bulk degradation of LD proteins and lipids (Schulze et al. 2017, Singh et al. 2009, Zechner et al. 2017).

Lipophagy is regulated by two small Rab GTPases, Rab10 and Rab7, that localize to LDs and are activated under autophagy-stimulating conditions (Li et al. 2016, Schroeder et al. 2015). Rab10 is required for the recruitment of LC3-positive autophagic membranes to LDs. It also forms a complex with the adaptor protein EH domain-binding protein 1 and the membrane-deforming ATP EH domain-containing 2, which together promote phagophore expansion and the engulfment of LDs within the autophagosome (Li et al. 2016). Rab7 regulates the trafficking of multivesicular bodies and lysosomes to LDs (Schroeder et al. 2015). The selective autophagy of many organelles involves organelle-specific receptors that bind to the autophagosome protein LC3 and initiate the assembly of the phagophore around the target organelle. The recruitment of autophagosome membranes can be mediated by organelle-resident receptors that contain Atg8 family-interacting motifs or LC3-interacting regions, such as Atg32, NIX, and BNIP, or by LIR-containing adapter proteins, such as OPTN, NBR1, TAX1B1, p62, and NDP52, that bind to ubiquitinated cargo (Pickles et al. 2018). Whether lipophagy employs a specific receptor or an alternative mechanism for the

recruitment of autophagic membranes remains unclear. P62 is one possible candidate for a lipophagy receptor, as it associates with LDs and is required for lipophagy (Lam et al. 2016, Tapia et al. 2019, Wang et al. 2017, Yan et al. 2019). To date no ubiquitination machinery has been implicated in either p62 recruitment to LDs or lipophagy. Deciphering what the mechanisms are that mediate the recruitment of autophagic machinery to LDs and if it is ubiquitin-dependent remain important questions.

Lipolysis and lipophagy can act synergistically in LD catabolism. The removal of PLIN2 by chaperone-mediated autophagy (CMA) (discussed in detail in the next section) is an early step in LD catabolism that is important for the recruitment of ATGL and autophagic membranes to LDs (Kaushik & Cuervo 2015). In hepatocytes, ATGL-dependent lipolysis functions upstream of lipophagy, reducing the size of existing LDs while liberating fatty acids that are reesterified and packaged into small LDs (Schott et al. 2019). LDs in hepatocytes can reach sizes $>5 \mu\text{m}$ in diameter. Lipolysis-mediated reductions in LD volume may be necessary to meet a size threshold compatible with autophagosome engulfment and fusion with the lysosome during lipophagy. The fatty acids released during lipolysis also function as signaling molecules to upregulate lipophagy gene programs through SIRT1 (Sathyanarayan et al. 2017). These studies indicate that LD size and fatty acid-regulated gene programs are important regulators of lipophagy, although their overall contribution to LD catabolism likely differs according to cell type and metabolic conditions.

In yeast, LDs are degraded by microlipophagy in response to nutrient-depleted conditions, including nitrogen starvation, stationary phase growth, and acute glucose restriction (Graef 2018). During microlipophagy, LDs are not enveloped within an autophagosome as in lipophagy. Instead, LDs are delivered directly to the vacuole (the yeast equivalent of the lysosome) for internalization and breakdown. LD transport to the vacuole may be facilitated by the biogenesis of LDs proximal to the vacuole under nutrient stress conditions and the physical tethering of the ER, vacuole, and LDs at the nuclear ER–vacuole junction by Mdm1p (Hariri et al. 2018, 2019). An essential event in microlipophagy is the striking partitioning of vacuolar lipids and proteins into microdomains, yielding large sterol- and sphingolipid-rich, raft-like liquid-ordered (Lo) domains that mediate LD docking and internalization (Toulmay & Prinz 2013, van Zutphen et al. 2014, Wang et al. 2014). LDs bound at these Lo domains are internalized by ESCRT-mediated inward budding of the vacuolar membrane, and subsequent membrane scission liberates LD-containing luminal vesicles for degradation by vacuolar hydrolases (Vevea et al. 2015). The catabolism of internalized LDs releases sterols that are required for the formation of the Lo domains, indicating a feedforward loop and a reciprocal relationship between LDs and microlipophagy (Seo et al. 2017, Wang et al. 2014). The NPC orthologs Ncr1p and Npc2p also play important roles in facilitating the formation and expansion of the Lo domains by transferring sterols from intraluminal vesicles formed by the multivesicular body pathway (Tsuji et al. 2017). Although LDs are not encapsulated within autophagosomes during microlipophagy, this process still requires core autophagy machinery and a subset of ATG proteins. Autophagy may impact microlipophagy indirectly due to its roles in the trafficking of Ncr1p, Npc2p, and sphingolipids to the vacuole for the formation of vacuolar Lo domains (Tsuji et al. 2017). However, some ATG proteins play direct roles in microlipophagy. For example, during acute glucose starvation, Atg14p transits from ER exit sites to Lo domains

in an AMPK/Snf1p-dependent manner (Seo et al. 2017). Atg14p localizes to the edges of the Lo domains and, together with Atg6p, drives the enlargement of Lo domains (Seo et al. 2017). Microlipophagy has also been observed to occur independently of core ATG machinery in response to defects in PC synthesis (Vevea et al. 2015) and the transition through diauxic growth (Oku et al. 2017). Whether microlipophagy occurs independently of Lo domains under these conditions or if Lo domains form through an alternate mechanism are not known. Furthermore, there is currently a lack of evidence demonstrating microlipophagy in mammalian cells.

5.2. Chaperone-Mediated Autophagy

CMA is a protein-degradation pathway that mediates the delivery of substrates to the lysosome for proteolysis (Kaushik & Cuervo 2018). In CMA, substrate proteins containing specific pentapeptide (KFERQ-like) motifs are recognized by the cytosolic chaperone heat shock-cognate 70-kDa protein (Hsc70) and shuttled to the lysosomal surface, where the Hsc70-substrate complex binds to lysosome-associated membrane protein type 2A (LAMP2A). Hsc70 facilitates substrate unfolding and translocation into the lysosomal lumen for proteolytic breakdown by lysosomal proteases. KFERQ-like CMA motifs were identified in PLIN2 (LDRLQ) and PLIN3 (SLKVQ), which target them for CMA (Kaushik & Cuervo 2015). Brief kiss-and-run as well as prolonged interactions between LDs and lysosomes have been observed; these organelle contact sites allow PLIN2 and PLIN3 to be transferred from LDs to lysosomes (Kaushik & Cuervo 2015, Schroeder et al. 2015, Valm et al. 2017). Mutation of the PLIN2 CMA-targeting motif reduces LD-lysosome contacts (Kaushik & Cuervo 2015), suggesting that a portion of these contacts may be mediated by interactions between PLINs and the CMA machinery. The small GTPase Rab7 has also been implicated in the formation of LD-lysosome contacts, the transfer of PLIN2 from LDs to lysosomes, and lipophagy (Schroeder et al. 2015), raising the possibility that Rab7 may regulate the CMA of LD proteins or a piecemeal autophagic event. The CMA-mediated degradation of PLIN2 is coupled to cellular energetics by 5'-AMP-activated protein kinase (AMPK), which phosphorylates PLIN2 and promotes its CMA during starvation (Kaushik & Cuervo 2016). Interestingly, the CMA-mediated removal of PLIN2 and PLIN3 facilitates the recruitment of lipolytic and lipophagic machinery (Kaushik & Cuervo 2015), suggesting that CMA degradation of PLINs is an upstream, initiating event for LD degradation by lipolysis and lipophagy. The disruption of CMA in cultured cells and in mouse models results in TAG and LD accumulation (Kaushik & Cuervo 2015), likely due to a combination of altered LD breakdown and reduced degradation of CMA substrates associated with lipid metabolism. Whether LD proteins other than PLIN2 and PLIN3 are targeted by CMA remains to be determined.

6. LIPID DROPLETS AND CELLULAR PROTEOSTASIS

While the canonical function of LDs is to store energy as TAG and suppress lipotoxicity, LDs have also been linked to less conventional roles in regulating cellular proteostasis and stress responses. The role of LDs in these processes is unclear but many possibilities have been suggested, including LDs sequestering aggregation-prone proteins prior to their degradation and regulating the ER lipid landscape to promote homeostasis.

6.1. Lipid Droplets and Endoplasmic Reticulum Stress

An imbalance between the ER protein-folding load and the capacity of quality control machinery can lead to the buildup of misfolded proteins in the ER lumen and membrane, a condition generally referred to as ER stress (Walter & Ron 2011). The cell is equipped with an adaptive homeostatic feedback mechanism, collectively referred to as the unfolded protein response (UPR), whose function is to alleviate ER stress (Walter & Ron 2011). The UPR encompasses three pathways initiated by ER transmembrane proteins (Ire1p in yeast; ATF6, PERK, and IRE1 in mammals) that sense the aberrant accumulation of unfolded ER proteins and activate downstream signaling cascades that lead to an expansion in the ER protein-folding capacity by coordinately inhibiting protein translation and upregulating the expression of ER folding machinery, ERAD factors, and lipid biosynthesis enzymes. The UPR can also be activated directly by alterations in the ER lipid composition that cause membrane bilayer stress independent of protein misfolding (Promlek et al. 2011, Volmer et al. 2013). Increased UPR signaling has been observed in yeast treated with saturated fatty acids or lacking key lipid metabolic enzymes (Jonikas et al. 2009, Pineau et al. 2009, Thibault et al. 2012), cultured mammalian cells treated with saturated fatty acids or following inhibition of steroyl-CoA desaturase (SCD1) (Ariyama et al. 2010, Cunha et al. 2008, Minville-Walz et al. 2010), animal models of metabolic diseases (Fu et al. 2011), and humans with obesity (Gregor et al. 2009).

The upregulation of LD biogenesis is a conserved response to ER stress. Yeast upregulate LD biogenesis under genetic or pharmacologically induced ER stress, such as treatment with the N-glycosylation inhibitor tunicamycin or the protein transport inhibitor Brefeldin A (Fei et al. 2009). Mammalian cells also increase LD production under pharmacologically induced ER stress (e.g., tunicamycin, thapsigargin, or bortezomib) (Hartman et al. 2010, Lee et al. 2012), as do the livers of transgenic mice that lack either Ire1 or ATF6 (Yamamoto et al. 2010, Zhang et al. 2011). Although LD biogenesis appears to be a conserved response to ER stress, it is not yet known why cells increase lipid storage under these conditions.

LDs may serve a protective role that suppresses ER stress. Consistent with this possibility, impairments in LD biogenesis are associated with an upregulation of UPR signaling. LD-deficient yeast strains (LD⁻) lacking the enzymes required for neutral lipid biosynthesis (*are1*, *are2*, *dga1*, and *lro1*) exhibit upregulated UPR signaling (Olzmann & Kopito 2011, Petschnigg et al. 2009) and sensitivity to oleic and palmitoleic acid but not palmitate (Garbarino et al. 2009, Petschnigg et al. 2009). In the absence of neutral lipid synthesis, oleate largely accumulates as a free fatty acid, whereas palmitate is shunted into PLs (Garbarino et al. 2009). Thus, the ability to buffer free fatty acids as neutral lipids or PLs is important for yeast to avoid fatty acid-mediated toxicity. Similarly, LDs are required to prevent ER stress and to provide a buffer for fatty acids to suppress lipotoxicity in mammalian cells. In adipocytes, protein kinase A-induced lipolysis releases fatty acids that are rapidly reesterified and sequestered as TAG in numerous small LDs (Chitraju et al. 2017). Inhibition of DGAT1 (but not DGAT2) during periods of lipolysis impairs LD biogenesis and is associated with increases in UPR markers (Chitraju et al. 2017). The role of DGAT1 in preventing ER stress is not limited to adipocytes, as inhibition of DGAT1 during oleate treatment in preadipocytes is accompanied by ER stress (D. Xu et al. 2018).

Disruptions in LD regulation that do not directly impact neutral lipid synthesis can also cause ER stress. For example, the reduction of ER–LD contacts by the depletion of Rab18 or NAG, a component of an ER–LD tethering complex, induces the upregulation of ER stress markers in preadipocytes treated with oleate (D. Xu et al. 2018). In addition, the loss of FIT2, an ER-resident protein that supports LD biogenesis, is associated with ER stress (Becuwe et al. 2018). However, it is unclear if the effect of FIT2 on ER stress is due to the regulation of LD biogenesis or ER morphology and membrane lipid composition through its putative activity as a lipid phosphatase (Becuwe et al. 2018, Hayes et al. 2017). Disruptions in LD biogenesis do not always lead to ER stress, and the importance of LDs in preventing ER stress is likely to be condition specific. For instance, the loss of DGAT1-dependent LD biogenesis during nutrient stress-induced autophagy has no effect on ER stress and instead causes mitochondrial dysfunction (Nguyen et al. 2017). It remains to be determined if the loss of LD biogenesis leads to UPR activation due to alterations in ER protein-folding capacity or in ER membrane lipid composition. The simplest explanation is that the inability to form LDs results in the accumulation of free fatty acids, which can then act as membrane detergents or be channeled into toxic lipid species in the ER. LD biogenesis may also be important to remove specific lipids from the ER to reestablish membrane homeostasis during periods of membrane stress.

6.2. Lipid Droplets as Mediators of Endoplasmic Reticulum Protein Degradation

LDs may mediate ER protein degradation through both proteasomal and autophagic pathways. One model suggests that LDs may temporarily sequester retrotranslocated ERAD substrates en route to the proteasome. Studies in yeast and mammalian cells have demonstrated that LD biogenesis is not generally required for ERAD (Nakatsukasa & Kamura 2016, Olzmann & Kopito 2011, To et al. 2017). Nonetheless, LDs have been reported to transiently associate with two different ERAD substrate proteins prior to their proteasomal degradation: apolipoprotein B-100 (Apo B-100) and HMG–CoA reductase. Polyubiquitinated Apo B-100 accumulates on the surface of LDs or in close proximity to LDs upon proteasome inhibition (Ohsaki et al. 2006, 2008; Suzuki et al. 2012). The close association of ubiquitinated Apo B-100 with LDs suggests that degradation-bound proteins may interact with LDs en route to the proteasome, potentially as a strategy to prevent their aggregation. However, it remains unclear whether LDs represent an on-pathway intermediate in Apo B-100 degradation or an off-pathway detour that is only evident under conditions of impaired proteasomal clearance. HMG–CoA reductase, the rate-limiting enzyme in cholesterol synthesis, localizes to LD-associated ER subdomains prior to its retrotranslocation to the cytosol (Hartman et al. 2010, Jo et al. 2013). Given that LDs are not generally required for the degradation of many ERAD substrates (Nakatsukasa & Kamura 2016, Olzmann & Kopito 2011, To et al. 2017), only a subset of ERAD substrates may associate with LDs or be degraded from LD-associated ER subdomains. Apo B-100 and HMG–CoA reductase are important regulators of lipid metabolism, and their association with LDs and/or degradation from LD-associated ER subdomains could be related to their cellular functions.

In yeast, LDs may facilitate ER protein degradation by mediating their delivery to the vacuole via microlipophagy (Vevea et al. 2015). In microlipophagy, LDs are delivered

directly to the vacuole at sterol-rich subdomains in an autophagosome-independent manner (Seo et al. 2017, van Zutphen et al. 2014, Wang et al. 2014). Yeast strains deficient in the synthesis of PC, the predominant membrane PL, exhibit alterations in their lipid profiles and upregulate LD biogenesis (Vevea et al. 2015). This increase in LDs likely reflects the channeling of excess lipids into TAG due to the inability to convert DAG into PC. The LDs formed under these conditions are degraded through microlipophagy (Vevea et al. 2015). Interestingly, LD fractions were enriched in both polyubiquitinated proteins and the ER chaperone Kar2p, leading Vevéa and colleagues (2015) to propose a role for LDs in physically removing unfolded proteins from the ER and facilitating their subsequent degradation by microlipophagy. However, the enrichment in polyubiquitinated proteins detected in the LD fraction may represent proteins found in regions of extensive LD-ER contacts, similar to the structures that harbor proteasome-bound HMG-CoA reductase (Hartman et al. 2010, Jo et al. 2013). Furthermore, the identities of substrate proteins that use this putative pathway for their degradation have yet to be identified.

6.3. Lipid Droplets and Toxic Protein Aggregates

Emerging findings implicate dysregulated lipid metabolism as a key factor in the etiology of Parkinson's disease (PD) and suggest a possible role for LDs in suppressing PD-associated lipotoxicity (Fanning et al. 2019, 2020; Imberdis et al. 2019; Soste et al. 2019; Vincent et al. 2018). α -Synuclein is a small, brain-enriched protein that is implicated in sporadic and familial forms of PD and is the primary aggregated protein within the PD-associated inclusion bodies known as Lewy bodies (Shahmoradian et al. 2019). α -Synuclein binds PLs and oleate (Sharon et al. 2001, Stöckl et al. 2008), and unsaturated fatty acids promote α -synuclein membrane binding, aggregation, and cytotoxicity (Fanning et al. 2019; Jo et al. 2002; Sharon et al. 2001, 2003). The expression of α -synuclein leads to alterations in the cellular lipid landscape, including increases in oleate, DAG, and TAG, as well as an increase in LD biogenesis (Fanning et al. 2019, Outeiro & Lindquist 2003, Vincent et al. 2018). Yeast strains deficient in LD biogenesis are more sensitive to α -synuclein-induced toxicity (Fanning et al. 2019), indicating a protective role for LDs under these conditions. LDs may act to sequester oleate and DAG as TAG, thereby mitigating a portion of α -synuclein toxicity and inhibiting its effects on membrane trafficking. SCD1 (Ole1 in yeast) inhibitors, which reduce the production of monounsaturated fatty acids such as oleate, strongly suppress α -synuclein toxicity in both yeast and cultured human induced pluripotent stem cell-derived neurons (Fanning et al. 2019, Imberdis et al. 2019, Vincent et al. 2018). The mechanism by which α -synuclein expression causes changes in lipid abundance is unknown, as is the mechanism by which oleate promotes α -synuclein toxicity. Neurons have a limited capacity to generate LDs, and recent findings demonstrate that these cells actively traffic damaged lipids via apolipoprotein E particles to astrocytes, which sequester these lipids in LDs prior to degradation (Ioannou et al. 2019, Liu et al. 2017a). Determining whether this neuronal lipid clearance pathway impacts α -synuclein toxicity and aggregation by reducing neuronal oleate concentrations will be important for future therapeutic research. LDs may also directly promote the clearance of inclusion bodies containing aggregated proteins in yeast, where LDs associate with inclusion bodies and increase their turnover via the release of an SE-derived detergent-like metabolite (Moldavski et al. 2015). Whether this is a general role for LDs remains unexplored.

7. PERSPECTIVES AND OPEN QUESTIONS

Numerous proteomic analyses of LDs have established a catalog of LD proteins in a variety of species, cell and tissue types, and metabolic conditions. The LD proteome composition reflects its role as a central player in cellular lipid and energy metabolism. Determining the composition and functions of heterogeneous LD subpopulations, possibly by leveraging proximity labeling methods, will be an important next step. While defining the LD proteome is crucial, we are still at the beginning of the process of understanding the mechanisms that regulate the LD proteome composition. Recent findings have established two classes of LD proteins defined by their initial insertion into the ER or LDs. The mechanisms that mediate their selective insertion into the ER or LDs and control their partitioning between the ER and LDs remain important areas of investigation. Similarly, while emerging data indicate that LD proteins can be degraded by CMA and the UPS, little is known about the degradation signals presented by substrates, the ubiquitination factors that recognize substrates and target them for degradation, and the mechanisms that regulate degradation under different metabolic states. Furthermore, while LDs clearly function in regulating cellular lipid homeostasis and responses to lipotoxicity, whether they play direct roles as mediators of protein quality control remains unresolved. Given the integral role of LD proteins in regulating all aspects of LD biology, defining the mechanisms that establish and remodel LD proteome composition is essential to understanding the cellular functions of LDs and their connections to human health and disease.

ACKNOWLEDGMENTS

We apologize to any authors whose important contributions were not cited in this review due to space limitations. We thank Alyssa Mathiowetz and Stefani Roberts for critical reading of this manuscript. Our work is supported by grants from the National Institutes of Health to J.A.O. (R01GM112948) and to M.A.R. (F31DK121477) and an American Cancer Society Research Scholar Grant award to J.A.O. (RSG-19-192-01-CCG). J.A.O. is a Chan Zuckerberg Biohub investigator.

LITERATURE CITED

- Abell BM, Holbrook LA, Abenes M, Murphy DJ, Hills MJ, Moloney MM. 1997 Role of the proline knot motif in oleosin endoplasmic reticulum topology and oil body targeting. *Plant Cell* 9(8):1481–93 [PubMed: 9286116]
- Ajjaji D, Ben M'barek K, Mimmack ML, England C, Herscovitz H, et al. 2019 Dual binding motifs underpin the hierarchical association of perilipins 1–3 with lipid droplets. *Mol. Biol. Cell* 30(5):703–16 [PubMed: 30649995]
- Ariyama H, Kono N, Matsuda S, Inoue T, Arai H. 2010 Decrease in membrane phospholipid unsaturation induces unfolded protein response. *J. Biol. Chem* 285(29):22027–35 [PubMed: 20489212]
- Bacle A, Gautier R, Jackson CL, Fuchs PFJ, Vanni S. 2017 Interdigitation between triglycerides and lipids modulates surface properties of lipid droplets. *Biophys. J* 112(7):1417–30 [PubMed: 28402884]
- Bartz R, Li W-H, Venables B, Zehmer JK, Roth MR, et al. 2007a Lipidomics reveals that adiposomes store ether lipids and mediate phospholipid traffic. *J. Lipid Res* 48(4):837–47 [PubMed: 17210984]
- Bartz R, Zehmer JK, Zhu M, Chen Y, Serrero G, et al. 2007b Dynamic activity of lipid droplets: protein phosphorylation and GTP-mediated protein translocation. *J. Proteome Res* 6(8):3256–65 [PubMed: 17608402]

- BasuRay S, Smagris E, Cohen JC, Hobbs HH. 2017 The PNPLA3 variant associated with fatty liver disease (I148M) accumulates on lipid droplets by evading ubiquitylation. *Hepatology* 66(4):1111–24 [PubMed: 28520213]
- BasuRay S, Wang Y, Smagris E, Cohen JC, Hobbs HH. 2019 Accumulation of PNPLA3 on lipid droplets is the basis of associated hepatic steatosis. *PNAS* 116(19):9521–26 [PubMed: 31019090]
- Becuwe M, Bond LM, Mejhert N, Boland S, Elliott SD, et al. 2018 FIT2 is a lipid phosphate phosphatase crucial for endoplasmic reticulum homeostasis. *bioRxiv* 291765 10.1101/291765
- Ben M'barek K, Ajjaji D, Chorlay A, Vanni S, Forêt L, Thiam AR. 2017 ER membrane phospholipids and surface tension control cellular lipid droplet formation. *Dev. Cell* 41(6):591–604.e7 [PubMed: 28579322]
- Bersuker K, Hendricks JM, Li Z, Magtanong L, Ford B, et al. 2019 The CoQ oxidoreductase FSP1 acts parallel to GPX4 to inhibit ferroptosis. *Nature* 575(7784):688–92 [PubMed: 31634900]
- Bersuker K, Olzmann JA. 2017 Establishing the lipid droplet proteome: mechanisms of lipid droplet protein targeting and degradation. *Biochim. Biophys. Acta Mol. Cell Biol. Lipids* 1862(10 Part B):1166–77 [PubMed: 28627435]
- Bersuker K, Olzmann JA. 2019 Identification of lipid droplet proteomes by proximity labeling proteomics using APEX2. *Methods Mol. Biol* 2008:57–72 [PubMed: 31124088]
- Bersuker K, Peterson CWH, To M, Sahl SJ, Savikhin V, et al. 2018 A proximity labeling strategy provides insights into the composition and dynamics of lipid droplet proteomes. *Dev. Cell* 44(1):97–112.e7 [PubMed: 29275994]
- Boeszoermenyi A, Nagy HM, Arthanari H, Phillip CJ, Lindermuth H, et al. 2015 Structure of a CGI-58 motif provides the molecular basis of lipid droplet anchoring. *J. Biol. Chem* 290(44):26361–72 [PubMed: 26350461]
- Boström P, Andersson L, Rutberg M, Perman J, Lidberg U, et al. 2007 SNARE proteins mediate fusion between cytosolic lipid droplets and are implicated in insulin sensitivity. *Nat. Cell Biol* 9(11):1286–93 [PubMed: 17922004]
- Brandt C, McFie PJ, Stone SJ. 2016 Diacylglycerol acyltransferase-2 and monoacylglycerol acyltransferase-2 are ubiquitinated proteins that are degraded by the 26S proteasome. *Biochem. J* 473(20):3621–37 [PubMed: 27531967]
- Brasaemle DL, Dolios G, Shapiro L, Wang R. 2004 Proteomic analysis of proteins associated with lipid droplets of basal and lipolytically stimulated 3T3-L1 adipocytes. *J. Biol. Chem* 279(45):46835–42 [PubMed: 15337753]
- Cermelli S, Guo Y, Gross SP, Welte MA. 2006 The lipid-droplet proteome reveals that droplets are a protein-storage depot. *Curr. Biol* 16(18):1783–95 [PubMed: 16979555]
- Chan SC, Lin S-C, Li P. 2007 Regulation of Cidea protein stability by the ubiquitin-mediated proteasomal degradation pathway. *Biochem. J* 408(2):259–66 [PubMed: 17711404]
- Chang C-L, Weigel AV, Ioannou MS, Pasolli HA, Xu CS, et al. 2019 Spastin tethers lipid droplets to peroxisomes and directs fatty acid trafficking through ESCRT-III. *J. Cell Biol* 218(8):2583–99 [PubMed: 31227594]
- Chen Y-C, Umanah GKE, Dephoure N, Andrabi SA, Gygi SP, et al. 2014 Msp1/ATAD1 maintains mitochondrial function by facilitating the degradation of mislocalized tail-anchored proteins. *EMBO J.* 33(14):1548–64 [PubMed: 24843043]
- Cheng J, Fujita A, Ohsaki Y, Suzuki M, Shinohara Y, Fujimoto T. 2009 Quantitative electron microscopy shows uniform incorporation of triglycerides into existing lipid droplets. *Histochem. Cell Biol* 132(3):281–91 [PubMed: 19557427]
- Chino H, Mizushima N. 2020 ER-phagy: quality control and turnover of endoplasmic reticulum. *Trends Cell Biol.* 30(5):384–98 [PubMed: 32302550]
- Chitraju C, Mejhert N, Haas JT, Diaz-Ramirez LG, Grueter CA, et al. 2017 Triglyceride synthesis by DGAT1 protects adipocytes from lipid-induced ER stress during lipolysis. *Cell Metab.* 26(2):407–18.e3 [PubMed: 28768178]
- Choi K, Kim H, Kang H, Lee S-Y, Lee SJ, et al. 2014 Regulation of diacylglycerol acyltransferase 2 protein stability by gp78-associated endoplasmic-reticulum-associated degradation. *FEBS J.* 281(13):3048–60 [PubMed: 24820123]

- Chorlay A, Monticelli L, Ferreira JV, Ben M'barek K, Ajjaji D, et al. 2019 Membrane asymmetry imposes directionality on lipid droplet emergence from the ER. *Dev. Cell* 50(1):25–42.e7 [PubMed: 31155466]
- Chorlay A, Thiam AR. 2020 Neutral lipids regulate amphipathic helix affinity for model lipid droplets. *J. Cell Biol* 219(4):e201907099 [PubMed: 32328636]
- opi A, Antoine-Bally S, Giménez-Andrés M, La Torre Garay C, Antonny B, et al. 2018 A giant amphipathic helix from a perilipin that is adapted for coating lipid droplets. *Nat. Commun* 9(1):1332 [PubMed: 29626194]
- Crunk AE, Monks J, Murakami A, Jackman M, Maclean PS, et al. 2013 Dynamic regulation of hepatic lipid droplet properties by diet. *PLOS ONE* 8(7):e67631 [PubMed: 23874434]
- Cunha DA, Hekerman P, Ladrière L, Bazarra-Castro A, Ortis F, et al. 2008 Initiation and execution of lipotoxic ER stress in pancreatic β -cells. *J. Cell Sci* 121(Part 14):2308–18 [PubMed: 18559892]
- Currie E, Guo X, Christiano R, Chitraju C, Kory N, et al. 2014 High confidence proteomic analysis of yeast LDs identifies additional droplet proteins and reveals connections to dolichol synthesis and sterol acetylation. *J. Lipid Res* 55(7):1465–77 [PubMed: 24868093]
- Czabany T, Wagner A, Zweytick D, Lohner K, Leitner E, et al. 2008 Structural and biochemical properties of lipid particles from the yeast *Saccharomyces cerevisiae*. *J. Biol. Chem* 283(25):17065–74 [PubMed: 18430725]
- Dai Z, Qi W, Li C, Lu J, Mao Y, et al. 2013 Dual regulation of adipose triglyceride lipase by pigment epithelium-derived factor: a novel mechanistic insight into progressive obesity. *Mol. Cell. Endocrinol* 377(1–2):123–34 [PubMed: 23850519]
- D'Aquila T, Zembroski AS, Buhman KK. 2019 Diet induced obesity alters intestinal cytoplasmic lipid droplet morphology and proteome in the postprandial response to dietary fat. *Front. Physiol* 10:180 [PubMed: 30890954]
- Deruyffelaere C, Purkrtova Z, Bouchez I, Collet B, Cacas J-L, et al. 2018 PUX10 is a CDC48A adaptor protein that regulates the extraction of ubiquitinated oleosins from seed lipid droplets in *Arabidopsis*. *Plant Cell* 30(9):2116–36 [PubMed: 30087208]
- Ding Y, Wu Y, Zeng R, Liao K. 2012 Proteomic profiling of lipid droplet-associated proteins in primary adipocytes of normal and obese mouse. *Acta Biochim. Biophys. Sin* 44(5):394–406 [PubMed: 22343379]
- Doll S, Freitas FP, Shah R, Aldrovandi M, Silva MCD, et al. 2019 FSP1 is a glutathione-independent ferroptosis suppressor. *Nature* 575(7784):693–98 [PubMed: 31634899]
- Du X, Zhou L, Aw YC, Mak HY, Xu Y, et al. 2020 ORP5 localizes to ER-lipid droplet contacts and regulates the level of PI(4)P on lipid droplets. *J. Cell Biol* 219(1):e201905162 [PubMed: 31653673]
- Dubey R, Stivala CE, Nguyen HQ, Goo Y-H, Paul A, et al. 2020 Lipid droplets can promote drug accumulation and activation. *Nat. Chem. Biol* 16(2):206–13 [PubMed: 31932720]
- Eastman SW, Yassaee M, Bieniasz PD. 2009 A role for ubiquitin ligases and Spartin/SPG20 in lipid droplet turnover. *J. Cell Biol* 184(6):881–94 [PubMed: 19307600]
- Edwards TL, Clowes VE, Tsang HTH, Connell JW, Sanderson CM, et al. 2009 Endogenous spartin (SPG20) is recruited to endosomes and lipid droplets and interacts with the ubiquitin E3 ligases AIP4 and AIP5. *Biochem. J* 423(1):31–39 [PubMed: 19580544]
- Fanning S, Haque A, Imberdis T, Baru V, Barrasa MI, et al. 2019 Lipidomic analysis of α -synuclein neurotoxicity identifies stearoyl CoA desaturase as a target for Parkinson treatment. *Mol. Cell* 73(5):1001–14.e8 [PubMed: 30527540]
- Fanning S, Selkoe D, Dettmer U. 2020 Parkinson's disease: proteinopathy or lipidopathy? *npj Parkinson's Dis.* 6:3 [PubMed: 31909184]
- Fei W, Wang H, Fu X, Bielby C, Yang H. 2009 Conditions of endoplasmic reticulum stress stimulate lipid droplet formation in *Saccharomyces cerevisiae*. *Biochem. J* 424(1):61–67 [PubMed: 19708857]
- Foresti O, Rodriguez-Vaello V, Funaya C, Carvalho P. 2014 Quality control of inner nuclear membrane proteins by the Asi complex. *Science* 346(6210):751–55 [PubMed: 25236469]

- Fu S, Yang L, Li P, Hofmann O, Dicker L, et al. 2011 Aberrant lipid metabolism disrupts calcium homeostasis causing liver endoplasmic reticulum stress in obesity. *Nature* 473(7348):528–31 [PubMed: 21532591]
- Fujimoto T, Parton RG. 2011 Not just fat: the structure and function of the lipid droplet. *Cold Spring Harb. Perspect. Biol* 3(3):a004838 [PubMed: 21421923]
- Gallardo-Montejano VI, Saxena G, Kusminski CM, Yang C, McAfee JL, et al. 2016 Nuclear Perilipin 5 integrates lipid droplet lipolysis with PGC-1 α /SIRT1-dependent transcriptional regulation of mitochondrial function. *Nat. Commun* 7:12723 [PubMed: 27554864]
- Garbarino J, Padamsee M, Wilcox L, Oelkers PM, D'Ambrosio D, et al. 2009 Sterol and diacylglycerol acyltransferase deficiency triggers fatty acid-mediated cell death. *J. Biol. Chem* 284(45):30994–1005 [PubMed: 19690167]
- Ghosh M, Niyogi S, Bhattacharyya M, Adak M, Nayak DK, et al. 2016 Ubiquitin ligase COP1 controls hepatic fat metabolism by targeting ATGL for degradation. *Diabetes* 65(12):3561–72 [PubMed: 27658392]
- Graef M 2018 Lipid droplet-mediated lipid and protein homeostasis in budding yeast. *FEBS Lett.* 592(8):1291–303 [PubMed: 29397034]
- Greenberg AS, Egan JJ, Wek SA, Garty NB, Blanchette-Mackie EJ, Londos C. 1991 Perilipin, a major hormonally regulated adipocyte-specific phosphoprotein associated with the periphery of lipid storage droplets. *J. Biol. Chem* 266(17):11341–46 [PubMed: 2040638]
- Greenwood DJ, Dos Santos MS, Huang S, Russell MRG, Collinson LM, et al. 2019 Subcellular antibiotic visualization reveals a dynamic drug reservoir in infected macrophages. *Science* 364(6447):1279–82 [PubMed: 31249058]
- Gregor MF, Yang L, Fabbrini E, Mohammed BS, Eagon JC, et al. 2009 Endoplasmic reticulum stress is reduced in tissues of obese subjects after weight loss. *Diabetes* 58(3):693–700 [PubMed: 19066313]
- Hariri H, Rogers S, Ugrankar R, Liu YL, Feathers JR, Henne WM. 2018 Lipid droplet biogenesis is spatially coordinated at ER-vacuole contacts under nutritional stress. *EMBO Rep.* 19(1):57–72 [PubMed: 29146766]
- Hariri H, Speer N, Bowerman J, Rogers S, Fu G, et al. 2019 Mdm1 maintains endoplasmic reticulum homeostasis by spatially regulating lipid droplet biogenesis. *J. Cell Biol* 218(4):1319–34 [PubMed: 30808705]
- Hartman IZ, Liu P, Zehmer JK, Luby-Phelps K, Jo Y, et al. 2010 Sterol-induced dislocation of 3-hydroxy-3-methylglutaryl coenzyme A reductase from endoplasmic reticulum membranes into the cytosol through a subcellular compartment resembling lipid droplets. *J. Biol. Chem* 285(25):19288–98 [PubMed: 20406816]
- Hayes M, Choudhary V, Ojha N, Shin JHH, Han G-S, et al. 2017 Fat storage-inducing transmembrane (FIT or FITM) proteins are related to lipid phosphatase/phosphotransferase enzymes. *Microb. Cell* 5(2):88–103 [PubMed: 29417057]
- Heckmann BL, Zhang X, Saarinen AM, Liu J. 2016 Regulation of G0/G1 switch gene 2 (G0S2) protein ubiquitination and stability by triglyceride accumulation and ATGL interaction. *PLOS ONE* 11(6):e0156742 [PubMed: 27248498]
- Hooper C, Puttamadappa SS, Loring Z, Shekhtman A, Bakowska JC. 2010 Spartins activate atrophin-1-interacting protein 4 (AIP4) E3 ubiquitin ligase and promotes ubiquitination of adipophilin on lipid droplets. *BMC Biol.* 8:72 [PubMed: 20504295]
- Hope RG, Murphy DJ, McLauchlan J. 2002 The domains required to direct core proteins of hepatitis C virus and GB virus-B to lipid droplets share common features with plant oleosin proteins. *J. Biol. Chem* 277(6):4261–70 [PubMed: 11706032]
- Hynynen R, Suchanek M, Spandl J, Bäck N, Thiele C, Olkkonen VM. 2009 OSBP-related protein 2 is a sterol receptor on lipid droplets that regulates the metabolism of neutral lipids. *J. Lipid Res* 50(7):1305–15 [PubMed: 19224871]
- Imberdis T, Negri J, Ramalingam N, Terry-Kantor E, Ho GPH, et al. 2019 Cell models of lipid-rich α -synuclein aggregation validate known modifiers of α -synuclein biology and identify stearyl-CoA desaturase. *PNAS* 116(41):20760–69 [PubMed: 31548371]

- Ingelmo-Torres M, González-Moreno E, Kassan A, Hanzal-Bayer M, Tebar F, et al. 2009 Hydrophobic and basic domains target proteins to lipid droplets. *Traffic* 10(12):1785–801 [PubMed: 19874557]
- Ioannou MS, Jackson J, Sheu S-H, Chang C-L, Weigel AV, et al. 2019 Neuron-astrocyte metabolic coupling protects against activity-induced fatty acid toxicity. *Cell* 177(6):1522–35.e14 [PubMed: 31130380]
- Jo E, Fuller N, Rand RP, St George-Hyslop P, Fraser PE. 2002 Defective membrane interactions of familial Parkinson's disease mutant A30P α -synuclein. *J. Mol. Biol.* 315(4):799–807 [PubMed: 11812148]
- Jo Y, Hartman IZ, DeBose-Boyd RA. 2013 Ancient ubiquitous protein-1 mediates sterol-induced ubiquitination of 3-hydroxy-3-methylglutaryl CoA reductase in lipid droplet-associated endoplasmic reticulum membranes. *Mol. Biol. Cell* 24(3):169–83 [PubMed: 23223569]
- Jonikas MC, Collins SR, Denic V, Oh E, Quan EM, et al. 2009 Comprehensive characterization of genes required for protein folding in the endoplasmic reticulum. *Science* 323(5922):1693–97 [PubMed: 19325107]
- Kamikubo K, Kato H, Kioka H, Yamazaki S, Tsukamoto O, et al. 2019 A molecular triage process mediated by RING finger protein 126 and BCL2-associated athanogene 6 regulates degradation of G0/G1 switch gene 2. *J. Biol. Chem* 294(40):14562–73 [PubMed: 31371451]
- Kaushik S, Cuervo AM. 2015 Degradation of lipid droplet-associated proteins by chaperone-mediated autophagy facilitates lipolysis. *Nat. Cell Biol* 17(6):759–70 [PubMed: 25961502]
- Kaushik S, Cuervo AM. 2016 AMPK-dependent phosphorylation of lipid droplet protein PLIN2 triggers its degradation by CMA. *Autophagy* 12(2):432–38 [PubMed: 26902588]
- Kaushik S, Cuervo AM. 2018 The coming of age of chaperone-mediated autophagy. *Nat. Rev. Mol. Cell Biol* 19(6):365–81 [PubMed: 29626215]
- Khan SA, Wollaston-Hayden EE, Markowski TW, Higgins L, Mashek DG. 2015 Quantitative analysis of the murine lipid droplet-associated proteome during diet-induced hepatic steatosis. *J. Lipid Res* 56(12):2260–72 [PubMed: 26416795]
- Khandelia H, Duelund L, Pakkanen KI, Ipsen JH. 2010 Triglyceride blisters in lipid bilayers: implications for lipid droplet biogenesis and the mobile lipid signal in cancer cell membranes. *PLOS ONE* 5(9):e12811 [PubMed: 20877640]
- Khmelniskii A, Blaszcak E, Pantazopoulou M, Fischer B, Omnus DJ, et al. 2014 Protein quality control at the inner nuclear membrane. *Nature* 516(7531):410–13 [PubMed: 25519137]
- Kimmel AR, Sztalryd C. 2016 The perilipins: major cytosolic lipid droplet-associated proteins and their roles in cellular lipid storage, mobilization, and systemic homeostasis. *Annu. Rev. Nutr* 36:471–509 [PubMed: 27431369]
- Kitamura T, Takagi S, Naganuma T, Kihara A. 2015 Mouse aldehyde dehydrogenase ALDH3B2 is localized to lipid droplets via two C-terminal tryptophan residues and lipid modification. *Biochem. J* 465(1):79–87 [PubMed: 25286108]
- Klemm EJ, Spooner E, Ploegh HL. 2011 Dual role of ancient ubiquitous protein 1 (AUP1) in lipid droplet accumulation and endoplasmic reticulum (ER) protein quality control. *J. Biol. Chem* 286(43):37602–14 [PubMed: 21857022]
- Kory N, Farese RV Jr., Walther TC. 2016 Targeting fat: mechanisms of protein localization to lipid droplets. *Trends Cell Biol.* 26(7):535–46 [PubMed: 26995697]
- Kory N, Thiam A-R, Farese RV Jr., Walther TC. 2015 Protein crowding is a determinant of lipid droplet protein composition. *Dev. Cell* 34(3):351–63 [PubMed: 26212136]
- Krahmer N, Farese RV Jr., Walther TC. 2013a Balancing the fat: lipid droplets and human disease. *EMBO Mol. Med* 5(7):973–83 [PubMed: 23740690]
- Krahmer N, Guo Y, Wilfling F, Hilger M, Lingrell S, et al. 2011 Phosphatidylcholine synthesis for lipid droplet expansion is mediated by localized activation of CTP:phosphocholine cytidyltransferase. *Cell Metab.* 14(4):504–15 [PubMed: 21982710]
- Krahmer N, Hilger M, Kory N, Wilfling F, Stoehr G, et al. 2013b Protein correlation profiles identify lipid droplet proteins with high confidence. *Mol. Cell Proteom* 12(5):1115–26
- Krahmer N, Najafi B, Schueder F, Quagliarini F, Steger M, et al. 2018 Organellar proteomics and phosphoproteomics reveal subcellular reorganization in diet-induced hepatic steatosis. *Dev. Cell* 47(2):205–21.e7 [PubMed: 30352176]

- Kretschmar FK, Mengel LA, Müller AO, Schmitt K, Blersch KF, et al. 2018 PUX10 is a lipid droplet-localized scaffold protein that interacts with CELL DIVISION CYCLE48 and is involved in the degradation of lipid droplet proteins. *Plant Cell* 30(9):2137–60 [PubMed: 30087207]
- Kumar N, Leonzino M, Hancock-Cerutti W, Horenkamp FA, Li P, et al. 2018 VPS13A and VPS13C are lipid transport proteins differentially localized at ER contact sites. *J. Cell Biol* 217(10):3625–39 [PubMed: 30093493]
- Lam T, Harmancey R, Vasquez H, Gilbert B, Patel N, et al. 2016 Reversal of intramyocellular lipid accumulation by lipophagy and a p62-mediated pathway. *Cell Death Discov.* 2:16061 [PubMed: 27625792]
- Lee J-S, Mendez R, Heng HH, Yang Z-Q, Zhang K. 2012 Pharmacological ER stress promotes hepatic lipogenesis and lipid droplet formation. *Am. J. Transl. Res* 4(1):102–13 [PubMed: 22347525]
- Li D, Zhao YG, Li D, Zhao H, Huang J, et al. 2019 The ER-localized protein DFCP1 modulates ER-lipid droplet contact formation. *Cell Rep.* 27(2):343–58.e5 [PubMed: 30970241]
- Li Z, Schulze RJ, Weller SG, Krueger EW, Schott MB, et al. 2016 A novel Rab10-EHBP1-EHD2 complex essential for the autophagic engulfment of lipid droplets. *Sci. Adv* 2(12):e1601470 [PubMed: 28028537]
- Li Z, Weller SG, Drizyte-Miller K, Chen J, Krueger EW, et al. 2019 Maturation of lipophagic organelles in hepatocytes is dependent upon a Rab10/dynamin-2 complex. *Hepatology* 72(2):486–502
- Liu L, MacKenzie KR, Putluri N, Maleti -Savati M, Bellen HJ. 2017a The glia-neuron lactate shuttle and elevated ROS promote lipid synthesis in neurons and lipid droplet accumulation in glia via APOE/D. *Cell Metab.* 26(5):719–37.e6 [PubMed: 28965825]
- Liu M, Ge R, Liu W, Liu Q, Xia X, et al. 2017b Differential proteomics profiling identifies LDPs and biological functions in high-fat diet-induced fatty livers. *J. Lipid Res* 58(4):681–94 [PubMed: 28179399]
- Liu P, Bartz R, Zehmer JK, Ying Y, Zhu M, et al. 2007 Rab-regulated interaction of early endosomes with lipid droplets. *Biochim. Biophys. Acta* 1773(6):784–93 [PubMed: 17395284]
- Liu P, Ying Y, Zhao Y, Mundy DI, Zhu M, Anderson RGW. 2004 Chinese hamster ovary K2 cell lipid droplets appear to be metabolic organelles involved in membrane traffic. *J. Biol. Chem* 279(5):3787–92 [PubMed: 14597625]
- Liu W, Morito D, Takashima S, Mineharu Y, Kobayashi H, et al. 2011 Identification of RNF213 as a susceptibility gene for moyamoya disease and its possible role in vascular development. *PLOS ONE* 6(7):e22542 [PubMed: 21799892]
- Luo H, Jiang M, Lian G, Liu Q, Shi M, et al. 2018 AIDA selectively mediates downregulation of fat synthesis enzymes by ERAD to retard intestinal fat absorption and prevent obesity. *Cell Metab.* 27(4):843–53.e6 [PubMed: 29617643]
- Mahamid J, Tegunov D, Maiser A, Arnold J, Leonhardt H, et al. 2019 Liquid-crystalline phase transitions in lipid droplets are related to cellular states and specific organelle association. *PNAS* 116(34):16866–71 [PubMed: 31375636]
- Masuda Y, Itabe H, Odaki M, Hama K, Fujimoto Y, et al. 2006 ADRP/adipophilin is degraded through the proteasome-dependent pathway during regression of lipid-storing cells. *J. Lipid Res* 47(1):87–98 [PubMed: 16230742]
- McFie PJ, Banman SL, Stone SJ. 2018 Diacylglycerol acyltransferase-2 contains a c-terminal sequence that interacts with lipid droplets. *Biochim. Biophys. Acta Mol. Cell Biol. Lipids* 1863(9):1068–81 [PubMed: 29902571]
- McFie PJ, Stone SL, Banman SL, Stone SJ. 2010 Topological orientation of acyl-CoA:diacylglycerol acyltransferase-1 (DGAT1) and identification of a putative active site histidine and the role of the n terminus in dimer/tetramer formation. *J. Biol. Chem* 285(48):37377–87 [PubMed: 20876538]
- Mejherth N, Kuruvilla L, Gabriel KR, Elliott SD, Guie M-A, et al. 2020 Partitioning of MLX-family transcription factors to lipid droplets regulates metabolic gene expression. *Mol. Cell* 77(6):1251–64.e9 [PubMed: 32023484]
- Minville-Walz M, Pierre A-S, Pichon L, Bellenger S, Fèvre C, et al. 2010 Inhibition of stearoyl-CoA desaturase 1 expression induces CHOP-dependent cell death in human cancer cells. *PLOS ONE* 5(12):e14363 [PubMed: 21179554]

- Miyanari Y, Atsuzawa K, Usuda N, Watashi K, Hishiki T, et al. 2007 The lipid droplet is an important organelle for hepatitis C virus production. *Nat. Cell Biol* 9(9):1089–97 [PubMed: 17721513]
- Mizushima N 2018 A brief history of autophagy from cell biology to physiology and disease. *Nat. Cell Biol* 20:521–27 [PubMed: 29686264]
- Moldavski O, Amen T, Levin-Zaidman S, Eisenstein M, Rogachev I, et al. 2015 Lipid droplets are essential for efficient clearance of cytosolic inclusion bodies. *Dev. Cell* 33(5):603–10 [PubMed: 26004510]
- Morito D, Nishikawa K, Hoseki J, Kitamura A, Kotani Y, et al. 2014 Moyamoya disease-associated protein mysterin/RNF213 is a novel AAA+ ATPase, which dynamically changes its oligomeric state. *Sci. Rep* 4:4442 [PubMed: 24658080]
- Nakatsukasa K, Kamura T. 2016 Subcellular fractionation analysis of the extraction of ubiquitinated polytopic membrane substrate during ER-associated degradation. *PLOS ONE* 11(2):e0148327 [PubMed: 26849222]
- Nguyen HP, Yi D, Lin F, Viscarra JA, Tabuchi C, et al. 2020 Aifm2, a NADH oxidase, supports robust glycolysis and is required for cold- and diet-induced thermogenesis. *Mol. Cell* 77(3):600–17.e4 [PubMed: 31952989]
- Nguyen KT, Lee C-S, Mun S-H, Truong NT, Park SK, Hwang C-S. 2019 N-terminal acetylation and the N-end rule pathway control degradation of the lipid droplet protein PLIN2. *J. Biol. Chem* 294(1):379–88 [PubMed: 30425097]
- Nguyen TB, Louie SM, Daniele JR, Tran Q, Dillin A, et al. 2017 DGAT1-dependent lipid droplet biogenesis protects mitochondrial function during starvation-induced autophagy. *Dev. Cell* 42(1):9–21.e5 [PubMed: 28697336]
- Nian Z, Sun Z, Yu L, Toh SY, Sang J, Li P. 2010 Fat-specific protein 27 undergoes ubiquitin-dependent degradation regulated by triacylglycerol synthesis and lipid droplet formation. *J. Biol. Chem* 285(13):9604–15 [PubMed: 20089860]
- Niyogi S, Ghosh M, Adak M, Chakrabarti P. 2019 PEDF promotes nuclear degradation of ATGL through COPI1. *Biochem. Biophys. Res. Commun* 512(4):806–11 [PubMed: 30926171]
- Ohsaki Y, Cheng J, Fujita A, Tokumoto T, Fujimoto T. 2006 Cytoplasmic lipid droplets are sites of convergence of proteasomal and autophagic degradation of apolipoprotein B. *Mol. Biol. Cell* 17(6):2674–83 [PubMed: 16597703]
- Ohsaki Y, Cheng J, Suzuki M, Fujita A, Fujimoto T. 2008 Lipid droplets are arrested in the ER membrane by tight binding of lipidated apolipoprotein B-100. *J. Cell Sci* 121(Part 14):2415–22 [PubMed: 18577578]
- Okreglak V, Walter P. 2014 The conserved AAA-ATPase Msp1 confers organelle specificity to tail-anchored proteins. *PNAS* 111(22):8019–24 [PubMed: 24821790]
- Oku M, Maeda Y, Kagohashi Y, Kondo T, Yamada M, et al. 2017 Evidence for ESCRT- and clathrin-dependent microautophagy. *J. Cell Biol* 216(10):3263–74 [PubMed: 28838958]
- Olzmann JA, Carvalho P. 2019 Dynamics and functions of lipid droplets. *Nat. Rev. Mol. Cell Biol* 20(3):137–55 [PubMed: 30523332]
- Olzmann JA, Kopito RR. 2011 Lipid droplet formation is dispensable for endoplasmic reticulum-associated degradation. *J. Biol. Chem* 286(32):27872–74 [PubMed: 21693705]
- Olzmann JA, Kopito RR, Christianson JC. 2013a The mammalian endoplasmic reticulum-associated degradation system. *Cold Spring Harb. Perspect. Biol* 5(9):a013185 [PubMed: 23232094]
- Olzmann JA, Richter CM, Kopito RR. 2013b Spatial regulation of UBXD8 and p97/VCP controls ATGL-mediated lipid droplet turnover. *PNAS* 110(4):1345–50 [PubMed: 23297223]
- Outeiro TF, Lindquist S. 2003 Yeast cells provide insight into α -synuclein biology and pathobiology. *Science* 302(5651):1772–75 [PubMed: 14657500]
- Patel S, Yang W, Kozusko K, Saudek V, Savage DB. 2014 Perilipins 2 and 3 lack a carboxy-terminal domain present in perilipin 1 involved in sequestering ABHD5 and suppressing basal lipolysis. *PNAS* 111(25):9163–68 [PubMed: 24927580]
- Petrasek J, Erhartova D, Levine B. 2019 Protective effect of SMAD-specific E3 ubiquitin protein ligase 1 in alcoholic steatohepatitis in mice. *Hepatol. Commun* 3(11):1450–58 [PubMed: 31701069]

- Petschnigg J, Wolinski H, Kolb D, Zellnig G, Kurat CF, et al. 2009 Good fat, essential cellular requirements for triacylglycerol synthesis to maintain membrane homeostasis in yeast. *J. Biol. Chem* 284(45):30981–93 [PubMed: 19608739]
- Pickles S, Vigié P, Youle RJ. 2018 Mitophagy and quality control mechanisms in mitochondrial maintenance. *Curr. Biol* 28(4):R170–85 [PubMed: 29462587]
- Pineau L, Colas J, Dupont S, Beney L, Fleurat-Lessard P, et al. 2009 Lipid-induced ER stress: synergistic effects of sterols and saturated fatty acids. *Traffic* 10(6):673–90 [PubMed: 19302420]
- Prévost C, Sharp ME, Kory N, Lin Q, Voth GA, et al. 2018 Mechanism and determinants of amphipathic helix-containing protein targeting to lipid droplets. *Dev. Cell* 44(1):73–86.e4 [PubMed: 29316443]
- Promlek T, Ishiwata-Kimata Y, Shido M, Sakuramoto M, Kohno K, Kimata Y. 2011 Membrane aberrancy and unfolded proteins activate the endoplasmic reticulum stress sensor Ire1 in different ways. *Mol. Biol. Cell* 22(18):3520–32 [PubMed: 21775630]
- Qi J, Gong J, Zhao T, Zhao J, Lam P, et al. 2008 Downregulation of AMP-activated protein kinase by Cidea-mediated ubiquitination and degradation in brown adipose tissue. *EMBO J.* 27(11):1537–48 [PubMed: 18480843]
- Qian H, Chen Y, Nian Z, Su L, Yu H, et al. 2017 HDAC6-mediated acetylation of lipid droplet-binding protein CIDEC regulates fat-induced lipid storage. *J. Clin. Invest* 127(4):1353–69 [PubMed: 28287402]
- Renne MF, Klug YA, Carvalho P. 2020 Lipid droplet biogenesis: a mystery “unmixing”? *Semin. Cell Dev. Biol* In press 10.1016/j.semcdb.2020.03.001
- Rowe ER, Mimmack ML, Barbosa AD, Haider A, Isaac I, et al. 2016 Conserved amphipathic helices mediate lipid droplet targeting of perilipins 1–3. *J. Biol. Chem* 291(13):6664–78 [PubMed: 26742848]
- Ruggiano A, Mora G, Buxó L, Carvalho P. 2016 Spatial control of lipid droplet proteins by the ERAD ubiquitin ligase Doa10. *EMBO J.* 35(15):1644–55 [PubMed: 27357570]
- Santinho A, Salo VT, Chorlay A, Li S, Zhou X, et al. 2020 Membrane curvature catalyzes lipid droplet assembly. *Curr. Biol* 30(13):2481–94.e6 [PubMed: 32442467]
- Sathyanarayan A, Mashek MT, Mashek DG. 2017 ATGL promotes autophagy/lipophagy via SIRT1 to control hepatic lipid droplet catabolism. *Cell Rep.* 19(1):1–9 [PubMed: 28380348]
- Schott MB, Weller SG, Schulze RJ, Krueger EW, Drizyte-Miller K, et al. 2019 Lipid droplet size directs lipolysis and lipophagy catabolism in hepatocytes. *J. Cell Biol* 218(10):3320–35 [PubMed: 31391210]
- Schroeder B, Schulze RJ, Weller SG, Sletten AC, Casey CA, McNiven MA. 2015 The small GTPase Rab7 as a central regulator of hepatocellular lipophagy. *Hepatology* 61(6):1896–907 [PubMed: 25565581]
- Schrul B, Kopito RR. 2016 Peroxin-dependent targeting of a lipid-droplet-destined membrane protein to ER subdomains. *Nat. Cell Biol* 18(7):740–51 [PubMed: 27295553]
- Schulze RJ, Sathyanarayan A, Mashek DG. 2017 Breaking fat: the regulation and mechanisms of lipophagy. *Biochim. Biophys. Acta Mol. Cell Biol. Lipids* 1862(10 Part B):1178–87 [PubMed: 28642194]
- Seo AY, Lau P-W, Feliciano D, Sengupta P, Le Gros MA, et al. 2017 AMPK and vacuole-associated Atg14p orchestrate μ -lipophagy for energy production and long-term survival under glucose starvation. *eLife* 6:e21690 [PubMed: 28394250]
- Shahmoradian SH, Lewis AJ, Genoud C, Hench J, Moors TE, et al. 2019 Lewy pathology in Parkinson’s disease consists of crowded organelles and lipid membranes. *Nat. Neurosci* 22(7):1099–109 [PubMed: 31235907]
- Shao S, Hegde RS. 2011 Membrane protein insertion at the endoplasmic reticulum. *Annu. Rev. Cell Dev. Biol* 27:25–56 [PubMed: 21801011]
- Sharon R, Bar-Joseph I, Frosch MP, Walsh DM, Hamilton JA, Selkoe DJ. 2003 The formation of highly soluble oligomers of α -synuclein is regulated by fatty acids and enhanced in Parkinson’s disease. *Neuron* 37(4):583–95 [PubMed: 12597857]

- Sharon R, Goldberg MS, Bar-Josef I, Betensky RA, Shen J, Selkoe DJ. 2001 α -Synuclein occurs in lipid-rich high molecular weight complexes, binds fatty acids, and shows homology to the fatty acid-binding proteins. *PNAS* 98(16):9110–15 [PubMed: 11481478]
- Singh R, Kaushik S, Wang Y, Xiang Y, Novak I, et al. 2009 Autophagy regulates lipid metabolism. *Nature* 458(7242):1131–35 [PubMed: 19339967]
- Soste M, Charmpi K, Lampert F, Gerez JA, van Oostrum M, et al. 2019 Proteomics-based monitoring of pathway activity reveals that blocking diacylglycerol biosynthesis rescues from alpha-synuclein toxicity. *Cell Syst.* 9(3):309–20.e8 [PubMed: 31521608]
- Spandl J, Lohmann D, Kuerschner L, Moessinger C, Thiele C. 2011 Ancient ubiquitous protein 1 (AUP1) localizes to lipid droplets and binds the E2 ubiquitin conjugase G2 (Ube2g2) via its G2 binding region. *J. Biol. Chem* 286(7):5599–606 [PubMed: 21127063]
- Stefanovic-Barrett S, Dickson AS, Burr SP, Williamson JC, Lobb IT, et al. 2018 MARCH6 and TRC8 facilitate the quality control of cytosolic and tail-anchored proteins. *EMBO Rep.* 19(5):e45603 [PubMed: 29519897]
- Stöckl M, Fischer P, Wanker E, Herrmann A. 2008 α -Synuclein selectively binds to anionic phospholipids embedded in liquid-disordered domains. *J. Mol. Biol* 375(5):1394–404 [PubMed: 18082181]
- Stone SJ, Levin MC, Farese RV Jr. 2006 Membrane topology and identification of key functional amino acid residues of murine acyl-CoA:diacylglycerol acyltransferase-2. *J. Biol. Chem* 281(52):40273–82 [PubMed: 17035227]
- Stone SJ, Levin MC, Zhou P, Han J, Walther TC, Farese RV Jr. 2009 The endoplasmic reticulum enzyme DGAT2 is found in mitochondria-associated membranes and has a mitochondrial targeting signal that promotes its association with mitochondria. *J. Biol. Chem* 284(8):5352–61 [PubMed: 19049983]
- Su W, Wang Y, Jia X, Wu W, Li L, et al. 2014 Comparative proteomic study reveals 17 β -HSD13 as a pathogenic protein in nonalcoholic fatty liver disease. *PNAS* 111(31):11437–42 [PubMed: 25028495]
- Sugihara M, Morito D, Ainuki S, Hirano Y, Ogino K, et al. 2019 The AAA+ ATPase/ubiquitin ligase mysterin stabilizes cytoplasmic lipid droplets. *J. Cell Biol* 218(3):949–60 [PubMed: 30705059]
- Suzuki M, Murakami T, Cheng J, Kano H, Fukata M, Fujimoto T. 2015 ELMOD2 is anchored to lipid droplets by palmitoylation and regulates adipocyte triglyceride lipase recruitment. *Mol. Biol. Cell* 26(12):2333–42 [PubMed: 25904333]
- Suzuki M, Otsuka T, Ohsaki Y, Cheng J, Taniguchi T, et al. 2012 Derlin-1 and UBXD8 are engaged in dislocation and degradation of lipidated ApoB-100 at lipid droplets. *Mol. Biol. Cell* 23(5):800–10 [PubMed: 22238364]
- Takahashi Y, Shinoda A, Kamada H, Shimizu M, Inoue J, Sato R. 2016 Perilipin2 plays a positive role in adipocytes during lipolysis by escaping proteasomal degradation. *Sci. Rep* 6:20975 [PubMed: 26876687]
- Tapia D, Jiménez T, Zamora C, Espinoza J, Rizzo R, et al. 2019 KDEL receptor regulates secretion by lysosome relocation- and autophagy-dependent modulation of lipid-droplet turnover. *Nat. Commun* 10(1):735 [PubMed: 30760704]
- Tauchi-Sato K, Ozeki S, Houjou T, Taguchi R, Fujimoto T. 2002 The surface of lipid droplets is a phospholipid monolayer with a unique fatty acid composition. *J. Biol. Chem* 277(46):44507–12 [PubMed: 12221100]
- Thibault G, Shui G, Kim W, McAlister GC, Ismail N, et al. 2012 The membrane stress response buffers lethal effects of lipid disequilibrium by reprogramming the protein homeostasis network. *Mol. Cell* 48(1):16–27 [PubMed: 23000174]
- To M, Peterson CWH, Roberts MA, Counihan JL, Wu TT, et al. 2017 Lipid disequilibrium disrupts ER proteostasis by impairing ERAD substrate glycan trimming and dislocation. *Mol. Biol. Cell* 28(2):270–84 [PubMed: 27881664]
- Tokunaga M, Shiheido H, Hayakawa I, Utsumi A, Takashima H, et al. 2013 Hereditary spastic paraplegia protein spartin is an FK506-binding protein identified by mRNA display. *Chem. Biol* 20(7):935–42 [PubMed: 23890011]

- Toulmay A, Prinz WA. 2013 Direct imaging reveals stable, micrometer-scale lipid domains that segregate proteins in live cells. *J. Cell Biol* 202(1):35–44 [PubMed: 23836928]
- Tsuji T, Fujimoto M, Tatematsu T, Cheng J, Orii M, et al. 2017 Niemann-Pick type C proteins promote microautophagy by expanding raft-like membrane domains in the yeast vacuole. *eLife* 6:e25960 [PubMed: 28590904]
- Valm AM, Cohen S, Legant WR, Melunis J, Hershberg U, et al. 2017 Applying systems-level spectral imaging and analysis to reveal the organelle interactome. *Nature* 546(7656):162–67 [PubMed: 28538724]
- Valverde DP, Yu S, Boggavarapu V, Kumar N, Lees JA, et al. 2019 ATG2 transports lipids to promote autophagosome biogenesis. *J. Cell Biol* 218(6):1787–98 [PubMed: 30952800]
- van Zutphen T, Todde V, de Boer R, Kreim M, Hofbauer HF, et al. 2014 Lipid droplet autophagy in the yeast *Saccharomyces cerevisiae*. *Mol. Biol. Cell* 25(2):290–301 [PubMed: 24258026]
- Varshavsky A 2019 N-degron and C-degron pathways of protein degradation. *PNAS* 116(2):358–66 [PubMed: 30622213]
- Vevea JD, Garcia EJ, Chan RB, Zhou B, Schultz M, et al. 2015 Role for lipid droplet biogenesis and microlipophagy in adaptation to lipid imbalance in yeast. *Dev. Cell* 35(5):584–99 [PubMed: 26651293]
- Vincent BM, Tardiff DF, Piotrowski JS, Aron R, Lucas MC, et al. 2018 Inhibiting stearyl-CoA desaturase ameliorates α -synuclein cytotoxicity. *Cell Rep.* 25(10):2742–54.e31 [PubMed: 30517862]
- Volmer R, van der Ploeg K, Ron D. 2013 Membrane lipid saturation activates endoplasmic reticulum unfolded protein response transducers through their transmembrane domains. *PNAS* 110(12):4628–33 [PubMed: 23487760]
- Walter P, Ron D. 2011 The unfolded protein response: from stress pathway to homeostatic regulation. *Science* 334(6059):1081–86 [PubMed: 22116877]
- Walther TC, Chung J, Farese RV Jr. 2017 Lipid droplet biogenesis. *Annu. Rev. Cell Dev. Biol* 33:491–510 [PubMed: 28793795]
- Wang C-W, Miao Y-H, Chang Y-S. 2014 A sterol-enriched vacuolar microdomain mediates stationary phase lipophagy in budding yeast. *J. Cell Biol* 206(3):357–66 [PubMed: 25070953]
- Wang L, Zhou J, Yan S, Lei G, Lee C-H, Yin X-M. 2017 Ethanol-triggered lipophagy requires SQSTM1 in AML12 hepatic cells. *Sci. Rep* 7(1):12307 [PubMed: 28951592]
- Wilfling F, Wang H, Haas JT, Kraemer N, Gould TJ, et al. 2013 Triacylglycerol synthesis enzymes mediate lipid droplet growth by relocalizing from the ER to lipid droplets. *Dev. Cell* 24(4):384–99 [PubMed: 23415954]
- Xie X, Yi Z, Sinha S, Madan M, Bowen BP, et al. 2016 Proteomics analyses of subcutaneous adipocytes reveal novel abnormalities in human insulin resistance. *Obesity* 24(7):1506–14 [PubMed: 27345962]
- Xu D, Li Y, Wu L, Li Y, Zhao D, et al. 2018 Rab18 promotes lipid droplet (LD) growth by tethering the ER to LDs through SNARE and NRZ interactions. *J. Cell Biol* 217(3):975–95 [PubMed: 29367353]
- Xu G, Sztalryd C, Londos C. 2006 Degradation of perilipin is mediated through ubiquitination-proteasome pathway. *Biochim. Biophys. Acta* 1761(1):83–90 [PubMed: 16448845]
- Xu G, Sztalryd C, Lu X, Tansey JT, Gan J, et al. 2005 Post-translational regulation of adipose differentiation-related protein by the ubiquitin/proteasome pathway. *J. Biol. Chem* 280(52):42841–47 [PubMed: 16115879]
- Xu S, Zhang X, Liu P. 2018 Lipid droplet proteins and metabolic diseases. *Biochim. Biophys. Acta Mol. Basis Dis* 1864(5 Part B):1968–83 [PubMed: 28739173]
- Yamamoto K, Takahara K, Oyadomari S, Okada T, Sato T, et al. 2010 Induction of liver steatosis and lipid droplet formation in ATF6 α -knockout mice burdened with pharmacological endoplasmic reticulum stress. *Mol. Biol. Cell* 21(17):2975–86 [PubMed: 20631254]
- Yan Y, Wang H, Wei C, Xiang Y, Liang X, et al. 2019 HDAC6 regulates lipid droplet turnover in response to nutrient deprivation via p62-mediated selective autophagy. *J. Genet. Genom* 46(4):221–29

- Yang X, Zhang X, Heckmann BL, Lu X, Liu J. 2011 Relative contribution of adipose triglyceride lipase and hormone-sensitive lipase to tumor necrosis factor- α (TNF- α)-induced lipolysis in adipocytes. *J. Biol. Chem* 286(47):40477–85 [PubMed: 21969372]
- Zechner R, Madeo F, Kratky D. 2017 Cytosolic lipolysis and lipophagy: two sides of the same coin. *Nat. Rev. Mol. Cell Biol* 18(11):671–84 [PubMed: 28852221]
- Zehmer JK, Bartz R, Liu P, Anderson RGW. 2008 Identification of a novel N-terminal hydrophobic sequence that targets proteins to lipid droplets. *J. Cell Sci* 121(11):1852–60 [PubMed: 18477614]
- Zhang C, Liu P. 2019 The new face of the lipid droplet: lipid droplet proteins. *Proteomics* 19(10):e1700223 [PubMed: 30216670]
- Zhang K, Wang S, Malhotra J, Hassler JR, Back SH, et al. 2011 The unfolded protein response transducer IRE1 α prevents ER stress-induced hepatic steatosis. *EMBO J.* 30(7):1357–75 [PubMed: 21407177]
- Zhang S, Wang Y, Cui L, Deng Y, Xu S, et al. 2016 Morphologically and functionally distinct lipid droplet subpopulations. *Sci. Rep* 6:29539 [PubMed: 27386790]
- Zhang X, Heckmann BL, Xie X, Saarinen AM, Liu J. 2014 Regulation of FSP27 protein stability by AMPK and HSC70. *Am. J. Physiol. Endocrinol. Metab* 307(11):E1047–56 [PubMed: 25315694]
- Zoni V, Khaddaj R, Campomanes P, Thiam AR, Schneiter R, Vanni S. 2020 Lipid droplet biogenesis is driven by liquid-liquid phase separation. *bioRxiv* 777466 10.1101/777466




Figure 1.

LD protein targeting. Class I LD proteins are inserted into the ER and are trafficked from the ER to nascent LDs. Class II LD proteins are inserted into LDs directly from the cytosol. These proteins include both amphipathic helix-containing proteins that recognize phospholipid packing defects in the LD monolayer and proteins that are modified by conjugated fatty acids. Abbreviations: ER, endoplasmic reticulum; LD, lipid droplet.




Figure 2. Mechanisms of LD protein degradation. LD proteins are present in multiple cellular compartments and have the potential to be degraded by ubiquitin-dependent pathways in the ER, cytosol, and LDs. Abbreviations: ERAD, endoplasmic reticulum-associated degradation; LD, lipid droplet; Ub, ubiquitin.