

AVERAGE LIFETIMES OF COLLECTIVE TRANSITIONS
IN THE SPIN 30-50 REGION

H. Hübel, U. Smilansky, R. M. Diamond,
F.S. Stephens, and B. Herskind

June 1978

RECEIVED
LAWRENCE
BERKELEY LABORATORY


OCT 6 1978

Prepared for the U.S. Department of Energy
under Contract W-7405-ENG-48

LIBRARY AND
DOCUMENTS SECTION

TWO-WEEK LOAN COPY

This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 6782 *



LBL-7750
c.2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

AVERAGE LIFETIMES OF COLLECTIVE TRANSITIONS
IN THE SPIN 30-50 REGION

H. Hübel,^{*} U. Smilansky,⁺ R. M. Diamond, and F. S. Stephens

Lawrence Berkeley Laboratory
University of California
Berkeley, California 94720

and

B. Herskind⁺⁺

The Niels Bohr Institute
University of Copenhagen, Denmark

ABSTRACT

A method is developed to determine average lifetimes of continuum gamma rays following heavy-ion compound-nucleus reactions. The resulting enhancement factors for E2 transitions depopulating states in the spin 30-50 region are of the same order of magnitude as for ground state rotational bands of deformed nuclei. This shows for the first time that these high-spin states decay through strongly collective bands.

^{*}Permanent address: Institut für Strahlen und Kernphysik, D-5300 Bonn, W.Germany. Supported by a Max Kade Foundation Fellowship.

⁺Permanent address: Department of Physics, The Weizmann Institute of Science, Rehovot, Israel

⁺⁺Supported in part by the Danish Natural Science Research Council.

The de-excitation γ -ray spectra of nuclei produced at very high angular momenta in heavy-ion compound-nucleus reactions are composed of three parts.¹ The intensity of the high-energy part of the spectra ($E_\gamma \gtrsim 1.5$ MeV for deformed nuclei in the heavy rare-earth region) falls exponentially with E_γ . In this region the angular correlation is roughly isotropic, suggesting about equal amounts of dipole and quadrupole transitions.^{2,3} Furthermore, the multiplicity associated with each transition is roughly independent of E_γ . This part of the spectrum is very likely produced by statistical transitions. The low-energy region of the spectrum ($\lesssim 0.7$ MeV) is dominated by discrete lines, the transitions between the well-separated and strongly populated low energy states. In this paper we will focus our attention on the intermediate range of γ -ray energies where the spectrum shows a broad structure (the "bump") with the following characteristics: i) The multiplicity of transitions at the high-energy edge of the bump is higher than the average multiplicity associated with either the statistical or discrete parts of the spectrum -- indicating that these transitions come from the highest spin states populated. ii) Angular distribution measurements show predominantly a stretched quadrupole component in these decays. These characteristics can be explained by a model which assumes that the quadrupole bump is composed of collective transitions within rotational bands parallel to the yrast line. The effective moments of inertia derived from the multiplicities and excitation energies of these transitions are close to the values for a rigid rotor. However, until now no measurements have shown that these transitions are indeed very highly collective. In the present experiment

an attempt is made to measure average lifetimes in the bump region by a Doppler shift method.

The experimental setup is similar to the one described in Ref. 3, with six 7.6×7.6 cm NaI detectors serving as a multiplicity filter. Two 7.6×7.6 cm NaI detectors were located at 0° and 90° with respect to the beam direction at a distance of 60 cm from the target to allow for neutron discrimination. Average lifetimes of the continuum γ -rays were deduced by comparing the Doppler shift for the spectra measured in the 0° detector with a thin self-supporting target and with a target on a thick backing.

The targets and compound nuclei studied using beams of 700 MeV ^{136}Xe from the LBL SuperHILAC are listed in Table I. In each case spectra were measured for a $\sim 1 \text{ mg/cm}^2$ self-supporting target and for a target of similar thickness on a 25 mg/cm^2 Au backing. In order to keep the conditions as equal as possible for the two spectra, a 25 mg/cm^2 Au foil was placed ~ 2 mm downstream from the thin foil targets. In addition, for the $^{27}\text{Al} + ^{136}\text{Xe}$ system, targets with Pb backings were used. In this case the stopping time is almost two times longer than in the Au stopper; this provides a check on the reliability of the method as discussed below.

As an example of our data, Fig. 1a shows the sum of the 2-6 fold³ spectra obtained in the 0° detector for the self-supporting ^{27}Al and the ^{27}Al on Au targets. Below ~ 600 keV the displacement of the two spectra results mainly from lifetimes of the discrete lines. In the region of 750 - 1500 keV, the lifetimes of the continuum γ -rays are responsible for the difference between the two spectra. At still higher γ -ray

energies, the lifetimes are much shorter than the characteristic stopping time, hence, no difference is observed between the backed and the unbacked targets. The different effects in these three regions are accentuated in the ratio of the two spectra which is displayed in Fig. 1b. The ratio for a different target, ^{28}Si , is shown in Fig. 1d.

Two methods were used to normalize the spectra obtained from the backed and the self-supporting targets: i) As was explained above, no difference is expected in the range of $E_\gamma \gtrsim 2$ MeV; therefore, the two spectra were matched in this energy region. ii) No Doppler shift is expected in the detector at 90° . Hence by taking the double ratio of spectra from the backed and from the self-supporting targets at 0° and 90° we were able to get a normalization which eliminates some of the possible systematic errors, at the expense of a larger statistical uncertainty.

The data were analyzed in the following way. First the pulse height spectra were unfolded⁴ to give the primary photon spectra. The spectrum for recoil into vacuum was then folded with the response function calculated for a Doppler-shifted distribution, and the calculated ratio of spectra for recoil into vacuum and for recoil into backing was compared to the experimental ratio. It should be noted that the theory provides only the Doppler response function. The result of convoluting the experimental spectra with the response function does not smooth the fluctuations in the original spectra as can be seen in Fig. 1c. The response function which generates the Doppler-broadened spectrum was obtained by following the population of the decaying levels during the slowing down of the emitting nuclei.

The population of levels in the continuum was calculated using a model in which the continuum is described by a series of rotational bands with an average moment of inertia \mathcal{I} . To each energy bin at a median energy E_γ a spin I is assigned using the relationship $E_\gamma = \frac{\hbar^2}{2\mathcal{I}}(4I-2)$ with the empirical^{2,3,5} value $\frac{2\mathcal{I}}{\hbar^2} = (150 \pm 20) \text{ MeV}^{-1}$. We assume that the level decays to the next member of the band by an E2 transition. The feeding of each level consists of two parts: the prompt feeding, and the cascading from upper members of the band. The prompt part is approximated by a σ_I -distribution of initial populations,² $\sigma_I \propto I / (1 + \exp((I - I_{\max})/2.5))$, where I_{\max} is the maximum angular momentum brought into the compound nucleus, determined for this case as $I_{\max} = 50$. For the cascading part, the transition lifetimes are determined according to the rotational model with a constant intrinsic quadrupole moment Q_0 [Refs. 6,7]

$$\tau = 8 \cdot 10^2 E_{(\text{keV})}^{-5} \frac{1}{\langle I_i 2 K 0 | I_i 2 I_f K \rangle^2 e^2 Q_0^2(\text{b})} \text{ sec.}$$

The population equations are solved for all members of the band which contribute to the feeding. To solve this system of 10-30 coupled linear equations, we derived a fast algorithm which made the analysis of the experiment tractable. The slowing-down process was calculated using the Northcliffe and Schilling⁸ electronic stopping power and the nuclear contribution was treated according to Blaugrund.⁹

The average Q_0 is obtained by fitting the calculated ratios of the spectra for the backed and unbacked targets to the experimental ratio with Q_0 as the only free parameter. The Q_0 values, corresponding enhancement factors (in Weisskopf units⁶), and deformation parameters⁷

obtained for the two systems investigated are given in Table I. Figure 1c shows the ratio of the unfolded spectra measured at 0° for a Au-backed and unbacked ^{27}Al target in the region of interest. The solid line is the best fitting calculated curve. The dashed lines show the ratios calculated for the two extreme values of Q_0 allowed by the errors given in Table I.

The uncertainties assigned to the results contain estimates of the following systematic errors: i) neglect of terms higher than first order in v/c ; of angular distribution effects due to hyperfine interaction (the deorientation effect is calculated to be insignificant); and of finite target thickness (we assume no transverse velocity even at the range of nuclear stopping); ii) uncertainty from insufficient knowledge of the stopping power; iii) an estimated five-fold increase in the statistical errors from unfolding the raw data.

There is a small amount of statistical γ -rays present in the bump region of the spectrum. Angular correlation experiments show,³ however, that the bump region consists of $\sim 90\%$ quadrupole radiation in these heavy rare earth nuclei. Furthermore, since the sensitivity of the method depends on the slope of the spectrum, and the slope of the statistical part of the spectrum is much less steep than that of the bump, our method is less sensitive to the lifetime of the statistical γ -rays.

With a moment of inertia parameter of $2\mathcal{I}/\hbar^2 = 150 \text{ MeV}^{-1}$ we can correlate a spin range of $I = 30-50$ to the γ -ray energy region of 0.8 to 1.4 MeV, in which we observe a measurable effect. We derive from the data that the transitions between these high spin states show an enhancement

of ~ 300 s.p.u. and a rather large deformation of $\beta_2 = 0.4$. The latter value is consistent with the moment of inertia observed in Refs. 2, 3 and 5. We conclude that the transitions in the E2 bump are of strongly collective nature, with as high or higher a degree of collectivity as in the ground state rotational bands of deformed nuclei.

As can be seen from Fig. 1c there is good agreement in both magnitude and slope for the calculated and measured curves. This indicates that for the investigated nuclei the model provides a satisfactory description of their behavior, suggesting no abrupt change in structure in the investigated region. Changes in structure corresponding to different lifetimes can give rise to regions where the fit with a single value of Q_0 is not good. This may provide a method for studying nuclear structure changes in the continuum region.

ACKNOWLEDGMENTS

We would like to thank Dr. D. Ward for helpful discussions on the stopping power calculations.

This work was supported by the Department of Energy.

TABLE I. Intrinsic quadrupole moments Q_0 , enhancement factors $B(E2)/B(E2)_w$, and deformation parameters β_2 for the $^{163}\text{Ho}^*$ and $^{164}\text{Er}^*$ compound systems.

Target	Compound nucleus	Stopper	$ Q_0 $		Average	$\frac{B(E2)}{B(E2)_w}$	β_2
			a)	b)			
^{27}Al	$^{163}\text{Ho}^*$	Au	9.3(2.5)	8.0(2.3)	8.2(2.2)	270(100)	0.37(11)
^{27}Al	$^{163}\text{Ho}^*$	Pb	7.0(2.6)	8.6(3.0)			
^{28}Si	$^{164}\text{Er}^*$	Au	8.5(3.0)	8.6(3.0)	9.8(3.3)	380(180)	0.43(15)
^{28}Si	$^{164}\text{Er}^*$	Au	12.2(4.9)	10.2(4.3)			

a) Determined from ratios of spectra at 0° as described in the text.

b) Determined from double ratios of spectra at 0° and 90° as described in the text.

REFERENCES

1. R.S.Simon, M.V.Banaschik, R.M.Diamond, J.O.Newton, and F.S.Stephens, Nucl. Phys. A290, 253 (1977).
2. M.A.Deleplanque, I.Y.Lee, F.S.Stephens, R.M.Diamond, and M.M.Aleonard, Phys. Rev. Letters 40, 629 (1978).
3. M.A.Deleplanque, Th.Byrski, R.M.Diamond, H.Hübel, F.S.Stephens, B.Herskind, and R.Bauer, to be published.
4. J.F.Mollenauer, Phys. Rev. 127, 867 (1962).
5. D.Hillis, B.Fernandez, F.Folkmann, J.Garrett, G.B.Hagemann, and B.Herskind, to be published.
6. A.Bohr and B.R.Mottelson, Nuclear Structure, Vol. I (W.A.Benjamin, New York & Amsterdam, 1969), p.389.
7. K.Alder and A.Winther, Coulomb Excitation (Academic Press, New York & London, 1966).
8. L.C.Northcliffe and R.F.Schilling, Nucl. Data A7, 233 (1970).
9. A.E.Blaugrund, Nucl. Phys. 88, 501 (1966).




Fig. 1. (a) Sum of 2-6 fold coincidence γ -ray spectra (raw data) measured at 0° for a self-supporting ^{27}Al target (solid curve) and for a ^{27}Al target on a Au backing (dotted curve).

(b) Ratio of the raw (not unfolded) spectra measured at 0° for a self-supporting ^{27}Al target and for a ^{27}Al target on a Au backing (the curve is drawn to guide the eye).

(c) Ratio of the unfolded spectra (dots) measured at 0° for a self-supporting ^{27}Al target and for a ^{27}Al target on a Au backing. The solid curve represents the best fit of the calculated ratio to the data. The dashed lines show the ratios calculated for the two extreme values of Q_0 allowed by the errors given in Table I.

(d) As Fig. 1b, but for a ^{28}Si target.

(XBL 786-7560)

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

TECHNICAL INFORMATION DEPARTMENT
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720