

# UC Berkeley

## UC Berkeley Previously Published Works

### Title

Gold-Catalyzed Hydrofluorination of Electron-Deficient Alkynes: Stereoselective Synthesis of  $\beta$ -Fluoro Michael Acceptors

### Permalink

<https://escholarship.org/uc/item/94b0v6bv>

### Journal

ACS Catalysis, 8(7)

### ISSN

2155-5435

### Authors

O'Connor, Thomas J  
Toste, F Dean

### Publication Date

2018-07-06

### DOI

10.1021/acscatal.8b01341

Peer reviewed



Published in final edited form as:

ACS Catal. 2018 July 6; 8(7): 5947–5951. doi:10.1021/acscatal.8b01341.

## Gold-Catalyzed Hydrofluorination of Electron-Deficient Alkynes: Stereoselective Synthesis of $\beta$ -Fluoro Michael Acceptors


Thomas J. O'Connor, F. Dean Toste\*

Department of Chemistry, University of California, Berkeley, California 04720, United States

### Abstract

The gold(I)-catalyzed, stereoselective hydrofluorination of electron-deficient alkynes with triethylamine trihydrogen fluoride ( $\text{Et}_3\text{N}\cdot 3\text{HF}$ ) is described. Fluorinated  $\alpha,\beta$ -unsaturated aldehydes, amides, esters, ketones, and nitriles were isolated in moderate to good yields as single diastereomers. In all but four cases, the (*Z*)-vinyl fluorides were initially formed in 97% diastereoselectivity. This work constitutes the first catalytic example of the diastereoselective preparation of a variety of  $\beta$ -alkyl,  $\beta$ -fluoro Michael acceptors from alkynes. Additionally, the described work expands access to  $\beta$ -aryl,  $\beta$ -fluoro Michael acceptors to the synthesis of  $\beta$ -fluoro- $\alpha,\beta$ -unsaturated amides and nitriles. The monofluoroalkenes formed through this strategy were readily transformed into other fluorine-containing compounds, and the developed method was applied to the synthesis of a fluorinated analogue of Exoderil, a topical antimycotic.

### Graphical Abstract



### Keywords

monofluoroalkenes; michael acceptors; hydrofluorination; gold catalysis; fluorine

New routes toward the selective fluorination of small molecules have been targeted in recent years due to the differences in the physical and biological properties between fluorinated compounds and those of their nonfluorinated analogues.<sup>1</sup> A fluorinated motif of particular interest is the monofluoroalkene. Monofluoroalkenes are isosteric with peptide bonds, and several bioactive compounds containing this motif have been reported.<sup>2</sup> Although several synthetic protocols exist to access  $\alpha$ -fluoro,  $\alpha,\beta$ -unsaturated carbonyl compounds—the

\*Corresponding Author: [fdtoste@berkeley.edu](mailto:fdtoste@berkeley.edu).

#### ASSOCIATED CONTENT

##### Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: [10.1021/acscatal.8b01341](https://doi.org/10.1021/acscatal.8b01341).

Experimental details and compound characterization data (PDF)

#### Notes

The authors declare no competing financial interest.

Horner–Wadsworth–Emmons reaction,<sup>3</sup> the Julia Olefination,<sup>4</sup> the Peterson Olefination,<sup>5</sup> and the Reformatsky reaction<sup>6</sup>—the stereoselective synthesis of  $\beta$ -fluoro,  $\alpha,\beta$ -unsaturated carbonyl compounds has proven to be a challenge, especially if  $\beta$ -alkyl substituents are desired.<sup>7</sup> Previous methods to access (*Z*)- $\beta$ -fluoro- $\alpha,\beta$ -unsaturated carbonyl compounds are limited by the formation of products with low diastereoselectivities or yields,<sup>8</sup> the requirement for prefunctionalized starting materials,<sup>9</sup> and narrow functional group tolerance.<sup>9b,c,10</sup> Because of these limitations, a stereoselective and functional-group-tolerant method to access (*Z*)- $\beta$ -alkyl,  $\beta$ -fluoro- $\alpha,\beta$ -unsaturated carbonyl compounds would be highly desirable.

The hydrofluorination of electron-deficient alkynes is perhaps the most direct method to generate (*Z*)- $\beta$ -fluoro  $\alpha,\beta$ -unsaturated carbonyl compounds from commercially available starting materials. Although some electron-deficient alkynes can undergo hydrofluorination in the absence of a catalyst, the diastereoselectivities of these reactions are generally moderate, especially for  $\beta$ -alkyl substrates.<sup>8a,b,10</sup> Traditional chromatographic techniques often fail to separate (*E*) and (*Z*) isomers of monofluoroalkenes; therefore, it is essential that the desired monofluoroalkenes are synthesized with high diastereomeric ratios.<sup>11</sup>

Since Sadighi's seminal report of the gold-catalyzed hydrofluorination of internal alkynes in 2007, other research groups have expanded the use of coinage metals for alkyne hydrofluorination.<sup>12</sup> Both Jiang, with excess AgF (Scheme 1a), and Nolan, with a catalytic amount of gold (Scheme 1b), prepared  $\beta$ -aryl,  $\beta$ -fluoro- $\alpha,\beta$ -unsaturated esters or ketones from electron-deficient, unsymmetrical alkynes.<sup>12c,e</sup> However, neither procedure reported the synthesis of  $\beta$ -alkyl,  $\beta$ -fluoro Michael acceptors or utilized alternative electron-withdrawing groups such as nitriles or amides. Alternative conditions were described by Hammond and Xu for the gold-catalyzed hydrofluorination of alkynes with a new DMPU/HF fluorinating reagent, but this procedure did not expand access to (*Z*)- $\beta$ -fluoro- $\alpha,\beta$ -unsaturated carbonyl compounds.<sup>12d</sup> The first gold-catalyzed synthesis of a  $\beta$ -alkyl- $\beta$ -fluoro Michael acceptor was demonstrated by Hammond and Xu in 2017 (Scheme 1c).<sup>12f</sup>

Although  $\beta$ -alkyl- $\beta$ -fluorovinylsulfones could be accessed in a (*Z*)-selective manner, alkynes that did not bear a sulfonyl group—such as aroyl and phosphonyl—failed to undergo hydrofluorination. Despite these advances in alkyne hydrofluorination by coinage metals, a general procedure to synthesize a variety of (*Z*)- $\beta$ -alkyl,  $\beta$ -fluoro Michael acceptors from electron-deficient alkynes is still an unsolved challenge.

Herein, we report a method for the preparation of a diverse array of  $\beta$ -alkyl,  $\beta$ -fluoro Michael acceptors from the gold-catalyzed hydrofluorination of electron-deficient alkynes. In addition to forming  $\beta$ -fluoro- $\alpha,\beta$ -unsaturated esters and ketones, this method is the first gold-catalyzed procedure to generate  $\beta$ -fluoro- $\alpha,\beta$ -unsaturated amides, nitriles, and aldehydes. A variety of  $\beta$ -alkyl as well as  $\beta$ -aryl substituents were tolerated; notably, 3° alkyl, alkenyl, and *o*-tolyl. Furthermore, we demonstrate that the monofluoroalkene products are synthetically versatile fluorinated building blocks.

The hydrofluorination of ethyl 2-butyrate (**1a**) with Et<sub>3</sub>N·3HF to form ethyl (*Z*)-3-fluorobut-2-enoate (**1b**) was selected as a model reaction. Monofluoroalkene **1b** formed in

moderate yields and low stereoselectivities under conditions similar to those reported by Sadighi (see Table 1, entry 1).<sup>12a</sup> Reactions employing AgBF<sub>4</sub> as the silver salt afforded alkene **1b** in greater chemical yield compared to reactions conducted in the presence of other silver salts (entry 1 and 2, see the Supporting Information for further details). Upon switching from gold catalysts bearing NHC-ligands to gold catalysts bearing phosphine ligands, modest improvements in both yield and stereoselectivity were observed (entries 3 and 4). Unfortunately, reactions conducted with several triaryl or trialkyl phosphine gold(I) complexes as catalysts generated a purple hue after several hours in the presence of Et<sub>3</sub>N·3HF, which has been reported by others as a visual indication of catalyst decomposition.<sup>13</sup>

Cationic-gold(I) complexes with dialkylbiarylphosphine ligands are known to be more stable toward decomposition pathways than cationic gold(I) complexes triaryl or trialkyl phosphines.<sup>14</sup> Upon switching the gold catalyst to CyJohnPhosAuCl, monofluoroalkene **1b** was generated in 84% yield. However, the stereoselectivity of the reaction conducted with CyJohnPhosAuCl decreased relative to the stereoselectivity of the reaction conducted with Cy<sub>3</sub>PAuCl as the catalyst (entry 3 and 4). Examination of a variety of dialkylbiaryl phosphinegold(I) complexes revealed that only reactions with RuPhos as the ligand afforded the greatest *Z:E* selectivity of **1b** (entries 6 and 7). For instance, in the presence of CyJohnPhos the yield of **1b** after 4 h was 85% but with a *Z:E* of 77:23.

In addition to the ligand effect on the reaction, both the solvent and additive were found to influence the yield and stereoselectivity of the hydrofluorination of alkynoate **1a**. Switching from potassium bisulfate to *p*-chlorobenzoic acid (*p*-Cl BA), a more soluble acid coadditive, resulted in a modest improvement in the yield of monofluoroalkene **1b** (entry 8). Reactions conducted with RuPhosAuCl and CH<sub>3</sub>CN as the solvent afforded the hydrofluorination product in a further improved yield while maintaining the *Z*-selectivity observed at shorter reaction times (entry 8 and 9). The change in solvent also ensured that the *Z:E* ratio did not decrease over time, permitting easier reaction monitoring as alkene isomerization was largely suppressed. Ultimately, reactions conducted in a solvent mixture of CH<sub>3</sub>CN:CH<sub>2</sub>Cl<sub>2</sub> maintained the high stereoselectivity of the hydrofluorination of alkyne **1a** while affording alkene **1b** in an improved yield (entry 9 and 10). The beneficial improvement in the yield of **1b** was observed with as little as 10 mol % *p*-Cl BA (entry 11 and 12). Other acid additives were examined, but benzoic acid derivatives appeared to provide an optimal p*K*<sub>a</sub> range (see Supporting Information, Table S4). Increasing the equivalents of Et<sub>3</sub>N·3HF did not have a significant influence on the reaction (entry 13); however, reactions with Et<sub>3</sub>N·2HF, Et<sub>3</sub>N·HF, and pyridine·HF (70% HF) failed to generate alkene **1b** (See Supporting Information).

Having identified suitable reactions conditions for the hydrofluorination of alkyne **1a**, we investigated the hydrofluorination of  $\beta$ -alkyl alkynoates, alkynones, alkynamides, and alkynenitriles (Table 2). Methyl, 1° alkyl, 2° alkyl, and vinyl  $\beta$ -substituted alkynoates underwent hydrofluorination in the presence of Et<sub>3</sub>N·3HF in a *Z*-selective manner in good yields. Notably, the final products were all isolated as a single diastereomer after standard silica gel column chromatography. Importantly, these results highlight this operationally simple, one-step route to  $\beta$ -alkyl,  $\beta$ -fluoro Michael acceptors from alkynes. The hydrofluorination reaction was also shown to be scalable, as fluoroalkenes **3b** and **5b** were

both prepared on a gram scale in good yield and with excellent *Z*-selectivity. For substrate **6a** with a bulky  $\beta$ -substituents, a higher reaction temperature was required to obtain the product in moderate yield (**6b**). The hydrofluorination of alkynoates bearing  $\beta$ -vinyl substituents provided straightforward access to fluorinated dienes **7b** and **8b**.

Hydrofluorination of the  $\gamma,\delta$ -alkene of either **7a** or **8a** was not detected by  $^{19}\text{F}$  NMR spectroscopy. The reaction conditions for the hydrofluorination of  $\beta$ -alkyl alkynoates were also suitable for the hydrofluorination of  $\beta$ -alkyl (hetero)aryl alkynones **9b** and **10b**.

Although methyl ketone **11a** proved to be a challenging substrate, **11b** was isolated in good yield with only a trace amount of the *E*-isomer. Both 2° and 3°  $\beta$ -alkyl alkynamides (**12–14a**) as well as alkynonitrile derivative **15a** underwent hydrofluorination to provide **12–15b** in moderate yields. Dec-2-ynal was the only substrate that did not undergo hydrofluorination in a diastereoselective manner under the standard conditions in Table 2 (72%, *Z:E* = 51:49). However, conducting the reaction at 5 °C did afford a *Z:E* ratio of >98:2 and 22% yield after 24 h. Unfortunately, after 96 h at 5 °C, the yield increased to 51% but the *Z:E* ratio decreased to 70:30.

To showcase the generality of this method, the hydrofluorination reactions of a variety of electron deficient alkynes bearing  $\beta$ -aryl substituents were also explored (Table 3).

Generally, the yields of  $\beta$ -aryl-monofluoroalkenes **16–30b** were comparable to those of their  $\beta$ -alkyl-analogues **2–15b**. In contrast to previous procedures, even a monofluoroalkene bearing an *ortho*-substituted aryl group (**19b**) was generated in modest yield.<sup>12e</sup> Compared with the esters and ketones, even the more electrophilic 2-phenylpropionaldehyde afforded **27b** in a *Z*-selective manner. Moreover, both  $\beta$ -aryl alkynonitriles and alkynamides were suitable substrates, generating otherwise difficult to access fluorinated motifs (**28–30b**). Finally, this methodology was found to be complementary to that reported by Hammond and Xu (See Supporting Information, Table S5)<sup>12f</sup>

The monofluoroalkenes generated from our catalytic process underwent a series of transformations demonstrating that  $\beta$ -fluoro Michael acceptors are valuable fluorinated building blocks (Scheme 2). For example, ester **3b** was reduced in the presence of DIBAL–H to yield the fluorinated allylic alcohol **1c** in high yield.<sup>15</sup> Aldehyde **27b** underwent Wittig olefination in modest yield to afford a 1-fluoro-2,4-diene **2c**.<sup>16</sup> In the presence of a suitable 1,3-ylide, ester **5b** underwent a regioselective [3 + 2] cycloaddition to generate a pyrrolidine with a quaternary fluorine center (**3e**).<sup>17</sup> Finally, amide **31b** was reduced in the presence of Meerwein's salt to furnish a fluorine-containing analogue of Exoderil **4c**.<sup>18</sup>

In conclusion, we have developed a stereoselective hydrofluorination of electron-deficient alkynes catalyzed by a RuPhos-ligated gold(I) complex. For the first time, direct access to a variety of (*Z*)- $\beta$ -alkyl,  $\beta$ -fluoro Michael acceptors was achieved. In addition, (*Z*)- $\beta$ -aryl,  $\beta$ -fluoro  $\alpha,\beta$ -unsaturated amides and nitriles were conveniently accessed with the disclosed method. The synthetic potential of the resulting monofluoroalkene was demonstrated with various transformations of the products without the loss of the newly installed fluorine atom, and with the synthesis of a fluorinated analogue of Exoderil.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## ACKNOWLEDGMENTS




We acknowledge the National Institute of General Medical Sciences (R35 GM118190) for financial support of this work. T.J.O. thanks the NSF (DGE 1752814) for a predoctoral fellowship.

## REFERENCES


- (1). (a) Bizet V; Besset T; Ma J-A; Cahard D Recent Progress in Asymmetric Fluorination and Trifluoromethylation Reactions. *Curr. Top. Med. Chem* 2014, 14, 901–940 [PubMed: 24484424] (b) Purser S; Moore PR; Swallow S; Gouverneur V Fluorine in Medicinal Chemistry. *Chem. Soc. Rev* 2008, 37, 320–330 [PubMed: 18197348] (c) Hagmann WK The Many Roles for Fluorine in Medicinal Chemistry. *J. Med. Chem* 2008, 51, 4359–4369 [PubMed: 18570365] (d) Muller K; Faeh C; Diederich F Fluorine in Pharmaceuticals: Looking Beyond Intuition. *Science* 2007, 317, 1881–1886 [PubMed: 17901324] (e) Yamazaki T; Taguchi T; Ojima I Fluorine in Medicinal Chemistry and Chemical Biology; John Wiley & Sons, Ltd, Chichester, U.K., 2009; pp 3–624.
- (2). (a) Landelle G; Bergeron M; Turcotte-Savard MO; Paquin JF Synthetic Approaches to Monofluoroalkenes. *Chem. Soc. Rev* 2011, 40, 2867–2908. [PubMed: 21399789] (b) Yanai H; Taguchi T Synthetic Methods for Fluorinated Olefins. *Eur. J. Org. Chem* 2011, 2011, 5939–5954.
- (3). Machleidt H; Wessendorf R Carbonyl-Fluorolefinierungen. *Justus Liebigs Ann. Chem* 1964, 674, 1–10.
- (4). Zajc B; Kake S Exceptionally Mild, High-Yield Synthesis of  $\alpha$ -Fluoro Acrylates. *Org. Lett* 2006, 8, 4457–4460. [PubMed: 16986924]
- (5). Welch JT; Herbert RW The Stereoselective Construction of (Z)-3-Aryl-2-fluoroalkenoates. *J. Org. Chem* 1990, 55 (16), 4782–4784.
- (6). Barma DK; Kundu A; Zhang H; Mioskowski C; Falck JR (Z)- $\alpha$ -Haloacrylates: An Exceptionally Stereoselective Preparation via Cr(II)-Mediated Olefination of Aldehydes with Trihaloacetates. *J. Am. Chem. Soc* 2003, 125, 3218–3219. [PubMed: 12630869]
- (7). (a) Yanai H; Taguchi T Synthetic Methods for Fluorinated Olefins. *Eur. J. Org. Chem* 2011, 2011 (30), 5939–5954. (b) Champagne PA; Desroches J; Hamel JD; Vandamme M; Paquin JF Monofluorination of Organic Compounds: 10 Years of Innovation. *Chem. Rev* 2015, 115, 9073–9174. [PubMed: 25854146] (c) Drouin M; Hamel J-D; Paquin J-F Synthesis of Monofluoroalkenes: A Leap Forward. *Synthesis* 2018, 50, 881–955.
- (8). (a) Albert P; Cousseau J Tetrabutylammonium and Polymer-supported Dihydrogentrifluoride: New Hydrofluorinating Reagents for Electrophilic Alkynes. *J. Chem. Soc., Chem. Commun* 1985, 961–962. (b) Gorgues A; Stéphan D; Cousseau J Mono-hydrofluorination of Electrophilic Alkynes by the Liquid Biphasic CsF-H<sub>2</sub>O-DMF System (DMF = N,N-dimethylformamide). *J. Chem. Soc., Chem. Commun* 1989, 1493–1494. (c) Sano K; Fukuhara T; Hara S Regioselective Synthesis of  $\beta$ -fluoro- $\alpha,\beta$ -unsaturated Ketones by the Reaction of  $\beta$ -diketones with DFMB. *J. Fluorine Chem* 2009, 130, 708–713.
- (9). (a) Zhang J; Liu L; Duan J; Gu L; Chen B; Sun T; Gong, Stereoselective One-Pot Sequential Dehydrochlorination/trans-Hydrofluorination Reaction of  $\beta$ -Chloro- $\alpha,\beta$ -unsaturated Aldehydes or Ketones: Facile Access to (Z)- $\beta$ -Fluoro- $\beta$ -arylenals/ $\beta$ -Fluoro- $\beta$ -arylenones. *Adv. Synth. Catal* 2017, 359, 4348–4358. (b) Yoshida M; Kawakami K; Hara S An Efficient Stereoselective Synthesis of (E)- $\beta$ -Fluoroalkenyliodonium Salts. *Synthesis* 2004, 2004, 2821–2824. (c) Yoshida M; Komata A; Hara S Stereoselective Synthesis of Fluoroalkenes via (Z)-2-Fluoroalkenyliodonium Salts. *Tetrahedron* 2006, 62, 8636–8645.
- (10). (a) McElroy KT; Purrington ST; Bumgardner CL; Burgess JP Lack of Polymerization of Fluorinated Acrylates. *J. Fluorine Chem* 1999, 95, 117–120. (b) Krishnan G; Sampson P Synthesis of  $\beta$ -Fluoro- $\alpha,\beta$ -unsaturated Esters and Nitriles via a Fluoro-Pummerer Rearrangement. *Tetrahedron Lett* 1990, 31, 5609–5612. (c) Patrick TB; Neumann J; Tatro A

- Cycloaddition Reactions of ethyl (E)- and (Z)-3-fluoropropenoate. *J. Fluorine Chem* 2011, 132, 779–782.
- (11). (a) Zhao Y; Jiang F; Hu J Spontaneous Resolution of Julia-Kocienski Intermediates Facilitates Phase Separation to Produce *Z*- and *E*-monofluoroalkenes. *J. Am. Chem. Soc* 2015, 137, 5199–5203. [PubMed: 25831969] (b) Thornbury RT; Toste FD Palladium-Catalyzed Defluorinative Coupling of 1-Aryl-2,2-Difluoroalkenes and Boronic Acids: Stereo-selective Synthesis of Monofluorostilbenes. *Angew. Chem., Int. Ed* 2016, 55, 11629–11632. (c) Sommer H; Fürstner A Stereospecific Synthesis of Fluoroalkenes by Silver-Mediated Fluorination of Functionalized Alkenylstannanes. *Chem. - Eur. J* 2017, 23, 558–562. [PubMed: 27883234] (d) Hu J; Han X; Yuan Y; Shi Z Stereoselective Synthesis of *Z* Fluoroalkenes through Copper-Catalyzed Hydrodefluorination of gem-Difluoroalkenes with Water. *Angew. Chem., Int. Ed* 2017, 56, 13342–13346.
- (12). (a) Akana JA; Bhattacharyya KX; Muller P; Sadighi JP Reversible C–F Bond Formation and the Au-Catalyzed Hydrofluorination of Alkynes. *J. Am. Chem. Soc* 2007, 129, 7736–7737. [PubMed: 17547409] (b) Gorske BC; Mbofana CT; Miller SJ Regio- and Stereoselective Synthesis of Fluoroalkenes by Directed Au(I) Catalysis. *Org. Lett* 2009, 11, 4318–4321. [PubMed: 19711904] (c) Li Y; Liu X; Ma D; Liu B; Jiang H Silver-Assisted Difunctionalization of Terminal Alkynes: Highly Regio- and Stereoselective Synthesis of Bromofluoroalkenes. *Adv. Synth. Catal* 2012, 354, 2683–2688. (d) Okoromoba OE; Han J; Hammond GB; Xu B Designer HF-Based Fluorination Reagent: Highly Regioselective Synthesis of Fluoroalkenes and gem-Difluoromethylene Compounds from Alkynes. *J. Am. Chem. Soc* 2014, 136, 14381–14384. [PubMed: 25260170] (e) Nahra F; Patrick SR; Bello D; Brill M; Obled A; Cordes DB; Slawin AM; O'Hagan D; Nolan SP Hydrofluorination of Alkynes Catalysed by Gold Bifluorides. *ChemCatChem* 2015, 7, 240–244. [PubMed: 26236406] (f) Zeng X; Liu S; Hammond GB; Xu B Divergent Regio- and Stereoselective Gold-catalyzed Synthesis of  $\alpha$ -Fluorosulfones and  $\beta$ -Fluorovinylsulfones from Alkynylsulfones. *Chem. - Eur. J* 2017, 23, 11977–11981 [PubMed: 28736856] (g) For reviews of gold catalysis and fluorine: Miro J; del Pozo C Fluorine and Gold: A Fruitful Partnership. *Chem. Rev* 2016, 116, 11924–11966. [PubMed: 27548659] (h) Hopkinson MN; Gee AD; Gouverneur V Gold Catalysis and Fluorine. *Isr. J. Chem* 2010, 50, 675–690.
- (13). Bartolomé C; Ramiro Z; Peñas-Defrutos MN; Espinet P Some Singular Features of Gold Catalysis: Protection of Gold(I) Catalysts by Substoichiometric Agents and Associated Phenomena. *ACS Catal* 2016, 6, 6537–6545.
- (14). (a) Malhotra D; Hammond GB; Xu B Ligand Design in Gold Catalysis and Chemistry of Gold-Oxonium Intermediates. *Top. Curr. Chem* 2014, 357, 1–24. (b) Gorin DJ; Sherry BD; Toste FD Ligand Effects in Homogeneous Au Catalysis. *Chem. Rev* 2008, 108, 3351–3378. [PubMed: 18652511]
- (15). Konno T; Ikemoto A; Ishihara T A New Entry to the Construction of a Quaternary Carbon Center having a Fluorine Atom-S(N)2' Reaction of  $\gamma$ -Fluoroallylic Alcohol Derivatives with Various Cyanocuprates. *Org. Biomol. Chem* 2012, 10, 8154–8163. [PubMed: 22965477]
- (16). Zhu S; Guo Z; Huang Z; Jiang H Bioinspired Intramolecular Diels-Alder Reaction: A Rapid Access to the Highly-Strained Cyclopropane-Fused Polycyclic Skeleton. *Chem. - Eur. J* 2014, 20, 2425–2430. [PubMed: 24501038]
- (17). (a) Mason JM; Murkin AS; Li L; Schramm VL; Gainsford GJ; Skelton BW A  $\beta$ -fluoroamine Inhibitor of Purine Nucleoside Phosphorylase. *J. Med. Chem* 2008, 51, 588020135884. (b) McAlpine I; Tran-Dube M; Wang F; Scales S; Matthews J; Collins MR; Nair SK; Nguyen M; Bian J; Alsina LM; Sun J; Zhong J; Warmus JS; O'Neill BT Synthesis of Small 3-Fluoro- and 3,3-Difluoropyrrolidines Using Azomethine Ylide Chemistry. *J. Org. Chem* 2015, 80, 7266–7274. [PubMed: 26057617]
- (18). Deiters A; Chen K; Eary CT; Martin SF Biomimetic Entry to the Sarpagan Family of Indole Alkaloids: Total Synthesis of (+)-Geissoschizine and (+)-N-Methylvellosimine. *J. Am. Chem. Soc* 2003, 125, 4541–4550. [PubMed: 12683825]


Previous Work:




This Work:

**Scheme 1.**Generation of  $\beta$ -Fluoro Michael Acceptors from Alkynes with Coinage Metals






**Scheme 2.**  
Diversification of Fluorinated Michael Acceptors

**Table 1.**Effect of the Reaction Conditions on the Hydrofluorination of **1a**


entry	L	solvent	additive	yield [%] (Z:E) <sup>b</sup>
1	IPr	CH <sub>2</sub> Cl <sub>2</sub>	KHSO <sub>4</sub>	50 (66:34)
2 <sup>c</sup>	IPr	CH <sub>2</sub> Cl <sub>2</sub>	KHSO <sub>4</sub>	43 (70:30)
3	PPh <sub>3</sub>	CH <sub>2</sub> Cl <sub>2</sub>	KHSO <sub>4</sub>	55 (60:40)
4	PCy <sub>3</sub>	CH <sub>2</sub> Cl <sub>2</sub>	KHSO <sub>4</sub>	64 (75:25)
5	CyJohnPhos	CH <sub>2</sub> Cl <sub>2</sub>	KHSO <sub>4</sub>	84 (55:45)
6	RuPhos	CH <sub>2</sub> Cl <sub>2</sub>	KHSO <sub>4</sub>	57 (56:44)
7 <sup>d</sup>	RuPhos	CH <sub>2</sub> Cl <sub>2</sub>	KHSO <sub>4</sub>	62 (97:3)
8 <sup>e</sup>	RuPhos	CH <sub>2</sub> Cl <sub>2</sub>	<i>p</i> -Cl BA <sup>e</sup>	66 (97:3)
9	RuPhos	CH <sub>3</sub> CN	<i>p</i> -Cl BA	70 (97:3)
10	RuPhos	1:4 CH <sub>2</sub> Cl <sub>2</sub> :CH <sub>3</sub> CN	<i>p</i> -Cl BA	76 (96:4)
11 <sup>f</sup>	RuPhos	1:4 CH <sub>2</sub> Cl <sub>2</sub> :CH <sub>3</sub> CN	<i>p</i> -Cl BA	71 (96:4)
12	RuPhos	1:4 CH <sub>2</sub> Cl <sub>2</sub> :CH <sub>3</sub> CN	none	65 (96:4)
13 <sup>g</sup>	RuPhos	1:4 CH <sub>2</sub> Cl <sub>2</sub> :CH <sub>3</sub> CN	<i>p</i> -Cl BA	80 (96:4)

<sup>a</sup>General reaction conditions: 0.2 mmol **1a**, plastic vial.<sup>b</sup>Yields and Z:E ratios were determined by <sup>19</sup>F NMR spectroscopy with 2,4-dinitrofluorobenzene as an internal standard.<sup>c</sup>5 mol % AgSbF<sub>6</sub>.<sup>d</sup>4 h.<sup>e</sup>*p*-chlorobenzoic acid.<sup>f</sup>10 mol % *p*-Cl BA.<sup>g</sup>3.0 equiv Et<sub>3</sub>N·3HF.

Table 2.

Scope of  $\beta$ -Alkyl,  $\beta$ -Fluoro Michael Acceptors

<sup>a</sup> Standard reaction conditions: 0.5 mmol **2-15a**, 3.0 equiv Et<sub>3</sub>N·3HF, 1.0 equiv *p*-Cl BA, 5 mol % RuPhosAuCl, 5 mol % AgBF<sub>4</sub>, 4:1 MeCN:CH<sub>2</sub>Cl<sub>2</sub> [0.7M], rt, 24 h.

<sup>b</sup> **2-15b** isolated as a single isomer except **11b**.

<sup>c</sup> Determined by <sup>19</sup>F NMR spectroscopy with PhF as an internal standard.

<sup>d</sup> 6.0 mmol scale.

<sup>e</sup> 5.0 mmol scale.

<sup>f</sup> 55 °C ginsoluble product.

*h*<sub>1.25 M, 4.0 equiv Et3N·HF.</sub>

*i*<sub>1.25 M, 4.0 equiv Et3N·3HF.</sub>

*j*<sub>1.43 M, 4.5 equiv Et3N·3HF, 50 °C.</sub>


Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 3.

Scope of  $\beta$ -aryl,  $\beta$ -fluoro Michael acceptors

<sup>a</sup>Standard reaction conditions: 0.5 mmol **16-30a**, 3.0 equiv Et<sub>3</sub>N·3HF, 1.0 equiv *p*-Cl BA, 5 mol % RuPhosAuCl, 5 mol % AgBF<sub>4</sub>, 4:1 MeCN:CH<sub>2</sub>Cl<sub>2</sub> [0.7M], rt, 24 h.

<sup>b</sup>**16-30b** isolated as a single isomer.

<sup>c</sup>Determined by <sup>19</sup>F NMR spectroscopy with PhF as an internal standard.

<sup>d</sup>1.43 M, 45 °C.

<sup>e</sup>45 °C.

<sup>f</sup>4.5 mmol.

<sup>g</sup>1.25 M, 55 °C, 4.0 equiv Et<sub>3</sub>N·3HF.

<sup>h</sup>45 °C, 48 h, 4.0 equiv Et<sub>3</sub>N·3HF.