

# UC Berkeley

## UC Berkeley Previously Published Works

### Title

Modeling the Effect of the 2018 Revised ACGIH® Hand Activity Threshold Limit Value® (TLV) at Reducing Risk for Carpal Tunnel Syndrome.

### Permalink

<https://escholarship.org/uc/item/94n03542>

### Journal

Journal of occupational and environmental hygiene, 16(9)

### ISSN

1545-9624

### Authors

Yung, Marcus  
Dale, Ann Marie  
Kapellusch, Jay  
[et al.](#)

### Publication Date

2019-09-01

### DOI

10.1080/15459624.2019.1640366

Peer reviewed



Published in final edited form as:

*J Occup Environ Hyg.* 2019 September ; 16(9): 628–633. doi:10.1080/15459624.2019.1640366.

## Modeling the Effect of the 2018 Revised ACGIH® Hand Activity Threshold Limit Value® (TLV) at Reducing Risk for Carpal Tunnel Syndrome

Marcus Yung, PhD<sup>1</sup>, Ann Marie Dale, PhD<sup>1</sup>, Jay Kapellusch, PhD<sup>2</sup>, Stephen Bao, PhD<sup>3</sup>, Carisa Harris-Adamson, PhD<sup>4,5</sup>, Alysha R Meyers, PhD<sup>6</sup>, Kurt T Hegmann, MD<sup>7</sup>, David Rempel, MD<sup>5,8</sup>, Bradley A. Evanoff, MD<sup>1</sup>

<sup>1</sup>Division of General Medical Sciences, Washington University School of Medicine, Saint Louis, MO, USA

<sup>2</sup>Department of Occupational Science and Technology, University of Wisconsin-Milwaukee, Milwaukee, Wisconsin, USA

<sup>3</sup>Safety and Health Assessment and Research for Prevention (SHARP) Program, Washington State Department of Labor and Industries, Olympia, Washington, USA

<sup>4</sup>Department of Environmental Health Sciences, University of California Berkeley, Berkeley, California, USA

<sup>5</sup>Department of Medicine, University of California at San Francisco, San Francisco, CA, USA

<sup>6</sup>Division of Surveillance, Hazard Evaluations, and Field Studies, National Institute for Occupational Safety and Health, CDC

<sup>7</sup>Department of Family and Preventive Medicine, University of Utah, Utah, USA

<sup>8</sup>Department of Bioengineering, University of California Berkeley, Berkeley, California, USA

### Abstract

Recent studies have shown the 2001 American Conference of Governmental Industrial Hygienists Threshold Limit Value for Hand Activity® was not sufficiently protective for workers at risk of carpal tunnel syndrome. These studies led to a revision of the Threshold Limit Value and Action Limit. This study compares the effect of applying the 2018 Threshold Limit Value® versus the 2001 Threshold Limit Value® to predict incident carpal tunnel syndrome within a large occupational cohort study (n = 4321 workers). Time from study enrollment to first occurrence of carpal tunnel syndrome was modeled using Cox proportional hazard regression. Adjusted and unadjusted hazard ratios for incident carpal tunnel syndrome were calculated using three exposure categories: below the Action Limit, between the Action Limit and Threshold Limit Value, and above the Threshold Limit Value. Workers exposed above the 2001 Action Limit demonstrated significant excess risk of carpal tunnel syndrome, while the 2018 Threshold Limit Value®

---

**Corresponding Author:** Bradley A. Evanoff, Washington University School of Medicine, Division of General Medical Sciences, 4523 Clayton Avenue, Campus Box 8005, Saint Louis, MO 63110, bevanoff@wustl.edu, Phone: 314-454-8638.

**Disclaimer.** The findings and conclusions in this paper are those of the authors and do not necessarily represent the views of the National Institute for Occupational Safety and Health.

demonstrated significant excess risk only above the Threshold Limit Value. Of 186 total cases of carpal tunnel syndrome, 52 cases occurred among workers exposed above the 2001 Threshold Limit Value®, versus 100 among those exposed above the 2018 value. Eliminating exposures above the 2001 Threshold Limit Value® might have prevented 11.2% of all cases of carpal tunnel syndrome seen in our cohort, versus 25.1% of cases potentially prevented by keeping exposures below the 2018 value. The 2018 revision of the Threshold Limit Value® better protects workers from carpal tunnel syndrome, a recognized occupational health indicator important to public health. A significant number of workers are currently exposed to forceful repetitive hand activity above these guidelines. Public health professionals should promulgate these new guidelines and encourage employers to reduce hand intensive exposures to prevent carpal tunnel syndrome and other musculoskeletal disorders.

## Keywords

Upper Extremity Musculoskeletal Disorder; MSD Prevention; Occupational Guidelines; Risk Assessment

---

## Introduction

Carpal tunnel syndrome (CTS) caused by frequent, forceful hand exertions is one of the most common and costly work-related musculoskeletal disorders (MSD).<sup>[1]</sup> The American Conference of Governmental Industrial Hygienists (ACGIH®) develops voluntary workplace exposure indices, thresholds, and limits to prevent occupational injuries and illnesses due to exposure to chemical and physical agents. In 2001, a threshold limit value (TLV®) for Hand Activity was published to prevent MSD among workers performing repetitive single task jobs.<sup>[2]</sup> The TLV is meant to represent conditions under which nearly all workers may be repeatedly exposed without adverse health effects; if exceeded, the risk of MSD is elevated, and control measures (e.g., engineering or administrative controls) should be employed to reduce exposure. A lower threshold, the Action Limit (AL), identifies a ‘moderate’ risk exposure (See Figure 1A) and should trigger increased monitoring or surveillance to ensure health. Recent large studies from the US and Italy examined the risk of new cases of CTS for exposures above and below the 2001 TLV and AL, and concluded that these standards were not sufficiently protective of workers.<sup>[3–5]</sup> The ACGIH subsequently revised the TLV for Hand Activity® (See Figure 1B).<sup>[6]</sup> This study summarizes the effect of applying the 2018 TLV® versus the 2001 TLV® to data from an occupational cohort study.

## Methodology

### Analyzed US Worker Cohort

The revised 2018 TLV® and the 2001 version were evaluated to compare risk prediction for the AL and TLV thresholds using the same source data as Kapellusch et al. (2014).<sup>[4]</sup> Pooled data were obtained from six prospective cohort studies that investigated workplace risk factors of upper extremity MSDs. In brief, 4321 workers, recruited by six research teams in the US, were followed between 2001 and 2010. All study participants were full-time

employees, older than 18 years of age, who were employed in jobs that involved hand-intensive, often repetitive and forceful activities, and were employed in industries such as Manufacturing, Healthcare and Social Assistance, Services, and Construction industries.

### Case Definition of Carpal Tunnel Syndrome (CTS)

Incident CTS was defined as both (1) symptoms of tingling, numbness, burning or pain in the thumb, index finger or long finger, and (2) an abnormal electrodiagnostic studies (EDS) consistent with median neuropathy at the wrist.<sup>[7]</sup> The criteria for median neuropathy included: (1) peak median sensory latency >3.7 ms or onset median sensory latency > 3.2 ms at 14 cm, (2) distal median motor latency > 4.5 ms, and (3) transcarpal sensory difference > 0.85 ms (difference between median and ulnar nerve sensory latency across wrist). All electrodiagnostic values were temperature adjusted to 32°C. Workers lost to follow-up prior to developing CTS were censored as a non-case on the date the worker departed the study.

### ACGIH TLV for Hand Activity

The TLV® considers both applied hand force and repetition of hand exertions. Trained analysts directly observed and videotaped the workers performing their usual jobs. Normalized peak hand force (NPF) was rated using the Borg CR-10 rating scale,<sup>[2]</sup> and frequency of hand exertions was rated using the hand activity level (HAL) 0–10 scale.<sup>[8]</sup> For workers who performed multiple tasks and/or changed jobs over the follow-up period, a time-weighted-average (TWA) exposure was calculated to create a single exposure value that accounts for the proportion of daily work time in each observed task and proportion of job time during the follow-up period.

For each worker, the 2018 TLV® equations were used to calculate a corresponding threshold NPF for TLV [ $NPF_{TLV}$  (Eq 1.1)] and for AL [ $NPF_{AL}$  (Eq 1.2)].

$$NPF_{TLV} = 0.56 \times (10 - HAL) \quad (\text{Eq. 1.1})$$

$$NPF_{AL} = NPF_{TLV} - 2 \quad (\text{Eq. 1.2})$$

A peak force index (PFI) for TLV [ $PFI_{TLV}$  (Eq 2.1)] and for AL [ $PFI_{AL}$  (Eq 2.2)] were then calculated for each worker (See Figure 1B). If PFI for AL and for TLV was greater than 1.0, the respective limit was exceeded; in the case for  $PFI_{AL}$ , a negative ratio (<0) also indicated that AL was exceeded. A negative  $PFI_{AL}$  ratio occurred when  $HAL > 6.4$  at any value for  $NPF_{AL}$ .

$$PFI_{TLV} = NPF_{OBS} / NPF_{TLV} \quad (\text{Eq. 2.1})$$

$$PFI_{AL} = NPF_{OBS} / NPF_{AL} \quad (\text{Eq. 2.2})$$

Using both  $PFI_{TLV}$  and  $PFI_{AL}$ , workers were categorized into: (1) *below AL*, (2) *between AL and TLV*, and (3) *above TLV*. To determine exposure classifications using the 2001

TLV®, a score (Eq. 3) was calculated and subsequently categorized workers into: *below AL* (score  $< 0.56$ ), *between AL and TLV* (score between 0.56 and 0.78), and *above TLV* (score  $> 0.78$ ).

$$Score = NPF_{OBS}/(10 - HAL) \quad (\text{Eq. 3})$$

### Statistical Analysis

After excluding prevalent CTS cases at baseline, subjects with no follow-up measurements, and subjects with missing biomechanical exposure data, the cohort was reduced to 2751. Descriptive statistics were calculated for the 2001 and the 2018 TLV and AL classifications. Person-years and CTS incidence density rates were calculated for each exposure category. Proportions of CTS cases was compared for the three TLV categories between 2001 and 2018 TLVs® with the chi-square test. To estimate differences in the number of cases of CTS attributable to work exposures above the TLV, we compared unadjusted incidence rates of CTS (cases/100person-years) for those exposed above the 2001 and 2018 TLVs to those exposed below the TLVs. Based on these incidence rates we then calculated the Attributable Proportion of CTS related to exposure for cases occurring in workers exposed above the TLV. Time from study enrollment to first occurrence of CTS was modeled using Cox proportional hazard (PH) regression. Both unadjusted and adjusted (co-variables: age, BMI, gender, and research site) hazard ratios (HR) were calculated for incident CTS using the three exposure categories. Co-variables were selected *a priori* as personal risk factors associated with increasing risk of developing CTS.<sup>[3-5, 7]</sup> All analyses were performed using SAS (version 9.4, SAS Institute, Cary, NC).

### Results

From the 2751 worker cohort, there were 6282 person-years of observation time over a maximum 6.4 years of follow-up (See Table 1). There were 186 incident cases of CTS with an overall incident density rate of 2.96 cases per 100 person-years. Application of the 2018 exposure recommendations markedly changed the categorization of exposures within our worker cohort. 42.4% of workers were classified as above the 2018 TLV vs. 23.2% above the 2001 TLV; 23.6% of workers were exposed below the 2018 AL vs. 57.6% of workers below the 2001 AL; and 34% vs. 19.2% classified as between the AL and TLV in 2018 and 2001, respectively. The 2001 and 2018 TLVs® had significantly different proportions of incident CTS cases within the three TLV categories ( $\chi^2(2) = 37.851$ ,  $p < 0.05$ ), reflecting fewer cases of CTS occurring below the 2018 TLV and AL, and more cases occurring above the TLV. For exposures below the AL, the 2018 threshold identified 52 fewer cases compared to the 2001 threshold (2001 TLV®: 86 CTS cases; 2018 TLV®: 34 cases). Below the TLV, the 2018 threshold identified 48 fewer cases compared to the 2001 threshold (2001 TLV®: 134 total CTS cases; 2018 TLV®: 86 total cases). Above the TLV, the 2018 threshold identified 48 more cases than the 2001 threshold (2001 TLV®: 52 CTS cases; 2018 TLV®: 100 CTS cases). Attributable proportion of CTS related to exposure above the TLV was 0.401 for the 2001 TLV and 0.467 for the 2018 TLV. This suggests that eliminating exposure above the TLV for all workers would have prevented 20.9 cases of CTS (11.2% of

all cases in the cohort) using the 2001 limit, and 46.7 cases (25.1% of cases) using the 2018 TLV. For both the 2001 and 2018 recommendations, further reductions would be achieved by eliminating exposures above the AL.

Proportional hazard models showed that risk of incident CTS decreased sharply for exposures below the TLV and exposures below the AL when comparing 2018 values to those from 2001 (see Table 2). In particular, large and statistically significant risk of CTS occurred above the AL but below the TLV when using the 2001 exposure recommendations (HR of 1.88, c.i. 1.30–2.72); under the 2018 TLV and AL, significant excess risk was seen only above the TLV.

## Discussion

Work-related MSDs remain a burden in the United States, with substantial employer and societal costs and impacts on the affected individual's quality of life. For example, estimated medical care costs of CTS exceed \$2 billion annually,<sup>[9]</sup> a number that does not include disability and other social costs. Importantly, CTS has the second highest rate of opioid prescribing by injury type among workers treated in under workers compensation.<sup>[10]</sup> Treatment of work-related disorders with prescription opioids has contributed to the current epidemic of opioid-related deaths,<sup>[11]</sup> with higher rates seen in industries with the highest injury rates.<sup>[12]</sup> Risk assessment and control methods exist for work-related MSDs, including ACGIH Threshold Limit Values® designed to prevent upper extremity MSDs such as CTS. This study evaluated the recommended 2018 TLV® as thresholds for risk factors of CTS using a large pooled cohort representing a diverse workforce in dozens of occupations and industries in the United States. The 2001 TLV® demonstrated little risk difference between exposures above the TLV and between the AL and the TLV when compared to exposures below the AL, whereas the 2018 TLV® demonstrated significant risk only above the TLV. Importantly, a substantial number of workers in our cohort were exposed above the 2001 and 2018 TLVs recommended by ACGIH, indicating an ongoing need to reduce exposures to prevent CTS and other upper extremity MSDs among workers in hand-intensive jobs.

In practical terms, if the 2018 TLV® was used for surveillance on this cohort, and if subsequent ergonomic interventions had been performed at the outset of the study to reduce exposure to those above the AL, then the 2018 TLV® might have prevented 28% of CTS cases that occurred below the 2001 AL. Similarly, 26% of workers classified as below TLV by the 2001 thresholds would have been classified as above TLV by the 2018 threshold and might also have been prevented. Thus, these hypothetical biomechanical exposure reducing interventions might have lowered the incidence rate of CTS for the cohort by 28% from 2.96 to 2.11 per 100 person-years.

The findings from this report are subject to at least two limitations. First, at baseline, workers reported an average of 7.6 years (SD = 8.6) of tenure within their company, and are likely representative of a survivor population. Therefore, study findings probably underestimate risks of exposure and the effect sizes may be somewhat lower than what would have been found in a population of newly hired workers. Second, workers who

changed jobs over the follow-up period were assigned a time-weighted-average for peak force and HAL measurements, which could have resulted in non-differential misclassification of exposure. In a sub-analysis, a single TWA value across multiple jobs over the follow-up period was found to be a suitable indicator of cumulative exposure. Job variability did not appear to affect our interpretation of the exposure-response associations.

## Conclusion

CTS is a recognized occupational health indicator of the working population important to public health.<sup>[13]</sup> The 2018 TLV® is demonstrated to improve the protection of workers who perform hand intensive tasks from risk of CTS. Adherence to the 2018 TLV® might have prevented 28% of CTS cases that occurred below the 2001 AL; 26% of CTS cases classified as below TLV by the 2001 thresholds would have been classified as above the TLV by the 2018 threshold. Many workers in hand intensive industries are still exposed above recommended limits.

## Recommendations

As part of a robust occupational health and safety program, adhering to the 2018 TLV® could be a key primary prevention strategy to reduce the public health burden of carpal tunnel syndrome and other upper extremity MSDs. The revised 2018 TLV® was an effective tool for predicting risk of CTS in our cohort, and should thus be used to trigger control measures when these limits are exceeded. When the AL or TLV are exceeded, employers should make changes (e.g., engineering controls and administrative controls) to the work environment to reduce workers' exposure to forceful, repetitive hand activities.

## Acknowledgments.


This study was supported by research funding from the Centers for Disease Control and Prevention/National Institute for Occupational Safety and Health (R01OH009712).

## References

1. Foley M, Silverstein B, and Polissar N: The economic burden of carpal tunnel syndrome: long-term earnings of CTS claimants in Washington State. *American Journal of Industrial Medicine* 50(3): 155–172 (2007). [PubMed: 17216630]
2. ACGIH: Threshold limit values for chemical substances and physical agents and biological exposure indices. Cincinnati, OH: American Conference of Governmental Industrial Hygienists, 2001.
3. Bonfiglioli R, Mattioli S, Armstrong T, Graziosi F, Marinelli F, A F et al.: Validation of the ACGIH TLV for hand activity level in the OCTOPUS cohort: a two-year longitudinal study of carpal tunnel syndrome. *Scandinavian Journal of Work, Environment and Health* 39(2): 155–163 (2013).
4. Kapellusch JM, Gerr FE, E. Malloy J, Garg A, Harris-Adamson C, Bao SS et al.: Exposure-response relationships for the ACGIH threshold limit value for hand-activity level: results from a pooled data study of carpal tunnel syndrome. *Scandinavian Journal of Work, Environment and Health* 40(6): 610–620 (2014).
5. Violante FS, Armstrong TJ, Fiorentini C, Graziosi F, Risi A, Venturi S et al.: Carpal tunnel syndrome and manual work: a longitudinal study. *Journal of Occupational and Environmental Medicine* 49(11): 1189–1196 (2007). [PubMed: 17993922]
6. Rempel D: 1631c Recent changes to the ACGIH hand activity level TLV. *Occupational and Environmental Medicine* 75(Suppl 2): A258–A258 (2018).

7. Dale AM, Harris-Adamson C, Rempel D, Gerr F, Hegmann K, Silverstein B et al.: Prevalence and incidence of carpal tunnel syndrome in US working populations: pooled analysis of six prospective studies. *Scandinavian Journal of Work, Environment and Health* 39(5): 495–505 (2013).
8. Latko WA, Armstrong TJ, Foulke JA, Herrin GD, Rabourn RA, and Ulin SS: Development and evaluation of an observational method for assessing repetition in hand tasks. *American Industrial Hygiene Association Journal* 58(4): 278–285 (1997). [PubMed: 9115085]
9. Falkiner S, and Myers S: When exactly can carpal tunnel syndrome be considered work-related? *ANZ Journal of Surgery* 72(3): 204–209 (2002). [PubMed: 12071453]
10. Thumula V, and Liu T-C: “Correlates of Opioid Dispensing”. Cambridge, MA: Workers Compensation Research Institute, 2018.
11. Cheng M, Sauer B, Johnson E, Porucznik C, and Hegmann K: Comparison of opioid-related deaths by work-related injury. *American Journal of Industrial Medicine* 56(3): 308–316 (2013). [PubMed: 23143851]
12. Massachusetts Department of Public Health, Bureau of Substance Addiction Services, Injury Surveillance Program, and Office of Special Analytic Projects: “Opioid-related Overdose Deaths in Massachusetts by Industry and Occupation, 2011–2015”, 2018.
13. NIOSH-CSTE Surveillance Planning Work Group: “The Role of the States in a Nationwide, Comprehensive Surveillance System for Work-Related Diseases, Injuries, and Hazards”: NIOSH Surveillance Coordinating Group, 2001.





**Figure 1.**  
ACGIH TLV for hand activity regression equations for threshold limit value and action limit for (A) 2001 guidelines and (B) revised 2018 guidelines

Author Manuscript  
Author Manuscript  
Author Manuscript  
Author Manuscript

**Table 1.**

Descriptive statistics and incidence density rates for ACGIH® categories for 2001 and 2018 guidelines.

| Variable                                | Mean | (SD)   | Subjects | CTS Cases | Person-Years | Incidence Rate Per 100 person years<br>(95% CI) |
|---|------|--------|----------|-----------|--------------|---|
| <b>Total Cohort</b>                     |      |        | 2751     | 186       | 6282         | 2.96 (2.55 – 3.42)                              |
| Age (years)                             | 39.6 | (11.5) |          |           |              |   |
| Gender                                  |      |        |          |           |              |   |
| Male                                    |      |        | 1351     | 71        |              |   |
| Female                                  |      |        | 1400     | 115       |              |   |
| BMI (kg/m <sup>2</sup> )                | 28.4 | (6.1)  |          |           |              |   |
| NPF <sub>Obs</sub> (Borg CR-10) – TWA   | 2.8  | (1.7)  |          |           |              |   |
| HAL (0–10) – TWA                        | 4.3  | (1.8)  |          |           |              |   |
| <b>ACGIH (2001) Exposure Categories</b> |      |        |          |           |              |   |
| < AL                                    |      |        | 1585     | 86        | 3879         | 2.22 (1.77 – 2.74)                              |
| AL and TLV                              |      |        | 529      | 48        | 1219         | 3.94 (2.90 – 5.22)                              |
| > TLV                                   |      |        | 637      | 52        | 1185         | 4.39 (3.28 – 5.76)                              |
| <b>ACGIH (2018) Exposure Categories</b> |      |        |          |           |              |   |
| < AL                                    |      |        | 649      | 34        | 1618         | 2.11 (1.46 – 2.94)                              |
| AL and TLV                              |      |        | 936      | 52        | 2261         | 2.30 (1.72 – 3.02)                              |
| > TLV                                   |      |        | 1166     | 100       | 2404         | 4.16 (3.39 – 5.06)                              |

**Table 2.**

Crude and adjusted associations between TLV for HAL® categories and incident CTS

| Variable  | 2001 ACGIH for Hand Activity® |             |         | 2018 ACGIH HA TLV® |             |         |
|---|-------------------------------|-------------|---------|--------------------|-------------|---------|
|   | HR                            | (95%CI)     | p>ChiSq | HR                 | (95%CI)     | p>ChiSq |
| <b>Unadjusted</b>                                   |                               |             |         |                    |             |         |
| < AL  | 1.00                          |             |         | 1.00               |             |         |
| AL and TLV  | 1.80                          | (1.27–2.57) | 0.0011  | 1.12               | (0.72–1.72) | 0.6171  |
| > TLV   | 2.01                          | (1.41–2.84) | <0.0001 | 2.03               | (1.37–3.00) | 0.0004  |
| <b>Adjusted for BMI, age, gender, research site</b> |                               |             |         |                    |             |         |
| < AL  | 1.00                          |             |         | 1.00               |             |         |
| AL and TLV  | 1.88                          | (1.30–2.72) | 0.0007  | 1.16               | (0.73–1.85) | 0.5225  |
| > TLV   | 1.73                          | (1.20–2.49) | 0.0034  | 1.99               | (1.28–3.10) | 0.0021  |

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript