

UCLA

UCLA Previously Published Works

Title

The Role of Xist in X-Chromosome Dosage Compensation

Permalink

<https://escholarship.org/uc/item/95k9v9vm>

Journal

Trends in Cell Biology, 28(12)

ISSN

0962-8924

Authors

Sahakyan, Anna

Yang, Yihao

Plath, Kathrin

Publication Date

2018-12-01

DOI

10.1016/j.tcb.2018.05.005

Peer reviewed



HHS Public Access

Author manuscript

Trends Cell Biol. Author manuscript; available in PMC 2019 December 01.

Published in final edited form as:

Trends Cell Biol. 2018 December ; 28(12): 999–1013. doi:10.1016/j.tcb.2018.05.005.

The role of *Xist* in X-chromosome dosage compensation

Anna Sahakyan, Yihao Yang, and Kathrin Plath*

David Geffen School of Medicine, Department of Biological Chemistry, Eli and Edythe Broad Center of Regenerative Medicine and Stem Cell Research, Jonsson Comprehensive Cancer Center, Molecular Biology Institute, University of California Los Angeles, Los Angeles, CA 90095, USA

Abstract

In each somatic cell of a female mammal one X chromosome is transcriptionally silenced via X-chromosome inactivation (XCI), initiating early in development. Although XCI events are conserved in mouse and human *post*-implantation development, regulation of X-chromosome dosage in *pre*-implantation development occurs differently. In pre-implantation development, *mouse* embryos undergo imprinted form of XCI, yet *humans* lack imprinted XCI and instead regulate gene expression of both X-chromosomes by dampening transcription. The long non-coding RNA *Xist/XIST* is expressed in mouse and human pre- and post-implantation development to orchestrate XCI, but its role in dampening is unclear. In this review, we discuss recent advances in our understanding of the role of *Xist* in X-chromosome dosage compensation in mouse and human.

Keywords

Xist; X-inactivation; embryonic stem cells; lncRNAs; dosage compensation

X chromosome dosage compensation

Male (XY) and female (XX) **eutherian** (see Glossary) mammals have equivalent expression levels of most X-chromosome genes despite the presence of an extra X chromosome in females. This X-chromosome dosage compensation is due to the phenomenon of X-chromosome inactivation (**XCI**), which refers to the transcriptional silencing and **heterochromatinization** of one of the two X chromosomes in females early in embryonic development [1]. Most of our knowledge of XCI is based on mouse studies, where two types of XCI exist: imprinted and random (Figure 1A).

In imprinted XCI, which initiates in all cells of the female mouse four- to eight-cell stage pre-implantation embryo, the paternally inherited X chromosome (X_p) undergoes

*Correspondence: kplath@mednet.ucla.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

inactivation, while the maternally inherited X chromosome (X_m) remains active [2,3]. As pre-implantation development progresses to form the blastocyst, cells of the trophectoderm layer, which give rise to extra-embryonic tissues (e.g. the placenta), maintain their imprinted XCI state. In contrast, epiblast cells of the blastocyst, which give rise to the embryo proper, reactivate the inactive X_p, re-establishing a state with two active X-chromosomes. The biallelic X-linked gene expression of epiblast cells is resolved again via XCI, but in this second wave of XCI either the X_p or the X_m is chosen at random for inactivation. Random XCI is maintained in all descendent somatic cells throughout life, resulting in adult mice that are a mosaic of cells expressing either maternal or paternal alleles of X-linked genes [4,5]. A group of X-linked genes express both the maternal and the paternal allele in each cell since these genes **escape** XCI and are thus the exception to the rule (reviewed by [6]). The chromosome-wide inactivation of the X chromosome, both in imprinted and random XCI, appears to always be governed by the **lncRNA** X inactive specific transcript (**Xist**), which is encoded in the X-inactivation center (XIC) of the X chromosome [7].

The occurrence of both imprinted and random XCI in the same species, as is the case in mouse, may not be very common. Most mammals studied utilize only one form of XCI for X-chromosome dosage compensation. In marsupials, only imprinted XCI is observed where the X_p is exclusively chosen for inactivation [8] (Figure 1C). Contrary to this, imprinted XCI does not occur in rabbit, pig, [9,10] horse, or human development [11,12,13] based on analysis of pre-implantation blastocysts [9,10,13] or placental tissues [11,12]. In human post-implantation development, both extra-embryonic and embryonic lineages dosage-compensate via random XCI [12,14] (Figure 1B). However, in the first week of human development, prior to implantation and XCI, the existence of a novel gene-dosage regulation has recently been uncovered [13]. Here, both X chromosomes remain active from the onset of zygotic gene activation until the blastocyst stage [9,13] (Figure 1B). However, transcription from both X chromosomes is tuned down, or dampened, resulting in a net reduction of X-linked gene expression in female blastocyst cells [13]. X-chromosome dampening (**XCD**) has not been observed in any other mammal yet, but it has been reported in the nematode *Caenorhabditis elegans* [15], although the underlying mechanism in human and nematode may differ. In the XX hermaphrodite *C. elegans*, the 3D conformation of the X chromosomes is remodeled to reduce chromosome-wide gene expression by half in order to achieve gene-expression balance between XX hermaphrodites and XO males [16]. 3D chromosome conformation also differs between the active and inactive X chromosomes in mammals [17], suggesting that chromosome conformation remodeling might also be at play in human XCD. However, unlike the mammalian inactive X-chromosome (**Xi**), lncRNAs have not been reported to regulate X-chromosome dose in the *C. elegans*. Instead, dampening in the nematode is carried out by the dosage compensation complex, a condensin-containing multi-subunit protein assembly that binds at multiple sites along the X chromosome and leads to chromosome-wide compaction and gene repression [18].

Whether XCD observed in human pre-implantation development and *C. elegans* are related at the molecular level needs further investigation. Moreover, when single cell RNA-sequencing data (Box 1) of pre-implantation human blastocysts are analyzed using different bioinformatics tools and approaches, the dosage compensation observed in human pre-

implantation embryos has been interpreted as initiation of XCI rather than dampening of both X chromosomes [19,20]. Fortunately, naïve human embryonic stem cells (hESCs), which are the *in vitro* counterparts of the pluripotent cells in the human pre-implantation embryo, exhibit XCD and thus can be used as a model system to address XCD and its relationship to the initiation of XCI further [21].

Box 1

Methods

Sox2 promoter-driven Cre recombinase: the Cre-recombinase enzyme, which recombines DNA at specific DNA sequences, coined loxP sites, is expressed only in cells where the *Sox2* gene is actively transcribed. This system is engineered by cloning the *Sox2* promoter and enhancer region upstream of the Cre recombinase gene. When the same cell contains DNA sequences flanked with loxP sites of the same orientation, the intervening sequence will be removed – or deleted.

Tetraploid complementation assay is a method for creating mice where all the cells of the embryo proper are derived from mouse pluripotent stem cells (PSCs) upon blastocyst injection of these cells. To accomplish this, cells of two-cell embryos are fused experimentally to form a tetraploid pre-implantation embryo. While the tetraploid cells of the blastocyst can develop into the extraembryonic tissues required for in utero development, they cannot contribute to the embryo proper. Hence, when such tetraploid blastocysts are injected with diploid mouse PSCs, all the tissues of the embryo proper come from these injected diploid cells, creating a non-chimeric mouse.

Single cell RNA-sequencing is a method of measuring global gene expression from individual cells to determine gene expression patterns unique to individual cells that would be otherwise be lost in bulk RNA-sequencing (from a group of cells) due to averaging of the data.

In this review, we discuss the experimental evidence examining the role of *Xist* in X-chromosome dosage compensation via imprinted and random XCI in mouse. We also consider *XIST* function in early human development and in human **pluripotent stem cells** (PSCs), reflecting on potential molecular mechanisms which might regulate context-dependent *XIST* function.

Long non-coding RNAs are key players in X-chromosome regulation

An intriguing fact about X-chromosome dosage regulation in all mammals is the utilization of long non-coding RNAs (lncRNAs) such as *Jpx*, *Ftx*, *Tsix*, and *XACT*, most of which are located in the XIC of the X chromosome [20,22,23–28] (Figure 2). When expressed, *Xist* is exclusively associated with the X-chromosome from which it is expressed, acting only *in cis* [29]. The lncRNAs *Jpx* and *Ftx* exert their function by acting as activators of *Xist* to fine-tune *Xist* expression and thus indirectly regulate XCI. The *Jpx* lncRNA product acts either *in cis* or *in trans* [23] and binds the *Xist* repressor CTCF, taking away repression of *Xist* transcription [22]. Contrary to this, the *Ftx* transcript itself is not required for *Xist*

regulation: it is the act of transcription of the *Ftx* locus that leads to *Xist* expression *in cis* [25]. Moreover, in mice, the lncRNA *Tsix*, which is transcribed in anti-sense orientation to *Xist*, represses *Xist* expression, thus ensuring *Xist* induction on the Xi and protecting the **Xa** (active X chromosome) from ectopic silencing by *Xist* [26,27]. *Tsix* is not expressed in human pre-implantation development [13], and the roles of *Jpx* and *Ftx* are yet to be examined in humans. A recently discovered lncRNA called **XACT** (X active coating transcript) that is unique to human PSCs, but does not reside in the XIC, may aid in maintaining transcriptional activity of the X chromosome from which it is expressed by counteracting *XIST* [20,28]. Interestingly, correlative studies suggest that *XIST* RNA, in addition to XCI, might also mediate the dampening of the transcriptional output of both X chromosomes of female human pre-implantation blastocysts [13,21] (discussed below). While *Xist* is unique to placental mammals, marsupials also use a cis-acting lncRNA encoded on the X chromosome, termed *Rsx* (RNA-on-the-silent X), which in many ways appears to act like *Xist* in XCI [8] (Figure 1). Taken together, it is rather interesting that different lncRNAs have evolved to regulate gene expression chromosome-wide *in cis*. This is perhaps due to the unique ability of lncRNAs to bind distant sites on chromatin while still tethered to their transcription loci. *Xist* [30] and other lncRNAs such as *HOTTIP* (HOXA transcript at the distal tip), which is encoded on mouse chromosome 6 and activates genes in its neighborhood [31], reach their target chromatin sites simply by proximity – by being close to these sites in 3D space due to the folding of chromatin within the nucleus (reviewed and illustrated in [32]). Understanding how X-chromosome dosage is regulated via lncRNAs will thus not only shed light onto X-chromosome biology but can serve as a starting point in understanding how lncRNAs localize to and act on chromatin in general.

***Xist* is required for XCI in mouse**

Xist is the best studied lncRNA to date. As the name suggests, *XIST* was discovered due to its association with the inactive X-chromosome – X inactive specific transcript [33–37]. The requirement of *Xist* for XCI was first directly implicated using female mouse embryonic stem cells (mESCs) with a mutated *Xist* gene lacking the first two thirds of exon 1 [38]. When induced to differentiate, both *in vitro* and *in vivo* (using aggregation chimeras), the X chromosome bearing the mutant truncated *Xist* gene was always spared from inactivation while *Xist* was expressed from the wild type X chromosome, causing non-random silencing [38]. This study demonstrated that *Xist* is required for choosing the chromosome for inactivation, and suggested, without direct evidence, that *cis*-expression of *Xist* is required for XCI.

Since mESCs are derived from the epiblast cells of the blastocyst, the role of *Xist* in pre-implantation development, from zygote to blastocyst formation, cannot be studied using these cells. Therefore, to further investigate the role of *Xist* in early development, mESCs were used to generate chimeric mice containing cells with a large deletion of *Xist*, which were then mated with wild type mice to generate hemizygote males or heterozygote females [39]. When the mutant *Xist* was inherited from the mother, both normal female and male pups were born. However, female embryos with a paternally-inherited mutant *Xist* had severe prenatal growth defects and survived until approximately embryonic day 10.5 (E10.5). In these embryos the extraembryonic tissues failed to develop due to the lack of

imprinted XCI of the paternally inherited X chromosome in these tissues [39] (Figure 3). Due to the pre-determined choice of the paternal X chromosome for silencing in imprinted XCI, this experiment is the first clear demonstration that XCI, specifically imprinted XCI, cannot be initiated without *Xist*. Note that there is no paternal inheritance of mutant *Xist* to be studied in males since males inherit a Y chromosome but no X chromosome from their father. The inability of the extraembryonic tissue to support the growth of the embryo in the absence of X-chromosome dosage compensation was investigated further and this failure was attributed to exhaustion of the extra-embryonic ectoderm due to premature cell differentiation [40].

A recent study used single cell RNA-sequencing to provide high temporal and chromosome-wide resolution of X-linked gene silencing in mouse pre-implantation development and its dependence on *Xist* RNA [3]. Comparisons of female wild type to mutant embryos carrying a paternal *Xist* deletion at the 8-, 16-, 32-cell, and blastocyst stages validated the need for *Xist* in initiating imprinted XCI [3], which was also demonstrated recently using RNA-sequencing of single embryos [2]. Such high-resolution data were important to clarify the role of *Xist* in imprinted XCI and to rule out prior arguments for an *Xist*-independent imprinted XCI [41]. The dynamics of silencing during the initiation of imprinted XCI revealed that genes are silenced with different kinetics. Genes silenced at an earlier stage of pre-implantation development were those near the XIC (where the *Xist* gene is located) in 3D space [3,30]. Intriguingly, this recapitulates the finding that at the initiation of random XCI, *Xist* first contacts the sites of the chromosome closest to its site of transcription in 3D, rather than linear space [30]. This correlation between the kinetics of imprinted XCI and the proximity to the *Xist* locus in 3D space independently supports the notion that imprinted XCI is mediated by *Xist*, and additionally suggests that the mechanism of *Xist* spreading *in cis* during the initiation of imprinted and random XCI is conserved.

Requirement of XCI in the development of the mouse embryo proper

Although the above-mentioned paternal *Xist* deletion experiment demonstrated that *Xist* is required for imprinted XCI, the early embryonic lethality due to malfunctioning extraembryonic tissues have made addressing the requirement of *Xist* in the embryo itself unfeasible with a germline mutation of *Xist*. A homozygous *Xist* knockout is required to address this question, which, when using a germline mutation, affects imprinted XCI as well (Figure 3). However, the maternal germline deletion of *Xist* allele was valuable to demonstrate the requirement of a functional *Xist* allele for choosing the X chromosome for XCI *in vivo* [42], extending the prior mESC study [38]. In the meantime, experiments demonstrating the sufficiency of *Xist* RNA for silencing was demonstrated. Particularly, ectopic expression of *Xist* cDNA from autosomes or the X chromosome [43,44], or activation of the endogenous *Xist* allele from the single X chromosome in male mESCs with an inducible promoter [30] demonstrated that *Xist* expression is sufficient to cause silencing *in cis*. These gain-of-function experiments also opened the way for dissection of *Xist* RNA for its functional units [43].

Studying the need for *Xist* in random XCI requires a unique approach that specifically deletes *Xist* in embryonic tissues. A recent study took on the challenge to assess the

importance of X-chromosome dosage compensation by random XCI in mouse embryonic development. To silence *Xist* specifically in the embryo while sparing the extra-embryonic tissue, *Xist* was conditionally deleted in the epiblast lineage [45]. While most mutant mice died *in utero*, surprisingly some mice survived to term, but exhibited growth retardation with reduced body size, dying within one month after birth (Figure 3). Only one mouse, which was a mosaic of XX and XO cells – cells that had lost one of their two X chromosomes – survived to adulthood. These data indicate that *Xist* and random XCI are required for normal embryonic development. However, given the survival to term, the phenotype appeared much weaker than expected and was a surprise – since XCI occurs soon after implantation the expectation was that *Xist* loss should have led to early embryonic lethality. This can perhaps be explained by another unexpected observation, namely that X-chromosome dosage compensation was not completely wiped out upon deletion of *Xist*. The authors reported partial X-chromosome dosage compensation in the absence of *Xist*, concluding that an *Xist*-independent mechanism was responsible [45].

This phenotype of partial X-chromosome dosage compensation in *Xist* mutant mice was observed due to less-than-expected increase in average expression levels of all genes by RNA-sequencing when compared to wild-type female mice, and due to the presence of mono-allelic gene expression in some, but not all cells at the single cell level of a few X-linked genes, assessed by fluorescent *in situ* hybridization [45]. However, understanding the *Xist* mutant mouse model system used [45] can perhaps better explain these above-mentioned observations. Deletion of *Xist* in the epiblast lineage was accomplished with the Sox2 promoter-driven Cre recombinase [45] (Box 1). However, the efficiency of *Xist* excision by the Sox2-driven Cre was not measured in the developmental interval around the induction of random XCI [46], as embryos were only harvested at E8.5 to confirm *Xist* deletion, a time point at which XCI has already occurred [45]. Originally, Sox2-driven Cre recombinase activity has been shown to occur in blastocyst outgrowths in culture and in all cells of the epiblast in E6.5 embryos *in vivo*, by assessing the removal of a ‘stop’ cassette in front of a beta-galactosidase reporter gene integrated into the ROSA26 locus [47,49]. Together, these findings suggest that the ROSA26 reporter recombined before E6.5 and possibly before XCI would be initiated *in vivo*, indicating that the Sox2-driven Cre-recombinase may be ideally suited to delete *Xist* before induction of random XCI. However, since chromatin accessibility may be different for the *Xist* locus in comparison to ROSA26, a region on mouse chromosome 6 identified because of its high recombination frequency [49], the Cre-mediated excision kinetics may differ for the two loci and thus need to be independently determined for the *Xist* locus. Additionally, the DNA segment flanked by LoxP sites, which need to come together for Cre-mediated recombination, is significantly longer in the *Xist* deletion construct [45] compared to the ROSA26 beta-galactosidase reporter system [48], and therefore potentially less favorable for deletion, again arguing for the need to establish *in vivo* *Xist* deletion kinetics in this mouse model. Hence, it cannot be ruled out that *Xist* deletion may have occurred *after* initiation/completion of random XCI.

Previous studies have shown that deletion of *Xist* has no dramatic *short-term* effect on the silent state of genes on the X chromosome when it occurs after the inactive X chromosome is fully established [50,51] (see below). Depending on the proportion of cells undergoing random XCI prior to *Xist* deletion, the embryos would then survive to term and demonstrate

incomplete dosage compensation at organism level due to the mixture of XaXa and XaXi cells (Xa is the active X chromosome, Xi is the inactive X chromosome). *Xist* deletion post XCI establishment would also explain the low number of pups surviving to term, since if not enough cells per embryo undergo X-chromosome dosage compensation prior to *Xist* deletion, the embryo would not be viable, arguing for the importance of *Xist* in XCI and embryonic development. Taken together, additional experiments with a temporally precise *Xist* deletion (and confirmation of *Xist* deletion before induction of random XCI) are required to dissect whether a novel embryonic X-chromosome dosage compensation mechanism in the absence of *Xist* or the delayed deletion of *Xist* relative to the onset of XCI explain the surprisingly weak consequences of the current *Xist* deletion in the embryo [45]. For instance, using homozygous *Xist* knockout mESCs in tetraploid complementation assays [52] (Box 1) could provide wild-type extraembryonic tissues capable of supporting normal development while all cells of the epiblast, derived from mESCs, would lack *Xist*, ruling out the possibility of random XCI occurring in any fraction of the cells. It would be interesting to see if/when development would fail in this scenario, but our hypothesis is that no viable pups would be obtained from such mice. Regardless, the current data argue that X-chromosome dosage compensation mediated by *Xist* is critical for embryonic development.

The influence of XCI on the developmental potential of female cells has been shown with mESCs, as the double dose of X-linked genes delays the differentiation of these cells due to its stabilizing effect on the naïve pluripotent state [55]. This stabilization is achieved via inhibition of the MAPK and Gsk3 pathways and stimulation of the Akt pathway, and XCI is needed to properly exit **naïve pluripotency** [53]. The delayed exit from **pluripotency** in the presence of two active X chromosomes may also occur *in vivo* [53], since embryos with a single X chromosome undergo accelerated development [54].

***Xist* executes XCI by recruiting a diverse set of proteins**

During initiation of XCI, *Xist* recruits numerous silencing factors to the X chromosome to establish **facultative heterochromatin**, also known as the Barr body [55]. This is accompanied by epigenetic changes including substitution of certain core histones, covalent modifications of histone tails, and promoter CpG methylation (reviewed by [1]). Although some of these *Xist*-induced epigenetic remodeling steps were discovered years ago, most of the proteins binding *Xist* directly and indirectly were identified only recently using mass spectrometry-based approaches and genetic screens [56–60]. While some of the *Xist*-binding proteins influence histone modifications (via the activation of the histone deacetylase HDAC3 through the engagement of SPEN by the 5' end of *Xist*) [57] or nuclear positioning of the *Xist*-coated Xi (through the binding of lamin B receptor (LBR) to *Xist*) [61], others induce RNA modification of adenosine methylation (m6A) on *Xist* to influence its silencing ability [62]. Identification of the *Xist* interactome – i.e. the proteins that directly or indirectly bind to *Xist* – has created a newfound appreciation of the multiple roles *Xist* plays in XCI, spanning from orchestrating chromosome-wide silencing, localizing itself to chromatin, altering chromatin state, and remodeling the 3D chromosome architecture, recently reviewed in [55]. While the primary sequence of *Xist* RNA is not well conserved between mouse and human, the gene structure (exons/introns) as well as the presence of key repeat regions [37] are conserved in the two species. Some of these repeat regions are important for *Xist*

function since they are the sites where proteins that directly interact with *Xist* bind to [56,57,63]. Therefore, although the mass-spectrometry based unbiased approaches have identified the *mouse Xist* interactome [56–58], it is safe to predict that the *humanXIST* interactome will largely overlap with that of mouse [64].

***Xist* is required for long-term maintenance of random XCI**

The Xi with *Xist* expression remains inactive in all somatic progeny of cells. In short-term *in vitro* studies (days – weeks), *Xist* does not seem to play a major role in the maintenance of the silent status of genes in random XCI in somatic cells [44,50]. This appears to also be the case in maintenance of imprinted XCI in the vole *Microtus Levis*, where ablation of *Xist* expression via deletion of its promoter region in trophoblast stem cells, which have already undergone imprinted XCI, does not lead to transcriptional reactivation or loss of repressive chromatin marks of the Xi [65]. However, a longer-term *in vivo* mouse study suggests that the prolonged absence of *Xist* in mice, initiated in the blood lineage using a tissue-specific Cre-recombinase, induces at least partial reactivation of genes on the X chromosome [51]. Notably, the experimentally induced deletion of *Xist* in hematopoietic cells in mice results in poor postnatal survival and development of **myelodysplasia** and various cancers of the blood with 100% penetrance [51] (Figure 3). The inevitable development of cancer in the absence of *Xist* clearly labels *Xist* as a potent tumor-suppressor, most likely due to its requirement in the maintenance of gene silencing in somatic cells. In agreement with this *in vivo* mouse study, abnormal reactivation of the Xi has also been reported in human breast cancer cells, although here an extra dose of X-linked genes is either due to Xi **erosion** or loss of an Xi combined with an Xa duplication [66]. However, the connection between *XIST*-dependent maintenance of XCI and cancer formation in humans needs to be further explored. Since the importance of *Xist* in maintaining XCI only became obvious from mouse *in vivo* studies, it is critical to address the role of *XIST* in human cancers with carefully designed experiments.

There are two instances in mouse development that require reactivation of the Xi: once in cells of the inner cell mass (ICM) of the blastocyst, when imprinted XCI needs to be reversed prior to induction of random XCI, and once more in the development of primordial germ cells, prior to meiosis [67]. In both cases, shutdown of *Xist* expression from the existing Xi precedes the removal of chromosome-wide transcriptional repression [68,69]. **Xi reactivation** is also observed *in vitro*, when female mouse somatic cells such as mouse embryonic fibroblasts are reprogrammed to form induced pluripotent stem cells (**iPSCs**). Whereas the necessity of *Xist* loss in Xi reactivation is difficult to address in the *in vivo* scenarios described above, the *in vitro* **reprogramming** system has allowed detailed studies of this relationship. Using ectopic maintenance of expression or the deletion of *Xist* in a reprogramming experiment, it was demonstrated that *Xist* loss is necessary, but not sufficient, for Xi reactivation in iPSC generation, reviewed in more detail in [67]. During reprogramming to iPSCs, Xi reactivation is one of the last steps of reprogramming, requiring DNA demethylation in addition to *Xist* RNA loss [70]. Similarly, in somatic cells, DNA demethylating agents, such as 5-aza-dC, induce reactivation of genes of the somatic Xi, albeit in a small proportion of cells, via induction of global DNA demethylation [71,72]. DNA methylation works in synergy with *Xist* RNA and **histone hypoacetylation** [71], as

well as the H3K9 trimethylation pathway [72,73] in maintaining the inactive state of the somatic Xi. In fact, it takes the synergism of triple-drug combinations targeting DNA methylation, topoisomerase activity (involved in relieving torsional stress during DNA replication and transcription) combined with knockdown of an *Xist*-interacting protein to obtain dramatic re-activation of the Xi, and even then the re-activation is not for all silenced genes [58]. Complete chromosome-wide reactivation of all silenced X-linked genes in somatic cells has not been reported thus far, highlighting the unbreachable nature of the multiple epigenetic layers protecting the Xi.

Xi reactivation in human pluripotent stem cells

Studies of the relationship between *XIST* and Xi reactivation in human cells are not as straightforward as in mouse, mainly because 1) there is no imprinted XCI to be reversed in human pre-implantation development [9], 2) reactivation of the Xi in human primordial germ cells is difficult to study due to the hurdles associated with obtaining appropriate tissue samples and the lack of a human germ cell culture system that recapitulates Xi reactivation, and 3) reprogramming of human somatic cells under standard conditions does not lead to Xi reactivation [74] as it results in iPSCs that are in a developmentally advanced – primed – pluripotent state [75]. However, when conventional human iPSCs and ESCs in the primed pluripotent state are expanded in culture, *XIST* expression becomes gradually lost, which is accompanied by methylation of the *XIST* promoter [74,76–78]. The *XIST* loss in these pluripotent cells is usually accompanied by partial reactivation of the Xi, where transcriptional repression of some, but not all genes on the Xi goes away, hence the overall inactive state of the Xi erodes, a phenomenon coined Xi erosion [77–80] (Figure 4). Importantly, erosion differs from escape of XCI as the genes undergoing erosion are initially silent on the Xi in early passage hPSCs and become reactivated upon extended passaging of these cells [80], whereas escape is not passage-dependent and rather includes genes on the Xi in chromosome regions with reduced *Xist* occupancy [30,81]. The extent of erosion of the Xi, i.e. the number of genes affected by this process, varies between individual human pluripotent stem cell (hPSC) lines; however, *XIST* loss occurs in nearly all hPSC lines studied over time in culture and often leads to Xi erosion [78,80] (Figure 4). Currently it remains to be tested whether loss of *XIST* expression causes Xi erosion, but the fact that no Xi erosion is observed while *XIST* is expressed in newly derived human iPSC lines [74] suggests such a causative relationship. While *XIST* may have a protective role in preventing Xi erosion, another X-linked lncRNA, *XACT* (X active coating transcript), has been implicated in driving Xi erosion in primed hPSCs [79]. While the eroded Xi does not interfere with hPSC growth or ability to differentiate, it does modulate these processes [80,82]. Moreover, when primed hPSCs with Xi erosion are differentiated, the reactivated genes on the Xi do not get re-silenced, resulting in somatic cells that at least partially lack dosage compensation of X-linked genes [80] (see [83] and [84] for detailed review) (Figure 4). Methods of repairing or preventing Xi erosion of female hPSCs are needed for their use in disease modeling [77] and regenerative medicine, particularly when considering X-linked diseases. For instance, iPSCs or iPSC-derived neurons from female patients with Lesch-Nyhan syndrome, a devastating disease affecting neurologic, cognitive, and behavioral functions [85], can be used to model the disease only in the presence of a faithfully silenced

Xi. This is because the disease phenotype is caused by a heterozygous mutation in the X-linked HPRT1 gene, leading to HPRT1 insufficiency in cells where the non-mutant HPRT1 resides on the Xi. When the region of the Xi harboring the HPRT1 gene undergoes erosion, it results in expression of the non-mutant HPRT1 gene product, overriding HPRT1 insufficiency. Thus, the Lesch-Nyhan diseases phenotypes can no longer be faithfully recapitulated with cultured iPSCs or iPSC-derived neurons in the presence of Xi erosion [77]. Additionally, in regenerative medicine such as cell replacement therapies, introducing cells with an Xi erosion into a patient may be treacherous because these cells lack proper dosage compensation of X-linked genes, a phenomenon observed in cancers [66].

The role of *XIST* in early human development

It is interesting to note that primed hESCs do not reflect the X-chromosome state of the human pre-implantation embryos from which they are derived: all cells of a female human blastocyst, including those of the epiblast lineage, have two active X chromosomes and simultaneously express *XIST* [9,13], (Figure 4). The recent discovery of this non-silencing *XIST* in early human development has intrigued many researchers who study X-chromosome dosage compensation, including us. The two immediate questions regarding this unusual X-chromosome state are, 1) what role, if any, does *XIST* have, and 2) what is the molecular mechanism disabling *XIST* from silencing the X chromosome(s). These and many other questions cannot be addressed with *in vitro* studies of conventional (primed) hESCs since their X-chromosome state is different from the cells of the blastocyst from which they are derived, most plausibly due to suboptimal cell culture composition used (reviewed in more detail in [83]). However, recently devised cell culture conditions, which have been formulated to support cells in a naïve (pre-implantation) pluripotent state, allow growth of hESCs that better resemble the pluripotent state of cells in the pre-implantation blastocyst from which they are derived [86,87]. Most importantly, the X-chromosome state of these naïve hESCs recapitulates many aspects of the human blastocyst, where female cells have two active X chromosome and express *XIST* [20,21]. While most of the cells in a pre-implantation female human blastocyst express *XIST* bi-allelically, this pattern is a minority in naïve hESCs which exhibit mostly mono-allelic *XIST* expression [21]. Hence, naïve hESCs resemble the blastocyst, but not perfectly, as there is still room for improvement in the naïve culture media formulation. The molecular mechanism behind the non-silencing *XIST* is currently not understood, but investigating *XIST*-interacting proteins and *XIST* RNA modifications, which have recently been demonstrated to be crucial for *Xist*'s silencing role in the mouse [55], warrant further investigation. Current naïve culture conditions will allow such studies since naïve hESCs exhibit non-silencing *XIST*, albeit mostly mono-allelically [21].

In addition to recapitulating the X-state of the pre-implantation blastocyst, naïve hPSCs allow *XIST*-mediated induction of XCI upon differentiation [21]. When primed hPSCs with large regions of Xi erosion are adapted to naïve pluripotency and then differentiated, the erosion is, for the first time, reversed and replaced with XCI [21] (Figure 4). Hence, the transition to the naïve state resets the X-chromosome abnormalities of the primed pluripotent state. However, when primed hPSCs are adapted to naïve pluripotency, the memory of the starting Xi does not get lost in the naïve transition, since upon differentiation

the starting Xi becomes silenced despite the presence of *de novo* XCI [21]. Therefore, although naïve hPSCs allow studies of *de novo* XCI in humans for the first time, they cannot be used to study choice of XCI since the process is non-random. The epigenetic memory of the starting Xi is unlikely due to DNA methylation, since the naïve state results in robust hypo-methylation of DNA [21,88,89], but may be due to the presence of histone modifications. For instance, it is possible that tri-methylation of histone H3 lysine 27 (H3K27me3), which was recently shown to regulate *Xist* imprinting in mice [90], marks the inactive or the active X chromosome through the transitions from primed to naïve pluripotency and eventually differentiation. It is, however, not clear whether naïve hESCs directly derived from the blastocyst or somatic cells directly reprogrammed to the naïve state can undergo random XCI upon differentiation.

Since there is no imprinted XCI in early human development, it has been unclear how X-chromosome dosage is compensated prior to onset of random XCI. Single cell RNA-sequencing of human pre-implantation embryos demonstrates gradual and time-dependent reduction of X-linked gene expression from both X chromosomes in embryonic days 4 to 7 in development [13]. This gradual dampening of X-linked gene expression correlates with upregulation of *XIST* [13]. X-chromosome dampening has also been observed in naïve *XIST*-expressing hESCs, further suggesting a novel role of *XIST* in human naïve pluripotency [21]. Independent analysis of the sequencing data from the pre-implantation blastocyst [13] and naïve hESC [21] studies has instead suggested the presence of XCI instead of XCD in human pre-implantation development [19]. If *XIST* is truly initiating XCI in the human blastocyst, given the fact that it is expressed from both X chromosomes in most cells, there must be a critical time-point at which point the cell decides to limit *XIST*'s silencing function to a single X chromosome, since silencing both X chromosomes is lethal due to phenotypic nullisomy of most X-chromosome genes [44,91]. Interestingly, blastocyst outgrowth studies demonstrated the presence of a *XIST*-negative transitional state between the *XIST*-expressing blastocyst cells and the *XIST*-expressing XCI cells [80]. In the transition from XCD to XCI in hPSCs an *XIST*-negative state is also observed [21]. These data suggest that X-chromosome dosage compensation via XCD does not lead to the initiation of XCI.

If *XIST* is responsible for XCD in naïve pluripotency, it might do so by mediating accumulation of some, but perhaps not all chromatin modifications that are also responsible for XCI. For instance, H3K27me3 [92] accumulates on the *XIST*-coated active X chromosome in naïve hESCs [21], which might be responsible for dampening of X-linked gene expression. Another hypothesis is that expression of the lncRNA *XACT* may counteract some but not all functions of *XIST*, thereby achieving dampening instead of silencing. Indeed, it has recently been shown that *XACT* prevents accumulation of *Xist* when ectopically expressed on the mouse X chromosome [20], consistent with the idea that *XACT* can limit *XIST*'s activity in naïve hPSCs. Interestingly, in a fraction of cells of rabbit blastocysts, *Xist* gets expressed from both X chromosomes, initiating silencing of both X-chromosomes before resolving to mono-allelic XCI via unknown mechanisms [9]. It is possible that the human scenario derives from such a mode of initiation of XCI and that *XACT* has evolved in primates to alleviate the detrimental consequences of inactivating both X-chromosomes for too long or in too many cells.

Regardless, naïve hPSCs, for the first time, allow detailed molecular studies of *XIST* and XCD, as well as the transition to XCI as cells exit pluripotency. Moreover, by studying these cells we can now gain insight into early human pre-implantation development and understand how it compares to what we already know in the mouse model organism.

Concluding Remarks and Future Directions

The biology of *Xist* unites researchers from multiple disciplines, including but not limited to those studying sex-chromosome dosage regulation, epigenetics (lncRNAs and chromatin remodeling), cell fate changes (reprogramming), cancer biology, disease pathogenesis (X-linked disorders), as well as development. *Xist*'s ability to recruit such a diverse group of researchers has enabled rapid advancement in understanding how *Xist* functions at the molecular level. We now understand that *Xist* acts as a scaffold to bring proteins to their site of action, effectively increasing the concentration of these proteins in a localized manner. Thanks to its very long-lived outcome and ability to be used in an allele-specific manner, *Xist*'s ability to silence genes can be mined for therapeutic purposes to balance gene expression in trisomic diseases such as Down Syndrome [93]. Furthermore, increased understanding of *Xist* can help engineer variants of this lncRNA to silence smaller and specific regions of a chromosome, increasing its therapeutic potential to silence mutant genes in an allele-specific manner.

Xist's unique expression pattern in human pre-implantation development (expression without silencing) is a great marker of human naïve pluripotency [21] which can be used to develop new and improved naïve culture conditions in the future. Lastly, from an evolutionary perspective, *Xist* is an interesting lncRNA to study since it carries out similar functions in mouse and human despite lack of sequence conservation, but also seems to have evolved extra functions in a context-dependent manner, which requires further investigation (see Outstanding Questions). Both mouse and human *Xist* serve as a wonderful model for expanding our knowledge on lncRNA function, while learning about development and dosage regulation.

Outstanding Questions

- Requirement of *Xist* in random X-chromosome inactivation (XCI) *in vivo* needs to be addressed with mouse models where deletion of *Xist* can be confirmed prior to induction of X-chromosome inactivation.
- Molecular mechanisms regulating X-chromosome dampening (XCD) in human pre-implantation embryos and naïve pluripotent stem cells are yet to be discovered.
- What is preventing *XIST* from silencing X-linked gene expression in human pre-implantation development and naïve pluripotent stem cells?
- Our knowledge of *Xist* biology from mouse studies needs to be extended to human *XIST*: what are the protein binding partners of human *XIST*? What are the steps in the initiation of *XCI*?

- Role of *XIST* and *XACT* in human pre-implantation development and naïve pluripotent stem cells needs to be examined with deletion studies.
- Why do naïve human pluripotent stem cells undergo non-random XCI? What is the epigenetic memory on the X-chromosomes in these cells?
- An improved culture condition for naïve human pluripotent stem cells that allows higher rate of bi-allelic *XIST* cells and random XCI upon differentiation needs to be devised.
- Are there other eutherians that exhibit XCD in pre-implantation development similar to humans?

Glossary

Erosion

loss of transcriptional silencing and DNA methylation for genes on the inactive X chromosome in cultured primed human pluripotent stem cells. Eroded regions of the inactive X chromosome usually contain more than one gene, and fail to re-inactivate upon exit from pluripotency.

ESC

embryonic stem cells are derived from the inner cell mass of the blastocyst before it implants in the uterus. ESCs are pluripotent and can self-renew indefinitely.

Escape

from X-chromosome inactivation happens in both human and mouse development when a small portion of X-linked genes do not undergo transcriptional silencing in early development despite residing on the inactive X-chromosome.

Eutheria

type of mammals whose fetal development requires a placenta (e.g. mouse and human). This excludes pouched mammals (marsupials, e.g. kangaroo) and egg-laying mammals (monotremes, e.g. platypus).

Facultative heterochromatin

a subtype of heterochromatin that can also be present as euchromatin (actively transcribed chromatin) in a different context, such as the X chromosome before and after X-chromosome inactivation.

Heterochromatin

regions of chromatin with low or no transcriptional activity.

Heterochromatinization

formation of heterochromatin via changes to the epigenome, such as covalent modifications of histones and methylation of DNA, which results in transcriptional inactivity.

Histone hypoacetylation

removal of the acetyl groups from histones, leading to reduced transcriptional activity.

LncRNA

long non-coding ribonucleic acid. These RNA molecules are more than 200 nucleotides long and do not encode proteins.

IPSCs

induced pluripotent stem cells derived from somatic cells by transcription factor overexpression-induced reprogramming.

Metatheria

also marsupials, these mammals give live birth after a rather short gestation time, after which development of the newborn continues in the mother's pouch.

Myelodysplasia

multiple types of cancers of blood progenitor cells in the bone marrow which interferes with normal maturation of blood cells, resulting in reduced platelet, or red or white blood cell count.

Naïve pluripotency

pluripotency state of cells of epiblast cells in the pre-implantation blastocyst.

Pluripotency

cell state that can give rise to any of the cell types in the body.

Primed pluripotency

pluripotency state of cells of epiblast cells in the post-implantation embryo.

PSC

pluripotent stem cells are obtained either from pre-implantation blastocysts (ESCs) or via reprogramming of somatic cells (iPSCs).

Reprogramming

forced expression of key transcription factors which remodel the epigenome of differentiated cells (i.e. fibroblasts) to transition these cells into the developmentally early, pluripotent state.

Xa

active X chromosome.

XACT

X active coating transcript is a recently discovered long non-coding RNA on the X chromosome, present in humans but not mice and expressed only in pluripotent cells from non-silent X chromosomes.

XCD

X-chromosome dampening is the tuning down, but not complete silencing of genes on the X chromosome. Transcriptional output from a dampened X chromosome is less than that from an active X but more than that from an inactive X-chromosome.

XCI

X-chromosome inactivation is the transcriptional silencing of most genes on the X chromosome in female mammalian cells.

Xi

inactive X chromosome with most genes not transcribed due to X-chromosome inactivation.

Xi reactivation

of the inactive X-chromosome refers to when all silenced genes become transcribed to resemble the state of the active X-chromosome. This is a chromosome-wide event and occurs in both mouse and human development during germ cell development (cells that give rise to gametes). In mouse development Xi reactivation also occurs in the transition from imprinted to random X-chromosome inactivation.

Xist

X inactive specific transcript is a lncRNA gene on the X chromosome. It does not leave the nucleus and stays associated with the chromosome from which it is transcribed. In mouse, studies have shown that it is required and sufficient for the induction of X-chromosome inactivation.

References

1. Dixon-McDougall T, Brown C. The making of a Barr body: the mosaic of factors that eXIST on the mammalian inactive X chromosome ¹. *Biochem Cell Biol.* 2016; 94:56–70. [PubMed: 26283003]
2. Wang F, et al. Regulation of X-linked gene expression during early mouse development by Rlim. *eLife.* 2016; 5
3. Borensztein M, et al. Xist-dependent imprinted X inactivation and the early developmental consequences of its failure. *Nat Struct Mol Biol.* 2017; 24:226–233. [PubMed: 28134930]
4. Lyon MF. Sex chromatin and gene action in the mammalian X-chromosome. *Am J Hum Genet.* 1962; 14:135–148. [PubMed: 14467629]
5. Wu H, et al. Cellular Resolution Maps of X Chromosome Inactivation: Implications for Neural Development, Function, and Disease. *Neuron.* 2014; 81:103–119. [PubMed: 24411735]
6. Balaton BP, Brown CJ. Escape Artists of the X Chromosome. *Trends Genet.* 2016; 32:348–359. [PubMed: 27103486]
7. Gendrel AV, Heard E. Noncoding RNAs and Epigenetic Mechanisms During X-Chromosome Inactivation. *Annu Rev Cell Dev Biol.* 2014; 30:561–580. [PubMed: 25000994]
8. Grant J, et al. Rxs is a metatherian RNA with Xist-like properties in X-chromosome inactivation. *Nature.* 2012; 487:254–258. [PubMed: 22722828]
9. Okamoto I, et al. Eutherian mammals use diverse strategies to initiate X-chromosome inactivation during development. *Nature.* 2011; 472:370–374. [PubMed: 21471966]
10. Hwang JY, Oh JN, Park CH, Lee DK, Lee CK. Dosage compensation of X-chromosome inactivation center-linked genes in porcine preimplantation embryos: Non-chromosome-wide initiation of X-chromosome inactivation in blastocysts. *Mech Dev.* 2015; 138:246–255. [PubMed: 26598280]
11. Wang X, Miller DC, Clark AG, Antczak DF. Random X inactivation in the mule and horse placenta. *Genome Res.* 2012; 22:1855–1863. [PubMed: 22645258]

12. Moreira de Mello JC, et al. Random X Inactivation and Extensive Mosaicism in Human Placenta Revealed by Analysis of Allele-Specific Gene Expression along the X Chromosome. *PLoS ONE*. 2010; 5:e10947. [PubMed: 20532033]
13. Petropoulos S, et al. Single-Cell RNA-Seq Reveals Lineage and X Chromosome Dynamics in Human Preimplantation Embryos. *Cell*. 2016; 165:1012–1026. [PubMed: 27062923]
14. Deng X, Berletch JB, Nguyen DK, Distèche CM. X chromosome regulation: diverse patterns in development, tissues and disease. *Nat Rev Genet*. 2014; 15:367–378. [PubMed: 24733023]
15. Meyer BJ. Targeting X chromosomes for repression. *Curr Opin Genet Dev*. 2010; 20:179–189. [PubMed: 20381335]
16. Crane E, et al. Condensin-driven remodelling of X chromosome topology during dosage compensation. *Nature*. 2015; 523:240–244. [PubMed: 26030525]
17. Pandya-Jones A, Plath K. The ‘lnc’ between 3D chromatin structure and X chromosome inactivation. *Semin Cell Dev Biol*. 2016; 56:35–47. [PubMed: 27062886]
18. Albritton SE, Ercan S. *Caenorhabditis elegans* Dosage Compensation: Insights into Condensin-Mediated Gene Regulation. *Trends Genet*. 2018; 34:41–53. [PubMed: 29037439]
19. Moreira de Mello JC, Fernandes GR, Vibranovski MD, Pereira LV. Early X chromosome inactivation during human preimplantation development revealed by single-cell RNA-sequencing. *Sci Rep*. 2017; 7
20. Vallot C, et al. XACT Noncoding RNA Competes with XIST in the Control of X Chromosome Activity during Human Early Development. *Cell Stem Cell*. 2017; 20:102–111. [PubMed: 27989768]
21. Sahakyan A, et al. Human Naive Pluripotent Stem Cells Model X Chromosome Dampening and X Inactivation. *Cell Stem Cell*. 2017; 20:87–101. [PubMed: 27989770]
22. Sun S, et al. Jpx RNA Activates Xist by Evicting CTCF. *Cell*. 2013; 153:1537–1551. [PubMed: 23791181]
23. Carmona S, Lin B, Chou T, Arroyo K, Sun S. LncRNA Jpx induces Xist expression in mice using both trans and cis mechanisms. *PLOS Genet*. 2018; 14:e1007378. [PubMed: 29734339]
24. Chureau C, et al. Ftx is a non-coding RNA which affects Xist expression and chromatin structure within the X-inactivation center region. *Hum Mol Genet*. 2011; 20:705–718. [PubMed: 21118898]
25. Furlan G, et al. The Ftx Noncoding Locus Controls X Chromosome Inactivation Independently of Its RNA Products. *Mol Cell*. 2018; 70:462–472.e8. [PubMed: 29706539]
26. Lee JT, Davidow LS, Warshawsky D. Tsix, a gene antisense to Xist at the X-inactivation centre. *Nat Genet*. 1999; 21:400–404. [PubMed: 10192391]
27. Gayen S, Maclary E, Buttigieg E, Hinten M, Kalantry S. A Primary Role for the Tsix lncRNA in Maintaining Random X-Chromosome Inactivation. *Cell Rep*. 2015; 11:1251–1265. [PubMed: 25981039]
28. Vallot C, et al. XACT, a long noncoding transcript coating the active X chromosome in human pluripotent cells. *Nat Genet*. 2013; 45:239–241. [PubMed: 23334669]
29. Jonkers I, et al. Xist RNA Is Confined to the Nuclear Territory of the Silenced X Chromosome throughout the Cell Cycle. *Mol Cell Biol*. 2008; 28:5583–5594. [PubMed: 18625719]
30. Engreitz JM, et al. The Xist lncRNA Exploits Three-Dimensional Genome Architecture to Spread Across the X Chromosome. *Science*. 2013; 341:1237973–1237973. [PubMed: 23828888]
31. Wang KC, et al. A long noncoding RNA maintains active chromatin to coordinate homeotic gene expression. *Nature*. 2011; 472:120–124. [PubMed: 21423168]
32. Engreitz JM, Ollikainen N, Guttman M. Long non-coding RNAs: spatial amplifiers that control nuclear structure and gene expression. *Nat Rev Mol Cell Biol*. 2016; 17:756–770. [PubMed: 27780979]
33. Brown CJ, et al. A gene from the region of the human X inactivation centre is expressed exclusively from the inactive X chromosome. *Nature*. 1991; 349:38–44. [PubMed: 1985261]
34. Brockdorff N, et al. Conservation of position and exclusive expression of mouse Xist from the inactive X chromosome. *Nature*. 1991; 351:329–331. [PubMed: 2034279]
35. Borsani G, et al. Characterization of a murine gene expressed from the inactive X chromosome. *Nature*. 1991; 351:325–329. [PubMed: 2034278]

36. Brown CJ, et al. The human XIST gene: analysis of a 17 kb inactive X-specific RNA that contains conserved repeats and is highly localized within the nucleus. *Cell*. 1992; 71:527–542. [PubMed: 1423611]
37. Brockdorff N, et al. The product of the mouse Xist gene is a 15 kb inactive X-specific transcript containing no conserved ORF and located in the nucleus. *Cell*. 1992; 71:515–526. [PubMed: 1423610]
38. Penny GD, Kay GF, Sheardown SA, Rastan S, Brockdorff N. Requirement for Xist in X chromosome inactivation. *Nature*. 1996; 379:131–137. [PubMed: 8538762]
39. Marahrens Y, Panning B, Dausman J, Strauss W, Jaenisch R. Xist-deficient mice are defective in dosage compensation but not spermatogenesis. *Genes Dev*. 1997; 11:156–166. [PubMed: 9009199]
40. Mugford JW, Yee D, Magnuson T. Failure of extra-embryonic progenitor maintenance in the absence of dosage compensation. *Dev Camb Engl*. 2012; 139:2130–2138.
41. Kalantry S, Purushothaman S, Bowen RB, Starmer J, Magnuson T. Evidence of Xist RNA-independent initiation of mouse imprinted X-chromosome inactivation. *Nature*. 2009; 460:647–651. [PubMed: 19571810]
42. Marahrens Y, Loring J, Jaenisch R. Role of the Xist gene in X chromosome choosing. *Cell*. 1998; 92:657–664. [PubMed: 9506520]
43. Wutz A, Rasmussen TP, Jaenisch R. Chromosomal silencing and localization are mediated by different domains of Xist RNA. *Nat Genet*. 2002; 30:167–174. [PubMed: 11780141]
44. Wutz A, Jaenisch R. A shift from reversible to irreversible X inactivation is triggered during ES cell differentiation. *Mol Cell*. 2000; 5:695–705. [PubMed: 10882105]
45. Yang L, Kirby JE, Sunwoo H, Lee JT. Female mice lacking Xist RNA show partial dosage compensation and survive to term. *Genes Dev*. 2016; 30:1747–1760. [PubMed: 27542829]
46. Rastan S. Timing of X-chromosome inactivation in postimplantation mouse embryos. *J Embryol Exp Morphol*. 1982; 71:11–24. [PubMed: 6185603]
47. Soriano P. Generalized lacZ expression with the ROSA26 Cre reporter strain. *Nat Genet*. 1999; 21:70–71. [PubMed: 9916792]
48. Hayashi S, Lewis P, Pevny L, McMahon AP. Efficient gene modulation in mouse epiblast using a Sox2Cre transgenic mouse strain. *Mech Dev*. 2002; 119(Suppl 1):S97–S101. [PubMed: 14516668]
49. Friedrich G, Soriano P. Promoter traps in embryonic stem cells: a genetic screen to identify and mutate developmental genes in mice. *Genes Dev*. 1991; 5:1513–1523. [PubMed: 1653172]
50. Csankovszki G, Panning B, Bates B, Pehrson JR, Jaenisch R. Conditional deletion of Xist disrupts histone macroH2A localization but not maintenance of X inactivation. *Nat Genet*. 1999; 22:323–324. [PubMed: 10431231]
51. Yildirim E, et al. Xist RNA Is a Potent Suppressor of Hematologic Cancer in Mice. *Cell*. 2013; 152:727–742. [PubMed: 23415223]
52. Nagy A, et al. Embryonic stem cells alone are able to support fetal development in the mouse. *Dev Camb Engl*. 1990; 110:815–821.
53. Schulz EG, et al. The two active X chromosomes in female ESCs block exit from the pluripotent state by modulating the ESC signaling network. *Cell Stem Cell*. 2014; 14:203–216. [PubMed: 24506884]
54. Thornhill AR, Burgoyne PS. A paternally imprinted X chromosome retards the development of the early mouse embryo. *Dev Camb Engl*. 1993; 118:171–174.
55. da Rocha ST, Heard E. Novel players in X inactivation: insights into Xist-mediated gene silencing and chromosome conformation. *Nat Struct Mol Biol*. 2017; 24:197–204. [PubMed: 28257137]
56. Chu C, et al. Systematic Discovery of Xist RNA Binding Proteins. *Cell*. 2015; 161:404–416. [PubMed: 25843628]
57. McHugh CA, et al. The Xist lncRNA interacts directly with SHARP to silence transcription through HDAC3. *Nature*. 2015; 521:232–236. [PubMed: 25915022]
58. Minajigi A, et al. A comprehensive Xist interactome reveals cohesin repulsion and an RNA-directed chromosome conformation. *Science*. 2015; 349:aab2276–aab2276.

59. Moindrot B, et al. A Pooled shRNA Screen Identifies Rbm15, Spen, and Wtap as Factors Required for Xist RNA-Mediated Silencing. *Cell Rep.* 2015; 12:562–572. [PubMed: 26190105]
60. Monfort A, et al. Identification of Spen as a Crucial Factor for Xist Function through Forward Genetic Screening in Haploid Embryonic Stem Cells. *Cell Rep.* 2015; 12:554–561. [PubMed: 26190100]
61. Chen CK, et al. Xist recruits the X chromosome to the nuclear lamina to enable chromosome-wide silencing. *Science.* 2016; 354:468–472. [PubMed: 27492478]
62. Patil DP, et al. m6A RNA methylation promotes XIST-mediated transcriptional repression. *Nature.* 2016; doi: 10.1038/nature19342
63. Pintacuda G, et al. hnRNPK Recruits PCGF3/5-PRC1 to the Xist RNA B-Repeat to Establish Polycomb-Mediated Chromosomal Silencing. *Mol Cell.* 2017; 68:955–969.e10. [PubMed: 29220657]
64. Van Nostrand EL, et al. Robust transcriptome-wide discovery of RNA-binding protein binding sites with enhanced CLIP (eCLIP). *Nat Methods.* 2016; 13:508–514. [PubMed: 27018577]
65. Shevchenko AI, et al. Impact of Xist RNA on chromatin modifications and transcriptional silencing maintenance at different stages of imprinted X chromosome inactivation in vole *Microtus levis*. *Chromosoma.* 2018; 127:129–139. [PubMed: 29151149]
66. Chaligné R, Heard E. X-chromosome inactivation in development and cancer. *FEBS Lett.* 2014; 588:2514–2522. [PubMed: 24937141]
67. Pasque V, Plath K. X chromosome reactivation in reprogramming and in development. *Curr Opin Cell Biol.* 2015; 37:75–83. [PubMed: 26540406]
68. Mak W, et al. Reactivation of the paternal X chromosome in early mouse embryos. *Science.* 2004; 303:666–669. [PubMed: 14752160]
69. de Napoles M, Nesterova T, Brockdorff N. Early Loss of Xist RNA Expression and Inactive X Chromosome Associated Chromatin Modification in Developing Primordial Germ Cells. *PLoS ONE.* 2007; 2:e860. [PubMed: 17848991]
70. Pasque V, et al. X Chromosome Reactivation Dynamics Reveal Stages of Reprogramming to Pluripotency. *Cell.* 2014; 159:1681–1697. [PubMed: 25525883]
71. Csankovszki G, Nagy A, Jaenisch R. Synergism of Xist RNA, DNA methylation, and histone hypoacetylation in maintaining X chromosome inactivation. *J Cell Biol.* 2001; 153:773–784. [PubMed: 11352938]
72. Minkovsky A, et al. The Mbd1-Atf7ip-Setdb1 pathway contributes to the maintenance of X chromosome inactivation. *Epigenetics Chromatin.* 2014; 7:12. [PubMed: 25028596]
73. Keniry A, et al. Setdb1-mediated H3K9 methylation is enriched on the inactive X and plays a role in its epigenetic silencing. *Epigenetics Chromatin.* 2016; 9
74. Tchieu J, et al. Female Human iPSCs Retain an Inactive X Chromosome. *Cell Stem Cell.* 2010; 7:329–342. [PubMed: 20727844]
75. Nichols J, Smith A. Naive and Primed Pluripotent States. *Cell Stem Cell.* 2009; 4:487–492. [PubMed: 19497275]
76. Shen Y, et al. X-inactivation in female human embryonic stem cells is in a nonrandom pattern and prone to epigenetic alterations. *Proc Natl Acad Sci.* 2008; 105:4709–4714. [PubMed: 18339804]
77. Mekhoubad S, et al. Erosion of Dosage Compensation Impacts Human iPSC Disease Modeling. *Cell Stem Cell.* 2012; 10:595–609. [PubMed: 22560080]
78. Nazor KL, et al. Recurrent Variations in DNA Methylation in Human Pluripotent Stem Cells and Their Differentiated Derivatives. *Cell Stem Cell.* 2012; 10:620–634. [PubMed: 22560082]
79. Vallot C, et al. Erosion of X Chromosome Inactivation in Human Pluripotent Cells Initiates with XACT Coating and Depends on a Specific Heterochromatin Landscape. *Cell Stem Cell.* 2015; 16:533–546. [PubMed: 25921272]
80. Patel S, et al. Human Embryonic Stem Cells Do Not Change Their X Inactivation Status during Differentiation. *Cell Rep.* 2017; 18:54–67. [PubMed: 27989715]
81. Simon MD, et al. High-resolution Xist binding maps reveal two-step spreading during X-chromosome inactivation. *Nature.* 2013; 504:465–469. [PubMed: 24162848]

82. Bruck T, Yanuka O, Benvenisty N. Human pluripotent stem cells with distinct X inactivation status show molecular and cellular differences controlled by the X-Linked ELK-1 gene. *Cell Rep.* 2013; 4:262–270. [PubMed: 23871667]
83. Sahakyan A, Plath K, Rougeulle C. Regulation of X-chromosome dosage compensation in human: mechanisms and model systems. *Philos Trans R Soc B Biol Sci.* 2017; 372:20160363.
84. Geens M, et al. Female human pluripotent stem cells rapidly lose X chromosome inactivation marks and progress to a skewed methylation pattern during culture. *Mol Hum Reprod.* 2016; 22:285–298. [PubMed: 26786180]
85. Bell S, Kolobova I, Crapper L, Ernst C. Lesch-Nyhan Syndrome: Models, Theories, and Therapies. *Mol Syndromol.* 2016; 7:302–311. [PubMed: 27920633]
86. Theunissen TW, et al. Systematic identification of culture conditions for induction and maintenance of naive human pluripotency. *Cell Stem Cell.* 2014; 15:471–487. [PubMed: 25090446]
87. Takashima Y, et al. Resetting Transcription Factor Control Circuitry toward Ground-State Pluripotency in Human. *Cell.* 2014; 158:1254–1269. [PubMed: 25215486]
88. Pastor WA, et al. Naive Human Pluripotent Cells Feature a Methylation Landscape Devoid of Blastocyst or Germline Memory. *Cell Stem Cell.* 2016; 18:323–329. [PubMed: 26853856]
89. Theunissen TW, et al. Molecular Criteria for Defining the Naive Human Pluripotent State. *Cell Stem Cell.* 2016; doi: 10.1016/j.stem.2016.06.011
90. Inoue A, Jiang L, Lu F, Zhang Y. Genomic imprinting of *Xist* by maternal H3K27me3. *Genes Dev.* 2017; 31:1927–1932. [PubMed: 29089420]
91. Guyochin A, et al. Live Cell Imaging of the Nascent Inactive X Chromosome during the Early Differentiation Process of Naive ES Cells towards Epiblast Stem Cells. *PLoS ONE.* 2014; 9:e116109. [PubMed: 25546018]
92. Plath K, Zhang Y, Panning B, Otte AP. Role of Histone H3 Lysine 27 Methylation in X Inactivation. *Science.* 2003; 300:131–135. [PubMed: 12649488]
93. Jiang J, et al. Translating dosage compensation to trisomy 21. *Nature.* 2013; 500:296–300. [PubMed: 23863942]

Trends

- X inactive specific transcript (*Xist*) is a long non-coding RNA that remains associated with the X chromosome from which it is expressed.
- *Xist* is unequivocally required for the imprinted form of X-chromosome inactivation (XCI) in mice *in vivo*, but demonstration of its indisputable requirement for random XCI *in vivo* is yet to be shown.
- Loss of *Xist* expression in mice *in vivo* and in conventional human pluripotent stem cells correlates with partial re-activation of genes residing on the inactive X chromosome, suggesting an important role of *Xist* in maintenance of the silent state of genes on the inactive X chromosome.
- Human pre-implantation embryos have a unique X-chromosome dosage compensation state called X-chromosome dampening (XCD), where transcriptional output is tuned down from both X chromosomes. Correlative observations suggest that XCD might be mediated by *XIST*.




Figure 1. X chromosome inactivation in different mammals

The X-chromosome states of eutherian (A and B) and **metatherian** (C) female mammals are shown in embryonic development. A) Imprinted XCI occurs in mouse pre-implantation development, but it is re-set in the cells that develop into the embryo to give way to random XCI, resulting in a mosaic adult female mouse. Both imprinted and random XCI in the mouse are regulated by the lncRNA *Xist*. B) Humans have evolved away from imprinted XCI as they dosage compensate in pre-implantation development by turning down transcription from both X chromosomes via XCD. Moreover, *XIST* is expressed on both dampened X chromosomes, where its functional role remains to be determined. In post-implantation development, similar to the mouse, human females display random XCI mediated by *XIST*. C) Metatherians, such as the marsupial opossum (*Monodelphis domestica*) dosage-compensate by inactivating the paternally-inherited X chromosome using the lncRNA *Rsx*. This imprinted dosage-compensation is maintained throughout marsupial development, resulting in a female adult with a transcriptionally inactive paternal X chromosome in all of its cells. XCI = X chromosome inactivation, *Xist/XIST* = X inactive specific transcript, X_a = active X chromosome, X_i = inactive X chromosome, X_m = maternal X chromosome, X_p = paternal X chromosome, XCD = X chromosome dampening, *Rsx* = RNA on the silent X.




Figure 2. Long non-coding RNAs involved in X chromosome dosage regulation in mouse and human

The X inactivation center (XIC) is located on the X chromosome and harbors the master regulator of XCI – the long non-coding RNA (lncRNA) *Xist*. A) In the mouse, *Xist* itself is positively regulated by the lncRNAs *Jpx* and *Ftx*, which are also encoded in the XIC, upstream of the *Xist* gene. The lncRNA *Tsix*, which is anti-sense to *Xist*, has a mutually exclusive expression pattern with *Xist*: it is expressed bi-allelically from both X chromosomes prior to XCI, and is thought to be a repressor of *Xist*. *Tsix* ‘protects’ the active X chromosome in pluripotency from being silenced by *Xist* upon induction of XCI, and is thought to be a repressor of *Xist*. B) The human *XIST* gene is also encoded in the XIC of the human X chromosome. The lncRNAs *JPX* and *FTX* are also upstream of *XIST* in the human XIC, similar to mouse. The role of *JPX* and *FTX* in regulating *XIST* expression in human is speculated based on mouse studies. Unlike mouse, the human XIC does not contain the *XIST* anti-sense lncRNA *TSIX*, since *TSIX* expression is not detected in pre-implantation blastocysts or human embryonic stem cells. A novel, human-specific lncRNA, X active coating transcript (*XACT*), however, is encoded about 40Mb upstream of the human XIC and seems to antagonize *XIST* in naïve pluripotency. Similar to mouse *Tsix*, expression of *XACT* is unique to pluripotent cells and not detected in somatic cells.




Figure 3. Lack of *Xist* at various developmental time-points highlights its importance in normal development

Summary of key studies addressing the role of *Xist* in mouse development from fertilization to birth and into adulthood. (i) When a zygote is formed with a maternally deleted *Xist* (inherited from the egg), mouse development progresses normally and results in non-mosaic adults where all cells inactivate the paternally inherited X chromosome (since only that X has the only functional *Xist* allele). (ii) However, when *Xist* is deleted from both X chromosomes (from the egg and the sperm), or from only the paternal X chromosome (sperm), extra-embryonic tissues fail to develop in the absence of imprinted XCI since this process requires paternally-inherited *Xist*, and thus mouse development halts 5–7 days post implantation. (iii) Conditional *Xist* deletion from both X chromosomes in epiblast cells that give rise to the embryo is often embryonically lethal, and if pups are born, they display partial loss of X-chromosome silencing and do not survive to adulthood. (iv) When *Xist* is deleted several days post-implantation, specifically in hematopoietic stem cells (HSCs), after the establishment of the Xi, pups are born but succumb to Multilineage Dysplasia as early as 1.5 months after birth. E = embryonic day, *Xist* = X inactive specific transcript, iXCI = imprinted X-chromosome inactivation, HSCs = hematopoietic stem cells.




Figure 4. The X-chromosome state of naïve and primed human pluripotent stem cells

(i) Female human somatic cells have an active and an *XIST*-expressing inactive X chromosome (Xa and Xi). (ii) Reprogramming of these cells to **primed pluripotency** does not change the X-chromosome state. (iii) Similarly, derivation of hPSCs from a pre-implantation blastocyst stabilizes the post-XCI state in primed pluripotent culture conditions. (iv) Over time in culture, the Xi loses expression of *XIST* and undergoes epigenetic erosion, resulting in partial reactivation and thus double-dose of the X-linked genes that fall in these eroded regions in primed hPSCs. (v) Although these cells can differentiate into somatic lineages, the resulting differentiated cells maintain the eroded X (Xe).

Female pre-implantation blastocysts have two active X chromosomes and express *XIST*, serving as a unique scenario where *XIST* expression does not cause XCI. (vi) When hESCs are derived under naïve pluripotent culture conditions, or when primed hPSCs are adapted to such naïve conditions, the X-chromosome state of resulting hPSCs resembles that of the pre-implantation blastocyst. (vii) Similar to normal development, differentiation of naïve hPSCs results in *XIST*-mediated XCI. *denotes the state found in majority of cells. hPSCs = human pluripotent stem cells, *XIST* = X inactive specific transcript, Xi = inactive X chromosome, Xa = active X chromosome, Xe = eroded X chromosome.