

UCLA

UCLA Previously Published Works

Title

Meta-analysis across Cohorts for Heart and Aging Research in Genomic Epidemiology (CHARGE) consortium provides evidence for an association of serum vitamin D with pulmonary function

Permalink

<https://escholarship.org/uc/item/9696h6m0>

Journal

British Journal Of Nutrition, 120(10)

ISSN

0007-1145

Authors

Xu, Jiayi

Bartz, Traci M

Chittoor, Geetha

et al.

Publication Date

2018-11-28

DOI

10.1017/s0007114518002180

Peer reviewed



Published in final edited form as:

Br J Nutr. 2018 November ; 120(10): 1159–1170. doi:10.1017/S0007114518002180.

Meta-Analysis across Cohorts for Heart and Aging Research in Genomic Epidemiology (CHARGE) Consortium Provides Evidence for an Association of Serum Vitamin D with Pulmonary Function

Jiayi Xu¹, Traci M. Bartz^{#2,3}, Geetha Chittoor^{#4,5}, Gudny Eiriksdottir^{#6}, Ani W. Manichaikul^{#7}, Fangui Sun^{#8}, Natalie Terzikhan^{#9,10}, Xia Zhou^{#11}, Sarah L. Booth¹², Guy G. Brusselle^{9,10,13}, Ian H. de Boer¹⁴, Myriam Fornage^{15,16}, Alexis C. Frazier-Wood¹⁷, Mariaelisa Graff⁵, Vilmundur Gudnason^{6,18}, Tamara B. Harris¹⁹, Albert Hofman^{10,20,21}, Ruixue Hou²², Denise K. Houston²³, David R. Jacobs Jr¹¹, Stephen B. Kritchevsky²³, Jeanne Latourelle^{24,25}, Rozenn N. Lemaitre^{3,14}, Pamela L. Lutsey¹¹, George O. Connor²⁴, Elizabeth C. Oelsner²⁶, James S. Pankow¹¹, Bruce M. Psaty^{3,14,27,28,29}, Rebecca R. Rohde⁵, Stephen S. Rich⁷, Jerome I. Rotter³⁰, Lewis J. Smith³¹, Bruno H. Stricker^{10,20}, V. Saroja Voruganti²², Thomas J. Wang³², M. Carola Zillikens^{20,33}, R. Graham Barr^{#26}, Josée Dupuis^{#8}, Sina A. Gharib^{#14,34}, Lies Lahousse^{#9,10,35}, Stephanie J. London^{#36}, Kari E. North^{#5}, Albert V. Smith^{#6,18}, Lyn M. Steffen^{#11}, Dana B. Hancock³⁷, and Patricia A. Cassano¹

¹Division of Nutritional Sciences, Cornell University, Ithaca, New York, United States ²Department of Biostatistics, University of Washington, Seattle, Washington, United States ³Cardiovascular Health Research Unit, University of Washington, Seattle, Washington, United States ⁴Department of Biomedical and Translational Informatics, Geisinger, Danville, Pennsylvania, United States ⁵Gillings School of Global Public Health, Department of Epidemiology, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, United States ⁶Icelandic Heart Association, Kopavogur, Iceland ⁷Center for Public Health Genomics, University of Virginia School of Medicine, Charlottesville, Virginia, United States ⁸Department of Biostatistics, Boston University School of

Corresponding author: Patricia A. Cassano, PhD, Division of Nutritional Sciences, Cornell University, 209 Savage Hall, Cornell University, Ithaca, NY 14853, pac6@cornell.edu, 607 255 7551 (PH); 607 255 2691 (FAX).

CONTRIBUTORS

PAC, DBH, and JX conceived and designed the study. RGB, JL, JD, SAG, LL, SJL, KEN, AVS, BMP, and LMS provided the data and supervised the data analysis in each cohort. JX, TMB, RRR, AVS, AWM, FS, NT, and XZ analyzed data within each cohort. JX, PAC and DBH meta-analyzed and interpreted the data. JX, PAC and DBH co-wrote and edited the first draft of the manuscript. PAC, DBH and JX had primary responsibility for final content. All authors provided data, analytic support and/or study design suggestions at all stages, critically reviewed the manuscript, and read and approved the final version.

Author Disclosure

Dr. Psaty serves on the DSMB of a clinical trial funded by the manufacturer (Zoll LifeCor) and on the Steering Committee of the Yale Open Data Access Project funded by Johnson & Johnson.

All other authors have no conflicts of interest. There is no commercial support or financial interest from the tobacco industry for the research presented.

The study sponsors were not involved in study design, data collection, data analysis, data interpretation, report writing, or decisions to submit the paper for publication. PAC and DBH had final responsibility for the decision to submit for publication.

Online Supporting Material

Supplemental table, figures, and methods are available.

Public Health, Boston, Massachusetts, United States ⁹Department of Respiratory Medicine, Ghent University Hospital, Ghent, Belgium ¹⁰Department of Epidemiology, Erasmus Medical Center, Rotterdam, the Netherlands ¹¹Division of Epidemiology and Community Health, University of Minnesota, Minneapolis, Minnesota, United States ¹²Jean Mayer-U.S. Department of Agriculture Human Nutrition Research Center on Aging, Tufts University, Boston, Massachusetts, United States ¹³Department of Respiratory Medicine, Erasmus Medical Center, Rotterdam, the Netherlands ¹⁴Department of Medicine, University of Washington, Seattle, Washington, United States ¹⁵Institute of Molecular Medicine, University of Texas Health Science Center at Houston, Houston, Texas, United States ¹⁶Human Genetics Center, School of Public Health, University of Texas Health Science Center at Houston, Houston, Texas, United States ¹⁷Children's Nutrition Research Center, Baylor College of Medicine, Houston, Texas, United States ¹⁸University of Iceland, Reykjavik, Iceland ¹⁹Laboratory of Epidemiology, Demography, and Biometry, National Institute on Aging, National Institutes of Health, Department of Health and Human Services, Bethesda, Maryland, United States ²⁰Netherlands Genomics Initiative (NGI)-sponsored Netherlands Consortium for Healthy Aging, Leiden, the Netherlands ²¹Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, Massachusetts, United States ²²Department of Nutrition and Nutrition Research Institute, University of North Carolina at Chapel Hill, Kannapolis, North Carolina, United States ²³Sticht Center on Aging, Department of Internal Medicine, Section on Gerontology and Geriatric Medicine, Wake Forest School of Medicine, Winston-Salem, North Carolina, United State ²⁴The Pulmonary Center, Department of Medicine, Boston University, Boston, Massachusetts, United State ²⁵Department of Neurology, Boston University, Boston, Massachusetts, United States ²⁶Department of Medicine, Columbia University, New York, New York, United States ²⁷Department of Epidemiology, University of Washington, Seattle, Washington, United States ²⁸Department of Health Services, University of Washington, Seattle, Washington, United States ²⁹Kaiser Permanente Washington Health Research Institute, Seattle, Washington, United States ³⁰Institute for Translational Genomics and Population Sciences, Los Angeles Biomedical Research Institute and Department of Pediatrics at Harbor-UCLA Medical Center, Torrance, California, United States ³¹Division of Pulmonary and Critical Care, Feinberg School of Medicine, Northwestern University, Chicago, Illinois, United States ³²Division of Cardiovascular Medicine, Department of Medicine, Vanderbilt University, Nashville, Tennessee, United States ³³Department of Internal Medicine, Erasmus Medical Center, Rotterdam, the Netherlands ³⁴Center for Lung Biology, University of Washington, Seattle, Washington, United States ³⁵Department of Bioanalysis, Ghent University, Ghent, Belgium ³⁶Division of Intramural Research, National Institute of Environmental Health Sciences, National Institutes of Health, Department of Health and Human Services, Research Triangle Park, North Carolina, United States ³⁷Behavioral and Urban Health Program, Behavioral Health and Criminal Justice Division, RTI International, Research Triangle Park, North Carolina, United States

These authors contributed equally to this work.

Abstract

The role that vitamin D plays in pulmonary function remains uncertain. Epidemiological studies reported mixed findings for serum 25-hydroxyvitamin D [25(OH)D]—pulmonary function association. We conducted the largest cross-sectional meta-analysis of the 25(OH)D–pulmonary

function association to date, based on nine European ancestry (EA) cohorts ($n=22,838$) and five African ancestry (AA) cohorts ($n=4,290$) in the CHARGE Consortium. Data were analyzed using linear models by cohort and ancestry. Effect modification by smoking status (current/former/never) was tested. Results were combined using fixed-effects meta-analysis. Mean (SD) serum 25(OH)D was 68 (29) nmol/L for EAs and 49 (21) nmol/L for AAs. For each 1 nmol/L higher 25(OH)D, forced expiratory volume in the first second (FEV₁) was higher by 1.1 mL in EAs (95% CI: 0.9,1.3; $P < 0.0001$) and 1.8 mL (95% CI: 1.1,2.5; $P < 0.0001$) in AAs ($P_{\text{race difference}}=0.06$), and forced vital capacity (FVC) was higher by 1.3 mL in EAs (95% CI: 1.0,1.6; $P < 0.0001$) and 1.5 mL (95% CI: 0.8,2.3; $P = 0.0001$) in AAs ($P_{\text{race difference}}=0.56$). Among EAs, the 25(OH)D–FVC association was stronger in smokers: per 1nmol/L higher 25(OH)D, FVC was higher by 1.7 mL (95% CI: 1.1,2.3) for current smokers and 1.7 mL (95% CI: 1.2,2.1) for former smokers, compared to 0.8 mL (95% CI: 0.4,1.2) for never smokers. In summary, the 25(OH)D associations with FEV₁ and FVC were positive in both ancestries. In EAs, a stronger association was observed for smokers compared to never smokers, which supports the importance of vitamin D in vulnerable populations.

Keywords

25-hydroxyvitamin D; vitamin D; forced expiratory volume; vital capacity; respiratory function tests; smoking; human; adult; whites; African Americans

INTRODUCTION

Chronic obstructive pulmonary disease (COPD), the third leading cause of mortality in the U.S.⁽¹⁾ and among the top 10 leading causes of total years of life lost in the world⁽²⁾, is characterized by progressive airway obstruction. Pulmonary function tests (PFTs), as performed by spirometry, are used to quantify pulmonary function parameters including forced expiratory volume in the first second (FEV₁) and forced vital capacity (FVC). Pulmonary function increases throughout childhood, plateaus in the 20s, and thereafter adults experience an age-related decline⁽³⁾. The majority of COPD cases (85%) are related to smoking⁽⁴⁾, which alters the trajectory in pulmonary function, by hindering growth, reducing peak function, and accelerating age-related decline⁽⁵⁾.

Vitamin D is proposed to have protective effects in the lungs via gene regulation⁽⁶⁾. *In vitro* studies found that 1,25-dihydroxyvitamin D, the active vitamin D metabolite, induced antimicrobial peptides for host defense in the lung and modulated airway remodeling⁽⁷⁾. In humans, 25-hydroxyvitamin D [25(OH)D] is the major vitamin D metabolite in serum, most of which forms a complex with vitamin D binding protein (~85–90% is DBP-bound)⁽⁸⁾, and then is metabolized to 1,25-dihydroxyvitamin D [1,25-(OH)₂D], the active steroid hormone form⁽⁹⁾. Total 25(OH)D is the commonly used biomarker of vitamin D status, and it is preferred to other vitamin D metabolites, such as non-DBP-bound 25(OH)D and 1,25-(OH)₂D, given that it is a comprehensive indicator for vitamin D stores, has a longer half-life (~3 weeks) and is less affected by calcium⁽¹⁰⁾. On average, African ancestry (AA) populations have lower serum 25(OH)D concentrations, due to multiple factors including genetics and skin pigmentation⁽¹¹⁾, but evidence exists that AA populations have higher 1,25-

(OH)₂D levels and greater bone mineral density compared to European ancestry (EA) populations⁰.

Previous observational cross-sectional studies of the vitamin D–pulmonary function association in the general population reported mixed findings. Most of these studies reported a positive association between 25(OH)D and pulmonary function⁽⁻⁾, although some reported a null or inverse association^(...), and two others reported a positive association under certain conditions, such as only in male current smokers⁰ or only in overweight and obese males⁰. The largest previous cross-sectional study, which included two Danish cohorts (total $n = 18,507$), reported positive associations of 25(OH)D with pulmonary function⁰. Only one prior cross-sectional study investigated serum 25(OH)D and pulmonary function in an ancestry group other than European, and it confirmed similar positive associations in the 3,957 AA participants studied⁰.

The current study investigated the hypothesis that serum 25(OH)D level is positively associated with pulmonary function. We leveraged the Cohorts for Heart and Aging Research in Genomic Epidemiology (CHARGE) Consortium to include population-based data on serum 25(OH)D and pulmonary function in a harmonized analysis. Additionally, we compared the association of serum 25(OH)D and pulmonary function across EA and AA groups and investigated effect modification by cigarette smoking.

MATERIALS AND METHODS

Cohorts and Participants

Nine prospective cohorts in the CHARGE Consortium were included (Table 1). All cohorts had EA participants, and five of the cohorts had AA participants. Only one cohort [Multi-Ethnic Study of Atherosclerosis (MESA)] has participants with other ancestries, and these other ancestries were not included in this study. Among the nine cohorts, the Framingham Heart Study (FHS) had two sub-cohorts analyzed separately: the Offspring and the Third-Generation (Gen3) cohorts. Our analysis pipeline harmonized the outcome and exposure definitions, the units on all variables, and the statistical modeling. The same exclusion criteria were applied to each cohort: missing PFTs, unacceptable PFTs using the American Thoracic Society and European Respiratory Society criteria for acceptability, missing serum 25(OH)D, serum 25(OH)D > 374.4 nmol/L (or 150ng/mL, leading to removal of a single outlier^(...)), or missing on other covariates (Supplemental Table 1).

Outcome and Exposure Assessment

Pre-bronchodilator pulmonary function outcomes (FEV₁, FVC, and FEV₁/FVC), which have similar accuracy as post-bronchodilator measures for long-term outcomes⁰, were measured in each cohort using standardized methods defined by the American Thoracic Society/European Respiratory Society criteria (Supplemental Table 2). The methods used to measure 25(OH)D varied by cohort (Supplemental Table 2). Three cohorts, including MESA, the Atherosclerosis Risk In Communities (ARIC) study, and the Cardiovascular Health Study (CHS), used the current reference method, liquid chromatography in tandem with mass spectrometry (LC-MS/MS); three cohorts, including FHS, the Coronary Artery

Risk Development in Young Adults (CARDIA) study, and the Health, Aging, and Body Composition (HABC) study, used radioimmunoassay (RIA); one cohort, the Age, Gene, Environment, Susceptibility Study—Reykjavik, Iceland (AGES), used chemiluminescence immunoassay (CLIA); and one cohort [the Rotterdam Study (RS)] used electro-CLIA. Only MESA calibrated the serum 25(OH)D measurement against the standard reference material 972⁰, which reflects the calendar time of the measurements in the cohorts, most of which occurred before the availability of the standard reference material (Supplemental Table 3). Measurements of the outcome and exposure variables were planned for either the full cohort (ARIC, CHS, FHS, HABC, and RS) or a subset of the cohort if the outcome or the exposure was only measured in an ancillary study (AGES, CARDIA, and MESA)⁽⁻⁾ (Supplemental Table 1). Continuous variables were used for serum 25(OH)D and pulmonary function to capture the association of 25(OH)D on PFTs across the broad distribution of ranges in the cohorts.^(,)

As shown in Table 1, among nine cohorts, four [AGES, CHS, FHS-Offspring, and FHS-Gen3] had a mean time difference of less than one year in the PFT measurements and the preceding 25(OH)D measurement, and the greatest mean time difference between 25(OH)D and PFT measurement was < 5 years [MESA]. Participants in ARIC and HABC had blood drawn for serum 25(OH)D after their PFT measure, but within 3 years.

Other covariates, including smoking status, pack-years (number of packs of cigarettes smoked per day times the number of years smoked), height, weight, and age, were measured concurrently with pulmonary function, except for CHS, which assessed covariates concurrent with the serum 25(OH)D measure, but within 1 year of the PFT measurement (Supplemental Table 3). All data collection and analysis was approved by the Institutional Review Board at each cohort's respective institution. Spirometry measures are available on the database of Genotypes and Phenotypes via accession numbers as follows: ARIC (phs000280), CARDIA (phs000285), CHS (phs000287), FHS (phs000007), and MESA (phs000209). Serum vitamin D measures are also available at the same accession numbers for CHS, FHS, and MESA.

Statistical Analysis in Individual Cohorts

All analyses were first conducted independently in each cohort, stratified by ancestry, given the lower mean serum 25(OH)D level in AA participants⁰. For FEV₁ and FEV₁/FVC, models were adjusted for smoking status, pack-years, height, height squared, age, age squared, sex, season of blood draw, and study center (if applicable); for FVC, the model was further adjusted for weight. Residual outliers, identified using the studentized residuals of the linear models (Supplemental Methods for more details), were excluded from all models (~0.3% of the total sample size). The model was extended to test the interaction between 25(OH)D and smoking status [never (reference group), former, and current smokers].

Meta-Analysis

We tested the association of serum 25(OH)D on each PFT outcome among individuals in each ancestry group and each cohort, separately, and then combined the effect estimates (also referred to as two-stage meta-analysis⁰), using inverse variance weighting and

assuming fixed-effects, with heterogeneity assessed via the I^2 statistic⁰. Random-effects meta-analysis was performed as a sensitivity analysis if there was potential heterogeneity ($I^2 > 30\%$). The comparison of meta-analyzed coefficients of the 25(OH)D–PFT associations for the two ancestry groups was conducted using a Z test⁰. Meta-analysis of the interaction terms of 25(OH)D with smoking status was also performed (Supplemental Methods for more details, Supplemental Table 4 and 5 for cohort-specific findings, and Supplemental Table 6 for meta-analyzed results).

Meta-regression was conducted to explore the potential causes of moderate heterogeneity in the meta-analysis of 25(OH)D on FEV₁ and FVC in the EA cohorts. Modifiers were tested individually in the meta-regression models to investigate heterogeneity; modifiers included factors that could vary between cohorts, such as proportion of ever, current, and former smokers, mean 25(OH)D level, assay method for serum 25(OH)D, time between 25(OH)D and PFT measures, and mean age of participants in each cohort. The two-sided type I error was examined at 0.05 for all analyses. Meta-analysis and meta-regression were conducted using the metafor package (version 1.9–8) in R (version 3.2.3., R Foundation for Statistical Computing, Vienna, Austria).

Regression coefficients (β) and standard errors (SE) calculated within each cohort per 1 nmol/L 25(OH)D are presented in the figures. Additionally, to put the magnitude of the 25(OH)D–PFT associations in terms relevant to public health, the meta-analyzed regression coefficients were multiplied by 10 nmol/L 25(OH)D, which is about half of the standard deviation (SD) of the 25(OH)D distribution.

RESULTS

We studied 22,838 EA and 4,290 AA participants. EA participants had higher FEV₁, FVC, and serum 25(OH)D than AA participants in each cohort, while FEV₁/FVC was similar across ancestry groups (Table 1 and Supplemental Figure 1). CARDIA and FHS-Gen3 were younger than the seven other cohorts, with consequently lower pack-years smoked in ever smokers. Across all cohorts, among EA participants, 17% were current smokers and 40% were former smokers; among AA participants, 22% were current smokers and 30% were former smokers. The serum 25(OH)D level was highest among never smokers [mean(SD) = 70(30) nmol/L], followed by former smokers [67 (29) nmol/L], and current smokers [64 (29) nmol/L] in EAs, while the trend was less obvious in AAs [49 (21) nmol/L in current smokers, 50 (21) nmol/L in former smokers, and 48 (21) nmol/L in never smokers]. The mean (SD) of serum 25(OH)D for EA participants across nine cohorts was 68 (29) nmol/L and for AA participants across five cohorts the mean (SD) was 49 (21) nmol/L.

Fixed-effects meta-analysis (Figure 1) revealed a consistently positive association of serum 25(OH)D with the PFT outcomes, FEV₁ and FVC, in both ancestry groups. To put these findings into context, a 10 nmol/L (~0.5 SD) higher 25(OH)D was associated with 11.1 mL higher FEV₁ in EAs ($P < 0.0001$) and 17.9 mL higher FEV₁ in AAs ($P < 0.0001$). Similarly, for a 10 nmol/L higher 25(OH)D, FVC was higher by 12.9 mL in EAs ($P < 0.0001$) and by 15.4 mL in AAs ($P = 0.0001$). The magnitudes of the 25(OH)D–PFT associations did not differ significantly between the two ancestry groups ($P = 0.06$ and $P = 0.56$ for FEV₁ and

FVC, respectively). The association of serum 25(OH)D with FEV₁/FVC reached statistical significance only in EAs (P = 0.0013), and the magnitude was negligible; a 10 nmol/L higher 25(OH)D was associated with a ratio being lower by 0.0055% (Supplemental Table 7 and Supplemental Figure 2 for ancestry- and cohort-specific findings).

In the main-effect meta-analysis of serum 25(OH)D on pulmonary function, EA cohorts had low to moderate heterogeneity, while AA cohorts had low heterogeneity (Figure 1, Supplemental Figure 2). We did a sensitivity analysis using random-effects meta-analysis among EA cohorts for the FEV₁ and FVC outcomes, and no substantial change was found in the meta-analyzed effect estimates and corresponding standard errors (coefficient (SE) of 1 nmol/L 25(OH)D on the FEV₁ outcome was 1.11 (0.12) mL in the fixed-effects model and 1.21 (0.19) mL in the random-effects model; coefficient (SE) on the FVC outcome was 1.29 (0.14) mL in the fixed-effects model and 1.31 (0.20) mL in the random-effects model). Meta-regression was also performed in the EA cohorts and we found that among these cohorts, cohorts with lower mean 25(OH)D concentration had stronger 25(OH)D–PFT associations (Figure 2). The proportion of ever smokers and of former smokers had significant linear associations with the 25(OH)D–PFT coefficients (Supplemental Figure 3), and these two variables were both highly correlated with mean 25(OH)D levels (Pearson's $r > 0.75$ for all pairwise correlations). The 25(OH)D–PFT association in EA cohorts varied by 25(OH)D assay method (meta-regression $p < 0.02$); the association was attenuated in cohorts using RIA compared to cohorts using LC-MS/MS (pairwise $p < 0.005$, Supplemental Figure 4). Mean age of each cohort was a significant positive modifier of the 25(OH)D–FEV₁ association, while time difference between 25(OH)D and spirometry measures did not affect the 25(OH)D–PFT association (Supplemental Figure 3).

To examine the potential impact of family relatedness between the FHS-Gen3 and the FHS-Offspring cohorts on the meta-analysis, sensitivity analysis confirmed that the findings were unchanged when either cohort was excluded (results not shown). In addition, the meta-analysis findings were not sensitive to exclusion of residual outliers.

In the EA cohorts, 25(OH)D had a greater positive association with FVC in current smokers than in never smokers ($\beta_{\text{current} \times 25(\text{OH})\text{D}} = 7.5$ mL for 10 nmol/L increment of 25(OH)D, $P = 0.047$). Similarly, 25(OH)D had a greater positive association with FVC in former smokers than in never smokers ($\beta_{\text{former} \times 25(\text{OH})\text{D}} = 7.9$ mL for 10 nmol/L increment of 25(OH)D, $P = 0.0065$) (Figure 3). For the FEV₁ outcome in the EA cohorts, the interaction coefficients for 25(OH)D and smoking status had the same positive direction as the coefficients for FVC, but were not statistically significant for either current ($P = 0.14$) or former smokers ($P = 0.14$). No statistical evidence of interaction of 25(OH)D and cigarette smoking was found in the AA cohorts for either outcome. To put the interaction finding into context, a 10 nmol/L higher serum 25(OH)D was associated with a 17.3 mL higher FVC in current smokers and a 16.6 mL higher FVC in former smokers, which was more than double the association magnitude in never smokers ($\beta = 7.8$ mL). A similar trend was found for the FEV₁ outcome in the EA cohorts. For 10 nmol/L higher serum 25(OH)D, FEV₁ was higher by 14.0 mL in current smokers, 12.0 mL in former smokers, and 8.0 mL in never smokers (Figure 4).

DISCUSSION

This study investigated the association of serum 25(OH)D with pulmonary function using multiple cohorts of different ancestries. We found a consistently positive association of serum 25(OH)D with FEV₁ and FVC across both EA and AA groups. In addition, in the EA group, a significantly stronger association was observed for current and former smokers, compared to never smokers.

A previous cross-sectional study in a European ancestry population (two Copenhagen cohorts: $n = 10,116$ and $n = 8,391$ respectively) similarly reported positive associations of 25(OH)D with FEV₁ percentage predicted and FVC percentage predicted, but not with FEV₁/FVC⁰. The magnitude of the association was about four times greater in the Copenhagen study, which may be due to the difference in the mean serum 25(OH)D (Danish median ~42 nmol/L vs. CHARGE median of ~65 nmol/L) given our finding that the 25(OH)D–PFT association was stronger in cohorts with lower serum 25(OH)D. Our finding for the serum 25(OH)D–FEV₁ association was similar in magnitude to the association reported in a British cohort of 6,789 participants with an average age of 45 years⁰, but weaker than a previous report from the FHS cohort⁰. Given that the rate of decline in FEV₁ at age 45 is increased by ~15 mL/year in current smokers⁰, we estimate that a 10 nmol/L higher 25(OH)D is similar to approximately 1 year of current smoking-related decline in FEV₁ for both ancestries, but in the opposite direction. Several putative biological mechanisms may support a causal association between low 25(OH)D levels and worse pulmonary function. First, lung tissue cells can locally convert 25(OH)D to 1,25-(OH)₂D⁰, the active form of vitamin D, which could improve the immune and anti-inflammatory responses in lungs via gene regulation⁽⁶⁾. If there is not enough circulating 25(OH)D, it is likely that the resolution of inflammation in lungs would be slower, which could have a negative impact on pulmonary function. In addition, 1,25-(OH)₂D in lungs, converted locally from 25(OH)D, can regulate the extracellular matrix homeostasis via the ERp60-mediated pathway⁰, and this is important for maintenance of lung structure. Furthermore, low vitamin D status could decrease circulating calcium status, which in turn can adversely affect thoracic skeleton mobility and respiratory muscle performance⁽⁶⁾.

Our findings show that the association of serum 25(OH)D with FEV₁ and FVC were stronger in magnitude in AA versus EA participants, although the difference by race did not reach statistical significance. The finding may reflect the lower serum 25(OH)D in AA participants, which is consistent with the meta-regression finding and with a previous study reporting attenuated associations at higher serum 25(OH)D (15). Future studies that investigate genetic variation in EAs and AAs in the context of serum 25(OH)D may help explain the differences.

In EA participants, the positive interaction terms between serum 25(OH)D and smoking status supported a stronger magnitude of association of serum 25(OH)D with FVC in current and former smokers than in never smokers, with a consistent, but not statistically significant, difference for FEV₁. The interaction finding is consistent with a prior cross-sectional National Health and Nutrition Examination Survey (NHANES) study, which reported a stronger 25(OH)D–FEV₁ association in current and former smokers than in never smokers

that was near statistical significance ($P = 0.06$)⁰. Given smokers have a higher level of oxidative stress and lower pulmonary function than never smokers, partly due to chronic inflammation in lung tissue, the stronger protective association of 25(OH)D on pulmonary function in smokers suggests a benefit for smokers. To explore this interaction, estimates of the 25(OH)D–PFT association were computed within each smoking category. In EA participants, the 25(OH)D–FEV₁ (or FVC) associations were statistically significant in all strata. Generally, in ever smokers of European ancestry, the coefficients for 25(OH)D were greater for FVC than for FEV₁.

Meta-regression provided additional evidence for effect modification by smoking. The proportion of ever smokers was a significant modifier of the association of serum 25(OH)D with FEV₁ and FVC. The higher the proportion of ever smokers, the greater the 25(OH)D–PFT association. More specifically, the proportion of former smokers explained the heterogeneity in the 25(OH)D–PFT association across cohorts more fully than the proportion of current smokers; this may be explained by a survival bias in older participants who were current smokers. The meta-regression, based on mean age of the cohorts, showed that cohorts with a higher mean age had a greater association magnitude of 25(OH)D with FEV₁. Given that meta-regression analysis uses cohort-level factors (e.g. mean age rather than age of each individual), ecological bias is possible⁰. Nevertheless, the age-related meta-regression finding was consistent with a prior NHANES study that showed the association of 25(OH)D and FEV₁ was stronger in people over age 60 compared to younger individuals⁰.

Several methodological considerations should be taken into account in interpreting the findings of this study. First, the meta-regression showed stronger 25(OH)D–PFT associations in cohorts with lower mean serum 25(OH)D, indicating a non-linear 25(OH)D–PFT association. This finding is consistent with a prior study in the FHS cohort, which reported a non-linear association and a stronger 25(OH)D–FEV₁ association in participants at risk of vitamin D deficiency (< 30 nmol/L)⁰. Second, serum 25(OH)D was measured by four different methods across the cohorts. For example, two cohorts with high mean 25(OH)D (> 90 nmol/L) used RIA methods. These same cohorts had a lower magnitude estimate of the 25(OH)D–PFT association; if the higher mean represents the ‘truth’ (and is not caused by measurement error in the RIA assay), then the lower 25(OH)D–PFT association may be primarily driven by the vitamin D distribution and not by the RIA method. Whether the assay method itself directly influences the estimate of the 25(OH)D–PFT association requires further data. Third, in this cross-sectional meta-analysis, minor differences were found in the time separation between the measurement of serum 25(OH)D and pulmonary function, but the meta-regression test for heterogeneity confirmed that time separation between measurements did not affect the 25(OH)D–PFT associations. Indeed, past studies with longitudinal measurements of serum 25(OH)D reported a high correlation of 25(OH)D measurements over a long period of time, with a correlation coefficient of 0.7 for measurements separated by 1 year, 0.5 for measurements separated by 5 years⁰, and 0.42–0.52 for measurements separated by 14 years⁰, which supports the use of a single 25(OH)D measurement to represent usual level. Fourth, residual confounding was unlikely given the consistent results across multiple cohorts in various settings. Weight was adjusted for the FVC outcome, given that higher weight and adiposity negatively affects lung volume (i.e., FVC)⁰; weight was not adjusted in the FEV₁ models, given FEV₁ is a measure of

airways obstruction and not physical restriction of lung volume. Physical activity was not adjusted because it is not a confounder in estimating the serum 25(OH)D–PFT association; while physical activity is known to contribute to oxygen utilization in lungs⁽¹⁾, little evidence and no biological rationale exists for a causal association of physical activity with either FEV₁ or FVC⁽¹⁾, which are markers for airways obstruction and lung volume, respectively. Finally, even though 3 cohorts (AGES, CARDIA, MESA) had the outcome or the exposure only measured in an ancillary study (random subset of the entire cohort), we do not expect selection bias to affect the estimate of the serum vitamin D–PFT association in this meta-analysis; indeed, the association magnitude and direction was consistent across all cohorts, regardless of the proportion of the original cohort contributing to the analysis. Thus, selection bias is expected to be negligible and would likely lead to an underestimated association, given the participants retained in the cohorts are expected to be, on average, healthier than those who were lost to follow-up.

This study meta-analyzed the serum 25(OH)D–PFT association across nine cohorts, according to a common pipeline that harmonized the variables and statistical analysis. The sample size comprised 17,569 EA participants from the United States; 5,269 EA participants from Iceland and the Netherlands; and 4,290 AA participants from the United States, all of whom were 19 to 95 years old. The sample provided excellent representation of the U.S. population, based on comparisons of demographic factors including sex, height, weight, smoking status, and COPD prevalence (~6.1%) to national surveys⁽¹⁾, which strengthens the external validity of the study's findings.

In summary, using meta-analysis, we estimated a positive association of serum 25(OH)D with the pulmonary function parameters FEV₁ and FVC in both EA and AA participants. Associations varied by smoking status in the EA group, with stronger serum 25(OH)D–PFT associations seen in current and former smokers. The observational design means we cannot infer a causal association, and future studies, such as randomized controlled trials or Mendelian randomization studies, are needed to further investigate the causality of 25(OH)D on pulmonary function.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Prof. James Booth in the Department of Biological Statistics and Computational Biology at Cornell University for providing advice on meta-analysis. We thank the staff (Lynn Johnson and Francoise Vermeylen) in Cornell Statistical Consulting Unit for their role in providing advice on methods. We also thank Hanfei Xu, Ph.D. student in the research group of Dr. Josée Dupuis for his assistance on residual model testing in the Framingham Heart Study.

Cohort Funding

This work was supported by National Institutes of Health (NIH) grant number R21 HL125574 funded by the National Heart, Lung, and Blood Institute (NHLBI) and the NIH Office of Dietary Supplements (ODS) (co-Principal Investigators [co-PIs]: DBH and PAC). The corresponding author (PAC) had full access to the data for the meta-analysis, and had final responsibility for the decision to submit for publication. No funding source had any role in the analysis of the data, the writing of the manuscript, or the decision to submit it. This work was also supported in part by R01HL077612 (PI: RGB) and by the Intramural Research Program of the National Institutes of

Health (NIH), National Institute of Environmental Health Sciences (ZO1 ES043012, PI: SJL). SJL is supported by the Intramural Research Program of NIH, National Institute of Environmental Health Sciences. Infrastructure for the CHARGE Consortium is supported in part by the NHLBI grant R01HL105756.

The Age, Gene/Environment Susceptibility (AGES)–Reykjavik Study has been funded by NIH contracts N01-AG-1–2100 and 271201200022C, the National Institute on Aging (NIA) Intramural Research Program, Hjartavernd (the Icelandic Heart Association), and the Althingi (the Icelandic Parliament). The study is approved by the Icelandic National Bioethics Committee, VSN: 00–063. The researchers are indebted to the participants for their willingness to participate in the study.

The Atherosclerosis Risk in Communities Study is carried out as a collaborative study supported by NHLBI contracts HHSN268201100005C, HHSN268201100006C, HHSN268201100007C, HHSN268201100008C, HHSN268201100009C, HHSN268201100010C, HHSN268201100011C, and HHSN268201100012C. 25(OH)D measurements were conducted with the support of R01 HL103706 from the NHLBI and R01 HL103706-S1 from the NIH ODS. The authors thank the staff and participants of the ARIC study for their important contributions.

This Cardiovascular Health Study (CHS) research was supported by NHLBI contracts HHSN268201200036C, HHSN268200800007C, N01HC55222, N01HC85079, N01HC85080, N01HC85081, N01HC85082, N01HC85083, N01HC85086; and NHLBI grants U01HL080295, R01HL085251, R01HL087652, R01HL105756, R01HL103612, R01HL120393, and R01HL130114 with additional contribution from the National Institute of Neurological Disorders and Stroke (NINDS). Additional support was provided through R01AG023629 from NIA. A full list of principal CHS investigators and institutions can be found at CHS-NHLBI.org. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. Vitamin D measurements were made possible by NHLBI (R01HL084443–01A2).

This work in Framingham Heart Study was supported by NHLBI's Framingham Heart Study contract (N01-HC-25195 and HHSN268201500001I). Vitamin D measurements in the Framingham study were made possible by NIA (R01 AG14759 to SLB.).

The Health Aging and Body Composition cohort study was supported by NIA contracts N01AG62101, N01AG2103, and N01AG62106, NIA grant R01-AG028050, NINR grant R01-NR012459, and in part by the Intramural Research Program of the NIA, NIH. This research was further supported by RC1AG035835, and the serum vitamin D assays were supported by R01AG029364.

The Multi-Ethnic Study of Atherosclerosis (MESA) study is conducted and supported by NHLBI in collaboration with MESA investigators. Support for MESA is provided by contracts HHSN268201500003I, N01-HC-95159, N01-HC-95160, N01-HC-95161, N01-HC-95162, N01-HC-95163, N01-HC-95164, N01-HC-95165, N01-HC-95166, N01-HC-95167, N01-HC-95168, and N01-HC-95169 from NHLBI, UL1-TR-000040, UL1-TR-001079, and UL1-TR-001881 from NCR, and DK063491 from the NIDDK. The MESA Lung study was supported by grants R01 HL077612, RC1 HL100543 and R01 HL093081 from NHLBI. Support for the Mineral Metabolite dataset was provided by grant HL096875.

The Rotterdam Study is funded by Erasmus Medical Center and Erasmus University, Rotterdam, the Netherlands; the Organization for the Health Research and Development (ZonMw); the Research Institute for Diseases in the Elderly (RIDE); the Dutch Ministry of Education, Culture, and Science; the Dutch Ministry for Health, Welfare, and Sports; the European Commission (DG XII), and the Municipality of Rotterdam. LL was a postdoctoral fellow of the Research Foundation—Flanders (FWO) in Brussels, Belgium. Part of this work was supported by a FWO-grant G035014N. DSM Nutritional Products AG, Kaiseraugst, Switzerland, sponsored the Vitamin D serum analyses. The authors are grateful to the study participants, the staff from the Rotterdam Study, and the participating general practitioners and pharmacists.

The Coronary Artery Risk Development in Young Adults Study (CARDIA) is supported by contracts HHSN268201300025C, HHSN268201300026C, HHSN268201300027C, HHSN268201300028C, HHSN268201300029C, and HHSN268200900041C from the National Heart, Lung, and Blood Institute (NHLBI), the Intramural Research Program of the National Institute on Aging (NIA), and an intra-agency agreement between NIA and NHLBI (AG0005).

Abbreviation Footnote

25(OH)D	25-Hydroxyvitamin D
AA	African Ancestry

AGES	Age Gene Environment Susceptibility Study—Reykjavik Iceland
ARIC	Atherosclerosis Risk in Communities Study
CARDIA	Coronary Artery Risk Development in Young Adults Study
CHARGE	Cohorts for Heart and Aging Research in Genomic Epidemiology Consortium
CHS	Cardiovascular Health Study
CLIA	Chemiluminescence Immunoassay
COPD	Chronic Obstructive Pulmonary Disease
EA	European Ancestry
FEV₁	Forced Expiratory Volume in the First Second
FHS (Offspring)	Framingham Heart Study—Offspring Cohort
FHS (Gen3)	Framingham Heart Study—Generation 3 Cohort
FVC	Forced Vital Capacity
HABC	Health Aging and Body Composition Study
LC-MS/MS	Liquid Chromatography in Tandem with Mass Spectrometry
MESA	Multi-Ethnic Study of Atherosclerosis
NHANES	National Health and Nutrition Examination Survey
PFT	Pulmonary Function Test
RIA	Radioimmunoassay
RS	Rotterdam (Netherlands) Study

REFERENCES

1. Minino AM, Murphy SL, Xu J, et al. (2011) Deaths: final data for 2008. *Natl Vital Stat Rep* 59, 1–126.
2. Collaborators GBDCoD (2017) Global, regional, and national age-sex specific mortality for 264 causes of death, 1980–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet* 390, 1151–210. [PubMed: 28919116]
3. Weiss ST (2010) Lung function and airway diseases. *Nat Genet* 42, 14–6. [PubMed: 20037613]
4. Forey BA, Thornton AJ, Lee PN (2011) Systematic review with meta-analysis of the epidemiological evidence relating smoking to COPD, chronic bronchitis and emphysema. *BMC Pulm Med* 11, 36. [PubMed: 21672193]
5. Guerra S, Stern DA, Zhou M, et al. (2013) Combined effects of parental and active smoking on early lung function deficits: a prospective study from birth to age 26 years. *Thorax* 68, 1021–8. [PubMed: 23847259]

6. Fletcher JM, Basdeo SA, Allen AC, et al. (2012) Therapeutic use of vitamin D and its analogues in autoimmunity. *Recent Pat Inflamm Allergy Drug Discov* 6, 22–34. [PubMed: 22122009]
7. Litonjua AA, ebrary Inc. (2012) Vitamin D and the lung mechanisms and disease associations. New York: Humana Press/Springer, <http://proxy.library.cornell.edu/login?url=http://link.springer.com/openurl?genre=book&isbn=978-1-61779-887-0>.
8. Bikle DD, Gee E, Halloran B, et al. (1986) Assessment of the free fraction of 25-hydroxyvitamin D in serum and its regulation by albumin and the vitamin D-binding protein. *J Clin Endocrinol Metab* 63, 954–9. [PubMed: 3745408]
9. Nykjaer A, Dragun D, Walther D, et al. (1999) An endocytic pathway essential for renal uptake and activation of the steroid 25-(OH) vitamin D3. *Cell* 96, 507–15. [PubMed: 10052453]
10. Zerwekh JE (2008) Blood biomarkers of vitamin D status. *Am J Clin Nutr* 87, 1087S–91S. [PubMed: 18400739]
11. Nielson CM, Jones KS, Chun RF, et al. (2016) Free 25-Hydroxyvitamin D: Impact of Vitamin D Binding Protein Assays on Racial-Genotypic Associations. *J Clin Endocrinol Metab* 101, 2226–34. [PubMed: 27007693]
12. Freedman BI, Register TC (2012) Effect of race and genetics on vitamin D metabolism, bone and vascular health. *Nat Rev Nephrol* 8, 459–66. [PubMed: 22688752]
13. Black PN, Scragg R (2005) Relationship between serum 25-hydroxyvitamin d and pulmonary function in the third national health and nutrition examination survey. *Chest* 128, 3792–8. [PubMed: 16354847]
14. Choi CJ, Seo M, Choi WS, et al. (2013) Relationship between serum 25-hydroxyvitamin D and lung function among Korean adults in Korea National Health and Nutrition Examination Survey (KNHANES), 2008–2010. *J Clin Endocrinol Metab* 98, 1703–10. [PubMed: 23533242]
15. Hansen JG, Gao W, Dupuis J, et al. (2015) Association of 25-Hydroxyvitamin D status and genetic variation in the vitamin D metabolic pathway with FEV1 in the Framingham Heart Study. *Respir Res* 16, 81. [PubMed: 26122139]
16. Afzal S, Lange P, Bojesen SE, et al. (2014) Plasma 25-hydroxyvitamin D, lung function and risk of chronic obstructive pulmonary disease. *Thorax* 69, 24–31. [PubMed: 23908128]
17. Berry DJ, Hesketh K, Power C, et al. (2011) Vitamin D status has a linear association with seasonal infections and lung function in British adults. *Br J Nutr* 106, 1433–40. [PubMed: 21736791]
18. Thuesen BH, Skaaby T, Husemoen LL, et al. (2015) The association of serum 25-OH vitamin D with atopy, asthma, and lung function in a prospective study of Danish adults. *Clin Exp Allergy* 45, 265–72. [PubMed: 24575884]
19. Tolppanen AM, Williams D, Henderson J, et al. (2011) Serum 25-hydroxy-vitamin D and ionised calcium in relation to lung function and allergen skin tests. *Eur J Clin Nutr* 65, 493–500. [PubMed: 21326268]
20. Shaheen SO, Jameson KA, Robinson SM, et al. (2011) Relationship of vitamin D status to adult lung function and COPD. *Thorax* 66, 692–8. [PubMed: 21653927]
21. Niruban SJ, Alagiakrishnan K, Beach J, et al. (2015) Association between vitamin D and respiratory outcomes in Canadian adolescents and adults. *J Asthma* 52, 653–61. [PubMed: 25563060]
22. Tolppanen AM, Sayers A, Granell R, et al. (2013) Prospective association of 25-hydroxyvitamin d3 and d2 with childhood lung function, asthma, wheezing, and flexural dermatitis. *Epidemiology* 24, 310–9. [PubMed: 23377091]
23. Lange NE, Sparrow D, Vokonas P, et al. (2012) Vitamin D deficiency, smoking, and lung function in the Normative Aging Study. *Am J Respir Crit Care Med* 186, 616–21. [PubMed: 22822023]
24. Khan S, Mai XM, Chen Y (2013) Plasma 25-hydroxyvitamin D associated with pulmonary function in Canadian adults with excess adiposity. *Am J Clin Nutr* 98, 174–9. [PubMed: 23656903]
25. O’Neal JD (2015) Vitamin D supplementation regimens for HIV-infected patients: a historical chart review. *Scholar Archive*.
26. Mannino DM, Diaz-Guzman E, Buist S (2011) Pre- and post-bronchodilator lung function as predictors of mortality in the Lung Health Study. *Respir Res* 12, 136. [PubMed: 21991942]

27. Phinney KW (2008) Development of a standard reference material for vitamin D in serum. *Am J Clin Nutr* 88, 511S–2S. [PubMed: 18689392]
28. Lederer DJ, Enright PL, Kawut SM, et al. (2009) Cigarette smoking is associated with subclinical parenchymal lung disease: the Multi-Ethnic Study of Atherosclerosis (MESA)-lung study. *Am J Respir Crit Care Med* 180, 407–14. [PubMed: 19542480]
29. Fujiyoshi A, Polgreen LE, Hurley DL, et al. (2013) A cross-sectional association between bone mineral density and parathyroid hormone and other biomarkers in community-dwelling young adults: the CARDIA study. *J Clin Endocrinol Metab* 98, 4038–46. [PubMed: 23966240]
30. Harris TB, Launer LJ, Eiriksdottir G, et al. (2007) Age, Gene/Environment Susceptibility-Reykjavik Study: multidisciplinary applied phenomics. *Am J Epidemiol* 165, 1076–87. [PubMed: 17351290]
31. Runarsdottir SB, Gudmundsson G, Aspelund T, et al. (2013) Prevalence of airflow obstruction in nonsmoking older individuals using different spirometric criteria: the AGES Reykjavik Study. *COPD* 10, 493–9. [PubMed: 23875743]
32. Morris TP, Fisher DJ, Kenward MG, et al. (2018) Meta-analysis of Gaussian individual patient data: Two-stage or not two-stage? *Stat Med* 37, 1419–38. [PubMed: 29349792]
33. Higgins JP, Thompson SG, Deeks JJ, et al. (2003) Measuring inconsistency in meta-analyses. *BMJ* 327, 557–60. [PubMed: 12958120]
34. Borenstein M, Wiley InterScience (Online service) (2009) Introduction to meta-analysis. Chichester, U.K.: John Wiley & Sons, http://encompass.library.cornell.edu/cgi-bin/checkIP.cgi?access=gateway_standard%26url=http://onlinelibrary.wiley.com/book/10.1002/9780470743386.
35. Mirabelli MC, Preisser JS, Loehr LR, et al. (2016) Lung function decline over 25 years of follow-up among black and white adults in the ARIC study cohort. *Respir Med* 113, 57–64. [PubMed: 26905512]
36. Hansdottir S, Monick MM, Hinde SL, et al. (2008) Respiratory epithelial cells convert inactive vitamin D to its active form: potential effects on host defense. *J Immunol* 181, 7090–9. [PubMed: 18981129]
37. Colotta F, Jansson B, Bonelli F (2017) Modulation of inflammatory and immune responses by vitamin D. *J Autoimmun* 85, 78–97. [PubMed: 28733125]
38. Dimeloe S, Richards DF, Urry ZL, et al. (2012) 1alpha,25-dihydroxyvitamin D3 promotes CD200 expression by human peripheral and airway-resident T cells. *Thorax* 67, 574–81. [PubMed: 22334534]
39. Boyan BD, Wong KL, Fang M, et al. (2007) 1alpha,25(OH)2D3 is an autocrine regulator of extracellular matrix turnover and growth factor release via ERp60 activated matrix vesicle metalloproteinases. *J Steroid Biochem Mol Biol* 103, 467–72. [PubMed: 17224270]
40. Herr C, Greulich T, Koczulla RA, et al. (2011) The role of vitamin D in pulmonary disease: COPD, asthma, infection, and cancer. *Respir Res* 12, 31. [PubMed: 21418564]
41. Schlaich C, Minne HW, Bruckner T, et al. (1998) Reduced pulmonary function in patients with spinal osteoporotic fractures. *Osteoporos Int* 8, 261–7. [PubMed: 9797911]
42. Thompson SG, Higgins JP (2002) How should meta-regression analyses be undertaken and interpreted? *Stat Med* 21, 1559–73. [PubMed: 12111920]
43. Hofmann JN, Yu K, Horst RL, et al. (2010) Long-term variation in serum 25-hydroxyvitamin D concentration among participants in the Prostate, Lung, Colorectal, and Ovarian Cancer Screening Trial. *Cancer Epidemiol Biomarkers Prev* 19, 927–31. [PubMed: 20332255]
44. Jorde R, Sneve M, Hutchinson M, et al. (2010) Tracking of serum 25-hydroxyvitamin D levels during 14 years in a population-based study and during 12 months in an intervention study. *Am J Epidemiol* 171, 903–8. [PubMed: 20219763]
45. Amara CE, Koval JJ, Paterson DH, et al. (2001) Lung function in older humans: the contribution of body composition, physical activity and smoking. *Ann Hum Biol* 28, 522–36. [PubMed: 11572518]
46. Burri PH, Gehr P, Muller K, et al. (1976) Adaptation of the growing lung to increased VO₂. I. IDPN as inducer of hyperactivity. *Respir Physiol* 28, 129–40. [PubMed: 988630]
47. Cheng YJ, Macera CA, Addy CL, et al. (2003) Effects of physical activity on exercise tests and respiratory function. *Br J Sports Med* 37, 521–8. [PubMed: 14665592]

48. Jamal A, Homa DM, O'Connor E, et al. (2015) Current cigarette smoking among adults - United States, 2005–2014. *MMWR Morb Mortal Wkly Rep* 64, 1233–40. [PubMed: 26562061]
49. Fryar CD, Gu Q, Ogden CL (2012) Anthropometric reference data for children and adults: United States, 2007–2010. *Vital Health Stat* 11, 1–48.
50. Ward BW, Nugent CN, Blumberg SJ, et al. (2017) Measuring the Prevalence of Diagnosed Chronic Obstructive Pulmonary Disease in the United States Using Data From the 2012–2014 National Health Interview Survey. *Public Health Rep* 132, 149–56. [PubMed: 28135423]




Figure 1. Forest plots of the meta-analysis of serum 25(OH)D on FEV1 and FVC across cohorts in the CHARGE Consortium, stratified by participant ancestry.

Associations are presented for serum 25(OH)D on (A) FEV1 in European ancestry cohorts ($n = 22,787$). (B) FEV1 in African ancestry cohorts ($n = 4,282$). (C) FVC in European ancestry cohorts ($n = 22,777$). (D) FVC in African ancestry cohorts ($n = 4,239$). β (unit: mL) denotes the coefficient from the fixed-effects meta-analysis for serum 25(OH)D on the pulmonary function outcome per 1 nmol/L increment of 25(OH)D, with its 95% confidence interval. Cohorts findings were ordered from the least to the most precise, and heterogeneity is presented (I^2).




Figure 2. Meta-regression of mean serum 25(OH)D levels against the association estimates of 25(OH)D with PFT in nine European ancestry cohorts in the CHARGE Consortium. (A) FEV_1 outcome (coefficient unit: mL per 1 nmol/L 25(OH)D), and (B) FVC outcome (coefficient unit: mL per 1nmol/L 25(OH)D). The modifier is mean serum 25(OH)D level of each nine cohorts. A linear regression line is present for each sub-figure, with a meta-regression p-value of 0.0006 for the FEV_1 outcome, and 0.005 for the FVC outcome. The figure also shows the measurement method for the serum 25(OH)D assay (legend shows symbols for each of the 4 assay methods).




Figure 3. Forest plots of the interaction meta-analysis of serum 25(OH)D and smoking status on FVC in the European ancestry cohorts in the CHARGE Consortium ($n = 22,777$). (A) Current Smokers and (B) Former Smokers. β (unit: mL) denotes the interaction term coefficient of 25(OH)D and smoking status on FVC from the fixed effects meta-analysis, per 1 nmol/L increment of 25(OH)D, with its 95% confidence interval. Cohorts were ordered from the least to the most precise, and heterogeneity is presented (I^2).




Figure 4. Meta-analysis of the association of serum 25(OH)D–PFT outcomes among current, former, and never smokers in the European ancestry cohorts in the CHARGE Consortium. FEV₁ and FVC are presented for each smoking status. β (unit: mL) denotes that 1 nmol/L higher serum 25(OH)D was associated with a β mL higher FEV₁ (or FVC), calculated from an analysis including the interaction of serum 25(OH)D and smoking status. The error bar represents ± 1 standard error. We used 22,787 EA participants for the FEV₁ outcome and 22,777 EA participants for the FVC outcome.

Table 1.

Cross-sectional participant characteristics of each cohort in the CHARGE Consortium (*n* = 27,128)*

European Ancestry Cohort	ARIC	CARDIA	CHS [†]	HABC [‡]	MESA	AGES	FHS (Offspring)	FHS (Gen3)	RS
Number of participants	8,327	172	1,297	1,411	1,113	1,685	1,639	3,610	3,584
Males, percentage	46.0	58.7	30.15	53.3	49.2	40.8	48.1	47.3	44.6
Current Smoker, percentage	23.4	11.6	9.4	6.5	8.4	9.8	14.3	15.3	16.0
Former Smoker, percentage	34.9	16.3	44.9	49.8	47.2	42.4	50.5	28.0	52.9
Pack-years [§]	28.0 (20.9)	6.2 (7.2)	28.1 (25.3)	36.4 (32.0)	30.1 (29.6)	24.6 (21.9)	26.5 (22.8)	12.4 (13.4)	22.9 (21.6)
Age, year	54.2 (5.7)	34.8 (3.1)	73.7 (4.4)	73.7 (2.8)	66.3 (9.9)	76.2 (5.6)	59.4 (9.3)	40.2 (8.7)	64.4 (9.7)
Height, m	1.69 (0.09)	1.73 (0.09)	1.63 (0.09)	1.67 (0.09)	1.69 (0.10)	1.67 (0.09)	1.68 (0.09)	1.71 (0.09)	1.69 (0.09)
Weight, kg [¶]	76.8 (16.2)	76.9 (17.0)	70.6 (14.2)	74.5 (14.5)	79.7 (17.3)	75.4 (14.7)	79.4 (17.2)	78.6 (18.4)	79.5 (14.6)
FEV ₁ , mL	2,946 (767)	3,881 (743)	2,010 (611)	2,324 (649)	2,556 (768)	2,142 (670)	2,724 (757)	3,592 (787)	2848 (866)
FVC, mL	3,987 (973)	4,967 (999)	2,881 (829)	3,118 (810)	3,492 (995)	2,877 (837)	3,711 (950)	4,621 (999)	3692 (1063)
FEV ₁ /FVC	0.739 (0.077)	0.785 (0.060)	0.700 (0.095)	0.745 (0.078)	0.734 (0.087)	0.744 (0.087)	0.733 (0.078)	0.779 (0.063)	0.771 (0.082)
Serum 25(OH)D, nmol/L^{**}	64.7 (21.8)	95.0 (35.3)	68.0 (27.9)	72.2 (25.6)	75.6 (28.2)	52.4 (23.5)	49.2 (18.9)	92.8 (36.0)	61.0 (27.4)
Never smoker	64.3 (21.0)	95.4 (34.4)	67.1 (25.1)	73.7 (25.9)	76.5 (27.7)	54.1 (22.8)	49.6 (18.6)	93.2 (35.4) ^{††}	59.7 (25.9)
Former smoker	67.1 (21.5)	94.5 (43.0)	69.4 (29.4)	71.7 (24.8)	76.2 (28.5)	52.3 (24.1)	49.8 (18.6)	93.5 (37.0)	62.3 (27.7)
Current smoker	61.8 (23.1)	92.7(29.5)	65.4 (33.2)	65.0 (28.1)	66.9 (28.2)	44.5 (22.7)	45.9 (20.6)	89.9 (36.3)	59.5 (29.4)
Method of 25(OH)D measurement	LC-MS/MS	RIA	LC-MS/MS	RIA	LC-MS/MS	CLIA	RIA	RIA	Electro-CLIA
Time from 25(OH)D to PFT, days ^{‡‡}	-1,073 (67)	1,122 (89)	363 (29)	-382 (39)	1,765 (112)	1 (5)	133 (377)	2 (61)	846 (808)
Season of 25(OH)D measurement, percentage^{§§}									
Spring	31.2	8.1	20.5	30.5	29.0	22.4	29.2	26.8	29.6
Summer	26.1	36.1	30.1	18.1	22.2	12.4	11.0	29.6	18.9
Fall	23.3	34.3	29.6	22.8	24.9	33.8	29.1	24.1	30.0
Winter	19.5	21.5	19.8	28.6	23.9	31.4	30.7	19.4	21.5
African Ancestry Cohort									
Number of participants	2,339	157	168	863	763				
Males, percentage	35.3	51.6	25.6	44.5	47.4				

European Ancestry Cohort	ARIC	CARDIA	CHS [†]	HABC [‡]	MESA	AGES	FHS (Offspring)	FHS (Gen3)	RS
Current Smoker, percentage	27.5	26.1	10.7	15.8	15.7				
Former Smoker, percentage	23.9	9.6	42.9	39.3	38.3				
Pack-years [§]	21.4 (20.7)	5.3 (4.6)	21.9 (18.3)	29.4 (23.4)	23.6 (21.8)				
Age, year	53.3 (5.7)	33.9 (3.2)	71.9 (4.5)	73.4 (2.9)	65.6 (9.7)				
Height, m	1.68 (0.09)	1.71 (0.10)	1.63 (0.08)	1.66 (0.09)	1.68 (0.10)				
Weight, kg [¶]	83.5 (17.1)	82.2 (16.9)	75.7 (13.3)	78.2 (15.1)	84.3 (16.8)				
FEV ₁ , mL	2,495 (638)	3,237 (709)	1,801 (508)	1,958 (566)	2,200 (667)				
FVC, mL	3,255 (806)	4,077 (920)	2,507 (706)	2,594 (712)	2,933 (869)				
FEV ₁ /FVC	0.768 (0.077)	0.799 (0.070)	0.723 (0.076)	0.757 (0.090)	0.755 (0.093)				
Serum 25(OH)D, nmol/L^{**}	47.4 (17.5)	69.4 (31.2)	44.6 (21.1)	51.8 (22.4)	47.9 (22.3)				
Never smoker	46.8 (16.7)	71.3 (30.1)	43.7 (19.2)	51.8 (22.7)	49.1 (22.3)				
Former smoker	48.5 (18.0)	69.2 (35.6)	47.2 (24.2)	52.3 (21.8)	49.3 (22.6)				
Current smoker	47.5 (18.4)	64.8 (32.4)	38.3 (14.9)	50.4 (23.2)	40.9 (20.0)				
Method of 25(OH)D measurement	LC-MS/MS	RIA	LC-MS/MS	RIA	LC-MS/MS				
Time from 25(OH)D to PFT, days ^{‡‡}	-1,054 (114)	1,101 (104)	350 (26)	-390 (53)	1,719 (115)				
Season of 25(OH)D measurement, percentage^{§§}									
Spring	30.0	10.2	58.9	35.6	34.6				
Summer	30.7	56.0	7.1	16.2	23.5				
Fall	20.7	23.6	8.9	24.9	19.7				
Winter	18.6	10.2	25.0	23.3	22.3				

* Data are presented as mean (SD) unless otherwise indicated; AGES, RS, and FHS only have participants of European ancestry; $n = 22,838$ for EAs, $n = 4,290$ for AAs, total $n = 27,128$.

[†]The number of participants used to compute descriptive statistics in CHS excluded those who had residual outliers based on the preliminary models ($n = 8$ for EAs and $n = 6$ for AAs); while other cohorts used the number of participants before applying residual exclusion for the descriptive statistics.

[‡]Numbers vary slightly for different outcomes in HABC (For the FVC outcome, $n = 1385$ for EAs and $n = 821$ for AAs; for the ratio outcome, $n = 1382$ for EAs and $n = 817$ for AAs). The numbers of participants for the FEV₁ outcome are used. However, the descriptive statistics is similar across different outcomes.

[§]Pack-years is calculated only among current and former smokers in each cohort.

[¶]We used 1,554 ever smokers here, instead of a total of 1,561 ever smokers in the Gen3 cohort, because the pack-years of seven ever smokers were so small that they were coded as 0. Therefore, these seven ever smokers do not contribute to the pack-years descriptive statistics here.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

[¶]The number of participants who have weight data is slightly different from the total number of participants in each cohort. However, the descriptive statistics of weight stays similar.

^{**} Mean (SD) of serum 25(OH)D level for all the participants in each cohort, and mean (SD) of 25(OH)D level in participants with each smoking status are shown here, stratified by ancestry.

^{††} We used 2,046 never smokers, rather than a total of 2,049 never smokers in the Gen3 cohort, to compute the 25(OH)D level in never smokers.

^{‡‡} The time difference is the interval between the time when pulmonary function was measured and the time when serum vitamin D was measured. The difference is positive, if the serum vitamin D was measured before the pulmonary function test; while the value is negative, if the serum vitamin D was measured after the pulmonary function test.

^{§§} The proportion of participants in each season when their serum was measured was rounded (thus rounding errors mean sums may not be exactly 100%).