

UC Riverside

UC Riverside Previously Published Works

Title

Selectively breeding for high voluntary physical activity in female mice does not bestow inherent characteristics that resemble eccentric remodeling of the heart, but the mini-muscle phenotype does.

Permalink

<https://escholarship.org/uc/item/96v0m1px>

Journal

Sports Medicine and Health Science, 5(3)

Authors

Leszczynski, Eric
Schwartz, Nicole
McPeck, Ashley
[et al.](#)

Publication Date

2023-09-01

DOI

10.1016/j.smhs.2023.07.003

Peer reviewed



Original Article

Selectively breeding for high voluntary physical activity in female mice does not bestow inherent characteristics that resemble eccentric remodeling of the heart, but the mini-muscle phenotype does

Eric C. Leszczynski^{a,*}, Nicole E. Schwartz^b, Ashley C. McPeck^a, Katharine D. Currie^a, David P. Ferguson^{a,*}, Theodore Garland Jr.^b

^a Department of Kinesiology, Michigan State University, East Lansing, MI, USA

^b Department of Evolution, Ecology, and Organismal Biology, University of California Riverside, Riverside, CA, USA

ARTICLE INFO

Keywords:

Echocardiography

Heart

Voluntary physical activity

Cardiovascular disease

Wheel running

ABSTRACT

Physical activity engagement results in a variety of positive health outcomes, including a reduction in cardiovascular disease risk partially due to eccentric remodeling of the heart. The purpose of this investigation was to determine if four replicate lines of High Runner mice that have been selectively bred for voluntary exercise on wheels have a cardiac phenotype that resembles the outcome of eccentric remodeling. Adult females (average age 55 days) from the 4 High Runner and 4 non-selected control lines were anaesthetized via vaporized isoflurane, then echocardiographic images were collected and analyzed for structural and functional differences. High Runner mice in general had lower ejection fractions compared to control mice lines (2-tailed $p = 0.0236$) and tended to have thicker walls of the anterior portion of the left ventricle ($p = 0.065$). However, a subset of the High Runner individuals, termed mini-muscle mice, had greater ejection fraction ($p = 0.0006$), fractional shortening percentage ($p < 0.0001$), and ventricular mass at dissection ($p < 0.0027$ with body mass as a covariate) compared to non-mini muscle mice. Mice from replicate lines bred for high voluntary exercise did not all have inherent positive cardiac functional or structural characteristics, although a genetically unique subset of mini-muscle individuals did have greater functional cardiac characteristics, which in conjunction with their previously described peripheral aerobic enhancements (e.g., increased capillarity) would partially account for their increased $\dot{V}O_{2max}$.

Introduction

Routinely engaging in moderate to vigorous physical activity is associated with reduced risk for non-communicable diseases,^{1–3} including the incidence of cardiovascular disease, hypertension, type II diabetes, and some types of cancer.^{1,4} In humans, the regulation of physical activity is under complex control by both environmental (access to sidewalks, availability of public transportation, perceived safety of the environment, and high socioeconomic status)⁵ and biological factors (ability of skeletal muscle to perform contractions, ability of the cardiovascular system to delivery oxygen to muscle, and neurobiological factors related to motivation, reward, and fatigue).^{4,6}

Given that genetic and genomic variants must underlie variation in some of the biological factors that regulate physical activity, the Garland Laboratory has used selective breeding for voluntary exercise to develop

a mouse model for the control of physical activity.⁷ Specifically, four replicate lines of High Runner (HR) mice have been bred for voluntary wheel-running behavior as young adults, while four non-selected control (C) lines are bred without regard to wheel running.⁸ Since reaching selection limits around generations 17–27 (depending on replicate line and sex), mice from the HR lines run ~2.5-3-fold more revolutions/day.⁹ Even without access to exercise wheels (i.e., in the absence of exercise training), the HR lines have a variety of co-segregating traits that are generally viewed as being associated with positive health outcomes and longevity, including greater activity when housed without wheels,¹⁰ reduced body fat,^{11,12} higher endurance on a motorized treadmill,¹³ and higher maximum aerobic capacity [i.e., maximal oxygen consumption, ($\dot{V}O_{2max}$)],^{14–17} as compared with the control lines.

One remarkable discovery in the HR mouse selection experiment is a muscle-mass polymorphism, in which some individuals have triceps surae and whole hindlimb muscles ~50% as heavy as in normal

* Corresponding author. Department of Kinesiology, Michigan State University, 308 W. Circle Dr. Room 39, East Lansing, MI, 48824, USA.

E-mail addresses: Leszczy8@msu.edu (E.C. Leszczynski), Fergu312@msu.edu (D.P. Ferguson).

<https://doi.org/10.1016/j.smhs.2023.07.003>

Received 20 December 2022; Received in revised form 18 May 2023; Accepted 5 July 2023

Available online 7 July 2023

2666-3376/© 2023 Chengdu Sport University. Publishing services by Elsevier B.V. on behalf of KeAi Communications Co. Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

List of abbreviations:

C	Control mouse
HR	High Runner mouse
$\dot{V} O_{2max}$	Maximal Oxygen Consumption
C-to-T	Cytosine to Thymine
Hsd	ICR Harlan-Sprague-Dawley
N	Sample size
CO₂	Carbon Dioxide
°C	Degrees Celsius
M-mode	Motion Mode
B-mode	Brightness Mode
REML	Restricted Maximum Likelihood
EKG	Electrocardiogram
DF	Degrees of Freedom
LSM	Least Squares Mean
SE	Standard Error
g	Grams
mg	Milligrams
mL	Milliliters
μL	Microliters
mm	Millimeters

individuals.¹⁸ This “mini-muscle” phenotype is caused by a single C-to-T base pair change (*Myh4*^{Minimsc}) between exon 11 and 12 of the *Myh4* skeletal muscle gene¹⁹ that behaves as a Mendelian recessive (i.e., heterozygotes have the normal phenotype). The reduced muscle mass is attributable to a strong reduction, and sometimes absence, of the fast glycolytic (type 2B) fiber type in locomotor muscles that normally contain this fiber type.^{20–22} Mini-muscle mice have a variety of correlated phenotypes (i.e., pleiotropic effects of the mini-muscle allele), including increased capillarity in medial gastrocnemius, increased mass-specific citrate synthase and myoglobin concentration in gastrocnemius, larger soleus muscles, larger hearts and other internal organs, a longer QRS complex in electrocardiograms, and a higher $\dot{V} O_{2max}$ even than other HR, non-mini mice.^{15,16,23–27} However, it is unknown if mini-muscle individuals have functional cardiovascular traits that would facilitate their high voluntary running speeds on wheels and their high $\dot{V} O_{2max}$. Along with the muscle phenotype displayed in the mini-muscle mice there could be inherent/genetic adaptations to the heart. In particular, eccentric remodeling of the heart is associated with increased left ventricle chamber size and modest increases in myocardial wall thickness, which increases cardiac health and reduces mortality from cardiovascular disease.²⁸

The purpose of the present investigation was to determine if selective breeding for high physical activity also results in an altered cardiac phenotype resembling cardiac eccentric remodeling when measured by echocardiography. If mice from the HR lines, or perhaps just the subset of mini-muscle HR mice, have inherent cardiac differences (i.e., in the absence of chronic exercise) that resemble eccentric remodeling, then further investigations could reveal potential genetically based cardio-protective mechanisms. Cardiac differences observed between mini-muscle and non-mini muscle mice would, specifically, suggest peripheral adaptations may assist with cardiac remodeling to facilitate greater oxygen delivery. However, if such phenotypes do not exist in untrained HR or mini-muscle mice, then the importance of regular engagement in physical activity to reduce the risk of cardiovascular disease would be reinforced. All mice included in this study were female, which we hypothesized would highlight inherent cardiac differences due to higher levels of wheel running in female mice.¹⁸

Methods*Ethical approval*

All animals were housed in a room temperature controlled at ~22 °C under a 12-h dark/light cycle. Food and water were provided *ad libitum* throughout the experimental procedures, and all procedures were approved by the University of California, Riverside Institutional Animal Care and Use Committee.

Selection experiment

The ongoing selection experiment was initiated in 1993 with 224 outbred Hsd:ICR (Harlan-Sprague-Dawley) mice. Mice were bred randomly for two generations and then separated into eight lines, with four to be bred for high activity (HR, $n = 31$) and four non-selected control (C, $n = 32$) lines.⁸ Briefly, at ~6–8 weeks of age, mice are housed individually in cages with attached wheels for 6 days. In the HR lines, breeders are chosen based on revolutions run on days 5 and 6; in the C lines, breeders are chosen without regard to running (for further details, see^{8,9}).

Originally present at low frequency in the base population (~7%), one of the C lines and two of the HR lines had individuals with hindlimb muscles that were ~50% smaller than normal-muscled individuals.^{18,24} This “mini-muscle” phenotype is no longer present in the C line, is fixed in HR line 3, and is polymorphic in HR line 6.^{15,29,30} Population-genetic modeling indicate that the mini-muscle trait was either neutral or under negative selection in the C lines, but favored in the HR lines.¹⁸ Of the 63 mice included in this study, all 7 in HR line 3 had the mini-muscle phenotype and 1 of the 8 mice in HR line 6 had it ($n = 55$ normal, $n = 8$ mini-muscle).

Experimental subjects and timeline

From generation 88, we sampled a total of 63 females (eight from each line, except seven from HR line 3). Each had experienced six days of wheel access as part of the routine testing protocol at an average starting age of 55 days (range = 48–65). Subsequently, each female had been a breeder, and all had given birth, with 60 of 63 successfully rearing their litter until weaning at 21 days of age. Echocardiography (see next section) was done 10–14 days after weaning (mean age = 137 days, range = 129–141), which was an average of 76 days (range = 68–84) after the end of the six-day period of wheel access. Given the amount of time that elapsed between wheel testing and echocardiography, any differences between the HR and C mice, or between mini- and normal-muscled individuals, should reflect inherent genetic differences, rather than possible effects of differential wheel running.

The following morning, mice were weighed, and body composition was measured by non-invasive quantitative magnetic resonance (EchoMRI-100; Echo Medical Systems LLC, Houston, Texas, USA), which independently calculated fat and lean mass. Mice were then euthanized via CO₂, and the heart ventricles were dissected free, blotted to remove blood, and weighed. As line HR line 6 is polymorphic for the mini-muscle phenotype (see above), we also collected and weighed the triceps surae muscle group for this line to identify mini-muscle individuals.¹⁸

Echocardiography

Investigators from Michigan State University traveled to the University of California, Riverside with a portable echocardiograph (Vivid IQ equipped with Hockey Stick probe and rodent analysis software, GE Healthcare, Chicago, IL, USA). As in previous studies,³¹ mice were anaesthetized using 2% isoflurane and placed in a supine position on a heated handling table. Limbs were secured to the table, and the hair over the trunk area was removed with clippers followed by application of hair

removal gel (Nair: Church & Dwight Co. Ewing, NJ, USA). Isoflurane concentration was then reduced to 1% and maintained throughout the rest of the measurement period. Body temperature was maintained at 37 °C via a supplemental heating lamp. Two dimensional echocardiographic images (M-mode short axis views at mid-papillary level and B-mode

parasternal long axis images) were collected for measures of cardiac structure and function in both short (left ventricular volume) and long axis (end diastolic volume) when applicable. The anesthesia process took less than 10 min, including hair removal and induction. Image analysis was conducted by a single researcher blinded to the mouse treatment

Table 1
Body mass, composition, and cardiovascular echocardiography results.

Variable Name, Units	N	C LSM	C SE	HR LSM	HR SE	Normal LSM	Normal SE	Mini-Muscle LSM	Mini-Muscle SE	Line Type DF	Line Type F	Line Type p	Mini-muscle DF	Mini-muscle F	Mini-Muscle p
Body Mass at Dissection (g)	62	34.66	1.78	31.07	1.57	34.71	1.10	31.02	1.93	6	2.64	0.155 1	53	3.89	<u>0.053 9</u>
Fat Mass (g)	59	0.67	0.03	0.62	0.03	0.53	0.02	0.76	0.04	6	1.89	0.218 1	49	22.81	<u>< 0.000 1</u>
Lean Mass (g)	59	1.45	0.02	1.41	0.02	1.45	0.01	1.41	0.02	6	3.16	0.125 6	50	3.81	<u>0.056 4</u>
Ejection Fraction (%)	36	1.87	0.01	1.82	0.01	1.81	0.01	1.89	0.02	6	9.09	<u>0.023 6</u>	24	15.72	<u>0.000 6</u>
Fractional Shortening (%)	35	36.64	1.78	32.71	1.28	29.98	0.96	39.37	2.29	6	4.10	0.09	25	13.56	<u>< 0.000 1</u>
Ventricle Mass at Dissection (mg)	60	152.80	5.06	158.30	4.30	145.30	2.90	165.70	6.20	6	0.83	0.396 1	50	9.94	<u>0.002 7</u>
Left Ventricle Mass (mg)	54	127.24	15.34	114.03	12.16	108.84	8.10	132.42	19.49	6	0.54	0.488 5	44	1.25	0.269 7
End Systolic Volume (μL)	37	-1.44	0.08	-1.36	0.06	-1.32	0.04	-1.49	0.10	6	0.71	0.431 2	26	2.36	0.136 6
End Diastolic Volume (μL)	37	-0.88	0.05	-0.88	0.04	-0.87	0.03	-0.89	0.06	6	0.00	0.999 7	26	0.13	0.722 9
Left Ventricle Volume Diastolic (μL)	54	142.93	22.85	97.97	18.10	105.17	12.01	135.73	29.06	6	2.83	0.143 5	43	0.94	0.337 4
Left Ventricle Volume Systolic (μL)	53	51.78	8.52	32.88	6.78	31.79	4.48	52.87	10.91	6	3.57	0.107 9	42	3.14	0.083 5
Anterior Wall Thickness (Diastole, mm)	53	0.86	0.05	0.99	0.04	0.88	0.03	0.97	0.06	6	5.09	<u>0.065 0</u>	43	1.80	0.186 4
Anterior Wall Thickness (Systole, mm)	54	1.32	0.08	1.45	0.06	1.37	0.04	1.41	0.11	6	1.71	0.239 1	44	0.12	0.728 3
Posterior Wall Thickness (Systole, mm)	53	1.21	0.08	1.25	0.06	1.18	0.04	1.27	0.09	6	0.18	0.688 8	43	0.68	0.414
Posterior Wall Thickness (Diastole, mm)	53	0.87	0.06	0.90	0.05	0.86	0.03	0.90	0.08	6	0.14	0.716 6	43	0.30	0.586 1
Relative Wall Thickness (mm)	37	-0.36	0.04	-0.34	0.03	-0.34	0.02	-0.37	0.06	6	0.15	0.710 2	28	0.28	0.599 6
Stroke Volume (μL)	53	91.69	15.07	63.88	12.21	68.88	8.20	86.69	18.98	6	2.48	0.166 3	42	0.77	0.386 5
Cardiac Output (mL/min)	53	54.52	9.12	38.65	7.40	41.55	4.98	51.63	11.46	6	2.21	0.188 0	42	0.67	0.416 1
Heart Rate (beats/min)	61	620.01	19.92	604.36	14.88	600.04	10.05	624.32	27.53	6	0.61	0.465 7	52	0.67	0.418 2

All data presented as Least Squares Mean and Standard Errors. Bold and underline p values denote significance (p < 0.05). N = Sample size, DF = degrees of freedom, LSM = least squares means, SE = standard error, C= Control mice, HR = high runner mice, g = grams, mg = milligrams, mL = milliliters, μL = microliters, mm = millimeters.

groups using dedicated software (EchoPAC; GE Healthcare, Horten, Norway). Standard measures of left ventricular structure and function were determined from the average of three cardiac cycles through EchoPAC software calculations.³²

Statistics

Images were evaluated for quality resulting in an N of 62 for M-mode images and N of 37 for B mode. Data were tested for normality using the Shapiro-Wilk test, and most residuals were found to be normally distributed, but the following variables were log-transformed: relative wall thickness, end systolic volume, left ventricle mass, posterior wall thickness (during diastole), left ventricle volume (during diastole), left ventricle internal diameter (during systole), left ventricle volume (during systole), ejection fraction, stroke volume, and cardiac output.

As in numerous previous studies of these lines of mice,^{10,13,15,17} data were analyzed with mixed models in SAS Procedure Mixed (SAS 9.4, SAS Institute, Inc., Cary, NC), using REML estimation and Type III tests of fixed effects. Line type (HR vs. C) and mini-muscle status were the main effects, and replicate line was nested within line type as a random effect. Degrees of freedom for testing the line type effect were always 1 and 6, whereas those for testing the mini-muscle effect varied with sample size. As another way to approach the data, we also used SAS Procedure Mixed with REML estimation and Type III tests of fixed effects with *a priori* contrasts to compare three groups: mice from the non-selected Control lines (1,2,4,5), those from the High Runner lines that had the mini-muscle (3, some of line 6), and those from High Runner lines that had normal muscles (some of line 6,7,8). Unlike the primary analyses presented in the text, these analyses did not use line as nested random effect.

Body mass and/or heart rate were used as covariates. Specifically, analyses of structural variables included body mass, whereas both body mass and heart rate were included for functional variables. Ejection fraction and fractional shortening percentage used only heart rate.

In all analyses, residuals were checked for skew and dependent variables were transformed as needed to improve normality (see Table 1). Outliers were removed based on criteria established *a priori*: when standardized values exceeded approximately 3 and/or were ≥ 1 unit from the next value. Significance was set at $p \leq 0.05$, and trends were considered at $0.05 < p \leq 0.10$.

Results

Table 1 presents significance levels as well as Least Squares Means and Standard Errors for both line type and mini-muscle status for all traits. HR and C mice did not differ statistically for body mass (Fig. 1A), fat mass (Fig. 1B), or lean mass (Fig. 1C). When comparing the data separated into three groups [mice from the non-selected Control lines (1,2,4,5), those from the High Runner lines that had the mini-muscle (3, some of line 6), and those from High Runner lines that had normal

muscles (some of line 6,7,8)] (Supplemental Table 1A), C mice had significantly greater body mass ($p < 0.0001$, Supplemental Fig. 1A) and lean mass ($p < 0.0001$, Supplemental Fig. 1C) compared to normal-muscle HR mice. No statistical differences in fat mass were observed between normal-muscle HR and C mice ($p = 0.1745$, Supplemental Fig. 1B).

HR mice had significantly lower ejection fractions ($p = 0.0236$, Fig. 2A) and tended to have a lower fractional shortening percentage ($p = 0.0900$, Fig. 2B) compared to the C lines. Furthermore, HR mice tended to have thicker anterior walls than C mice during diastole ($p = 0.0650$, Fig. 3A). When separating mice into the three groups, the ejection fraction difference ($p = 0.0052$, Supplemental Fig. 2A) and fractional shortening percentage trend ($p = 0.0515$, Supplemental Fig. 2B) remained. However, normal-muscle HR mice had significantly thicker anterior walls during diastole compared to C mice ($p = 0.0286$, Supplemental Fig. 3A).

As compared with normal-muscle mice, the mini-muscle phenotype was associated with reduced total body ($p = 0.0539$, Fig. 1A) and lean mass ($p = 0.0564$, Fig. 1C), but significantly higher fat mass ($p < 0.0001$ with lean mass as a covariate, Fig. 1B). Furthermore, mini-muscle mice had significantly greater ejection fraction ($p = 0.0006$, Fig. 2A) and fractional shortening percentage ($p < 0.0001$, Fig. 2B) compared to non-mini muscle mice.

When separating the normal-muscled and mini-muscle HR mice, the latter had significantly lower body mass ($p = 0.002$) and lean mass ($p < 0.0001$) compared to C mice. Mini-muscle mice also had significantly greater fat mass (with lean as a covariate) compared to both C mice ($p = 0.0017$) and normal-muscle HR mice ($p < 0.0001$, Supplemental Fig. 1B). Mini-muscle mice had greater ejection fraction ($p = 0.0004$) than normal-muscle HR mice, and greater fractional shortening percentage than both C ($p = 0.0316$) and normal-muscle HR mice ($p = 0.0009$). Mini-muscle mice had significantly thicker anterior walls ($p = 0.0026$, Supplemental Fig. 3A) compared to C mice.

Finally, mini-muscle mice had significantly greater ventricular mass weighed at dissection ($p = 0.0027$ with body mass as a covariate, Fig. 4A) and tended to have greater left ventricular volumes during systole as estimated by echocardiography ($p = 0.0835$). We observed no statistical differences in the other cardiac traits between HR and C mice or between mini- and normal-muscle mice. Differences in ventricular mass at dissection remained after separating the normal-muscle HR mice, with mini-muscle mice having greater ventricular mass than both C ($p < 0.0001$) and normal-muscle HR mice ($p < 0.0001$, Supplemental Fig. 4A) at dissection.

Discussion

Work with rodent models demonstrates that physical activity engagement is regulated by both environmental^{33–36} and biological^{4,7,37–44} factors, as well as epigenetic mechanisms.^{45,46} Over the past 30 years, the Garland laboratory has bred mice for high physical




Fig. 1. 1A) Body mass, 1B) fat (lean mass as covariate), and 1C) lean mass of control (C), high runner (HR), normal, and mini muscle mice. LS Means and associated standard errors from SAS Procedure Mixed (see text). g = Grams.




Fig. 2. 2A) Ejection fraction and 2B) fractional shortening percentage of control (C), high runner (HR), normal, and mini muscle mice (heart rate as a covariate). LS Means and associated standard errors from SAS Procedure Mixed (see text).




Fig. 3. Anterior wall thickness during 3A) Diastole and 3B) Systole of control (C), high runner (HR), normal, and mini muscle mice (body mass as a covariate). LS Means and associated standard errors from SAS Procedure Mixed (see text). mm = millimeters.




Fig. 4. Left ventricular mass measured via 4A) Dissection and 4B) Echocardiography of control (C), high runner (HR), normal, and mini muscle mice (body mass as a covariate). LS Means and associated standard errors from SAS Procedure Mixed (see text). mg = milligrams.

activity (as measured by voluntary wheel running).^{7,47,48} Over the course of almost 100 generations, various neural, anatomical, and functional adaptations that promote or support high levels of endurance running have evolved in the HR lines,^{12,13,17,18,20,24,25} and some of the underlying genetic and genomic changes have been identified.^{49–51}

Regular engagement in physical activity is viewed as cardioprotective and is known to elicit favorable changes in cardiac structure and function, including left ventricular eccentric remodeling. Whether individuals, genetic strains or even species that regularly engage in physical activity have innate (i.e., genetically “programmed”) cardiac features that might resemble those of eccentric remodeling is unknown (see also Kay et al., 2018²⁹). We tested this hypothesis by use of the unique, selectively bred High Runner (HR) lines of mice. We observed no statistically significant differences in any cardiovascular structures between HR mice and those from the non-selected control lines; rather, HR mice had reduced ejection fraction and tended to have thicker anterior

walls, contrary to our initial hypotheses. These results are counterintuitive and may indicate that even though HR mice are bred for high activity during a 6-day period of wheel access, they need to engage in endurance activity for some length of time to exhibit characteristics of cardiac remodeling. Although eccentric remodeling has not previously been examined in the HR model, a previous study²⁵ reported that chronic wheel access (13–14 weeks) led to a greater degree of ventricular hypertrophy in HR than in C mice, which could be explained statistically by the greater amount of running by the former (i.e., “more pain, more gain”). Similarly, HR mice have enhanced trainability of cardiac function as compared with C mice over six days of wheel access, as indicated by their longer PR duration (measured via electrocardiogram) afterwards.²³ Indeed, an initial investigation into cardiac gene expression in the HR and control lines indicated that chronic exercise (20–33 months) offset many age-related gene expression changes observed in the ventricles of sedentary animals.⁵²

A primary limitation to aerobic exercise capacity is an adequate cardiovascular system to provide necessary oxygen to the skeletal muscle mitochondria for sustained contraction (e.g., see^{53–56}). Among various changes, consistent engagement in endurance activity elicits eccentric remodeling of the heart that primarily consists of increased ventricular volume, modest increases in wall thickness, and a reduction in relative wall thickness.²⁸ Previous research has observed minimal differences between the HR lines bred for high activity and the non-selected control lines, despite HR lines previously demonstrating greater $\dot{V}O_{2\max}$, greater lipid utilization during exercise, increased myoglobin concentration in their ventricles, and faster speeds throughout wheel running sessions compared to non-HR mice.^{25,26,57,58} Although the reduction in ejection fraction of HR mice, in the present investigation, was surprising, cardiovascular measurements in elite cyclists showed similar reductions when compared to sedentary volunteers,⁵⁹ and professional basketball players had lower ejection fraction than the general population while maintaining normal systolic function.⁶⁰ Therefore, despite the traditional association between lower ejection fraction and systolic dysfunction, we propose the HR mice rather have evolved cardiac traits (lower left ventricular ejection fraction, thicker anterior walls) similar to those observed in the “athlete’s heart”,^{61–64} see also Kay et al., 2019.²³

Although we did not observe many universal differences between the HR and control-line mice, the subset of HR individuals with the mini-muscle phenotype had cardiac differences (when compared to non-mini muscle mice) indicative of health-positive adaptations, including significantly greater ejection fraction, fractional shortening percentage, and left ventricular mass (the last also shown in previous studies, e.g.,^{13,15,17,18,65}). The elevated ejection fraction and fractional shortening percentage may be explained by a longer duration QRS complex (measured via live EKG analysis:²³), which is positively correlated with left ventricular size in humans^{23,66} but does not predict athletic performance.^{23,67,68} Previous work shows that mini-muscle mice have greater $\dot{V}O_{2\max}$ (greater than other HR mice in some studies:^{15,26,58}) and peripheral adaptations in greater capillary density and capillary-to-fiber ratio of their gastrocnemius.^{15,27,58} Mini-muscle mice also have greatly reduced hindlimb muscle masses in conjunction with alterations in skeletal muscle enzyme concentrations and fiber types that appear beneficial for prolonged aerobic activity.^{23,25,27,36} The greater capillary density, in addition to smaller skeletal muscle (and more resistance to flow through them), may require mini-muscle mice to generate a higher blood pressure to facilitate blood flow through these tissues and ensure perfusion of the working tissue, although higher blood pressures were not observed in tail-cuff blood pressure measurements.⁶⁹ The greater activity of skeletal muscle enzymes in the mini-muscle mice may also necessitate greater blood delivery, and result in further cardiac remodeling to facilitate the increased need. Therefore, in addition to the peripheral alterations previously observed, the mini-muscle mice also have altered cardiovascular characteristics to facilitate blood flow to smaller tissues that are ultimately conducive to prolonged, aerobically supported physical activity.

The inherent cardiac functional and structural differences between the mini-muscle and non-mini muscle HR mice demonstrate the idea of multiple solutions for high voluntary activity from a given starting point⁵⁷; however, it is important to note we investigated adult mice that were not exposed to more than 6 days of running, and even that short exposure occurred an average of 76 days prior (see Methods). Therefore, although the HR mice were genetically bred to be highly active, they were not regularly participating in physical activity at the time of investigation, so the cardiac phenotype observed in the HR mice should be representative of their untrained or baseline state. Specifically, the HR mice may have a reduced ejection fraction at baseline, but an increased capacity to adapt, which may result in an improved cardiac phenotype later in life if trained. Evidence of increased plasticity to training has been demonstrated in the HR mice previously,^{23,25} and thus structural changes in response to training should be investigated in future studies.

Although the present results are interesting, we note several limitations. First, data collection was conducted by investigators traveling with a portable echocardiography. The machine was not equipped with the stationary mounting system common on most mouse echocardiographs, which reduced quality of the images and consequently sample size for some measures. Second, measurements were conducted after anesthetization, so more differences could emerge during a pharmacological stress test on the mice to examine maximal cardiac function. Thirdly, this study included only females, and as such, inherent genetic differences could be present in males that are absent in females. Finally, cardiac differences might be observed at the single myocyte level, as both cardiac calcium transit kinetics and contractile functioning differences were observed between rats bred for high- and low endurance capacity during forced treadmill exercise⁷⁰ and which also differ in voluntary wheel running.⁷¹

Conclusions

In summary, even though mice were bred for high activity, it may be necessary for mice to engage in moderate to vigorous physical activity to observe positive eccentric remodeling of the heart. Additional studies would be needed to test this hypothesis. Although mice from the HR lines bred for high activity showed limited structural or functional differences from the non-selected control lines, the subset of HR mini-muscle mice have high ejection fraction and fractional shortening percentage than non-mini muscle mice, which could aid $\dot{V}O_{2\max}$ and endurance activity engagement.³¹

Submission statement

All authors have read and agree with the contents of the manuscript. While this manuscript is being reviewed for this journal, the manuscript will not be submitted elsewhere for review and publication.

Authors’ contribution

TG and DF devised the study. TG and DF led data collection, assisted by NS and EL. EL led analysis of data, assisted by AM, KD, and DF. TG ran statistical analysis with assistance from DF and NS. EL, TG, and DF drafted the manuscript. All authors read and approved the manuscript.

Ethical approval statement

All animals were housed in a room temperature controlled at 22 °C under a 12-h dark/light cycle. All food and water was provided ad libitum throughout the experimental procedures, and all experimental procedures were approved by the University of California, Riverside Institutional Animal Care and Use Committee.

Funding

The study was funded by the Department of Kinesiology at Michigan State University and NSF grant IOS-2038528 to T.G. and Natalie Holt.

Data availability

Data generated or analyzed during this study are available from the corresponding author upon reasonable request.

Conflict of interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

The authors wish to thank all members of the Neonatal Nutrition and Exercise Research Lab and ECHO Lab at Michigan State University and the Garland Laboratory at the University of California, Riverside.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.smhs.2023.07.003>.

References

- Kraus WE, Powell KE, Haskell WL, et al. Physical activity, all-cause and cardiovascular mortality, and cardiovascular disease. *Med Sci Sports Exerc.* 2019;51(6):1270–1281. <https://doi.org/10.1249/MSS.0000000000001939>.
- Zhao M, Veeranki SP, Li S, Steffen LM, Xi B. Beneficial associations of low and large doses of leisure time physical activity with all-cause, cardiovascular disease and cancer mortality: a national cohort study of 88,140 US adults. *Br J Sports Med.* 2019;53(22):1405–1411. <https://doi.org/10.1136/bjsports-2018-099254>.
- Williams PT. Dose-response relationship of physical activity to premature and total all-cause and cardiovascular disease mortality in walkers. *PLoS One.* 2013;8(11):e78777. <https://doi.org/10.1371/journal.pone.0078777>.
- Lightfoot JT, De Geus EJC, Booth FW, et al. Biological/genetic regulation of physical activity level: consensus from GenBioPAC. *Med Sci Sports Exerc.* 2018;50(4):863. <https://doi.org/10.1249/mss.0000000000001499>.
- Humpel N, Owen N, Leslie E. Environmental factors associated with adults' participation in physical activity: a review. *Am J Prev Med.* 2002;22(3):188–199. [https://doi.org/10.1016/s0749-3797\(01\)00426-3](https://doi.org/10.1016/s0749-3797(01)00426-3).
- Rowland T. *Biologic Regulation of Physical Activity.* Human Kinetics; 2016.
- Garland Jr T, Schutz H, Chappell MA, et al. The biological control of voluntary exercise, spontaneous physical activity and daily energy expenditure in relation to obesity: human and rodent perspectives. *J Exp Biol.* 2011;214(Pt 2):206–229. <https://doi.org/10.1242/jeb.048397>.
- Swallow JG, Carter PA, Garland Jr T. Artificial selection for increased wheel-running behavior in house mice. *Behav Genet.* 1998;28(3):227–237. <https://doi.org/10.1023/a:1021479331779>.
- Careau V, Wolak ME, Carter PA, Garland Jr T. Limits to behavioral evolution: the quantitative genetics of a complex trait under directional selection. *Evolution.* 2013;67(11):3102–3119. <https://doi.org/10.1111/evo.12200>.
- Copes LE, Schutz H, Dlugosz EM, Acosta W, Chappell MA, Garland Jr T. Effects of voluntary exercise on spontaneous physical activity and food consumption in mice: results from an artificial selection experiment. *Physiol Behav.* 2015;149:86–94. <https://doi.org/10.1016/j.physbeh.2015.05.025>.
- Swallow JG, Koteja P, Carter PA, Garland Jr T. Food consumption and body composition in mice selected for high wheel-running activity. *J Comp Physiol B.* 2001;171(8):651–659. <https://doi.org/10.1007/s003600100216>.
- Hiramatsu L, Garland Jr T. Mice selectively bred for high voluntary wheel-running behavior conserve more fat despite increased exercise. *Physiol Behav.* 2018;194:1–8. <https://doi.org/10.1016/j.physbeh.2018.04.010>.
- Meek TH, Lonquich BP, Hannon RM, Garland Jr T. Endurance capacity of mice selectively bred for high voluntary wheel running. *J Exp Biol.* 2009;212(18):2908–2917. <https://doi.org/10.1242/jeb.028886>.
- Swallow JG, Garland Jr T, Carter PA, Zhan WZ, Sieck GC. Effects of voluntary activity and genetic selection on aerobic capacity in house mice (*Mus domesticus*). *J Appl Physiol (1985).* 1998;84(1):69–76. <https://doi.org/10.1152/jappl.1998.84.1.69>.
- Cadney MD, Hiramatsu L, Thompson Z, et al. Effects of early-life exposure to Western diet and voluntary exercise on adult activity levels, exercise physiology, and associated traits in selectively bred High Runner mice. *Physiol Behav.* 2021;234:113389. <https://doi.org/10.1016/j.physbeh.2021.113389>.
- Hiramatsu L, Kay JC, Thompson Z, et al. Maternal exposure to Western diet affects adult body composition and voluntary wheel running in a genotype-specific manner in mice. *Physiol Behav.* 2017;179:235–245. <https://doi.org/10.1016/j.physbeh.2017.06.008>.
- Kolb EM, Kelly SA, Middleton KM, Sermsakdi LS, Chappell MA, Garland Jr T. Erythropoietin elevates but not voluntary wheel running in mice. *J Exp Biol.* 2010;213(3):510–519. <https://doi.org/10.1242/jeb.029074>.
- Garland Jr T, Morgan MT, Swallow JG, et al. Evolution of a small-muscle polymorphism in lines of house mice selected for high activity levels. *Evolution.* 2002;56(6):1267–1275. <https://doi.org/10.1111/j.0014-3820.2002.tb01437.x>.
- Kelly SA, Bell TA, Selitsky SR, et al. A novel intronic single nucleotide polymorphism in the myosin heavy polypeptide 4 gene is responsible for the mini-muscle phenotype characterized by major reduction in hind-limb muscle mass in mice. *Genetics.* 2013;195(4):1385–1395. <https://doi.org/10.1534/genetics.113.154476>.
- Bilodeau GM, Guderley H, Joannis DR, Garland Jr T. Reduction of type IIB myosin and IIB fibers in tibialis anterior muscle of mini-muscle mice from high-activity lines. *J Exp Zool A Ecol Genet Physiol.* 2009;311(3):189–198. <https://doi.org/10.1002/jez.518>.
- McGillivray DG, Garland Jr T, Dlugosz EM, Chappell MA, Syme DA. Changes in efficiency and myosin expression in the small-muscle phenotype of mice selectively bred for high voluntary running activity. *J Exp Biol.* 2009;212(7):977–985. <https://doi.org/10.1242/jeb.026625>.
- Guderley H, Houle-Leroy P, Diffie GM, Camp DM, Garland Jr T. Morphometry, ultrastructure, myosin isoforms, and metabolic capacities of the “mini muscles” favoured by selection for high activity in house mice. *Comp Biochem Physiol B Biochem Mol Biol.* 2006;144(3):271–282. <https://doi.org/10.1016/j.cbpb.2006.02.009>.
- Kay JC, Claghorn GC, Thompson Z, Hampton TG, Garland Jr T. Electrocardiograms of mice selectively bred for high levels of voluntary exercise: effects of short-term exercise training and the mini-muscle phenotype. *Physiol Behav.* 2019;199:322–332. <https://doi.org/10.1016/j.physbeh.2018.11.041>.
- Houle-Leroy P, Guderley H, Swallow JG, Garland Jr T. Artificial selection for high activity favors mighty mini-muscles in house mice. *Am J Physiol Regul Integr Comp Physiol.* 2003;284(2):R433–R443. <https://doi.org/10.1152/ajpregu.00179.2002>.
- Kelly SA, Gomes FR, Kolb EM, Malisch JL, Garland Jr T. Effects of activity, genetic selection and their interaction on muscle metabolic capacities and organ masses in mice. *J Exp Biol.* 2017;220(6):1038–1047. <https://doi.org/10.1242/jeb.148759>.
- Rezende EL, Garland Jr T, Chappell MA, Malisch JL, Gomes FR. Maximum aerobic performance in lines of *Mus* selected for high wheel-running activity: effects of selection, oxygen availability and the mini-muscle phenotype. *J Exp Biol.* 2006;209(1):115–127. <https://doi.org/10.1242/jeb.01883>.
- Wong LE, Garland Jr T, Rowan SL, Hepple RT. Anatomic capillarization is elevated in the medial gastrocnemius muscle of mighty mini mice. *J Appl Physiol.* 2009;106(5):1660–1667. <https://doi.org/10.1152/jappphysiol.91233.2008>.
- Shave R, Howatson G, Dickson D, Young L. Exercise-induced cardiac remodeling: lessons from humans, horses, and dogs. *Vet Sci.* 2017;4(1):9. <https://doi.org/10.3390/vetsci4010009>.
- Kay JC, Ramirez J, Contreras E, Garland Jr T. Reduced non-bicarbonate skeletal muscle buffering capacity in mice with the mini-muscle phenotype. *J Exp Biol.* 2018;221(10):jeb172478. <https://doi.org/10.1242/jeb.172478>.
- Castro AA, Garland T, Ahmed S, Holt NC. Trade-offs in muscle physiology in selectively bred high runner mice. *J Exp Biol.* 2022;225(23):jeb244083. <https://doi.org/10.1242/jeb.244083>.
- Ferguson DP, Monroe TO, Heredia CP, et al. Postnatal undernutrition alters adult female mouse cardiac structure and function leading to limited exercise capacity. *J Physiol.* 2019;597(7):1855–1872. <https://doi.org/10.1111/JP277637>.
- West CR, Crawford MA, Poormasjedi-Meibod MS, et al. Passive hind-limb cycling improves cardiac function and reduces cardiovascular disease risk in experimental spinal cord injury. *J Physiol.* 2014;592(8):1771–1783. <https://doi.org/10.1113/jphysiol.2013.268367>.
- Schmitt EE, Vellers HL, Porter WW, Lightfoot JT. Environmental endocrine disruptor affects voluntary physical activity in mice. *Med Sci Sports Exerc.* 2016;48(7):1251–1258. <https://doi.org/10.1249/MSS.0000000000000908>.
- Eclarinal JD, Zhu S, Baker MS, et al. Maternal exercise during pregnancy promotes physical activity in adult offspring. *FASEB J.* 2016;30(7):2541–2548. <https://doi.org/10.1096/fj.201500018r>.
- Leszczynski EC, Visker JR, Ferguson DP. The effect of growth restriction on voluntary physical activity engagement in mice. *Med Sci Sports Exerc.* 2019;51(11):2201–2209. <https://doi.org/10.1249/mss.0000000000002040>.
- Acosta W, Meek TH, Schutz H, Dlugosz EM, Vu KT, Garland Jr T. Effects of early-onset voluntary exercise on adult physical activity and associated phenotypes in mice. *Physiol Behav.* 2015;149:279–286. <https://doi.org/10.1016/j.physbeh.2015.06.020>.
- Ferguson DP, Dangott LJ, Vellers HL, Schmitt EE, Lightfoot JT. Differential protein expression in the nucleus accumbens of high and low active mice. *Behav Brain Res.* 2015;291:283–288. <https://doi.org/10.1016/j.bbr.2015.05.035>.
- Ferguson DP, Dangott LJ, Schmitt EE, Vellers HL, Lightfoot JT. Differential skeletal muscle proteome of high- and low-active mice. *J Appl Physiol (1985).* 2014;116(8):1057–1067. <https://doi.org/10.1152/jappphysiol.00911.2013>.
- Knab AM, Bowen RS, Hamilton AT, Lightfoot JT. Pharmacological manipulation of the dopaminergic system affects wheel-running activity in differentially active mice. *J Biol Regul Homeost Agents.* 2012;26(1):119–129.
- Bowen RS, Knab AM, Hamilton AT, McCall JR, Moore-Harrison TL, Lightfoot JT. Effects of supraphysiological doses of sex steroids on wheel running activity in mice. *J Steroids Horm Sci.* 2012;3(2):110. <https://doi.org/10.4172/2157-7536.1000110>.
- Lightfoot JT, Leamy L, Pomp D, et al. Strain screen and haplotype association mapping of wheel running in inbred mouse strains. *J Appl Physiol (1985).* 2010;109(3):623–634. <https://doi.org/10.1152/jappphysiol.00525.2010>.
- Knab AM, Lightfoot JT. Does the difference between physically active and couch potato lie in the dopamine system? *Int J Biol Sci.* 2010;6(2):133–150. <https://doi.org/10.7150/ijbs.6.133>.
- Knab AM, Bowen RS, Hamilton AT, Gulleddge AA, Lightfoot JT. Altered dopaminergic profiles: implications for the regulation of voluntary physical activity. *Behav Brain Res.* 2009;204(1):147–152. <https://doi.org/10.1016/j.bbr.2009.05.034>.
- Kelly SA, Nehrenberg DL, Hua K, Garland Jr T, Pomp D. Functional genomic architecture of predisposition to voluntary exercise in mice: expression QTL in the brain. Research Support, N.I.H. *Extramural Genetics.* 2012;191(2):643–654. <https://doi.org/10.1534/genetics.112.140509>.
- MacKay H, Scott CA, Duryea JD, et al. DNA methylation in AgRP neurons regulates voluntary exercise behavior in mice. *Nat Commun.* 2019;10(1):1–11. <https://doi.org/10.1038/s41467-019-13339-3>.
- Latchney SE, Cadney MD, Hopkins A, Garland T. DNA methylation analysis of imprinted genes in the cortex and hippocampus of cross-fostered mice selectively bred for increased voluntary wheel-running. *Behav Genet.* 2022;1–17. <https://doi.org/10.1007/s10519-022-10112-z>.
- Wallace IJ, Garland Jr T. Mobility as an emergent property of biological organization: insights from experimental evolution. *Evol Anthropol.* 2016;25(3):98–104. <https://doi.org/10.1002/evan.21481>.

48. Swallow JG, Hayes JP, Koteja P, Garland Jr T. Selection experiments and experimental evolution of performance and physiology. *Experiment Evolut: Concepts, Methods Appl Select Experiments*. 2009;301–351. <https://doi.org/10.1525/california/9780520247666.003.0012>.
49. Saul MC, Majdak P, Perez S, Reilly M, Garland Jr T, Rhodes JS. High motivation for exercise is associated with altered chromatin regulators of monoamine receptor gene expression in the striatum of selectively bred mice. *Gene Brain Behav*. 2017;16(3):328–341. <https://doi.org/10.1111/gbb.12347>.
50. Hillis DA, Yadgary L, Weinstock GM, et al. Genetic basis of aerobically supported voluntary exercise: results from a selection experiment with house mice. *Genetics*. 2020;216(3):781–804. <https://doi.org/10.1534/genetics.120.303668>.
51. Nguyen QAT, Hillis D, Katada S, et al. Coadaptation of the chemosensory system with voluntary exercise behavior in mice. *PLoS One*. 2020;15(11):e0241758. <https://doi.org/10.1371/journal.pone.0241758>.
52. Bronikowski AM, Carter PA, Morgan TJ, et al. Lifelong voluntary exercise in the mouse prevents age-related alterations in gene expression in the heart. *Physiol Genom*. 2003;12(2):129–138. <https://doi.org/10.1152/physiolgenomics.00082.2002>.
53. Hillman SS, Hedrick MS. A meta-analysis of in vivo vertebrate cardiac performance: implications for cardiovascular support in the evolution of endothermy. *J Exp Biol*. 2015;218(8):1143–1150. <https://doi.org/10.1242/jeb.118372>.
54. Scott GR, Dalziel AC. Physiological insight into the evolution of complex phenotypes: aerobic performance and the O₂ transport pathway of vertebrates. *J Exp Biol*. 2021;224(16):jeb210849. <https://doi.org/10.1242/jeb.210849>.
55. van der Zwaard S, de Ruiter CJ, Noordhof DA, et al. Maximal oxygen uptake is proportional to muscle fiber oxidative capacity, from chronic heart failure patients to professional cyclists. *J Appl Physiol*. 2016;121(3):636–645. <https://doi.org/10.1152/jappphysiol.00355.2016>.
56. Gifford JR, Garten RS, Nelson AD, et al. Symmorphosis and skeletal muscle: in vivo and in vitro measures reveal differing constraints in the exercise-trained and untrained human. *J Physiol*. 2016;594(6):1741–1751. <https://doi.org/10.1113/jp271229>.
57. Garland Jr T, Kelly SA, Malisch JL, et al. How to run far: multiple solutions and sex-specific responses to selective breeding for high voluntary activity levels. *Proc Biol Sci*. 2011;278(1705):574–581. <https://doi.org/10.1098/rspb.2010.1584>.
58. Templeman NM, Schutz H, Garland Jr T, McClelland GB. Do mice bred selectively for high locomotor activity have a greater reliance on lipids to power submaximal aerobic exercise? *Am J Physiol Regul Integr Comp Physiol*. 2012;303(1):R101–R111. <https://doi.org/10.1152/ajpregu.00511.2011>.
59. Abergel E, Chatellier G, Hagege AA, et al. Serial left ventricular adaptations in world-class professional cyclists: implications for disease screening and follow-up. *J Am Coll Cardiol*. 2004;44(1):144–149. <https://doi.org/10.1016/j.jacc.2004.02.057>.
60. Paterick TE, Gordon T, Spiegel D. Echocardiography: profiling of the athlete's heart. *J Am Soc Echocardiogr*. 2014;27(9):940–948. <https://doi.org/10.1016/j.echo.2014.06.008>.
61. De Innocentis C, Ricci F, Khanji MY, et al. Athlete's heart: diagnostic challenges and future perspectives. *Sports Med*. 2018;48(11):2463–2477. <https://doi.org/10.1007/s40279-018-0985-2>.
62. Weeks KL, McMullen JR. The athlete's heart vs. the failing heart: can signaling explain the two distinct outcomes? *Physiology*. 2011;26(2):97–105. <https://doi.org/10.1152/physiol.00043.2010>.
63. Prior DL, La Gerche A. The athlete's heart. *Heart*. 2012;98(12):947–955. <https://doi.org/10.1136/heartjnl-2011-301329>.
64. Maron BJ. Distinguishing hypertrophic cardiomyopathy from athlete's heart: a clinical problem of increasing magnitude and significance. *Heart*. 2005;91(11):1380. <https://doi.org/10.1136/heartjnl-2011-301329>.
65. Swallow JG, Rhodes JS, Garland Jr T. Phenotypic and evolutionary plasticity of organ masses in response to voluntary exercise in house mice. *Integr Comp Biol*. 2005;45(3):426–437. <https://doi.org/10.1093/icb/45.3.426>.
66. Carlsson MB, Trägårdh E, Engblom H, et al. Left ventricular mass by 12-lead electrocardiogram in healthy subjects: comparison to cardiac magnetic resonance imaging. *J Electrocardiol*. 2006;39(1):67–72. <https://doi.org/10.1016/j.jelectrocard.2005.07.005>.
67. Physick-Sheard PW, Hendren CM. Heart score: physiological basis and confounding variables. *Equine Exerc Physiol*. 1983;1:121–134.
68. Hanson CM, Kline KH, Foreman JH. Measurements of heart scores and heart weights in horses of two different morphic body types. *Comp Biochem Physiol A Comp Physiol*. 1994;108(2–3):175–178. [https://doi.org/10.1016/0300-9629\(94\)90083-3](https://doi.org/10.1016/0300-9629(94)90083-3).
69. Kolb EM, Kelly SA, Garland Jr T. Mice from lines selectively bred for high voluntary wheel running exhibit lower blood pressure during withdrawal from wheel access. *Physiol Behav*. 2013;112:49–55. <https://doi.org/10.1016/j.physbeh.2013.02.010>.
70. Palpant NJ, Szatkowski ML, Wang W, et al. Artificial selection for whole animal low intrinsic aerobic capacity co-segregates with hypoxia-induced cardiac pump failure. *PLoS One*. 2009;4(7):e6117. <https://doi.org/10.1371/journal.pone.0006117>.
71. Swallow JG, Wroblewska AK, Waters RP, Renner KJ, Britton SL, Koch LG. Phenotypic and evolutionary plasticity of body composition in rats selectively bred for high endurance capacity. *J Appl Physiol*. 2010;109(3):778–785. <https://doi.org/10.1152/jappphysiol.01026.2009>.