

# Lawrence Berkeley National Laboratory

## LBL Publications

### Title

Showers Initiated by 335 Mev Bremsstrahlung

### Permalink

<https://escholarship.org/uc/item/9721n62z>

### Authors

Blocker, Wade

Kenney, Robert W

### Publication Date

2024-03-06

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UNCLASSIFIED

Copy 2

UCRL-516  
Abstract

Showers Initiated by 335 Mev Bremsstrahlung. WADE BLOCKER AND ROBERT W. KENNEY. Radiation Laboratory, University of California, Berkeley, California.

The variation of cascade shower intensity with depth has been determined for showers produced in lead, copper, aluminum, and carbon by the 335 Mev bremsstrahlung from the Berkeley synchrotron.

The collimated beam from the synchrotron passed through a monitor ionization chamber and fell normally upon closely stacked plates of the element under study. The shower particles from various thicknesses of this element were then observed by a detector ionization chamber situated behind the plates. An essentially infinite amount of the element under study was placed behind the detector chamber to provide backscattered radiation. This contribution to the total intensity was very considerable in the case of lead.

Shower intensities in the initial half radiation length allow the thin layer pair production and Compton contributions to the shower to be derived. The ordinates and abscissae of the maxima of the shower curves are compared with theory. The exponential tails of the shower curves give the gamma ray absorption coefficients at the energy of minimum absorption cross-section. These values are compared with theory. A thin sheet of lead observed by an ionization chamber gave ionization values which are used to derive the total energy of the X-ray beam. This work was sponsored by the A.E.C.