

UC Davis

UC Davis Previously Published Works

Title

Complete genome sequence of Halorhabdus utahensis type strain (AX-2T)

Permalink

<https://escholarship.org/uc/item/97d8t4kh>

Journal

Environmental Microbiome, 1(3)

ISSN

1944-3277

Authors

Anderson, Iain

Tindall, Brian J

Pomrenke, Helga

et al.

Publication Date

2009-11-01

DOI

10.4056/sigs.31864

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Complete genome sequence of *Halorhabdus utahensis* type strain (AX-2^T)

Iain Anderson¹, Brian J. Tindall², Helga Pomrenke², Markus Göker², Alla Lapidus¹, Matt Nolan¹, Alex Copeland¹, Tijana Glavina Del Rio¹, Feng Chen¹, Hope Tice¹, Jan-Fang Cheng¹, Susan Lucas¹, Olga Chertkov^{1,3}, David Bruce^{1,3}, Thomas Brettin^{1,3}, John C. Detter^{1,3}, Cliff Han^{1,3}, Lynne Goodwin^{1,3}, Miriam Land^{1,4}, Loren Hauser^{1,4}, Yun-Juan Chang^{1,4}, Cynthia D. Jeffries^{1,4}, Sam Pitluck¹, Amrita Pati¹, Konstantinos Mavromatis¹, Natalia Ivanova¹, Galina Ovchinnikova¹, Amy Chen⁵, Krishna Palaniappan⁵, Patrick Chain^{1,6}, Manfred Rohde⁷, Jim Bristow¹, Jonathan A. Eisen^{1,8}, Victor Markowitz⁵, Philip Hugenholtz¹, Nikos C. Kyrpides¹, and Hans-Peter Klenk^{2*}

¹ DOE Joint Genome Institute, Walnut Creek, California, USA

² DSMZ - German Collection of Microorganisms and Cell Cultures GmbH, Braunschweig, Germany

³ Los Alamos National Laboratory, Bioscience Division, Los Alamos, New Mexico, USA

⁴ Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA

⁵ Biological Data Management and Technology Center, Lawrence Berkeley National Laboratory, Berkeley, California, USA

⁶ Lawrence Livermore National Laboratory, Livermore, California, USA

⁷ HZI - Helmholtz Centre for Infection Research, Braunschweig, Germany

⁸ University of California Davis Genome Center, Davis, California, USA

*Corresponding author: Hans-Peter Klenk

Keywords: halophile, free-living, non-pathogenic, aerobic, euryarchaeon, *Halobacteriaceae*

Halorhabdus utahensis Wainø *et al.* 2000 is the type species of the genus, which is of phylogenetic interest because of its location on one of the deepest branches within the very extensive euryarchaeal family *Halobacteriaceae*. *H. utahensis* is a free-living, motile, rod shaped to pleomorphic, Gram-negative archaeon, which was originally isolated from a sediment sample collected from the southern arm of Great Salt Lake, Utah, USA. When grown on appropriate media, *H. utahensis* can form polyhydroxybutyrate (PHB). Here we describe the features of this organism, together with the complete genome sequence, and annotation. This is the first complete genome sequence of a member of halobacterial genus *Halorhabdus*, and the 3,116,795 bp long single replicon genome with its 3027 protein-coding and 48 RNA genes is part of the *Genomic Encyclopedia of Bacteria and Archaea* project.

Introduction

Strain AX-2^T (= DSM 12940 = JCM 11049) is the type strain of the species *Halorhabdus utahensis*, and represents one of only two species currently assigned to the genus *Halorhabdus* [1]. Strain AX-2^T was first described by Wainø *et al.* in 2000 [1] as Gram-negative, motile and extremely pleomorphic organism. The organism is of interest because of its position in the tree of life, where the genera *Halorhabdus* and *Halomicrobium* constitute one of the deepest branches within the large euryarchaeal family *Halobacteriaceae*. Here we present a summary classification and a set of

features for *H. utahensis* strain AX-2^T together with the description of the complete genomic sequencing and annotation.

Classification and features

Only one other 16S rRNA encoding sequence has been deposited in the INSDC databases with a similarity of greater than 97% to that of strain AX-2^T. That sequence belongs to the other species classified in the genus *Halorhabdus*, *H. tiamatea*, which was isolated from a sample of the brine-sediment interface of the Shaban Deep in the




Figure 2. Scanning electron micrograph of *H. utahensis* strain AX-2^T

Table 1. Classification and general features of *H. utahensis* strain AX-2^T according to the MGS recommendations [9]

MIGS ID	Property	Term	Evidence code
	Classification	Domain <i>Archaea</i>	TAS [10]
		Phylum <i>Euryarchaeota</i>	TAS [11,12]
		Class <i>Halobacteria</i>	TAS [13]
		Order <i>Halobacteriales</i>	TAS [14]
		Family <i>Halobacteriaceae</i>	TAS [15]
		Genus <i>Halorhabdus</i>	TAS [1]
		Species <i>Halorhabdus utahensis</i>	TAS [1]
		Type strain AX-2	TAS [1]
	Gram stain	negative	TAS [1]
	Cell shape	rod to pleomorphic	TAS [1]
	Motility	motile by a single flagellum	TAS [1]
	Sporulation	nonsporulating	TAS [1]
	Temperature range	mesophile, 15-55°C	TAS [1]
	Optimum temperature	50°C	TAS [1]
	Salinity	halophile, at least 9% (w/v) NaCl, maximum 30%, with an optimum at 27%	TAS [1]
MIGS-22	Oxygen requirement	primarily aerobic; facultatively anaerobic, growth via glucose fermentation	TAS [1]
	Carbon source	glucose, xylose and fructose	TAS [1]
	Energy source	carbohydrates	NAS
MIGS-6	Habitat	aquatic	TAS [1]
MIGS-15	Biotic relationship	free living	NAS
MIGS-14	Pathogenicity	none	NAS

Table 1. Classification and general features of *H. utahensis* strain AX-2^T according to the MIGS recommendations [9] (cont.)

MIGS ID	Property	Term	Evidence code
	Biosafety level	1	TAS [16]
	Isolation	sediment of Great Salt Lake, Utah	TAS [1]
MIGS-4	Geographic location	sediment of Great Salt Lake, Utah	TAS [1]
MIGS-5	Sample collection time	before 2000	TAS [1]
MIGS-4.1	Latitude, Longitude	41.177, -112.502	NAS
MIGS-4.2			
MIGS-4.3	Depth	sea level	TAS [1]
MIGS-4.4	Altitude	not reported	

Evidence codes - IDA: Inferred from Direct Assay (first time in publication); TAS: Traceable Author Statement (i.e., a direct report exists in the literature); NAS: Non-traceable Author Statement (i.e., not directly observed for the living, isolated sample, but based on a generally accepted property for the species, or anecdotal evidence). These evidence codes are from the Gene Ontology project [17]. If the evidence code is IDA, then the property was directly observed for a living isolate by one of the authors or an expert mentioned in the acknowledgements.

Genome sequencing and annotation

Genome project history

This organism was selected for sequencing on the basis of each phylogenetic position, and is part of the *Genomic Encyclopedia of Bacteria and Archaea* project. The genome project is deposited in the Genome OnLine Database [7] and the complete

genome sequence in GenBank. Sequencing, finishing and annotation were performed by the DOE Joint Genome Institute (JGI). A summary of the project information is shown in Table 2.

Table 2. Genome sequencing project information

MIGS ID	Property	Term
MIGS-31	Finishing quality	Finished
MIGS-28	Libraries used	Three genomic libraries: two Sanger libraries (8 kb pMCL200 and fosmid pcc1Fos) and one 454 pyrosequence standard library
MIGS-29	Sequencing platforms	ABI3730, 454 GS FLX
MIGS-31.2	Sequencing coverage	8.3x Sanger; 21.2x pyrosequence
MIGS-30	Assemblers	Newbler version 1.1.02.15, phrap
MIGS-32	Gene calling method	Prodigal, GenePRIMP
	INSDC ID	CP001687
	Genbank Date of Release	August 27, 2009
	GOLD ID	Gc01053
	NCBI project ID	29305
	Database: IMG-GEBA	2501416929
MIGS-13	Source material identifier	DSM 12940
	Project relevance	Tree of Life, GEBA

Growth conditions and DNA isolation

H. utahensis strain AX-2^T, DSM 12940, was grown in DSMZ medium 927 (*H. utahensis* medium) [18] at 40°C. DNA was isolated from 1-1.5 g of cell paste using a Qiagen Genomic 500 DNA Kit

(Qiagen, Hilden, Germany) as described in Wu *et al.* [19].

Genome sequencing and assembly

The genome was sequenced using a combination of Sanger and 454 sequencing platforms. All general aspects of library construction and

sequencing performed at the JGI can be found on the JGI website (<http://www.jgi.doe.gov/>). 454 Pyrosequencing reads were assembled using the Newbler assembler, version 1.1.02.15 (Roche). Large Newbler contigs were broken into 3,474 overlapping fragments of 1,000 bp and entered into assembly as pseudo-reads. The sequences were assigned quality scores based on Newbler consensus q-scores with modifications to account for overlap redundancy and to adjust inflated q-scores. A hybrid 454/Sanger assembly was made using the parallel phrap assembler (High Performance Software, LLC). Possible mis-assemblies were corrected with Dupfinisher or transposon bombing of bridging clones [20]. Gaps between contigs were closed by editing in Consed, custom primer walk or PCR amplification. A total of 212 Sanger finishing reads were produced to close gaps, to resolve repetitive regions, and to raise the quality of the finished sequence. The final assembly consists of 26,545 Sanger and 382,722 pyrosequence (454) reads. Together all sequence types provided 29.5× coverage of the genome. The error rate of the completed genome sequence is less than 1 in 100,000.

Genome annotation

Genes were identified using Prodigal [21] as part of the Oak Ridge National Laboratory genome annotation pipeline, followed by a round of manual curation using the JGI GenePRIMP pipeline (<http://geneprimp.jgi-psf.org/>) [22]. The predicted CDSs were translated and used to search the National Center for Biotechnology Information (NCBI) nonredundant database, UniProt, TIGRFam, Pfam, PRIAM, KEGG, COG, and InterPro databases. Additional gene prediction analysis and functional annotation was performed within the Integrated Microbial Genomes Expert Review platform (<http://img.jgi.doe.gov/er>) [23].

Genome properties

The genome is 3,116,795 bp long and comprises one main circular chromosome with a 62.9% GC content (Table 3 and Figure 3). Of the 3,075 genes predicted, 3,027 were protein coding genes, and 48 RNAs; 29 pseudogenes were also identified. The majority of the protein-coding genes (60.5%) were assigned with a putative function, while those remaining were annotated as hypothetical proteins. The distribution of genes into COGs functional categories is presented in Table 4.

Table 3.

Attribute	Value	% of Total
Genome size (bp)	3,116,795	100.00%
DNA Coding region (bp)	2,768,833	88.83%
DNA G+C content (bp)	1,960,463	62.90%
Number of replicons	1	
Extrachromosomal elements	0	
Total genes	3,075	100.00%
RNA genes	48	1.59%
rRNA operons	1	
Protein-coding genes	3,027	98.30%
Pseudo genes	29	1.90%
Genes with function prediction	1,860	60.47%
Genes in paralog clusters	473	15.38%
Genes assigned to COGs	1,946	63.28%
Genes assigned Pfam domains	1,918	62.37%
Genes with signal peptides	705	22.93%
Genes with transmembrane helices	782	25.43%
CRISPR repeats	2	




Figure 3. Graphical circular map of the genome. From outside to the center: Genes on forward strand (color by COG categories), Genes on reverse strand (color by COG categories), RNA genes (tRNAs green, rRNAs red, other RNAs black), GC content, GC skew.

Table 4. Number of genes associated with the general COG functional categories

Code	Value	% age	Description
J	157	5.2	Translation, ribosomal structure and biogenesis
A	1	0.0	RNA processing and modification
K	120	3.9	Transcription
L	115	3.8	Replication, recombination and repair
B	3	0.0	Chromatin structure and dynamics
D	26	0.8	Cell cycle control, mitosis and meiosis
Y	0	0.0	Nuclear structure
V	41	1.3	Defense mechanisms
T	121	4.0	Signal transduction mechanisms
M	82	2.7	Cell wall/membrane biogenesis
N	33	1.0	Cell motility
Z	0	0.0	Cytoskeleton
W	0	0.0	Extracellular structures
U	25	0.8	Intracellular trafficking and secretion
O	95	3.1	Posttranslational modification, protein turnover, chaperones
C	147	4.8	Energy production and conversion

Table 4. Number of genes associated with the general COG functional categories

Code	Value	% age	Description
G	107	3.5	Carbohydrate transport and metabolism
E	165	5.4	Amino acid transport and metabolism
F	65	2.1	Nucleotide transport and metabolism
H	106	3.5	Coenzyme transport and metabolism
I	42	1.4	Lipid transport and metabolism
P	122	4.0	Inorganic ion transport and metabolism
Q	24	0.8	Secondary metabolites biosynthesis, transport and catabolism
R	0	10.9	General function prediction only
S	214	7.1	Function unknown
-	1,081	35.7	Not in COGs

Acknowledgements

We gratefully acknowledge the help of Susanne Schneider (DSMZ) for DNA extraction and quality analysis. This work was performed under the auspices of the US Department of Energy's Office of Science, Biological and Environmental Research Program, and by the University of California, Lawrence Berkeley

National Laboratory under contract No. DE-AC02-05CH11231, Lawrence Livermore National Laboratory under Contract No. DE-AC52-07NA27344, and Los Alamos National Laboratory under contract No. DE-AC02-06NA25396, as well as German Research Foundation (DFG) INST 599/1-1.

References

- Wainø M, Tindall BJ, Ingvorsen K. *Halorhabdus utahensis* gen. nov., sp. nov., an aerobic, extremely halophilic member of the *Archaea* from Great Salt Lake, Utah. *Int J Syst Evol Microbiol* 2000; **50**:183-190. [PubMed](#)
- Antunes A, Taborda M, Huber R, Moissl C, Nobre MF, da Costa MS. *Halorhabdus tiamatea* sp. nov., a non-pigmented, extremely halophilic archaeon from a deep-sea, hypersaline anoxic basin of the Red Sea, and emended description of the genus *Halorhabdus*. *Int J Syst Evol Microbiol* 2008; **58**:215-220. [PubMed](#) [doi:10.1099/ijs.0.65316-0](https://doi.org/10.1099/ijs.0.65316-0)
- McGenity TJ, Gemmell RT, Grant WD, Stan-Lotter H. Origins of halophilic microorganisms in ancient salt deposits. *Environ Microbiol* 2000; **2**:243-250. [PubMed](#) [doi:10.1046/j.1462-2920.2000.00105.x](https://doi.org/10.1046/j.1462-2920.2000.00105.x)
- Lee C, Grasso C, Sharlow MF. Multiple sequence alignment using partial order graphs. *Bioinformatics* 2002; **18**:452-464. [PubMed](#) [doi:10.1093/bioinformatics/18.3.452](https://doi.org/10.1093/bioinformatics/18.3.452)
- Castresana J. Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Mol Biol Evol* 2000; **17**:540-552. [PubMed](#)
- Stamatakis A, Hoover P, Rougemont J. A rapid bootstrap algorithm for the RAxML web-servers. *Syst Biol* 2008; **57**:758-771. [PubMed](#) [doi:10.1080/10635150802429642](https://doi.org/10.1080/10635150802429642)
- Liolios K, Mavromatis K, Tavernarakis N, Kyrpides NC. The Genomes OnLine Database (GOLD) in 2007: status of genomic and metagenomic projects and their associated metadata. *Nucleic Acids Res* 2008; **36**:D475-D479. [PubMed](#) [doi:10.1093/nar/gkm884](https://doi.org/10.1093/nar/gkm884)
- Tindall BJ, Schneider S, Lapidus A, Copeland A, Glavina Del Rio T, Nolan M, Lucas S, Cheng F, Saunders E, Bruce D., et al. Complete genome sequence of *Halomicrobium mukohataei* type strain (arg-2T). *Stand Genomic Sci* 2009; **1**:
- Field D, Garrity G, Gray T, Morrison N, Selengut J, Sterk P, Tatusova T, Thomson N, Allen MJ, Angiuoli SV, et al. Towards a richer description of our complete collection of genomes and metagenomes: the "Minimum Information about a Genome Sequence" (MIGS) specification. *Nat Biotechnol* 2008; **26**:541-547. [PubMed](#) [doi:10.1038/nbt1360](https://doi.org/10.1038/nbt1360)
- Woese CR, Kandler O, Wheelis ML. Towards a natural system of organisms: proposal for the domains *Archaea*, *Bacteria*, and *Eucarya*. *Proc Natl Acad Sci USA* 1990; **87**:4576-4579. [PubMed](#) [doi:10.1073/pnas.87.12.4576](https://doi.org/10.1073/pnas.87.12.4576)
- Garrity GM, Holt JG. Phylum All. *Euryarchaeota* phy. nov. In *Bergey's Manual of Systematic Bacteriology*, vol. 1. 2nd ed. Edited by: Garrity, GM, Boone, DR and Castenholz, RW. Springer, New York; 2001:211-355.

12. List Editor. Validation of publication of new names and new combinations previously effectively published outside the IJSEM. Validation List no. 85. *Int J Syst Evol Microbiol* 2002; **52**:685-690. [PubMed doi:10.1099/ijms.0.02358-0](#)
13. Garrity GM, Lilburn TG, Cole JR, Harrison SH, Euzéby J, Tindall BJ. "Part 1- The Archaea, Phyla Crenarchaeota and Euryarchaeota" Taxonomic Outline of the *Bacteria* and *Archaea*. 2007. www.taxonomicoutline.org
14. Grant WD, Larsen H. Group III. Extremely halophilic archaeobacteria. Order *Halobacteriales* ord. nov. In: JT Staley, MP Bryant, N Pfennig & JG Holt (eds) *Bergey's Manual of Systematic Bacteriology*, first edition, Vol 3, The Williams & Watkins Co., Baltimore, 1989, pp 2216-2228.
15. Gibbons NE. Family V. *Halobacteriaceae* fam. nov. In: RE Buchanan & NE Gibbons (eds) *Bergey's Manual of Determinative Bacteriology*, eighth edition, The Williams & Watkins Co., Baltimore, 1974, pp 279.
16. Anonymous. Biological Agents: Technical rules for biological agents. www.baua.de TRBA 466.
17. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, *et al.* Gene ontology: tool for the unification of biology. *Nat Genet* 2000; **25**:25-29. [PubMed doi:10.1038/75556](#)
18. List of growth media used at DSMZ: http://www.dsmz.de/microorganisms/media_list.php
19. Wu M, Hugenholtz P, Mavromatis K, Pukall R, Dalin E, Ivanova N, Kunin V, Goodwin L, Wu M, Tindall BJ, *et al.* A phylogeny-driven genomic encyclopedia of *Bacteria* and *Archaea*. *Nature* (In press).
20. Sims D, Brettin T, Detter JC, Han C, Lapidus A, Copeland A, Glavina Del Rio T, Nolan M, Chen F, Lucas S, *et al.* Complete genome of *Kytococcus sedentarius* type strain (541^T). *Stand Genomic Sci* 2009; **1**:12-20. [doi:10.4056/sigs.761](#)
21. Anonymous. Prodigal Prokaryotic Dynamic Programming Genefinding Algorithm. Oak Ridge National Laboratory and University of Tennessee 2009 <http://compbio.ornl.gov/prodigal/>
22. Pati A, Ivanova N, Mikhailova N, Ovchinnikova G, Hooper SD, Lykidis A, Kyrpides NC. GenePRIMP: A Gene Prediction Improvement Pipeline for microbial genomes. 2009 (Submitted)
23. Markowitz VM, Mavromatis K, Ivanova NN, Chen IMA, Chu K, Kyrpides NC. Expert Review of Functional Annotations for Microbial Genomes. *Bioinformatics* 2009; **25**:2271-2278. [PubMed doi:10.1093/bioinformatics/btp393](#)